

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Magma extrusion during the Ubinas 2013-2014 eruptive crisis based on satellite thermal imaging (MIROVA) and ground-based monitoring**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1566585> since 2016-06-15T11:59:50Z

*Published version:*

DOI:10.1016/j.jvolgeores.2015.07.005

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published as

Diego Coppola<sup>1</sup>, Orlando Macedo<sup>2</sup>, Domingo Ramos<sup>3</sup>, Anthony Finizola<sup>4</sup>, Dario Delle Donne<sup>5-6</sup>, José del Carpio<sup>2</sup>, Randall White<sup>7</sup>, Wendy McCausland<sup>7</sup>, Riky Centeno<sup>2</sup>, Marco Rivera<sup>3</sup>, Fredy Apaza<sup>3</sup>, Beto Ccallata<sup>3</sup>, Wilmer Chilo<sup>3</sup>, Corrado Cigolini<sup>1,8,9</sup>, Marco Laiolo<sup>6</sup>, Ivonne Lazarte<sup>3</sup>, Roger Machaca<sup>3</sup>, Pablo Masias<sup>3</sup>, Mayra Ortega<sup>3</sup>, Nino Puma<sup>2</sup>, Edú Taipe<sup>3</sup> (2015)

**Magma extrusion during the Ubinas 2013-2014 eruptive crisis based on satellite thermal imaging (MIROVA) and ground-based monitoring.**

JOURNAL OF VOLCANOLOGY AND GEOTHERMAL RESEARCH, Volume: 302, Pages: 199-210

DOI: 10.1016/j.jvolgeores.2015.07.005

Published: SEP 1 2015

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ *Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform*]

32

33 **Magma extrusion during the Ubinas 2013-2014 eruptive crisis based on**  
34 **satellite thermal imaging (MIROVA) and ground-based monitoring**

35

36 **Diego Coppola<sup>1</sup>, Orlando Macedo<sup>2</sup>, Domingo Ramos<sup>3</sup>, Anthony Finizola<sup>4</sup>, Dario**  
37 **Delle Donne<sup>5-6</sup>, José del Carpio<sup>2</sup>, Randall White<sup>7</sup>, Wendy McCausland<sup>7</sup>, Riky**  
38 **Centeno<sup>2</sup>, Marco Rivera<sup>3</sup>, Fredy Apaza<sup>3</sup>, Beto Ccallata<sup>3</sup>, Wilmer Chilo<sup>3</sup>, Corrado**  
39 **Cigolini<sup>1,8,9</sup>, Marco Laiolo<sup>6</sup>, Ivonne Lazarte<sup>3</sup>, Roger Machaca<sup>3</sup>, Pablo Masias<sup>3</sup>,**  
40 **Mayra Ortega<sup>3</sup>, Nino Puma<sup>2</sup>, Edú Taipe<sup>3</sup>,**

41

42

43 <sup>1</sup> *Dipartimento di Scienze della Terra, Università di Torino, Italy.*

44 <sup>2</sup> *Observatorio Vulcanológico del Sur, Instituto Geofísico del Perú (OVS-IGP), Perú.*

45 <sup>3</sup> *Observatorio Vulcanológico del Ingemmet (OVI), Perú..*

46 <sup>4</sup> *Université de La Réunion, IPGP, UMR 7154, Sorbonne Paris Cité, La Réunion,*  
47 *France.*

48 <sup>5</sup> *Dipartimento di Scienze della Terra e del Mare, Università di Palermo, Italy.*

49 <sup>6</sup> *Dipartimento di Scienze della Terra, Università di Firenze, Italy.*

50 <sup>7</sup> *Volcano Disaster Assistance Program, USGS, USA.*

51 <sup>8</sup> *NatRisk, Centro Interdipartimentale sui Rischi Naturali in Ambiente Montano e*  
52 *Collinare, Università degli Studi di Torino, Italy*

53 <sup>9</sup> *Present address: Institute for Geothermal Sciences, University of Kyoto, Beppu, Japan*

54

55

56

57

58 **Email contacts:**

59 Diego Coppola: [diego.coppola@unito.it](mailto:diego.coppola@unito.it)

60 Orlando Macedo: [orlando.macedo@igp.gob.pe](mailto:orlando.macedo@igp.gob.pe)

61 Domingo Ramos: [dramos@ingemmet.gob.pe](mailto:dramos@ingemmet.gob.pe)

62 Anthony Finizola: [anthony.finizola@gmail.com](mailto:anthony.finizola@gmail.com)

63 Dario Delle Donne: [dario.d.donne@gmail.com](mailto:dario.d.donne@gmail.com)

64 Randall White: [rwhite@usgs.gov](mailto:rwhite@usgs.gov)

65 Wendy McCausland: [wmccausland@usgs.gov](mailto:wmccausland@usgs.gov)

66 José del Carpio: [joseadelcarpio@hotmail.com](mailto:joseadelcarpio@hotmail.com)

67 Riky Centeno: [riky.centeno@igp.gob.pe](mailto:riky.centeno@igp.gob.pe)

68 Marco Rivera: [mrivera@ingemmet.gob.pe](mailto:mrivera@ingemmet.gob.pe)

69 Fredy Apaza: [fapaza@ingemmet.gob.pe](mailto:fapaza@ingemmet.gob.pe)

70 Beto Callata: [beto.ccallata@gmail.com](mailto:beto.ccallata@gmail.com)

71 Wilmer Chilo: [wchilom@gmail.com](mailto:wchilom@gmail.com)

72 Corrado Cigolini: [corrado.cigolini@unito.it](mailto:corrado.cigolini@unito.it)

73 Marco Laiolo: [marco.laiolo@unito.it](mailto:marco.laiolo@unito.it)

74 Ivonne Lazarte: [ilazarte@ingemmet.gob.pe](mailto:ilazarte@ingemmet.gob.pe)

75 Roger Machaca: [rmachacca@ingemmet.gob.pe](mailto:rmachacca@ingemmet.gob.pe)

76 Pablo Masias: [pmasias@ingemmet.gob.pe](mailto:pmasias@ingemmet.gob.pe)

77 Mayra Ortega: [mortega@ingemmet.gob.pe](mailto:mortega@ingemmet.gob.pe)

78 Nino Puma: [npuma@igp.gob.pe](mailto:npuma@igp.gob.pe)

79 Edú Taipe: [edtaipe@ingemmet.gob.pe](mailto:edtaipe@ingemmet.gob.pe)

80

81

82

83 **ABSTRACT**

84

85 After three year of mild gases emissions, the Ubinas volcano entered in a new eruptive  
86 phase on September 2<sup>nd</sup>, 2013. The MIROVA system (a space-based volcanic hot-spot  
87 detection system), allowed us to detect in near real time the thermal emissions  
88 associated with the eruption and provided early evidence of magma extrusion within the  
89 deep summit crater. By combining IR data with plume height, sulfur emissions, hot  
90 spring temperatures and seismic activity, we interpret the thermal output detected over  
91 Ubinas in terms of extrusion rates associated eruption. We suggest that the 2013-2014  
92 eruptive crisis can be subdivided into three main phases: (i) shallow magma intrusion  
93 inside the edifice, (ii) extrusion and growing of a lava plug at the bottom of the summit  
94 crater coupled with increasing explosive activity and finally (iii) disruption of the lava  
95 plug and gradual decline of the explosive activity. The occurrence of the 8.2 Mw  
96 Iquique (Chile) earthquake (365 km away from Ubinas) on April 1<sup>st</sup>, 2014, may have  
97 perturbed most of the analyzed parameters, suggesting a prompt interaction with the  
98 ongoing volcanic activity. In particular, the analysis of thermal and seismic datasets  
99 shows that the earthquake may have promoted the most intense thermal and explosive  
100 phase that culminated in a major explosion on April 19<sup>th</sup>, 2014.

101 These results reveal the efficiency of space-based thermal observations in detecting the  
102 extrusion of hot magma within deep volcanic craters and in tracking its evolution. We  
103 emphasize that, in combination with other geophysical and geochemical datasets,  
104 MIROVA is an essential tool for monitoring remote volcanoes with rather difficult  
105 accessibility, like those of the Andes that reach remarkably high altitudes.

106

107

## 108 **INTRODUCTION**

109 The Andes, one of Earth's highest subaerial mountain ranges, host more Holocene  
110 active volcanoes than any other volcanic region in the world (Tilling, 2009) but less  
111 than 25 of the ~200 potentially active volcanoes are continuously monitored (Stern,  
112 2004). Within the last decades, population growth and economic development within  
113 the Andean countries drastically increased volcanic risk within the areas surrounding  
114 active volcanoes.

115 The highly elevated (>4000 m) region of southern Peru is a unique example because it  
116 hosts seven active volcanoes located at less than 160 km from Arequipa, the 2<sup>nd</sup> most  
117 important city of Peru (with nearly one million inhabitants). The same area was also the  
118 site of largest explosive eruption in historical times within the Andes (Huaynaputina  
119 volcano; AD 1600; Thouret et al., 1999) and, in terms of earthquakes and volcanic  
120 eruptions, may be considered one of the most hazardous regions in South America  
121 (Degg and Chester, 2005). Progress and refinement of volcano monitoring techniques is  
122 therefore strategic to mitigate future volcanic crises.

123 Space-based thermal observations of volcanic activity (Harris, 2013) represent a useful,  
124 safe and inexpensive tool that may strongly improve volcano surveillance, especially at  
125 active volcanoes with difficult and dangerous access. A survey performed by Jay et al.  
126 (2013) on central and southern Andes reveals that low-amplitude volcanic hotspots  
127 detectable from space are effectively more common than expected, especially at  
128 volcanoes characterized by low level thermal anomalies such as fumaroles and geysers.  
129 For example, by using a high resolution thermal sensor (i.e. ASTER), the authors found  
130 hot spots at 4 Peruvian volcanos, Sabancaya (5967 m), El Misti (5822 m), Ubinas (5672  
131 m) and Huaynaputina (4850 m), with pixel-integrated thermal anomalies spanning from

132 6 to 13 °C above background. However, due to the low temporal coverage of the high  
133 resolution imagers, a systematic monitoring of these volcanoes is still not sufficient to  
134 track daily or weekly variations that may accompany thermal unrest or ongoing eruptive  
135 activity.

136 A new volcanic hot spot detection system, named MIROVA (Middle InfraRed  
137 Observations of Volcanic Activity), combines an high sensitivity for detecting small  
138 thermal anomalies with the improved temporal coverage typical of moderate-resolution  
139 sensors, both factors necessary for near real time monitoring applications (Coppola et  
140 al., 2015). The system is based on the analysis of infrared data acquired by the Moderate  
141 Resolution Imaging Spectroradiometer sensor (MODIS), and uses the Middle InfraRed  
142 Radiation (MIR) recorded at 1 km<sup>2</sup> resolution in order to detect, locate and measure the  
143 heat radiated from volcanic activity (hereby called Volcanic Radiative Power; VRP in  
144 MW). In particular, MIROVA provides thermal maps (50 x 50 km) and VRP time-  
145 series within 1 to 4 hours of each satellite overpass, thus enabling thermal monitoring of  
146 a volcanic target approximately 4 times per day (Coppola et al., 2015).

147 Since July 2013 MIROVA's observations became operational at 3 Peruvian volcanoes  
148 ([www.mirovaweb.it](http://www.mirovaweb.it)) including Ubinas (5672 m), among the most active volcanoes in  
149 Peru (Thouret et al., 2005). Ubinas entered into a new eruptive crisis a few months later,  
150 on September 2013, and the whole eruption was monitored both by IGP and  
151 INGEMMET Peruvian institutions. This gives us the unique opportunity to relate the  
152 thermal flux detected by MIROVA to a series of field observations and other  
153 geophysical datasets. After presenting the data acquisition and the chronology of the  
154 eruption we will describe the analyzed parameters in terms of eruptive dynamics, with  
155 particular emphasis on the contribution of MIROVA in tracking the extrusion of magma  
156 within the deep crater of Ubinas. Finally, we will discuss the interaction between the

157 Iquique earthquake (Mw 8.2), that struck the coasts of Chile and south Peru on April 1<sup>st</sup>  
158 2014, and the volcanic activity observed at Ubinas.

159

160 Figure – 1

161

## 162 **2 - UBINAS VOLCANO**

163

164 Ubinas volcano (16.355°S -70.903°W; 5672 m a.s.l.) is considered the most active  
165 volcano of Peru, with an average of 7 eruptions (VEI 2-3) per century (cf. Thouret et al.,  
166 2005; Rivera et al., 2014). The volcano is located beyond the main arc of the Central  
167 Andean Volcanic Zone (CVZ), approximately 70 km East of Arequipa city (Fig. 1).  
168 More than 5000 people live within 12 km from the crater. Together with Huaynaputina  
169 and Ticsani volcanoes, Ubinas forms the Ubinas–Huaynaputina–Ticsani Volcanic  
170 Group (UHTVG; Lavallée et al., 2009). Volcanism of this area has been inferred to be  
171 strongly ruled by the structural setting, being dominated by a N165 trending normal  
172 fault and a sinistral, N130 strike-slip fault. (Thouret et al., 2005; Lavallée et al., 2009;  
173 Fig. 1). The edifice lies along the western margin of the Rio Tambo graben (E–W  
174 extensional regime) with a 2,000 m altitude gradient between the low-relief high  
175 plateau, to West, and the Ubinas valley, to E and SE (cf. Byrdina et al., 2013; Gonzales  
176 et al., 2014; Lavallée et al., 2009).

177 The summit area of the volcano is represented by a steep-wall 1.4 km wide caldera  
178 whose floor is ash covered. An ash cone occupies the central portion of the caldera and  
179 is itself truncated to the south by the most recent vent; a triangular funnel-shaped pit-  
180 crater 400 m in diameter and ~300 m deep (Rivera, 1997; Thouret et al., 2005; see Fig.  
181 1c). The recent eruptions occurred within the southern pit crater, where fumarolic areas,



182 emitting volcanic gas and steam have been commonly observed during inter-eruptive  
183 periods (cf. Gonzales et al., 2014). Before the 2013-2014 episode, the last eruption of  
184 Ubinas refers to the 2006-2009 crisis (Rivera et al., 2010; 2014). As previously  
185 occurred, this event produced an increasing number of fumaroles, strong degassing,  
186 phreatic to phreatomagmatic activity, magma extrusion and vulcanian explosions (2-4  
187 km height columns). The eruptive crisis lasted 3 years, showing a general decline since  
188 2007. The ash fallout affected the 5,000 inhabitants and involved the hydrology and the  
189 cultivation in an area of 100 km<sup>2</sup> (Rivera et al., 2010). As a whole about 7 Mm<sup>3</sup> of ash  
190 has been emitted reaching a distance of 80 km from the summit (cf. Rivera et al., 2014).  
191 According to Rivera et al. (2014) the 2006-2009 activity represents an “archetype of  
192 Ubinas’s eruptions” in the last 500 years: repeated ascent of small-volume magma  
193 batches (< 10 Mm<sup>3</sup>) from a shallow reservoir (4-7 km) that interact during their ascent  
194 with sectors of the hydrothermal system in the shallowest portion of the conduit  
195 (Gonzales et al., 2014).  
196 Between 2010 and 2013 the activity was characterized by weak gas emissions (BGVN  
197 38:08) accompanied by mild degassing from the summit crater. On September 2013 a  
198 new eruptive episode started (BGVN 38:08).

199

200

### 201 **3 - DATA ACQUISITION**

202 We present more than one year of continuous monitoring data acquired between July 1<sup>st</sup>,  
203 2013 and September 1<sup>st</sup>, 2014. Five datasets are considered and discussed in the  
204 following sections. These consist of: (1) Volcanic Radiative Power; (2) SO<sub>2</sub> density; (3)  
205 Plume Elevation; (4) Temperature of thermal water, (5) Cumulative daily energy of  
206 hybrid earthquakes and (6) daily number of seismic events.

207

### 208 **3.1 – MIROVA - Volcanic Radiative Power (VRP)**

209 The MIROVA system is based on MODIS Level 1B data provided in near real time by  
210 the LANCE-MODIS system (<http://lance-modis.eosdis.nasa.gov/>). MODIS images have  
211 a nominal spatial resolution of 1 km (on IR bands) and allow a target volcano to be  
212 imaged approximately four times per day. Level 1B granules are analysed automatically  
213 according to five principal steps. These are: (i) data extraction, (ii) cropping and  
214 resampling, (iii) definition of Region of Interest (ROIs), (iv) hot-spot detection and  
215 finally (v) calculation of the “excess” of MIR radiance and Volcanic Radiative Power  
216 (VRP) (see Coppola et al., 2015, for details of the processing scheme of MIROVA).

217 Volcanic Radiative Power (VRP) is calculated by using the MIR method (Wooster et  
218 al., 2003) according to which for any individual “alerted” pixel, the VRP is calculated  
219 as:

220

$$221 \quad VRP_{PIX} = 18.9 \times A_{PIX} \times (L_{4alert} - L_{4bk}) \quad (1)$$

222

223 where  $A_{PIX}$  is the pixel size (1 km<sup>2</sup> for the resampled MODIS pixels), and  $L_{4alert}$  and  
224  $L_{4bk}$  are the 4 μm (MIR) radiance of the alerted pixel/s and background, respectively.

225 When two or more pixels (a cluster of pixels) are alerted, the total radiative power is  
226 calculated as being the sum of each single  $VRP_{PIX}$ .

227 As discussed by Coppola et al., 2015, a number of issues must be taken into account  
228 when using the data automatically provided by MIROVA. In particular, the presence of  
229 clouds and the viewing geometry angle may strongly affect the thermal signal detected  
230 by MODIS. While the clouds may completely absorb the IR radiation from the ground,  
231 the radiating source located at the bottom of a deep crater (as in the case of Ubinas) may

232 remain undetected when the satellite zenith angle is very high (i.e.  $> 30^\circ$ ). For these  
233 reasons the visual inspection of each image was performed routinely during the eruption  
234 allowing to interpret the thermal signal case by case, and to discard one hotspot which  
235 was related to a fire that occurred on September 19<sup>th</sup>, 2013 on the south flank of the  
236 volcano.

237

238 Between July 2013 and September 2014 MIROVA detected hotspots in 62 images, with  
239 VRP ranging from 1 to 37 MW (Fig. 3a). Based on the analyzed data we estimated that  
240 the 2013-2014 eruption of Ubinas radiated approximately  $5.5 \times 10^{13}$  J into the  
241 atmosphere. An example of selected images processed by MIROVA throughout the  
242 eruption is given in Fig. 2

243

244 Figure 2

245

### 246 **3.2 - SO<sub>2</sub> density (OMI)**

247 SO<sub>2</sub> density in Dobson Units (DU) is calculated from Ozone Monitoring Instrument  
248 (OMI) images and published daily through the Aura Validation Data Center website  
249 (AVDC, <http://avdc.gsfc.nasa.gov/>). For this study we used the daily average of SO<sub>2</sub>  
250 density values (Fig. 3b) estimated for the Planetary Boundary Layer PBL-SO<sub>2</sub> column  
251 with center of mass altitude of 0.9 km (<http://so2.gsfc.nasa.gov/docs.html>), just above  
252 the crater area of Ubinas volcano and within 50 km radius. As noted on this website,  
253 some of the values may be potentially biased by noise due to South Atlantic Anomaly  
254 (SAA)

### 255 **3.3 - Plume elevation**

256 The estimation of the daily maximum plume elevation (Fig. 3c) have been performed  
257 during the period July 1<sup>st</sup>, 2013 to March 9<sup>th</sup>, 2014 by a local assistant from Ubinas  
258 village and after March 10<sup>th</sup>, 2014 by way of a video camera installed 28 km to the west  
259 of the active vent.

260 The video camera (Campbell Scientific CC5MPX) records one picture of 5 Megapixels  
261 every 30 seconds. These images are sent to Arequipa by telemetry.

262

### 263 **3.4 - Temperature of thermal waters**

264 Temperature in the thermal spring of “Ubinas Thermal”, located ~5.6 km SE from the  
265 volcano’s summit (UB1 in Fig 1; Gonzales et al., 2014), has been measured every 5  
266 minutes by a data logger (Tinytag Aquatic 2). In this survey, we plot the temperatures  
267 measured every day at 4.00 AM (local time) in order to avoid any fluctuations due to  
268 variations of solar radiation (Fig. 3d).

269

### 270 **3.5 - Cumulative daily energy of hybrid earthquakes**

271 The Ubinas volcano seismic network is a telemetric network maintained by the OVS-  
272 IGP. This network consists of 4 permanent digital telemetric stations (UB1, UB2, UB3  
273 and UB4) distributed over the entire volcanic cone, between 4850 and 5000 masl. Two  
274 stations (UB1 and UB2) are equipped with Guralp CMG-40T, 3C broadband sensors,  
275 and the other two (UB3 and UB4) with Lennartz LE-3Dlite short period sensors. All  
276 digitizers are RefTek-130.

277 The analysis and classification of seismic signals recorded by the network, the  
278 determination of the seismic energy, the location of events, etc., is performed daily  
279 using mainly the UB1 seismic data, UB1 being the most reliable station, situated 2.5 km  
280 NW of the active vent.

281 The energy has been calculated by using the following equation:

$$282 \quad E_{seismic} = 2\pi r^2 \rho c \frac{1}{A} \int S^2 U(t)^2 dt \quad (2)$$

283 (Johnson and Aster, 2005) where  $r$  is the source-station distance,  $\rho$  is the density,  $c$  is  
284 the P wave velocity,  $A$  is the attenuation correction,  $S$  the seismic site response  
285 correction and  $U(t)$  is the particle velocity. The source is fixed below the active vent. By  
286 setting  $\rho = 2600 \text{ kg m}^{-3}$ ,  $c = 3000 \text{ m s}^{-1}$ ,  $A$  and  $S$  were fixed at 1.

287 During the current eruption all types of seismic events have been observed like  
288 Volcano-tectonic (VT), Long period (LP), hybrid (events having high and low  
289 frequencies), tremors, explosions and exhalations. During the seven months with  
290 maximum activity, February to August 2014, these events were recorded with mean  
291 rates of 44 VTs/month, 5223 LPs/month, 1104 hybrids/month, 164 explosions or  
292 exhalations/month and 6 hours daily of tremor. All these events occurred near the  
293 surface by the crater zone, except VTs which were located between 1 and 3 km deep  
294 below the crater zone.

295 During the stage of magmatic eruption, characterized by intense eruptions of tephra, the  
296 most important signals are earthquakes of the hybrid type, which have been detected  
297 and registered from the second week of February onwards (Figure 3). Since these  
298 features have been associated with magma ascent to the crater (White et al., 1998;  
299 White, 2011) the OVS decided to conduct a surveillance of the eruption based primarily  
300 on the cumulative daily energy (Fig. 3e) and on the daily rate (Fig. 3f) of such hybrid  
301 type earthquakes.

302

#### 303 **4 - RESULTS**

304 Based on the recorded parameters and the observed phenomena the 2013-2014 eruption  
305 of Ubinas has been subdivided into three main phases (Fig. 3).

306 On September 02<sup>nd</sup> 2013 at 03:46 UTC, a phreatic explosion generated an ash plume  
307 that rose 1.5 to 2 km above the crater. A few minutes after the explosion, at 04:00 UTC,  
308 the MIROVA system detected the first thermal anomaly since the beginning of real time  
309 observations (Fig. 3a). This explosion was the first of a short sequence of phreatic  
310 events that occurred between September 2<sup>nd</sup> and 7<sup>th</sup>, and marks the beginning of the  
311 Phase I. In the following months, occasional puffs of steam and gases rose typically 500  
312 m above the crater without producing any thermal anomaly detectable by MIROVA.  
313 However, between September and December 2013, the “Ubinas Thermal” (UT) hot  
314 spring started to increase its temperature (Fig. 3d). On January 2014 plume height was  
315 persistently below 500 m although anomalous SO<sub>2</sub> concentrations (above 10 DU) were  
316 detected by OMI on a few occasions (Fig. 3b and 3c).

317 An increase in seismic activity started on February 03<sup>rd</sup> 2014 and marks the beginning  
318 of the Phase II (Fig. 3e and 3f). Daily energy of hybrid earthquakes suddenly increased  
319 on 9<sup>th</sup> February (rising for the first time above 5MJ) and was followed, on February 10<sup>th</sup>  
320 , by the first thermal anomaly detected by MIROVA after September 2013 (Fig. 3a).  
321 Note than the picture taken on February 11<sup>th</sup>, 2014, does not allow seeing the bottom of  
322 the crater (Fig. 4a). However, the MODIS image of February 10<sup>th</sup> was acquired with a  
323 very low satellite zenith angle (Fig. 2b) allowing the detection of magmatic anomalies  
324 better than those collected by means of visual observations from the caldera summit.

325 After a peak on February 14<sup>th</sup> (114 MJ; Fig. 3e) the energy of hybrid earthquakes started  
326 to decline, although thermal anomalies were regularly detected and increased over time,  
327 reaching about 6-7 MW on March 1<sup>st</sup>. These thermal detections occurred only under  
328 near-zenithal viewing geometry (satellite zenith < 35°) thus suggesting that the  
329 “radiating source” was located at the bottom of the deep summit crater and likely  
330 corresponded to a growing lava body. A field survey of March 1<sup>st</sup> confirmed, for the

331 first time, the presence of an elongated body of incandescent lava at the bottom of  
332 Ubinas crater that, by that time, measured about 30-40 m in diameter (Fig. 4b). During  
333 the following days the VRP continued to increase due to the growth of the lava body  
334 within the deep crater, reaching about 12 MW on March 12<sup>th</sup>. At the same time a  
335 moderate growth of sulfur emissions and plume height was recorded by OMI (Fig. 3b  
336 and 3c). Throughout March 2014 the explosivity of the eruption slightly increased  
337 (plume height reached more than 2000 m and sulfur emission between 10 and 35 DU)  
338 but a clear reduction in thermal emissions was detected by MIROVA between 14 and 31  
339 March (VRP of 2-7 MW, Fig. 3a, 3b, 3c). The visual inspection of the MODIS images  
340 allowed us to discard the presence of clouds or volcanic plumes as the origin of this  
341 decrease (Fig. 2). Hence, we infer that a noticeable reduction of magma flux entering  
342 the crater characterized this period, although incandescence was still visible at the  
343 surface (Fig. 4c). Interestingly, this sharp decrease of the thermal output in mid-March  
344 2014 was almost coeval with a marked escalation of seismicity (Fig. 3f) with hybrid's  
345 energy release reaching 580 MJ on March 31<sup>st</sup>, 2014 (Fig. 3e).

346

347 Figure 3

348

349 On April 1<sup>st</sup>, 2014 at (23.46 UTC) a 8.2 Mw earthquake struck off the coast of Chile,  
350 365 km south of Ubinas volcano (Fig. 1). The earthquake triggered a tsunami of up to  
351 2.1 m (that hit the town of Iquique, 95 km south east from the epicenter) and affected  
352 regions of Tacna, Moquegua and Arequipa, in South Peru. The Iquique earthquake was  
353 felt in Arequipa town (intensity of III-IV) and eventually affected the plumbing system  
354 of Ubinas as suggested by the perturbation of the monitored parameters (Fig. 3).

355 The hot spring at “Ubinas Thermal” station (UT) recorded this event with a sharp drop  
356 of the water temperature (Fig. 3d). Evaluation of precipitation data recorded at Ubinas  
357 village (courtesy of Meteorological Service of Peru) allowed us to rule out that the  
358 decrease of temperature resulted from the rainfall. More likely the temperature drop  
359 may have resulted from a temporary mixing of fresh and thermal water, as well as from  
360 a permeability change induced by transient stresses (Manga et al., 2012). Temperature  
361 drop is typical of the response of many wells and springs in a variety of environments to  
362 the seismic waves from distant earthquakes (Hills et al., 2002) as for example observed  
363 in the past years at Ubinas and Misti volcano (IGP internal Reports for June, 23<sup>th</sup> 2001).  
364 A major change in the eruptive behavior of Ubinas was also recorded by seismic and  
365 MIROVA data. Indeed, after several days of increasing trend, seismicity inverted its  
366 tendency just after the earthquake, starting to decline in both hybrid’s energy release  
367 (Fig. 3e) and number of events (Fig. 3f). At the same time the thermal activity within  
368 the crater had strongly intensified reaching 37 MW on April 4th, 2014. This was the  
369 highest thermal anomaly recorded during the eruption and occurred only 3 days after the  
370 8.2 Mw earthquake (Fig 3a). In the following days several thermal detections reached  
371 more than 20 MW and were accompanied by the most explosive phase, characterized by  
372 multiple explosions ejecting incandescent tephra and blocks around the crater. In this  
373 period the ash plumes reached altitudes of more than 4000 m and were coupled with  
374 sulfur emissions as high as 45 DU (Fig. 3b, 3c).

375 A major explosive event occurred on April 19<sup>th</sup> 2014, where blocks of fresh basaltic  
376 andesite magma, 40-50 cm in size, were ejected up to 2.6 km from the active vent. The  
377 volcanic plume rose more than 6 km above the crater, and a block of 5 x 4 x 2m has  
378 been found inside the summit caldera, at 660 m from the active vent.



379 This explosion marks the end of the Phase II and the transition into the Phase III.  
380 Thermal output at the summit crater dropped abruptly after the major explosion, and the  
381 detection of weak thermal anomalies (about 1 MW) became sporadic between May and  
382 September 2014 (Fig. 3a). At the same time, the explosive activity gradually declined  
383 although it was punctuated by distinct explosive events of decreasing intensity. On May  
384 9, at 03:50 (UTC), an isolated thermal anomaly of 30 MW was detected (Fig. 3a) and  
385 resulted from the hot material ejected during one coeval explosion.

386 The abrupt drop of thermal emissions after the explosion of April 19<sup>th</sup> was likely  
387 associated with the massive disruption of the magma plug, extruded within the crater  
388 during the previous explosive phase. Actually, the explosion cut the head of the magma  
389 column (strongly deepening the thermal source) which however continued to feed tens  
390 of minor explosive events during Phase III. Field observations on July 31<sup>st</sup> confirmed  
391 that the lava body at the bottom of the crater had disappeared (Fig. 4d), even though  
392 moderate explosions and SO<sub>2</sub> emissions were still significant (Fig. 3b and 3c). On the  
393 other hand, after the earthquake-induced drop of temperature, the hot spring UT  
394 gradually returned to pre-earthquake temperatures (yet showing multiple minor  
395 fluctuations, Fig. 3d). As a whole the Phase III consisted of a waning eruptive period  
396 and by September 2014 the activity of Ubinas consisted only of weak plume emissions  
397 and sporadic low altitude explosions (Fig. 3b and 3c).

398

399 Figure 4 -

400

401 **5 - DISCUSSION**

402 The phreatic explosions of September 2013 mark the beginning of the Phase I and  
403 possibly reflect the first interaction between a new ascending magma batch and the  
404 large hydrothermal system located under the Ubinas crater (Gonzales et al., 2014).

405 It is worth noting that the temperature of the “Ubinas thermal” hot spring increased  
406 significantly during this phase (Fig 3d), probably reflecting a period of shallow magma  
407 intrusion inside the edifice and its interaction with the thermal waters stored within the  
408 hydrothermal system. On the other hand, the increase in the energy of hybrid events on  
409 February 3<sup>rd</sup>, 2014 (Fig. 3e), just followed by summit thermal anomalies (on February  
410 10<sup>th</sup>, 2014), clearly indicate that 166 days after the first phreatic explosion, the magma  
411 had reached the bottom of the summit crater. By assuming that the base of the  
412 hydrothermal system is located 1500-3000 m below the summit (Gonzales et al., 2014),  
413 we may infer that during this intrusive stage the andesitic magma batch rose to very  
414 shallow depth with an average velocity of  $1-2 \times 10^{-4} \text{ m s}^{-1}$ .

415 Overall, during the most intense “eruptive” period (Phase II), we observe a good  
416 correlation of the peaks and trends between thermal output, SO<sub>2</sub> emissions, and plume  
417 elevation (Fig. 2). Notably, our datasets reveal that since the first detection of magma at  
418 the bottom of the crater (on February 10<sup>th</sup>), all the above parameters share a general  
419 increasing trend throughout the Phase II (Fig. 3) that was apparently amplified by the  
420 Iquique earthquake (discussed below).

421 Although thermal, SO<sub>2</sub>, and plume elevation sample distinct eruptive processes (i.e.  
422 heat, gas and ash release), it’s interesting to note that all these parameters are somehow  
423 related to rate of magma ascent within the shallow volcanic conduit (Gonnermann and  
424 Manga, 2012). For example, under certain assumptions, space-based thermal flux  
425 released by active lava bodies may be converted into estimates of Time Averaged lava  
426 Discharge Rates (TADR; Harris et al., 2009; Dragoni and Tallarico 2009, Coppola et

427 al., 2013) thus providing a way to calculate volumetric flux of effused-extruded lavas  
 428 (see next section). On the other hand, the SO<sub>2</sub> emissions have been widely used to  
 429 calculate magma degassing rates (i.e. Allard et al., 1994) and in some cases may provide  
 430 constraints into the magma supply rate (by using the so called “Petrologic Method”;  
 431 Shinoara, 2008). Nonetheless, during explosive eruptions, plume height has been  
 432 commonly correlated to Mass Eruption Rates (MER) (Morton et al., 1956; Sparks et al.,  
 433 1997; Wilson et al., 1978; Mastin et al., 2009) thus providing a further way to evaluate  
 434 magma production rates occurring during explosive events. Accordingly, we may infer  
 435 that the Phase II was characterized by a general acceleration of magma eruption rates  
 436 that culminated into the paroxysmal phase of April 19<sup>th</sup>, 2014.

437

## 438 **5.1 - EXTRUSION RATES AND VOLUMES FROM THERMAL DATA**

439

440 Coppola et al. (2013) proposed a simple method to provide first-order estimates of time-  
 441 averaged lava discharge rates (TADR) from MODIS-derived VRP. This is based on an  
 442 empirical parameter called “radiant density” ( $c_{rad}$ ):

$$443 \quad TADR = \frac{VRP}{c_{rad}} \quad (3)$$

444 where TADR is in m<sup>3</sup>s<sup>-1</sup>, VRP is in W, and  $c_{rad}$  is the radiant density, expressed in J m<sup>-3</sup>

445 The radiant density approach relies on the fact that under a given discharge rate, basic,  
 446 intermediate and acidic lava bodies radiate thermal energy differently because of their  
 447 different bulk rheology. Based on the analysis of several distinct worldwide eruptions,  
 448 the authors suggest that the radiant density of a lava body can be predicted ( $\pm 50\%$ ) on  
 449 the basis of the silica content of the erupted products:

$$450 \quad c_{rad} = 6.45 \times 10^{25} \times (X_{SiO_2})^{-10.4} \quad (4)$$

451 where  $X_{\text{SiO}_2}$  is the silica content (wt%) of the erupted magma. For Ubinas andesite, here  
452 we used a silica content of 56 wt% (Rivera pers. comm.), and we calculated a radiant  
453 density comprised between  $2.1 \times 10^7$  and  $8.5 \times 10^7 \text{ J m}^{-3}$ . Accordingly, we estimated  
454 that  $1.5 \pm 0.75 \text{ Mm}^3$  of magma have been extruded between its first appearance at the  
455 surface, on February 10<sup>th</sup>, and the major explosion of April 19<sup>th</sup>, 2014 (mean output rate  
456 of  $0.25 \pm 0.12 \text{ m}^3 \text{ s}^{-1}$ ).

457 A more detailed analysis reveals that, although characterized by a general increasing  
458 trend, the extrusion of magma was somehow cyclic, with at least three distinct stages  
459 (Fig 5a). The first stage (Phase IIb) was recorded between February 10<sup>th</sup> and March  
460 12<sup>nd</sup>, when the extrusion rate gradually increased from  $0.1 \pm 0.05$  to  $0.31 \pm 0.15 \text{ m}^3 \text{ s}^{-1}$ .  
461 The second stage (Phase IIc) followed a drastic reduction of magma extrusion on March  
462 14<sup>th</sup>, and was characterised by output rates that remained below  $0.2 \text{ m}^3 \text{ s}^{-1}$  up to the end  
463 of March (Fig. 5a). The third stage started after April 1<sup>st</sup>, and was characterised by the  
464 sudden increase of extrusion rate (up to  $1.4 \pm 0.02 \text{ m}^3 \text{ s}^{-1}$  on April 4<sup>th</sup>) that persisted  
465 above  $1 \text{ m}^3 \text{ s}^{-1}$  up to the major explosion of April 19<sup>th</sup> (Phase IId). It is worth noting that  
466 during this intense extrusive and explosive phase the magma plug was probably  
467 continuously extruded and disrupted by the recurrent explosions so that by the end of  
468 May 2014 no magma body was present at the bottom of the crater (Fig. 4d). Sporadic  
469 thermal emissions were also recorded during the Phase III. However, due to the poor  
470 continuity of the thermal signal we don't consider this phase as extruding significant  
471 magma volume within the pit crater. Instead, we regard the above values as a minimum  
472 estimate for the erupted magma volumes, essentially extruded during the Phase II.  
473 Accordingly, we may estimate that a minimum of  $1.5 \pm 0.75 \text{ Mm}^3$  of ash and tephra  
474 should have been produced by the continuous disruption of the magma body at the  
475 bottom of the pit crater, especially throughout the most intense Phase IId. By

476 comparison, about 7 Mm<sup>3</sup> of ash tephra was erupted during the long-lasting 2006-2009  
477 eruptive crisis throughout hundreds of vulcanian explosions (Rivera et al., 2014).

478

## 479 **5.2 - THERMAL AND HYBRID SEISMIC ACTIVITY COMPARISON**

480 The comparison between hybrid seismic and thermal activity recorded during the Phase  
481 II reveals a more complex eruptive dynamic and outlines some mutual relationships  
482 between these parameters (Fig. 5). In fact, the increase of hybrid's energy and the  
483 heightening of daily seismicity seem to have systematically "preceded" the cycles of  
484 major magma extrusion at the summit crater. For example, in early February 2014,  
485 when the conduit was still "closed" (no signs of thermal activity), the occurrence of few  
486 but energetic hybrids suggests that the magma had to overcome a high resistance before  
487 eventually appearing on the surface (Phase IIa; Fig. 5). On the other hand, after the first  
488 "opening" of the conduit (i.e. after February 10<sup>th</sup>), the magma rose almost easily  
489 (increasing extrusion rate) and hybrid's energy decreased accordingly (Phase IIb; Fig.  
490 5). Once again these trends inverted since March 12<sup>th</sup>, when the reduction of the  
491 extrusion rate was coupled with a renewed and stronger increase in seismic activity.  
492 This suggest that the ascent of magma in this period was somehow "reduced" (or  
493 blocked), because of obstruction(s) within the conduit or, eventually, because the  
494 magma was too viscous to flow out. In our view, the Phases IIa and IIc might reflect  
495 stages of pressure build-up within the plumbing system where the slow ascending  
496 magma had to push hard (increasing daily hybrid's energy) to open the conduit (Phases  
497 IIa) or to extrude magma at low rates (Phases IIc). The earthquake of April 1<sup>st</sup> occurred  
498 at this critical stage and may have contributed to unblocking the conduit and promoting  
499 the most intense activity at the vent (Phase 2d; see also next section). It is worth noting  
500 that, after the earthquake, both the energy of hybrids and the number of seismic events

501 decreased regularly, thus suggesting a general depressurization of the shallow magmatic  
502 system during the highest extrusive phase as well as during the following waning phase  
503 (Fig. 5b and 5c).

504 Ottemöller (2008) found similar correlations during the 2003 extrusive eruption at  
505 Soufrière Hills Volcano (SHV), whereby higher energy releases of hybrid events  
506 reflected increased pressurization during periods of low extrusion rates. Conversely,  
507 lower energy releases have been inferred to be associated with rapid extrusion and  
508 reduced pressurization. Thus, the cyclic magma extrusion observed at Ubinas volcano is  
509 not uncommon, and may be explained by non-linear processes related to degassing,  
510 crystallization and rheological stiffening of magma, as observed in many other dome-  
511 building eruptions (Denlinger and Hoblitt 1996; Melnik and Sparks 2006; Costa et al.,  
512 2007).

513

### 514 **5.3 – POSSIBLE INTERACTION BETWEEN THE IQUIQUE EARTHQUAKE** 515 **AND UBINAS VOLCANO**

516

517 Regional earthquakes have been proved to be able to interact with volcanoes by  
518 triggering new volcanic unrest (Hill et al., 2002), by perturbing the medium in the  
519 vicinity of a volcanic conduit (Manga et al., 2012; Battaglia et al., 2012; Lesage et al.,  
520 2014), or, by causing several-fold increases in thermal output (Delle Donne et al.,  
521 2010), eruption rates (Harris et al., 2007) and/or gas transfer (Cigolini et al. 2007), at  
522 already erupting volcanoes.

523 Our datasets suggest that at least four parameters recorded during the Ubinas eruption  
524 (water temperature at Ubinas thermal station, thermal flux at the summit crater, plume  
525 height and hybrid seismicity) appear to have been perturbed by the M 8.2 Iquique

526 earthquake (Fig. 3). While the temperature drop at the Ubinas Thermal station (Fig. 3d)  
527 may represents a local response of the shallow hydrothermal system to the seismic  
528 waves, it is possible that the other three parameters reflect a response of the magmatic  
529 system to the megathrust earthquake. In particular, the response of Ubinas volcano  
530 consisted on a 2-3 fold increase of volumetric flux and plume height, coupled with a  
531 decrease in the amplitude and rate of hybrid earthquakes (Fig. 5).

532 Seismic waves travel to great distances without losing much of their energy (Hill et al.,  
533 2002; Delle Donne et al. 2010) so that dynamic stress induced by their passage may  
534 have effectively influenced activity at Ubinas volcano. Transient stress perturbations  
535 may in fact promote nucleation, growth and ascent of gas bubbles, acting as a  
536 vesiculation pump (Manga and Brodsky, 2006; Harris and Ripepe, 2007) thus causing  
537 the enhanced magma extrusion and explosive activity as effectively observed at Ubinas  
538 just after the Iquique earthquake (Fig.5).

539 Delle Donne et al. (2010) statistically constrain the maximum distance for triggering an  
540 eruption at a given volcano, which is strictly dependent on the magnitude of the  
541 earthquake. These authors suggest that the orientation of the seismogenic faults, in  
542 respect to the location of the volcano, can also play a role in facilitating the triggering  
543 mechanism (i.e., by means of focusing the radiated energy in a strike-parallel direction).

544 The Iquique megathrust earthquake ( $M_w = 8.2$ ; Distance=365 km; fault strike:  $348.9^\circ$ ;  
545 Lai et al., 2014) fully meet the cited conditions, since the seismogenic fault that  
546 generated the earthquake is characterized by a directivity perfectly compatible with the  
547 location of Ubinas (Fig. 6). This support the hypothesis that the transient stress changes  
548 may have accelerated the eruptive processes at Ubinas, already operative at the time of  
549 the triggering earthquake.

550 On the other hand Bonali et al. (2013), among others, suggest that earthquake-induced  
551 “unclamping” (normal stress reduction within the magma pathway) may also promote  
552 an increase in magma flux at volcanoes that are already in a critical state. To test this  
553 possibility, we estimated the normal stress change and volumetric dilatation induced by  
554 Mw 8.2 Iquique earthquake, by using Coulomb 3.3 software [e.g., Toda et al., 2005]. As  
555 input fault model we used the Finite Fault Results (Ji et al., 2002; Bassin et al., 2000)  
556 computed by G. Hayes\_(NEIC-USGS) available at:

557 [http://comcat.cr.usgs.gov/earthquakes/eventpage/usc000nzvd#scientific\\_finite-fault](http://comcat.cr.usgs.gov/earthquakes/eventpage/usc000nzvd#scientific_finite-fault).

558 To calculate the normal stress change we assumed a shallow (~2 km deep), nearly  
559 vertically dipping and 165°N oriented dyke (consistent with the geology) and a deep  
560 magma chamber located within 5 and 10 km below the surface (Lavallée et al., 2009;  
561 Rivera et al., 2010). Results of the analysis suggest that the Iquique earthquake may  
562 have produced a weak clamping, in the order of only 0.03 bar (Fig. 7a), at the shallow  
563 feeder dyke, coupled with a minor volumetric compression of deep magma chamber  
564 (Fig. 7b). Accordingly, we may conclude that the earthquake did not cause a normal  
565 stress reduction (Bonali et al., 2013) but, conversely, it may have induced only a very  
566 small compression of the Ubinas plumbing system. However, given its low amplitude, it  
567 is unlikely that this weak “clamping” could have promoted the “squeezing” of the  
568 magma filled pathway throughout a small but stable deformation (e.g. Nostro et al,  
569 1998; Bautista et al., 1996).

570 On the other hand our results suggest that Ubinas volcano may have responded  
571 promptly to the Iquique earthquake (< 3 days) by increasing the eruption rate at the vent  
572 and by decreasing the shallow seismicity surrounding the conduit (Fig. 5).

573 At Soufrière Hills Volcano, faster extrusion rates were systematically observed during  
574 deflation periods, the latter being characterized by decreasing hybrid earthquakes



575 (Green & Neuberg 2006). Accordingly, we suggest that the lowering in the hybrid  
576 energy release was not directly related to the earthquake itself, but more likely to the  
577 acceleration of the extrusive-explosive processes occurring at the vent. In turn the  
578 increase in discharge rates induced a pressure drop within the magma plug and caused  
579 the gradual waning of seismicity.

580 Besides, several other Peruvian volcanoes, in a critical or quiescent state, satisfy the  
581 criteria described by Delle Donne et al., 2010 (distance and strike alignment) and might  
582 have been also perturbed by stress changes induced by the Iquique earthquake (Fig. 6).  
583 These include Sabancaya and El Misti, among the most active volcanoes of Peru in the  
584 last century. Further investigations will better clarify if the seismic waves caused by the  
585 Iquique earthquake had sufficient energy to trigger a response in fumarolic emission or  
586 volcano seismicity at these volcanoes.

587

## 588 **CONCLUSIONS**

589

590 The combined use of satellite and field monitoring techniques allowed us to track the  
591 evolution of the 2013-2014 eruptive crisis of Ubinas volcano. The eruption has been  
592 subdivided into three main phases based on a set of 6 parameters and the observed  
593 activity. While Phase I is related to a period of shallow magma intrusion, the other two  
594 phases were ascribed to stages of waxing (Phase II) and waning (Phase III) eruptive  
595 activity, respectively.

596 In particular, the space-based observations performed by the MIROVA system provided  
597 the first evidences of magma extrusion within the bottom of the summit pit-crater and  
598 allowed to constrain extrusion rates during the most intense eruptive phase (Phase II).  
599 We estimated that between February and April 2014, at least  $1.5 \text{ Mm}^3$  of magma were

600 extruded during week-long cycles and were abruptly disrupted during the major  
601 explosion of April 19<sup>th</sup>, 2014. Nonetheless, during the Phase II we recognized a general  
602 acceleration of all the eruptive processes (magma extrusion, plume height, SO<sub>2</sub>  
603 emission) which were apparently perturbed by the occurrence of the Iquique earthquake  
604 (Mw 8.2) on April 1st 2014. A preliminary analysis suggests that the prompt 3-fold  
605 increase of the extrusion rate was principally a response to the dynamic stress changes  
606 induced by the earthquake, and may have favored a general depressurization of the  
607 shallow magmatic system associated to the decreasing hybrids seismic activity.

608 The observations provided by MIROVA demonstrate the capability to track the  
609 presence of magma within deep craters, and allow us to better understand the extrusive  
610 processes by correlating MODIS data with other datasets, such as plume emissions and  
611 seismic activity (i.e., the energy released by hybrid events). In conclusion, we hereby  
612 demonstrate that MIROVA is an additional and efficient tool for monitoring safely, and  
613 in near real time, extrusive-explosive volcanoes with hard and dangerous access.

614

## 615 **ACKNOWLEDGEMENTS**

616

617 MIROVA is a collaborative project between the Universities of Turin and Florence  
618 (Italy), and is supported by the Italian Civil Protection Department. Additional funds  
619 were provided by MIUR, Fondazione Cassa di Risparmio di Torino and Fondazione  
620 Compagnia di San Paolo di Torino. The seismic study was financed by the APNOP  
621 Meta 022 and the PP068 Meta 007 of the OVS- Instituto Geofísico del Perú. We  
622 acknowledge L. Wilson for the editorial handling. We thank, S. Byrdina and J.F. Lénat  
623 for their constructive comments. We are grateful to D. Hill for discussions and  
624 suggestions on an early version of this manuscript. We particularly thank M. Alvarez

625 and J. Acosta for field observations. We acknowledge the LANCE-MODIS system  
626 (<http://lance-modis.eosdis.nasa.gov/>) for providing Level 1B MODIS data. This is IPGP  
627 contribution number: 3644. Any use of trade, firm, or product names is for descriptive  
628 purposes only and does not imply endorsement by the U.S. Government.

629

630

## 631 **REFERENCES**

632

633 Allard, P., Carbonelle, J., Mètrich, N., Loyer, H., Zettwoog, P., 1994. Sulphur output  
634 and magma degassing budget of Stromboli volcano. *Nature* 368, 326-330,  
635 doi:10.1038/368326a0.

636

637 Bassin, C., Laske, G., Masters, G., 2000. The Current Limits of Resolution for Surface  
638 Wave Tomography in North America, *EOS Trans AGU*, 81, F897.

639

640 Battaglia, j., Métaxian, J-P., Garaebiti, E., 2012. Earthquake-volcano interaction imaged  
641 by coda wave interferometry. *Geophys. Res. Lett.*, 39 (L11309).

642

643 Bonali, F.L., Tibaldi, A., Corazzato, C., Tormey, D.R., Lara, L.E., 2013. Quantifying  
644 the effect of large earthquakes in promoting eruptions due to stress changes on  
645 magma pathway: The Chile case. *Tectonophysics* 583, 54–67. doi:  
646 10.1016/j.tecto.2012.10.025.

647

648 Byrdina, S., Ramos, D., Vandemeulebrouck, J., Masias, P., Revil, A., Finizola, A.,  
649 Gonzales Zuñiga, K., Cruz, V., Antayhua, Y., Macedo, O. 2013. Influence of the

650 regional topography on the remote emplacement of hydrothermal systems with  
651 examples of Ticsani and Ubinas volcanoes, Southern Peru. *Earth Planet. Sci. Lett.*  
652 365, 152-164.

653

654 Cigolini, C., Laiolo, M., Coppola, D., 2007. Earthquake–volcano interactions detected  
655 from radon degassing at Stromboli (Italy). *Earth Planet. Sci. Lett.* 257 (3-4), 511–  
656 525.

657

658 Coppola, D., Laiolo, M., Piscopo, D., Cigolini, C. 2013. Rheological control on the  
659 radiant density of active lava flows and domes. *J. Volcanol. Geotherm. Res.*, 249,  
660 39–48, doi:10.1016/j.jvolgeores.2012.09.005.

661

662 Coppola D., Laiolo M., Cigolini C., Delle Donne D., Ripepe M., 2015. Enhanced  
663 volcanic hot-spot detection using MODIS IR data: results from the MIROVA  
664 system. In: Harris, AJL., De Groeve, T., Garel, F., & Carn, S.A. (eds) *Detecting,*  
665 *Modelling and Responding to Effusive Eruptions.* Geological Society, London,  
666 Special Publications, 426, first published on May 14, 2015, doi:10.1144/SP426.5

667 Degg, M.R., Chester, D.K., 2005. Seismic and volcanic hazards in Peru: Changing  
668 attitudes to disaster mitigation: *The Geographical Journal*, 171(2), 125-145.

669

670 Denlinger, R.P., Hoblitt, R.P., 1996. Cyclic eruptive behavior of silicic volcanoes.  
671 *Geology* 27(5), 459-462.

672

673 de Silva, S.L., Francis, P., 1990. Potentially active volcanoes of Perú - Observations  
674 using the Landsat Thematic Mapper and space shuttle photography. *Bull. Volcanol.*,  
675 52, 286-301.

676

677 Costa, A., Melnik, O., Sparks, R.S.J., Voight, B., 2007. Control of magma flow in dykes  
678 on cyclic lava dome extrusion. *Geophys. Res. Lett.* 34 (L02303).

679

680 Delle Donne, D., Harris, A.J.L., Ripepe, M., Wright, R., 2010. Earthquake-induced  
681 thermal anomalies at active volcanoes. *Geology* 38, 71-74, doi: 10.1130/G30984.1.

682

683 Dragoni, M., Tallarico, A., 2009. Assumptions in the evaluation of lava effusion rates  
684 from heat radiation. *Geophys. Res. Lett.* 36 (8), L08302, DOI:  
685 10.1029/2009GL037411.

686

687 Gonnermann, H.M., Manga, M., 2013. Magma ascent in the volcanic conduit. In:  
688 *Modeling volcanic processes: The physics and mathematics of volcanism*, Eds.  
689 Fagents, S.A., Gregg, T.K.P., Lopez, R.C., Cambridge University Press, 55-84.

690

691 Gonzales, K., Finizola, A., Lénat, J.F., Macedo, O., Ramos, D., Thouret, J.C., Fournier,  
692 N., Cruz, V., Pistre, K., 2014. Asymmetrical structure, hydrothermal system and  
693 edifice stability: The case of Ubinas volcano, Peru, revealed by geophysical surveys.  
694 *J. Volcanol. Geotherm. Res.*, 276, 132-144.

695

696 Harris, A.J.L., 2013 *Thermal remote sensing of active volcanoes: a user's manual*.  
697 Cambridge University Press, pp. 736

698

699 Harris, A.J.L., Ripepe, M., 2007. Regional earthquake as a trigger for enhanced  
700 volcanic activity: Evidence from MODIS thermal data. *Geophys. Res. Lett.* 34,  
701 L02304, doi: 10.1029/2006GL028251.

702

703 Harris, A.J.L., Baloga, S.M., 2009. Lava discharge rates from satellite-measured heat  
704 flux. *Geophys. Res. Lett.* 36 (19), doi: 10.1029/2009GL039717.

705

706 Hill, D.P., Pollitz, F., Newhall, C., 2002. Earthquake-volcano interactions. *Physics*  
707 *Today* 55, 41-47.

708

709 Jay, J.A., Welch, M., Pritchard, M.E., Mares, P.J., Mnich, M.E., Melkonian, K.,  
710 Aguilera, F., Naranjo, J.A., Sunagua, M., Clavero, J., 2013. Volcanic hotspots of the  
711 central and southern Andes as seen from space by ASTER and MODVOLC between  
712 the years 2000 and 2010. *Geol. Soc. London Special Publication* 380, p.

713

714 Ji, C., D.J. Wald, Helmberger, D.V., 2002. Source description of the 1999 Hector Mine,  
715 California earthquake; Part I: Wavelet domain inversion theory and resolution  
716 analysis, , *Vol* 92, No. 4. pp. 1192-1207.

717

718 Johnson J.B., Aster R.C., 2005. Relative partitioning of acoustic and seismic energy  
719 during strombolian eruptions. *J. Volcanol. Geotherm. Res.*, 148, 334-354.

720

721 Lavallée, Y., de Silva, S.L., Salas, G., Byrnes, J.M., 2009. Structural control on  
722 volcanism at the Ubinas, Huaynaputina, and Ticsani Volcanic Group (UHTVG),  
723 southern Peru. *J. Volcanol. Geotherm. Res.*, 186 (3–4), 253–264.  
724

725 Lay, T., Yue, H., Brodsky, E. E. , An, C. , 2014. The April 1, 2014 Iquique, Chile Mw  
726 8.1 earthquake rupture sequence. *Geophys. Res. Lett.*, 41 (11), 3818–3825, doi:  
727 10.1002/2014GL060238.  
728

729 Lesage, P., G. Reyes-Dávila, and R. Arámbula-Mendoza (2014), Large tectonic  
730 earthquakes induce sharp temporary decreases in seismic velocity in Volcán de  
731 Colima, Mexico, *J. Geophys. Res. Solid Earth*, 119, 4360–4376,  
732 doi:10.1002/2013JB010884.

733 Manga, M., Brodsky, E., 2006. Seismic triggering of eruption in the far field: Volcanoes  
734 and Geysers. *Annual Review of Earth and Planetary Sciences*, 34, 263-291.  
735

736 Manga, M., I. Beresnev, E. E. Brodsky, J. E. Elkhoury, D. Elsworth, S. E. Ingebritsen,  
737 D. C. Mays, and C.-Y. Wang (2012), Changes in permeability caused by transient  
738 stresses: Field observations, experiments, and mechanisms, *Rev. Geophys.*, 50,  
739 RG2004, doi:10.1029/2011RG000382.  
740

741 Mastin, L.G., Guffanti, M., Servranckx, R., Webleyd, P., Barsotti, S., Dean, K., Durant,  
742 A., Ewert, J.W., Neri, A., Rose, W.I., Schneider, D., Siebert, L., Stunder, B.,  
743 Swanson, G., Tupper, A., Volentik, A., Waythomas, C.F., 2009. A multidisciplinary  
744 effort to assign realistic source parameters to models of volcanic ash-cloud transport  
745 and dispersion during eruptions. *J. Volcanol. Geotherm. Res.*, 186 (1–2), 10–21.

746

747 Melnik, O., Sparks, R.S.J., 1999. Nonlinear dynamics of lava dome extrusion, *Nature*,  
748 402, 37 – 41.

749

750 Morton, B.R., Taylor, G.T., Turner, J.S., 1956. Turbulent gravitational convection from  
751 maintained and instantaneous sources. *Transaction of the Royal Society London ser.*  
752 *A* 234, 1-23.

753

754 Ottemöller, L., 2008. Seismic hybrid swarm precursory to a major lava dome collapse:  
755 9-12 July 2003, Soufrière Hills Volcano, Montserrat. *J. Volcanol. Geotherm. Res.*,  
756 177 (4), 903-910.

757

758 Rivera, M., Thouret, J.C. , Mariño, J., Berolatti, R., Fuentes, J. 2010. Characteristics and  
759 management of the 2006-2008 volcanic crisis at the Ubinas volcano (Peru). *J.*  
760 *Volcanol. Geotherm. Res.*, 198 (1-2), 19-34.

761

762 Rivera, M., Thouret, J.C., Samaniego, P., Le Pennec, J.L. 2014. The 2006-2009 activity  
763 of the Ubinas volcano (Peru): Petrology of the 2006 eruptive products and insights  
764 into genesis of andesite magmas, magma recharge and plumbing system. *J. Volcanol.*  
765 *Geotherm. Res.*, 270, 122-141.

766

767 Shinohara, H., 2008. Excess degassing from volcanoes and its role on eruptive and  
768 intrusive activity. *Review Geophys.* 46 (4), RG4005, doi:10.1029/2007RG000244.

769



770 Sparks, R.S.J., Bursik, M.I., Carey, S.N., Gilbert, J., Glaze, L.S., Sigurdsson, H.,  
771 Woods, A., 1997. Volcanic Plumes. John Wiley and Sons, p. 574.  
772

773 Stern, C.R., 2004. Active Andean volcanism: its geologic and tectonic setting. Revista  
774 Geologica del Chile, Vol. 31(2), p. 161-206  
775

776 Thouret, J.C., Rivera, M., Worner, G., Gerbe, M.C., Finizola, A., Fornari, M., Gonzales,  
777 K., 2005. Ubinas : the evolution of the historically most active volcano in southern  
778 Peru. Bull. Volcanol. 67(6), 557-589.  
779

780 Thouret, J.-C.; Davilla, J.; Eissen, J.-P. (May 1999). Largest explosive eruption in  
781 historical times in the Andes at Huaynaputina volcano, A.D. 1600, southern Peru.  
782 Geology 27 (5): 435–438.  
783

784 Tilling, R.I., 2009. Volcanism and associated hazards: the Andean perspective. Adv.  
785 Geosci. 22, p. 125-137  
786

787 White R., Miller A., Lynch L., and Power J. (1998). Observations of hybrid seismic  
788 events at Soufriere Hills Volcano, Montserrat: July 1995 to September 1996.  
789 Geophys Res Lett, Vol 25, N°19, 3657-3660.  
790

791 White R. (2011). Monitoring volcanoes and forecasting eruptions. Volcano Observatory  
792 Best Practices Workshop: Eruption Forecasting, 11-15 September 2011, Erice, Italy.  
793

794 Wilson, L., Sparks, R.S.J., Huang, T.C., Watkins, N.D., 1978. The control of volcanic  
795 column heights by eruption energetics and dynamics. *J. Geophys. Res.*, 83, 1829-  
796 1836.

797

798 Wooster, M.J., Zhukov, B., Oertel, D., 2003. Fire radiative energy for quantitative study  
799 of biomass burning: Derivation from the BIRD experimental satellite and comparison to  
800 MODIS fire products. *Remote Sensing of Environment* 86 (1), 83-107.

## 801 **FIGURES**

802

803 **Figure 1.** (a) Location of Ubinas Volcano and Iquique earthquake (b) Snapshot of  
804 MIROVA-derived IR image (February 10<sup>th</sup>, 2014) overlapped on Google Earth. Red  
805 pixels at the volcano's summit indicate the presence of sub-pixels thermal anomalies,  
806 related to a new incandescent magma body. UB-1 refer to the location of the Broadband  
807 seismic station and UT refer to the location of "Ubinas thermal" hot spring. (c) Details  
808 of the Ubinas caldera with the deep southern pit-crater (image from Google Earth).

809

810

811 **Figure 2** – Selected thermal images elaborated by MIROVA system over Ubinas  
812 volcano. The images (50 x 50 km) are centered on the summit of the volcano and  
813 draped over a shaded relief map. For more information on thermal maps produced by  
814 MIROVA, please refer to Coppola et al., 2015).

815

816

817 **Figure 3.** Correlation between (a) MIROVA, (b) SO<sub>2</sub> OMI, (c) plume elevation, (d)  
818 temperature of "Ubinas thermal" hot spring, (e) seismic energy released by hybrid

819 events, (f) number of hybrid seismic events. “A”, “B”, “C” and “D” stands for pictures  
820 shown in figure 4.

821

822

823

824

825

826

827 **Figure 4.** Pictures of the active vent taken from inside the summit caldera on A)  
828 February 11<sup>th</sup> 2014, B) March 1<sup>st</sup> 2014, C) March 19<sup>th</sup> 2014, and D) July 31<sup>st</sup> 2014. Note  
829 the fresh lava in the bottom on the crater in figure C.

830

831

832 **Figure 5.** Detail of datasets records between January and June 2014. (a) Plume height  
833 (above crater rim); (b) Volcanic Radiative Power (left axis) has been converted into  
834 extrusion rates (only mean values are represented) using the radiant density approach  
835 (see the text for more details). (c) Cumulative daily energy released by hybrid’s events.  
836 Note how the phases of increased energy (IIa and IIc) anticipate the phases of major  
837 magma extrusion (IIb and IId). (d) Number of daily hybrid seismic events. The  
838 occurrence of the Iquique earthquake (yellow star) is followed by an increase of the  
839 extrusion rate coupled with the gradual reduction of seismic activity represented by the  
840 hybrid’s energy and number of daily hybrid events. Dotted horizontal lines in (a) and  
841 (b) represent pre- and post- earthquake mean values outlining a 2-3 fold increase in the  
842 extrusive-explosive activity.

843

844

845 **Figure 6.** (a) Geographical relationship between the Iquique earthquake (black star) and  
846 Ubinas volcano (red circle). Focal mechanism and parameters related to the April 1<sup>st</sup>,  
847 2014 mainshock are from Lai et al. (2014). Note how the strike of Iquique faults is  
848 aligned ( $\pm 15^\circ$ ) with several Peruvian volcanoes including Ubinas, Sabancaya and El  
849 Misti volcanoes. (b) Magnitude–Distance and (c) Magnitude–Azimuth relationship of  
850 volcano-earthquake interactions with the Ubinas case represented by the red star  
851 (modified after Delle Donne et al., 2010).

852

853

854 **Figure 7.** Normal stress change along a  $165^\circ\text{N}$  oriented vertical dyke (a) and volumetric  
855 dilatation (b) produced by Iquique Mw 8.2 Earthquake, calculated at a depth of 2 and 10  
856 km, respectively. Ubinas Volcano is located in the area characterized by a weak  
857 volumetric compression and a weak increase of horizontal normal stress along a  
858 hypothetical  $165^\circ\text{N}$  oriented feeding vertical dyke.