



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Geographic variation in air temperature leads to intraspecific variability in the behavior and productivity of a eusocial insect

 This is a pre print version of the following article:

 Original Citation:

 Availability:

 This version is available http://hdl.handle.net/2318/140785

 since 2016-06-23T11:20:04Z

 Published version:

 DOI:10.1007/s10905-013-9436-y

 Terms of use:

 Open Access

 Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright

(Article begins on next page)

protection by the applicable law.



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on: Questa è la versione dell'autore

5 dell'opera: [**Journal of Insect Behavior** (DOI: 10.1007/s10905-013-9436-y]

The definitive version is available at:

La versione definitiva è disponibile alla URL: [http://link.springer.com/article/10.1007%2Fs10905-013-

10 **9436-y**]

Geographic variation in air temperature

leads to intraspecific variability in the

behavior and productivity of a eusocial insect

15

Stefania Fucini¹, Alessia Uboni¹, M. Cristina Lorenzi^{1,*}

¹ University of Turin, Department of Life Sciences and Systems Biology, via Accademia Albertina 13, 10123 Torino, Italy

20

Running head: Intraspecific differences in behavior and colony

productivity

25 * Corresponding author: M. Cristina Lorenzi, University of Turin, Department of Life Sciences and Systems Biology, Via Accademia Albertina 13, 10123 Torino, Italy, tel: +39 011 6704512, fax: +39 011 670 4508, e-mail: cristina.lorenzi@unito.it

Abstract

- 30 In primitively eusocial insects, air temperature is the environmental factor that primarily affects colony cycle. Several studies demonstrated interspecific differences in the adaptation of eusocial insects to local air temperature. Nevertheless, studies on intraspecific adaptations are rare. In this study, we investigate the influence of air temperature on local adaptations in behavior and colony productivity of *Polistes biglumis*
- 35 foundresses living in warm and cold temperate zones. We hypothesized that foundresses from warm temperate zones would show a higher activity level compared to those from cold temperate zones before brood emergence, based on differences in air temperature between the two zones. After brood emergence, we expected a reduced foundress activity level in the warm climate zone, due to workers' help. In contrast, foundresses
- 40 living in the cold-climate zone, which do not produce workers, were expected to remain active throughout the nesting season. We also hypothesized that colony productivity was higher in warm-climate colonies. As expected, warm-climate foundresses reduced their activity level after brood emergence and, with their relatively large number of workers, continued egg production throughout the nesting season. Further studies are necessary
- 45 to assess if these intraspecific differences are attributable to phenotypic plasticity or genetic divergence.

KEY WORDS: SOCIAL INSECTS; *POLISTES BIGLUMIS*; SOCIAL TRAITS; PRODUCTIVITY; INTRASPECIFIC VARIABILITY

Introduction

55

Climate is a physical force that can shape numerous life history traits of an organism and thus delineate distinct characteristics that define a species. Nevertheless, sometimes different climate regimes also generate trait differences among individuals or populations belonging to the same

species (e.g., Husby et al. 2010; Oliveira et al. 2013). Understanding the effects that climate has on the ecology and phenology of a species, as well as determining the ability of a species to adapt to different climate regimes, is essential because it gives us the ability to foreseen how

60 ecosystems will react to climate changes in the long-term (Somero 2010). In eusocial insects, air temperature can impact the activity level of colony members, their social roles, the colony cycle, and the colony productivity. Differences in local air temperatures cause, for example, marked interspecific differences in social structure and behavior in the

- 65 genus *Polistes* (Hymenoptera, Vespidae; Yamane 1996). *Polistes* paper wasps are independent-founding eusocial insects. Species living in temperate zones share common features in their colony cycle: mated females hibernate in winter, then found a nest solitarily or in association, and incur very expensive processes, like provisioning of a first brood of
- workers. After first-brood emergence, namely during the post-emergence phase (Reeve 1991), these workers take over foraging and provision a second brood that consists typically of reproductives (males and gynes).
 However, species living at different climatic conditions present differences in their colony cycle and social structure. Lorenzi and Turillazzi (1986)

75 documented that *P. biglumis* foundresses living in cool temperate zones in

the Alps were the most active females in their colonies, foraging and feeding larvae both before and after brood emergence. Foundress active behavior was most likely due to the harsh climate conditions they experienced. Low annual mean air temperature (3.7 °C) limited the length

- 80 of the colony activity to 3.5-4 months and the number of brood produced annually to less than 30 individuals. In contrast, *Polistes* spp. from warm temperate zones, living at relatively lower altitudes and higher temperatures, have a longer nesting season (approximately 6 months; Reeve 1991) and foundresses produce relatively larger broods (e.g., about
- 85 250 adults produced annually in *P. chinensis* colony; Miyano 1980). Up until now, most studies have focused on interspecific variation in *Polistes* colony productivity in response to climatic conditions, while studies on intraspecific variation have been rare (e.g., Yamane 1972; Yamane and Kawamichi 1975; Miyano 1980; Inagawa et al. 2001).
- 90 In the present study, we investigated the influence of climate on activity level and productivity of *P. biglumis* foundresses living in warm and cold temperate zones. We expected that foundresses living in warm temperate zones would exhibit higher activity level than foundresses living in cold climate during the pre-emergence phase, and that they would
- 95 reduce their activity in the post-emergence phase. We also expected that warm-climate foundresses would produce higher numbers of brood compared to their cold-climate conspecifics.

Materials and methods

100 Study sites and populations

105

During the summers of 2006 and 2007, we studied in the field foundress behavior and colony productivity of three *Polistes biglumis* populations living in the Alps and one living in the Apennines. For details on the species distribution and colony cycle, see Lorenzi and Turillazzi (1986) and Fucini et al. (2009).

Alpine populations – Alpine populations were located near Montgenèvre (Hautes Alpes, France, 1850 m a.s.l.), Ferrere (Valle Stura di Demonte, Cuneo, Italy, 1900 m a.s.l), and Carì (Val Leventina, Cantone Ticino, Switzerland, 1650 m a.s.l). The mean monthly air temperature in

110 the nesting season ranged between 8 and 13 °C. Wasps were active for about 6 hours/day (Fucini et al. 2009).

Apennine population - The Apennine population was located near Monte Mare (Abruzzo National Park, Lazio and Molise, Isernia, Italy, 1740 m a.s.l). The climate was relatively milder than the Alpine climate, with

115 mean monthly air temperatures 6-7 °C higher. The length of the wasp daily activity period was about 9 hours (Fucini et al. 2009).

A total of 478 colonies was found in 2006 (Ferrere: n = 129, Montgenèvre: n = 139, Carì: n = 71 and Monte Mare: n = 139) and a total of 349 in 2007 (Ferrere: n = 104, Montgenèvre: n = 96, Carì: n = 72 and

120 Monte Mare: n = 77). We individually marked each nest with a number and each foundress with a unique color combination using enamel paint.

Foundress activity

We conducted behavioral observations in the field on a total of 79 125 colonies in 2006 and 106 in 2007. We collected behavioral data for one/two continuous hours per nest (depending on weather conditions) between 0830 and 1700 hours local time in the Apennines and between 1000 and 1600 hours local time in the Alps, for a total of about 320 hours of observation time (149 hours in 2006 and 171 in 2007), during both the

130 pre- and post-emergence phase (see Table 1 for details).

We measured time spent off nest (i.e., foraging) by noting departure and arrival time of the foundresses. We also measured the time spent resting on the nest. Other activities performed by the foundresses during the observation time were noted but not included in the analyses.

135 Behavioral data were analyzed as percentage of the observation time spent foraging or resting on the nest.

Colony productivity

We measured nest size and colony productivity by counting the total 140 number of cells and immature brood in each nest approximately once a week throughout the nesting season. During pre-emergence, we classified large larvae and pupae as first brood, while eggs and small larvae were defined as second brood. Cells that were not occupied by immature brood (usually peripheral cells) were classified as empty.

Statistical analyses

We tested data from different years (2006 and 2007; two-sample Kolmogorov-Smirnov test) and populations in the Alps (Montgenèvre, Ferrere, and Carì; Kruskall-Wallis test) for homogeneity, before pooling

150 them (all P > 0.05). Where needed, we checked whether the data accommodated the assumptions of normality and homogeneity of variance.

Foundress activity - We used Generalized linear models (GZLMs) for binomial distributions (logit link) to compare foundress activity (i.e.

- 155 foraging/[foraging + resting]) between brood-emergence phases (pre- and post-emergence) and zones (Apennines *vs* Alps). In the GZLMs foraging was the dependent variable, zone and phase were categorical factors, and the number of larvae in the nest and the mean monthly temperature were covariates.
- 160 *Colony productivity* We used MANOVA to compare colony productivity in the Apennines and in the Alps at the end of the preemergence phase, since most of the foundress reproductive success results from eggs laid during this phase and nest size usually does not increase significantly during the post-emergence phase (Lorenzi and
- 165 Turillazzi 1986). Number of cells, number of empty cells, and first and second brood were the dependent variables and zone was the categorical factor.

Sample sizes were not consistent throughout the study period, because we discovered some nests later in the season, nest predation occurred at different stages during colony cycle, and a few nests were

missed on some surveys. Statistical tests were performed using SPSS 14.0 statistical package for Windows (SPSS Inc, Chicago, IL).

Results

175 Foundress activity

The proportion of time that foundresses spent foraging *vs* resting changed in different ways between zones relatively to brood-emergence phases (GZLM on foraging: factor zone: Wald χ^2 = 1840.271, df = 1, P < 0.0001; factor phase: Wald χ^2 = 29088.193; covariate number of larvae:

- 180 Wald χ^2 = 1242.234, df = 1, P < 0.0001; covariate temperature: Wald χ^2 = 2713.181, df = 1, P < 0.0001). Indeed, during the pre-emergence phase Apennine foundresses foraged significantly longer and rested significantly less than Alpine foundresses. These differences were significantly associated to the zone (and expectedly they were associated to
- temperature and number of larvae in the nests) (GZLM on foraging: factor zone: Wald χ^2 = 736.023, df = 1, P < 0.0001; covariate number of larvae: Wald χ^2 = 2232.533, df = 1, P < 0.0001; covariate temperature: Wald χ^2 = 157.889, df = 1, P < 0.0001). During the post-emergence phase, the difference between zones in the time foundresses spent foraging *vs*
- 190 resting reverted: Apennine foundresses spent significantly less time foraging (and rested longer) than Alpine foundresses (GZLM on foraging: factor zone: Wald χ^2 = 6894.287, df = 1, P < 0.0001; covariate number of larvae: Wald χ^2 = 576.529, df = 1, P < 0.0001; covariate temperature: Wald χ^2 = 5511.595, df = 1, P < 0.0001).

Pre-emergence colony productivity in the Apennines and in the Alps

In late pre-emergence, the number of cells built by the foundresses in the Apennine zone and Alpine zones did not vary significantly (MANOVA:

- 200 $F_{1,128} = 3.107$, P = 0.08). However, fewer cells were left empty in Apennine nests than in Alpine nests ($F_{1,128} = 26.081$, P < 0.0005). Apennine foundresses produced significantly more first brood and second brood than Alpine foundresses (first brood: $F_{1,128} = 12.429$, P = 0.001; second brood: $F_{1,128} = 20.862$, P < 0.0005; see Table 2 for mean values ±
- standard deviation).

Discussion

Our results provide evidence that cold-temperate (Alpine) and warmtemperate (Apennine) populations of *P. biglumis* differ in terms of

- 210 foundress behavior and colony productivity. Apennine foundresses were more active outside their nests, and produced a higher number of both first and second brood than those from the Alpine populations. We attribute these differences mainly to the different climate conditions of the two zones and to how foundresses allocate resources to brood production and
- 215 energetically costly tasks. Fucini et al. (2009) demonstrated that in the Apennine population the first female offspring emerges as workers, i.e., as females that are active both on and outside the nests. These workers take care of energetically costly tasks, such as nest construction, foraging and provisioning of larvae. Apennine foundresses take advantage of the
- 220 elevated number of workers emerged during the post-emergence phase, and save energy that they allot to egg production. Indeed, after the emergence of the first brood, Apennine foundresses rested longer on their nests and foraged less (Fig. 1). Foraging is an energetically costly task that reduces the opportunity of direct reproduction in social insects in
- 225 general (Schmid-Hempel and Wolf 1988), and in *Polistes* wasps in particular (Markiewicz and O'Donnell 2001). Thus, *P. biglumis* foundresses living in the Apennines had brood production patterns similar to those of other *Polistes* species, which live at lower altitude in warm climate conditions (e.g., Pardi 1942, 1946; Miyano 1986).
- 230 In the Alpine populations, the situation was strikingly different. Cold climatic conditions reduced the foundress activity during the pre-

emergence phase, and thus limited the production of first and second brood with respect to the Apennine foundresses. Alpine foundresses were relatively active during the post-emergence phase, compared to the

- 235 Apennine foundresses. The difference was not only due to the small numbers of first brood produced in the Alps, but also to the fact that this brood emerged as future gynes (Fucini et al. 2009). A behavioral comparison between the pre- and post-emergence phase reveals that Alpine foundresses spent about 30% of the time out of their nests both
- 240 before and after brood emergence (Fig. 1). Resource-consuming activities, such as foraging, may reduce the resources for egg production (Markiewicz and O'Donnell 2001). The cold weather contributes to reducing fertility as well (Karsai 1997). The strategy of Alpine *P. biglumis* foundresses consists in forgoing worker production and continuing to be
- 245 active over the whole nesting season. By doing so, they ensure the production of reproductive brood in an extremely short daily and seasonal activity period (Fucini et al. 2009).

The nests in the Apennines had less peripheral empty cells than those in the Alps. Peripheral empty cells may serve for thermoregulation, i.e. to 250 protect immature brood from cold weather (Yamane 1969, 1972; Yamane and Kawamichi 1975; Lorenzi and Turillazzi 1986; Hozumi et al. 2008). Our results support this hypothesis.

In conclusion, we observed that *P. biglumis* foundresses can adapt to local conditions, exhibiting high flexibility in their social traits. Air

temperature seems to play a key role in influencing the activity level of *P. biglumis* foundresses (indeed it was a significant covariate in our models).
 Future studies are needed to assess if the observed variability in *P.*

biglumis populations is determined either by phenotypic plasticity or by genetic differentiation.

Acknowledgments

We are grateful to the many students who helped with collecting field data. We thank Valeria Di Bona for helpful discussions. The *P. biglumis* population located in Valle Stura di Demonte was found thank to Prof. Augusto Vigna-Taglianti. We thank C. Sulli, Servizio

265 Scientifico Ambientale, for permission to work in the Parco Nazionale d'Abruzzo, Lazio e Molise. Funding for this work was obtained from the MURST ex 60% (to M. C. L.).

References

270	Fucini, S., Di Bona, V., Mola, F., Piccaluga, C. and Lorenzi, M. (2009). Social wasps without workers: geographic variation of caste expression in the paper wasp Polistes biglumis. Insectes Soc. 56(4): 347-358.
275	Hozumi, S., Yamane, S. and Katakura, H. (2008). Building of extra cells in the nests of paper wasps (Hymenoptera; Vespidae; Polistes) as an adaptive measure in severely cold regions. Sociobiology 51(2): 399-414.
280	Husby, A., Nussey, D. H., Visser, M. E., Wilson, A. J., Sheldon, B. C. and Kruuk, L. E. (2010). Contrasting patterns of phenotypic plasticity in reproductive traits in two great tit (Parus major) populations. Evolution 64(8): 2221-2237.
285	Inagawa, K., Kojima, J., Sayama, K. and Tsuchida, K. (2001). Colony productivity of the paper wasp Polistes snelleni: Comparison between cool-temperate and warm-temperate populations. Insectes Soc. 48(3): 259-265.
	Karsai, I. (1997). Brood patterns in wasp combs: the influence of brood on egg-laying and building by adults. Ethol. Ecol. Evol. 9(1): 27-44.
290	Lorenzi, M. and Turillazzi, S. (1986). Behavioural and ecological adaptations to the high mountain environment of Polistes biglumis bimaculatus. Ecol. Entomol. 11(2): 199-204.
295	Markiewicz, D. A. and O'Donnell, S. (2001). Social dominance, task performance and nutrition: implications for reproduction in eusocial wasps. J. Comp. Physiol., A 187(5): 327-333.
300	Miyano, S. (1980). Life tables of colonies and workers in a paper wasp, Polistes chinensis antennalis, in central Japan (Hymenoptera: Vespidae). Res. Popul. Ecol. 22(1): 69-88.

Miyano, S. (1986). Colony development, worker behavior and male production in orphan colonies of a Japanese paper wasp, Polistes chinensis antennalis Pérez (Hymenoptera: Vespidae). Res. Popul. Ecol. 28(2): 347-361.

Oliveira, R., Carvalho, A. and Schlindwein, C. (2013). Plasticity in male territoriality of a solitary bee under different environmental conditions. J. Insect Behav., 10.1007/s10905-013-9385-5.

310

305

Pardi, L. (1942). Ricerche sui Polistini V. La poliginia iniziale di Polistes gallicus (L.). Bollettino dell'Istituto di Entomologia dell'Università di Bologna 14: 1-106.

315 Pardi, L. (1946). Ricerche sui Polistini VI. La "dominazione" e il ciclo ovarico annuale in Polistes gallicus (L.). . Bollettino dell'Istituto di Entomologia dell'Università di Bologna 15: 25-84.

Reeve, H. K. (1991). Polistes. In K. G. Ross and R. W. Matthews(eds),The social biology of wasp, Cornell University Press, Ithaca, pp. 99-148.

Schmid-Hempel, P. and Wolf, T. (1988). Foraging effort and life span of workers in a social insect. J. Anim. Ecol. 57: 509-521.

325 Somero, G. (2010). The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. J. Exp. Biol. 213(6): 912-920.

Yamane, S. (1969). Preliminary observations on the life history of two
 Polistine wasps, Polistes snelleni and P. biglumis in Sapporo. Journal of the Faculty of Science, Hokkaido University, Series VI, Zoology 17(1): 78-105.

Yamane, S. (1972). Life cycle and nest architecture of Polistes wasp in the Okushiri Island, northern Japan (Hymenoptera, Vespidae). Journal of the Faculty of Science, Hokkaido University, Series VI, Zoology 18: 440-458.

Yamane, S. and Kawamichi, T. (1975). Bionomic comparison of Polistes biglumis (Hymenoptera, Vespidae) at two different localities in Hokkaido,

340 northern Japan, with reference to its probable adaptation to cold climate. The Entomological Society of Japan, Kontyû, Tokio 43(2): 214-232.

345

Yamane, S. (1996). Ecological factors influencing the colony cycle of Polistes wasps. In S. Turillazzi and M. J. West-Eberhard(eds), Natural history and evolution of paper-wasps, Oxford University Press, 75, pp. 97. Table 1. Total number of colonies observed, divided by study site, and total observation hours.

	Pre-emergence		Post-emergence	
	2006	2007	2006	2007
Ferrere	17	9	2	6
Montgenèvre	12	9	1	1
Carì	10	15	3	4
Monte Mare	27	40	7	22
Total colonies	66	73	13	33
Total observation hours	125	121	24	50

Table 2. Colony productivity in Apennine and Alpine colonies in the late pre-

355	emergence phase.	Values are means	± standard deviations.
-----	------------------	------------------	------------------------

	Pre-emergence		
	Apennine	Alpine	
Total cells	48.45 ± 7.63	44.75 ± 11.20	
Empty cells	9.15 ± 6.45	17.66 ± 8.79	
First brood	14.42 ± 2.97	10.80 ± 5.63	
Second brood	24.91 ± 6.91	16.29 ± 10.05	

Fig 1. Proportion of time that Apennine and Alpine foundresses spent foraging (grey) and resting (black) during the pre- and post-emergence phase. Column heights represent mean percentage values. Vertical bars represent ± standard deviation.