



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Screening and evolution of volatile compounds during ripening of 'Nebbiolo', 'Dolcetto' and 'Barbera' (*Vitis vinifera* L.) neutral grapes by SBSE-GC/MS.**This is the author's manuscript***Original Citation:**Availability:*This version is available <http://hdl.handle.net/2318/1571310> since 2020-04-01T15:40:41Z*Published version:*

DOI:10.1007/s00217-015-2626-4

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**Screening and evolution of volatile compounds during ripening of ‘Nebbiolo’, ‘Dolcetto’ and
‘Barbera’ (*Vitis vinifera* L.) neutral grapes by SBSE-GC/MS.**

Antonio Carlonmagno^{1)*}, Andrea Schubert²⁾, Alessandra Ferrandino¹⁾

¹⁾ Centro di Ricerche in Viticoltura ed Enologia, Università di Torino

C.so Enotria 2/C

12051 Alba (CN), Italy

²⁾ DISAFA, Dipartimento di Scienze Agrarie, Forestali, Alimentari, Università di Torino

Largo P. Braccini 2

10095 Grugliasco (TO), Italy

*Corresponding author:

Tel.: +39 0173-441486; fax: +39 0173-441349

antonio.carlonmagno@unito.it




Figure 1

Evolution of total pre-fermentative volatiles, C6 compounds, aldehydes, alcohols, benzenoids, monoterpenes, C13-norisoprenoids and sesquiterpenes during ripening in Nebbiolo, Dolcetto and Barbera grape berries.
(DPV = Days Post Véraison).