

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Pliocene and Early Pleistocene carpological records of terrestrial plants from the southern border of the Po Plain (northern Italy).**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/151087> since 2016-06-29T14:36:25Z

*Published version:*

DOI:10.1016/j.revpalbo.2014.10.007

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in *REVIEW OF PALAEOBOTANY AND PALYNOLOGY*, 218, 2015, 10.1016/j.revpalbo.2014.10.007.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.revpalbo.2014.10.007

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0034666714001511>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/151087>

Pliocene and Early Pleistocene carpological records of terrestrial plants from the southern border  
of the Po Plain (N-Italy)

Edoardo Martinetto <sup>a\*</sup>, Giovanni Monegato <sup>b</sup>, Andrea Irace <sup>b</sup>, Stefano Claudio Vaiani <sup>c</sup>, Elena  
Vassio <sup>a</sup>

<sup>a</sup> Dipartimento di Scienze della Terra, Università degli Studi di Torino. Via Valperga Caluso 35, I-  
10125 Torino, Italy; edoardo.martinetto@unito.it

<sup>b</sup> CNR - IGG Institute of Geosciences and Earth Resources. Via Valperga Caluso 35, I-10125  
Torino, Italy

<sup>c</sup> Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università degli Studi di  
Bologna. Via Zamboni 67, I-40126 Bologna, Italy, stefano.vaiani@unibo.it

**KEY WORDS** - Fruits and seeds, biochronology, Italy, Pliocene and Early Pleistocene.

**ABSTRACT** - The analysis of fruit and seed assemblages in Pliocene and Early Pleistocene  
successions of the southern border of the Po Plain showed that these deposits contain an  
interesting record of the terrestrial palaeoflora which integrates the information provided by the  
extensive pollen record available in this area. We could fix in the geological time scale, locally  
with a precision of 100 ka, 14 carpoflora-bearing layers spanning from 5.1 to 0.9 Ma, thus  
obtaining a framework which is not available in any other parts of Europe. This permitted us to  
provide precise ages for the occurrence of fruits and seeds of a few plant groups with a proven late  
Cenozoic differentiation (*Carex*, *Hypericum*, *Thymelaea*), as an important support to future  
phylogenetic analyses. However, our main aim is to evaluate the usefulness of our dataset for

biochronological analyses. At first glance, the fossil record of several species showed an interesting distribution, often limited to one to three chronostratigraphic stages among the Zanclean, Piacenzian, Gelasian and Calabrian. An elementary palaeoclimatic characterisation of the individual species, but also of the plant assemblages, suggests that a temperature decrease is one of the causes of the observed palaeofloral differences. The input of the new data into the CENOFITA database led to an improvement of the chart of the distribution of carpological taxa in the Pliocene and Early Pleistocene of northern Italy. Within this new chart, the carpological species were clustered in groups with a similar chronologic distribution, which are useful for the biochronological interpretation of undated carpological assemblages.

## 1. Introduction

The Pliocene and Early Pleistocene (Zanclean-Calabrian) continental palaeoflora of northern Italy, as well as its ecological and climatic interpretation, has been documented by several papers dealing with rather long pollen sequences (Lona and Bertoldi, 1973; Ravazzi and Rossignol Strick, 1995; Bertini, 2001 and 2010) and by macrofloral records referred to more limited chronostratigraphic intervals (Gregor, 1990; Mai, 1995; Basilici et al., 1997; Martinetto, 1995, 1999, 2001b and 2003; Martinetto and Sami, 2001; Martinetto et al., 2007 and 2012; Ghiotto, 2010).

Due to the possibility to precisely identify several carpological taxa at species level much detailed macrofloral data are presently provided by fruits and seeds (carpological remains), whose Zanclean-Calabrian records from northern Italy (34 localities in Fig. 1) have been listed into the CENOFITA database (Martinetto and Vassio, 2010). The richest carpological assemblages are reported mainly from shallow marine and continental sections of NW Italy (Fig. 1), whose precise

chronological framing is usually problematic (Bertoldi and Martinetto, 1996; Martinetto, 1999; Cavallo and Martinetto, 2001; Martinetto et al., 2007).

Since the first analyses on the Pliocene fruit and seed assemblages in northern Italy (Gregor, 1986; Martinetto, 1995) it was noticed that the species composition varied conspicuously from older to younger strata, and was able to provide useful information on the floral change from 5 to 1 Ma. Therefore, a long-lasting research for fruit-bearing layers in stratigraphically well-framed sections has been carried out in northern and central Italy, discontinuously, for the last 20 years (Martinetto, 1994, 1999 and 2001a; Martinetto et al., 2007). Recently, new interest for the examination of the material from the studied sections was raised by studies on continental sections of NW-Italy, where the concomitant occurrence of some carpological taxa seems to be crucial for the chronologic framing of the deposits (Irace et al., 2012).

The first aim of this paper is to point out the presence of rather rich assemblages of carpological remains of terrestrial plants in Pliocene and Early Pleistocene successions exposed along the southern border of the Po Plain, thus improving the preliminary reports by Gregor (1986 and 1990). Additionally, we wanted to obtain an updated scheme reporting the occurrence of such plant taxa in the Zanclean-Calabrian of northern Italy and we tried to recognize in this fossil record the signal of changes in the terrestrial flora (and vegetation). We also wanted to discuss the possible causes of the appearance and disappearance of the plant species in the fossil record, by considering environmental changes, migrations as well as phylogeny of those plant taxa with a proven diversification in the last 5 Ma (e.g., Thymelaeaceae: Galicia-Herbada, 2006; Cyperaceae: Jiménez-Mejías and Martinetto, 2013).

## 2. Geological setting and material

## 2.1 Geological setting

During the Pliocene-Early Pleistocene interval the southern side of the Po Plain area was affected by the compressional N-NE-verging Apennine tectonic activity, which was responsible for the formation of coeval but distinct and structurally separated thrust-top and foredeep basins (e.g., Ricci Lucchi, 1986; Ghielmi et al., 2010).

At the western termination of the Po Plain, the Plio-Pleistocene Piedmont Basin evolved as a wide wedge top basin, bounded to the north by the westernmost arc of the Padane thrust front (Fig. 2). To the south-east, the Castell'Arquato Basin acted as a closed piggy-back basin on top of Apennine thrust fronts. The Adriatic Foredeep developed in the south-eastern sector of the Po Plain.

During the Piacenzian-Calabrian time span, transitional to continental successions were deposited in the Piedmont Basin, while deep water to shelfal units filled the Castell'Arquato and the Adriatic Foredeep basins (Fig. 3). In these basins paralic and continental facies appeared only in the Calabrian.

## 2.2 Stratigraphic sections and carpological material

The fossil fruit and seed material studied in this work comprises both previously unpublished material from freshly analysed sediment samples (East area samples in Fig. 4) and taxonomically revised collection material (West area samples, Fig. 4). All these carpological samples were prepared starting from bulk sediment samples collected in well constrained sections along the southern side of the Po Plain (Fig. 2). Seven Pliocene-Lower Pleistocene successions of marine to transitional environment were taken into consideration in the East area, and compared with the existing records of 3 sections from the West area (Fig. 3). In both areas, several successions, straddling the Pliocene-Early Pleistocene time interval (Barbieri, 1967; Colalongo, 1969;

95 Cremonini et al., 1969; Iaccarino and Monegatti, 1996; Rio et al., 1997; Monegatti et al., 2002;  
96 Roveri and Taviani, 2003; Calabrese and Di Dio, 2009), are particularly suitable for  
97 palaeobotanical studies due to the frequent occurrence of layers with terrestrial plant remains.  
98 Recent magnetostratigraphic analyses (Gunderson et al., 2012, 2014) fixed to the end of the Early  
99 Pleistocene some layers with rich palaeofloras in the Stirone (Gregor, 1990) and Enza (Iaccarino  
100 and Monegatti, 1996) sections; the same age has been proposed by Muttoni et al. (2011) for the  
101 leaf-bearing Oriolo section (Martinetto and Sami, 2001), which is not reconsidered here because  
102 of the scarcity of carpological remains.

103 The palaeocarpological analyses of samples collected in the West area (Fig. 3) concerned the  
104 record of two beds from the Argille Azzurre Fm. (BR3, PO1), already described by Martinetto  
105 (1995) and Cavallo and Martinetto (1996). Micropalaeontological (Casnedi, 1971; Pavia et al.,  
106 1989; Violanti, 2012) and stratigraphic (Vigna et al., 2010) studies assigned the marine sediments  
107 of this formation, in the studied area, to the Zanclean, and precisely to biozones MPL2 and MPL3  
108 (5.1-3.8 Ma). Three other layers (RDB1, RDB6, RT1), sampled in the Villafranchian type area  
109 (Fig. 2), represent the continental deposits of the “Villafranchian lower complex” (*sensu* Carraro,  
110 1996), which were ascribed to the middle Piacenzian (Boano et al., 1999; Napoleone et al., 2003;  
111 Martinetto et al., 2007). In this basin dated sites with Gelasian or Calabrian carpological  
112 assemblages are lacking.

113 In the East area, seven sections (1-7 in Fig. 3) were sampled for carpological analysis. Among  
114 these, sections 1-4 belong to the Castell’Arquato Basin (Roveri and Taviani, 2003), which  
115 includes the historical Piacenzian stratotype (e.g., Mayer-Eymar, 1858; Pareto, 1865; Barbieri,  
116 1967; Raffi et al., 1989). Sections 5-7 belong to the Adriatic Foredeep (Colalongo et al., 1982a and  
117 1982b; Ricci Lucchi, 1986; Iaccarino and Monegatti, 1996).

The samples are located in stratigraphic logs (Fig. 3) that represent a portion of longer stratigraphic successions. As a whole, 21 samples of fruit and seed-bearing deposits were collected, 9 of which have been so-far analysed, as reported below:

1) **Chiavenna Rocchetta section** (Roveri and Taviani, 2003). This succession was deposited in a shelf environment during the Piacenzian. The sample CVE2 comes from a silty sand layer, rich in marine mollusc shells and with scattered terrestrial plant remains, cropping out on the left bank of the Chiavenna Creek, below the first calcarenitic body. According to the magnetostratigraphic data reported by Roveri and Taviani (2003) for a neighbouring drill-core, the CVE2 bed was attributed to the subchron Kaena.

2) **Lugagnano Quarry section** (Barbieri, 1967). This stratigraphic interval is assigned to the lower part of the Zanclean (MPL2 Zone) because of the occurrence of *Globorotalia margaritae* (listed as *Globorotalia hirsuta* by the author) and the absence of *Globorotalia puncticulata*. Benthic foraminiferal assemblages are highly diversified and include mainly *Bulimina aculeata minima*, *Bulimina costata*, *Cibicides pseudoungerianus*, *Siphonina reticulata*, *Sphaeroidina bulloides*, *Uvigerina peregrina*, *Uvigerina pigmea* and *Uvigerina rutila* (Barbieri, 1967). These species are considered to reflect deposition within an outer shelf or upper slope environment (e.g. Blanc-Vernet, 1969; Rasmussen, 2005; Schweizer, 2006). The presence of rare shallow marine taxa, specifically *Ammonia beccarii*, *Criboelphidium semistriatum*, *Elphidium macellum* and *Elphidium complanatum*, suggests downslope transport during deposition. The sample LU1 was collected in a sandy silt layer rich in plant remains, in the middle portion of Barbieri's (1967) section 2. Gregor (1990) reported a list of carpological taxa gathered by hand-picking on the outcrop of the Lugagnano Quarry section, which included the species *Visnea germanica*.



141 3) **Castell'Arquato composite section** (Monegatti et al., 1997 and 2002; Roveri and Taviani,  
142 2003). This composite section includes Zanclean to Calabrian marine sediments. Carpological  
143 samples have been collected in subsections Monte Falcone and Arda.

144 3a) **Monte Falcone subsection** (Monegatti et al., 2002). This succession includes silt and sand  
145 passing upward to dominant calcarenite, deposited within a shallow marine to coastal  
146 environment. Magnetostratigraphic analyses reveal the occurrence of normal polarity interval  
147 2An.1n (Gauss Chron) of late Piacenzian age (Monegatti et al., 1997; 2002). The sample CQ1 was  
148 collected in the lower portion of the outcropping succession, within shallow marine silty sands rich  
149 in mollusc shells and with scattered terrestrial plant remains, deposited at 30-60 m water depth  
150 (Monegatti et al., 2002).

151 3b) **Arda subsection** (Dominici, 2001; Roveri and Taviani, 2003). The succession was deposited  
152 in a shoreface to inner shelf environment (5-40 m water depth). This represents the youngest part  
153 of the composite section, assigned to the Gelasian-Calabrian transition (Calabrese and Di Dio,  
154 2009). The collection of samples from as much as four layers (AD3, AD6, AD7, AD9 in Fig. 3)  
155 was stimulated by the detailed stratigraphic and palaeoenvironmental analyses of Dominici (2001);  
156 however only the sample from the AD3 layer has been so-far analysed.

157 4) **Stirone section** (Papani and Pelosio, 1963). The part of the historical Stirone River succession  
158 (Cremaschi, 1982; Dominici, 2001; Gunderson et al., 2012) considered in this work includes  
159 Gelasian and Calabrian sediments, showing shelf deposits passing upward to a cyclic alternation  
160 of shallow marine to continental deposits (Fig. 3). In the Calabrian portion (Stirone-Laurano in  
161 Martinetto, 1999), two samples were collected in layers rich of terrestrial plant remains (Fig. 3):  
162 LA1 comes from freshwater dark grey massive muds, immediately preceding the base of the  
163 Jaramillo subchron (Gunderson et al., 2012), which were rich in terrestrial plant remains  
164 associated to tree stumps in growth position; LA3 was collected in shallow-marine sands referable  
165 to the last 200 ka of the Matuyama chron (C1r.1r; Cremaschi, 1982; Gunderson et al., 2012).

These samples correspond, respectively, to beds S3 and S7-8 of Gregor (1986, 1990), from which this author reported abundant carpological remains, not yet subjected to a detailed taxonomic treatment. Mai (1994) reported a cone of *Picea florschuetzii* Van der Hammen from “Millepioppi”, the locality where Gregor’s bed S3 (our LA1) was in outcrop during the 1980s.

5) **Enza Creek section** (Iaccarino and Monegatti, 1996). The lower part of this succession consists of shelf clays locally interbedded with calcarenites and sands. The upper part includes mainly coastal sands passing upward to continental sediments (Iaccarino and Monegatti, 1996). An age model for the section has been recently provided by Gunderson et al. (2014), combining magnetobiostratigraphic data with cosmogenic radionuclide burial dating and cyclostratigraphy. According to these authors, the section of Fig. 3 is almost entirely attributed to the Calabrian. The boundary between Calabrian and “Ionian” has been traced within continental sediments. Seven samples were collected (Fig. 3), however, only sample EZ38, providing a rich carpological assemblage, has been analysed so-far. Magnetostratigraphic analyses reveal that this sample has been collected in proximity of the upper boundary of Jaramillo subchron (Gunderson et al., 2014), allowing an age attribution of about 1.0 Ma. The depositional environment of plant-bearing samples has been investigated through analyses of foraminiferal assemblages in the whole section (Appendix A), showing an overall regressive tendency from inner shelf (EZ20) to coastal (EZ30-40) and continental environments (EZ50-51).

6) **Santerno River section** (Colalongo, 1969; Cremonini et al., 1969). The part of the Santerno River succession considered in this work, includes open marine clays with interbedded sands (Cremonini et al., 1969). The Gelasian-Calabrian boundary occurred in the lowermost portion of the analyzed section (Fig. 3; Colalongo, 1969). The sample SNT1 comes from a sandy layer, rich in phytoclasts (Gastaldo, 1994), which cropped out in the year 2001, close to the Gelasian-Calabrian boundary (Fig. 3). Foraminiferal assemblages of a 2.6 m-thick succession including the

sampled layer, have been investigated (Appendix A), in order to improve the biostratigraphic and palaeoenvironmental framework of plant-bearing sediments. Foraminiferal associations reveal an outer shelf depositional environment attributed to the *G. cariacensis* biozone, according to comparable assemblages observed in the nearby Lamone section by Vaiani (2000).

7) **Marecchia River section** (Colalongo et al., 1982a; Rio et al., 1997). The part of the Marecchia River succession considered in this work, consists of slope sediments, represented by clays and sapropels, with subordinate sands. These deposits are ascribed to the late Piacenzian (Gauss Chron, normal polarity interval 2An.1n) according to the stratigraphic scheme reported by Rio et al. (1997). Three samples were collected in sandy beds below (MR1) and above (MR2, MR3) the sapropel bearing intervals M1 and M2 of Rio et al. (1997), possibly corresponding to sapropel cluster O of Roveri and Taviani (2003). Only the sample MR3 has been so-far analysed.

As a whole the analyzed sections are substantially well-calibrated within a detailed chronostratigraphic frame. This allows us to propose a rather precise chronological framing (locally with resolution of ca. 100 ka) of the plant-bearing layers (Tab. 1) and to place the East area analyzed material into a definite chronological order (Fig. 4).

### 3. Methods

This research was not intended to complete the analyses in a short time, since several weeks of analytic work are needed for each sample; rather, the aim was to sample and preserve for long time the carpological record provided by a few layers, often exposed in ephemeral sections, which had been calibrated, more or less precisely, from a chronostratigraphic point of view. In order to improve our knowledge of the biochronologic range of some key-species found in the late Cenozoic of northern Italy, we decided to analyse their occurrences in ten chronostratigraphically framed sections, cropping out along the southern border of the Po Plain (Fig. 2), where we

distinguished a West area (Piedmont Basin in Fig. 3) from an East area (Castell'Arquato Basin and Adriatic Foredeep in Fig. 3). The stratigraphic position of the sampled layers is indicated in synthetic logs (Fig. 3), with the aim to favour the connection between palaeobotanical data and the results of other, already rich, stratigraphic records (mainly magnetostratigraphy, foraminifer and nannoplankton biostratigraphy).

Methodological approaches to the late Cenozoic biochronology of terrestrial plant fossils are discussed by Martinetto and Ravazzi (1997), Martinetto (1999), and Martinetto et al. (2007), who evidenced the scattered occurrence of plant macrofossils in the stratigraphic record. The biochronological analysis carried out in this paper proceeded with the input of newly obtained occurrence data into the CENOFITA database (Martinetto and Vassio, 2010), in which the proved or inferred age of each fossil site is reported. Those species whose chronostratigraphic distribution in N-Italy appeared to be restricted to one or a few stages (Zanclean, Piacenzian, Gelasian, Calabrian) were singled out in order to evaluate their biochronologic potential, depending on biological and ecological features (habitat, growth form, dispersal mechanism, etc.), which are discussed below.

The collection of carpological samples was carried out outside the frame of specific projects, but profited from the results of stratigraphic studies carried out by other authors (Barbieri, 1967; Cremonini et al., 1969; Colalongo, 1969; Iaccarino and Monegatti, 1996; Rio et al., 1997; Dominici, 2001; Monegatti et al., 2002; Roveri and Taviani, 2003), which often permitted a correct lithostratigraphic positioning of the plant-bearing layers. Furthermore, in two of the studied sections a new sampling for foraminifer assemblages has been carried out in the attempt to improve the biostratigraphic framing of the plant-bearing beds (see Appendix A).

The labels used for the sampled layers are indicated in Table 1, and they are also used for indicating carpological samples, with the addition of a conventional code of 3 characters which designates all the carpological records of each individual sample in the CENOFITA database

(Letter-numer-Letter, e.g. A0B). The sediments (usually 4-30 dm<sup>3</sup>) were treated according to the method described by Basilici et al. (1997), and floating material was collected with final mesh size of 0.3 mm. As for the sinking material, the removal of the clastic matrix left a very limited volume of plant-bearing residue (maximum 20% of the original volume). When completely dry, this was stored for long-term preservation, and analysed after variable times, up to 25 years after preparation (Stirone section). For the picking of fruit and seeds the residue of each sample was separated in 2 fractions: > 3 mm and < 3 mm: the coarser fraction was examined at naked eye, and the finer one under a Wild M3B stereomicroscope. Only a few selected carpological species with characteristic morphology (those listed in Fig. 4) were picked out, and we took care to proceed in a uniform way for all the samples. The identification of these species has been carried out by comparison to the CENOFITA collection of late Cenozoic carpological material (Martinetto and Vassio, 2010), which had been the subject of several taxonomical studies (Basilici et al, 1997; Cavallo and Martinetto, 2001; Martinetto, 2001a and 2001c; Mai and Martinetto, 2006; Jacques et al., 2011; Vassio, 2012; Jiménez-Mejías and Martinetto, 2013; Martinetto and Festa, 2013). In a few cases comparison to modern material has been necessary, and it was carried out in the carpological collection of the Museum für Naturkunde of Berlin (label MFN, without number) and in the Modern Carpological Collection of the Turin University (Martinetto et al., 2014), whose samples are indicated by the label MCC and designated by a collection number (e.g. MCC0001). Images of the fossil fruits and seeds (Plate I) were obtained by using a Leica M205 C optical microscope integrated with Leica Microsystems' Leica DFC290 HD digital camera and Leica Application Suite software.

For the ecological and climatic characterisation of the fossil taxa we conducted a search for the nearest living relatives (NLRs), and we consulted the data contained in the ClimGrim (Grimm and Denk, 2012) and PALAEOFLORA databases (Utescher and Mosbrugger, 2013), as well as those reported in the supplementary material of Denk et al. (2013). The reconstructed thermal

requirements of the species treated in this paper have been indicated by boundary values of Mean Annual Temperature (MAT in °C), as explained in Fig. 4. Furthermore, we determined the Köppen signature (of several species as explained by Denk et al. (2013). When the "potential modern analogs" (our NLRs) were not reported by these authors, we analysed the modern geographic distribution and we checked their Köppen climatic zone (see Supplementary material). We consulted Boufford (1997) for *Boehmeria*, Galicia-Herbada (2006) for *Thymelaea*, Fang et al. (2011) for Chinese plants, Hansen and Sunding (1993) for Macaronesian plants, Stevens (2014) for *Phytolacca* and *Stemona*, and Thomson et al. (1999a, 1999b) for North American plants.

#### 4. Taphonomy

As discussed above, 10 of the 14 studied fruit and seed assemblages were formed in a marine or nearshore palaeoenvironment. Unfortunately there is no actuopalaeobotanical study dealing with the taphonomy fruits and seeds of terrestrial plants deposited on marine bottoms, so we do not know the degree of fidelity of such fossil record with respect to the contemporary flora and vegetation of the terrestrial environment. Zavala et al. (2012) illustrated, with specific reference to the Plio-Pliostocene sediments of the Eastern area, that the origin of the concentration of terrestrial plant remains in marine successions can be explained by a massive transport of terrestrial phytoclasts to the marine bottoms, carried out by the rivers. This allows us to interpret such marine assemblages in the light of the present knowledge on the relationships between vegetation and fruit and seed assemblages in fluvial deposits (Cappers, 1993; Gee, 2005; Vassio and Martinetto, 2012). These studies indicate that fruit and seed assemblages provide a record of the flora and vegetation of a broad area of the river's catchment, although with several mismatches. For our

purposes, the above-mentioned studies also suggest that, given a fixed type of terrestrial flora, we must assume that the list of plants obtained by analysing fruits and seeds in different samples of marine sediment (with plant phytoclasts of terrestrial provenance) would be rather different from one sample to the other, just due to taphonomical processes. For this reason a large number of well-calibrated marine samples would be needed to reliably assess when a terrestrial plant species appeared, disappeared or became more or less frequent in the surrounding terrestrial areas.

In addition, the geological context of the studied marine successions suggests that the possibility of reworking of fossil plant remains has to be considered. In fact the progressive uplift of the northern Apennine foothills determined the exposure and erosion of the Pliocene sediments, and the recycling of their plant phytoclasts into the Pleistocene marine and continental basins is theoretically possible. There are very few reports in the literature about fruits and seeds in modern or fossil assemblages for which reworking from older sediments could be demonstrated (e.g., Thomasson, 1991). In light of the poor available evidence we suggest that reworked fossil fruits and seeds seem to be extremely rare.

## 5. Palaeobotanical results

### 5.1 Sample analysis

The analysis of the 9 palaeocarpological samples from the East area sections showed that fruits and seeds made up an extremely subordinate component of the phytoclasts (ca. 1 fruit/seed per 100-1000 wood fragments), and their taxonomic analysis indicated a prevalence of terrestrial plants, although fruits of the seagrass *Cymodocea* cf. *nodosa* occurred in several samples (see below). The palaeocarpological analysis of these samples did not permit so far to obtain a

complete list of taxa, but the presence/absence of the species listed in Fig. 4 was carefully checked into the residues of all samples.

The number of carpological species and specimens represented in different samples varies greatly. Samples CQ1, CVE2, EZ38, LA1, LU1 yielded poor assemblages in terms of number of both species (<20) and fruit/seed specimens (<50). Rich carpological assemblages, certainly with more than 30 species and more than 50 (but often >200) carpological specimens, were obtained from layers AD3, LA3, MR3 and SNT1.

## 5.2 Carpological data analysis

By analysing the whole CENOFITA dataset for N-Italy (Fig. 5) we could gather most species into six groups on the basis of the similar biochronologic distribution: Group 1, only Zanclean records; Group 2, records in the Zanclean and in the late Piacenzian (from 3.0 to 2.8 Ma); Group 3, records in the Zanclean and in the middle-late Piacenzian (from 3.3 to 2.8 Ma); Group 4, definite records in the (Zanclean-) Piacenzian, and clues for the persistence after the Piacenzian/Gelasian transition given by occurrences in localities under investigation (Irace et al., 2012); Group 5, records from (Zanclean-) Piacenzian to Calabrian, but not from the Middle Pleistocene onwards; Group 6, records in the (Gelasian-) Calabrian, and partly also from the Middle Pleistocene onwards. In general, the East area record (Fig. 4) fitted within the biochronologic distribution of each species provided by the analysis of the N-Italian CENOFITA dataset (Fig. 5), and only two species showed a broader distribution in the East area record (Fig. 4): *Eurya stigmosa* and *Sparganium nanum*, occurring here newly in the Gelasian. As an alternative explanation to the longer survival of these two species in the East area, we must consider the possible reworking of fruit and seed specimens from older sediments, already discussed above. Additionally, the East area samples did not contain any record of a few key-species which, according to the data of the N-Italian



336 CENOFITA database, would seem to be potentially useful for the biochronologic framing of the  
337 West area fluvial-alluvial successions: *Actinidia faveolata*, *Boehmeria lithuanica*, *Carex*  
338 *flagellata*, *Cryptomeria rhenana*, *Datisca gratioloides*, *Meliosma wetteraviensis*, *Styrax maximus*,  
339 *Schoenoplectus isolepioides*, *Symplocos schereri*. All of these occur in the West area samples  
340 reconsidered in this work (Fig. 4).

341 We also observe the lack in the East area samples of any record of Group 1 ("Zanclean") elements  
342 (Fig. 4), even in the Zanclean assemblage of Lugagnano (LU1). However, Gregor (1990) reported  
343 the occurrence in this locality of *Visnea germanica*, which suggests that we missed the Group 1  
344 elements due to relatively small size of our sediment sample. The West area record shows that  
345 Group 1 elements are well represented in the Zanclean sample of Pocapaglia (PO1, over 100 dm<sup>3</sup>  
346 of sediment analysed: Cavallo and Martinetto, 1996; Repetto, 1997), whereas only 4 out of 10  
347 species occur in the less intensively sampled layer of Breolungi (BR3, 19 dm<sup>3</sup>).

348 The climatic characterisation of the species treated in this paper is shown by the boundary values  
349 of Mean Annual Temperature (MAT in °C) reported in Fig. 4, which permit to point out several  
350 thermophile species (e.g. MAT>9°C or >15°C, etc.), as well as cool-tolerant species (e.g.  
351 MAT>5°C or >3°C, etc.) and a few warm-intolerant ones (e.g. MAT<14°C or <17°C).  
352 Furthermore, the Köppen climate type for the Pliocene plant assemblages (BR3, PO1, RDB1,  
353 RDB6, RT1, LU1, CVE2, MR3, CQ1) can be easily determined as Cfa, being this the single type  
354 shared by all the occurring plant taxa (see Supplementary material). For the Early Pleistocene  
355 assemblages (AD3, EZ38, LA1, LA3 and SNT1) the main Köppen signature is for the Cfa and Cfb  
356 climate types. Thus, our elementary climatic analysis does not point out a definite change of the  
357 Köppen climate types within the Pliocene and the Early Pleistocene assemblages studied.  
358 However, the boundary values of MAT (Fig. 4) distinctly show that the Pliocene floras are rich in  
359 thermophile species, that disappeared or became rare in the Early Pleistocene floras, which are  
360 mainly made up by cool-tolerant and warm-intolerant species. This situation suggests that a

decrease in temperatures is one of the causes of the palaeofloral differences observed between Pliocene and Early Pleistocene assemblages.

## 6. Occurrences of selected plant taxa

The complete list of taxa which have been studied in the West and East area samples is reported in Fig. 4, and in this section we provide additional information on selected species. The description of each species displays its occurrence only in the studied sections, expressed by the label of the plant-bearing layer, as reported in Tab. 1. Other known records of each species, even in the studied region, are reported under "*General distribution and characteristics*".

### Family Betulaceae

#### *Carpinus betulus* L.

1996 *Carpinus betulus* L. - Cavallo and Martinetto: pl. 5, fig. 2 (Pocapaglia, layer PO1).

*Occurrence:* West area: BR3, PO1 (Zanclean); RDB1, RDB6 (Piacenzian); East area: LU1 (Zanclean); CQ1, MR3 (Piacenzian); AD3 (Gelasian); SNT1 (early Calabrian), LA1, EZ38, LA3 (late Calabrian).

*General distribution and characteristics:* *C. betulus* has been selected for this analysis in order to evaluate the pattern of the fossil record of this plant, whose habit, ecology, and diaspore taphonomy are well known. This is a medium-sized deciduous tree, still living in the Po Plain nowadays, and characterizing its few relics of semi-natural forest vegetation (Pignatti, 1982). It is also a species frequently found in modern fruit and seed accumulations, where its wind-dispersed

fruits are overrepresented with respect to the frequency in the surrounding vegetation (Gee, 2005; Vassio and Martinetto, 2012). Its fossil record in northern Italy covers all the intervals from latest Miocene (Messinian: Cavallo et al., 1986), through Pliocene (Pavia, 1970; Martinetto, 1994; Cavallo and Martinetto, 1996; 2001), Early Pleistocene (Martinetto and Sami, 2001; Ghiotto, 2010), Middle Pleistocene (Martinetto, 2009), to the Holocene (Vassio, 2012).

The lack of *C. betulus* in three Piacenzian samples (Fig. 4) can be explained by two possibilities: i) it was absent or very rare in the time slices corresponding to the deposition of the layers where its fruits are lacking; ii) it was present in the time slices corresponding to the deposition of the layers where its fruits are lacking, but has not been recorded. We would propend for the second hypothesis and conclude that our data contribute to the assessment of a long-lasting presence of *Carpinus betulus* in the studied area during Pliocene and Early Pleistocene.

#### Family Cucurbitaceae

##### *Trichosanthes fragilis* E.M. Reid

#### Plate I, 1

*Occurrence:* East area: MR3 (Piacenzian).

*General distribution and characteristics:* this species had been reported in Italy only for the Front site (Martinetto, 1995; Martinetto et al., 2007) and is figured here for the first time (Plate I, 1). From the analysis of the distribution of its N-Italian occurrences it would seem an index fossil of the Piacenzian age, but this is due to the rarity of this species in the fossil record, which does not permit to use it in biochronological analyses.

405 Family Cymodoceaceae

406 *Cymodocea* cf. *nodosa* (Ucria) Ascherson

407 Plate I, 2

408 *Occurrence:* West area: PO1 (Zanclean); East area: LU1 (Zanclean); CVE2, MR3, CQ1  
409 (Piacenzian); AD3 (Gelasian); SNT1 (Calabrian).

410 *General distribution and characteristics:* The occurrence of fossil fruits of *Cymodocea* in other  
411 Zanclean sites of the West area had been previously reported by Pavia (1976) and Basilici et al.  
412 (1997). The fossil fruits from the studied samples seem to be identical to the modern ones  
413 (reference material: MCC1662) of *Cymodocea nodosa* (Ucria) Ascherson, a common seagrass in  
414 the Mediterranean; yet we prefer to adopt the open nomenclature (*Cymodocea* cf. *nodosa*) because  
415 the fossils did not yet undergo a thorough morphological characterisation, and possible differences  
416 between older (Zanclean) and younger (early Calabrian) remains (Fig. 5) must be verified.

417 The record of *Cymodocea* fruits in the studied samples shows very well how the occurrence data  
418 coincide with the sediment facies. In fact, by examining Fig. 5 it appears that *Cymodocea* cf.  
419 *nodosa* has a very constant occurrence in the Zanclean, Piacenzian and the oldest Calabrian, but it  
420 is lacking in the late Calabrian assemblages. The "disappearance" of this seagrass can be attributed  
421 to the unsuitable palaeoenvironment (coastal to continental) of the younger Calabrian deposits.

422

423 Family Cyperaceae

424 *Carex* cf. *elata* All.

425 Plate I, 3

426 *Occurrence:* East area: LA1 (Calabrian).

427 *General distribution and characteristics:* The importance of the fossil record of *C. elata* and  
428 related forms in the Plio-Pleistocene has been recently pointed out by Martinetto et al. (2012) and  
429 Jiménez-Mejías and Martinetto (2013); therefore a careful search for the tiny fruits has been  
430 carried out. However, the record in the studied sections turned out to be extremely scanty, possibly  
431 due to the prevailing marine-coastal facies. In fact, *C. cf. elata* was only detected in the  
432 palustrine-lacustrine assemblage LA1 (Fig. 4).

433 The modern *C. elata* is a rather tall herb forming tussocks in various non-saline waterlogged  
434 environments of western Eurasia. Actuopalaeobotanical analyses (Vassio, 2012; Vassio and  
435 Martinetto, 2012) showed that its tiny fruits are only abundant in the sediments that are directly  
436 deposited in the places where the plant grows, whereas they are rare in the adjoining fluvial  
437 sediments (Bertolotto et al., 2012). So, the absence of the fruits in marine-coastal sediments, in  
438 which the abundance of plant material depends on the fluvial input (Zavala et al., 2012), does not  
439 mean that the plant was absent in the continental palustrine environments.

440

441 *Schoenoplectus cf. litoralis* (Schrad.) Palla

442 Plate I, 4

443 *Occurrence:* East area: AD3 (Gelasian), LA3 (Calabrian).

444 *Taxonomical remarks:* Recent work on Cyperaceae showed the great taxonomic importance of  
445 fruit morphology (e. g. Pignotti, 2003; Ercole et al., 2012; Jiménez-Mejías and Martinetto, 2013).  
446 Some fruits found in the studied sections can easily be assigned to the genus *Schoenoplectus* for  
447 the following characters: achenes with a narrowly obovate outline, acute base and largely obtuse  
448 apex, flat on one side (Plate I, 4a) and convex on the other (Plate I, 4b); apex with a tubercle, but  
449 style-remain lacking; epidermal cells small-sized and subrectangular. Only five modern European  
450 species bear, as the fossil specimens, smooth fruits (Jiménez-Mejías et al., 2007): *S. pungens*, *S.*

451 *lacustris*, *S. litoralis*, *S. tabaernamontani* and *S. triqueter*. Another taxon, *S. carinatus* has been  
452 found to be a hybrid (Ercole et al., 2012). The morphologically similar *S. corymbosum* has fruits  
453 with small irregularly distributed wrinkles (Jiménez-Mejías et al., 2007).

454 The nutlets of *S. pungens* (MCC1515), *S. lacustris* (MCC1983), *S. tabaernamontani* (MCC1995)  
455 and *S. triqueter* (MCC2005) are always longer than 2.3 mm, so the smaller dimensions of the  
456 fossils (LA3: 1.5-1.9; AD3: 1.5-1.8 mm), suggest assignment to *S. litoralis*, whose modern nutlets  
457 are 1.6-1.9 [2.1] mm-long. However, Pignotti (2003) reported that also the hybrid *S. carinatus* has  
458 comparably small fruits, and this is one of the reason for keeping the open nomenclature  
459 *Schoenoplectus* cf. *litoralis*. Another reason is the total lack of the typical, feather-like perianth  
460 bristles (not preserved).

461 *General distribution and characteristics:* We are not aware of previous fossil records of *S.*  
462 *litoralis*. Today this is an herbaceous plant of rather limited distribution, found in coastal marshes.  
463 The time of diversification of the five above-mentioned species of *Schoenoplectus* was not  
464 discussed in the recent phylogenetic analyses (Muasya et al., 2009; Jung and Choi, 2011), however  
465 some divergence times should note date back more than 5 Ma, in consideration of the patterns of  
466 diversification pointed out in the Cyperaceae clades (Escudero et al., 2012; Escudero and Hipp,  
467 2013).

468 Our Gelasian fossil record of *S.* cf. *litoralis*, as well as the “upper Pliocene” (but possibly  
469 Gelasian) ones of *S. lacustris* (Mai and Walther, 1988) and *S. lacustroides* (Velichkevich and  
470 Zastawniak, 2003) may become important for dating the phylogenetic tree, but need a more  
471 detailed morphological characterisation. As *S. lacustris* is broadly distributed in lakes, marshes  
472 and rivers, its time of diversification could be used for the biochronology of continental sediments.  
473 The time of diversification of *S. litoralis*, which developed an adaptation to survive in salty soils,  
474 could be useful for the biochronology of coastal successions.

475

476 Family Ehretiaceae

477 *Ehretia europaea* E.M. Reid

478 1996 *Ehretia* sp. - Cavallo and Martinetto: pl. 3, figs. 2, 3 (Pocapaglia, layer PO1).

479 *Occurrence*: West area: PO1 (Zanclean).

480 *General distribution and characteristics*: The two endocarps figured by Cavallo and Martinetto  
481 (1996) are 2.0 mm broad and 2.1-2.5 mm long. These dimensions are similar to those of the two  
482 type specimens of *E. europaea* (1.8-1.25 x 1.6-2.75), described by Reid (1923) from the Late  
483 Miocene locality of Pont-de-Gail in France. Also the diagnostic morphological characters  
484 (Gottschling et al., 2002) show a good agreement between the Italian and the French specimens.

485 Gottschling et al. (2002) place *E. europaea* in the clade *Ehretia* II, but do not indicate a precise  
486 Nearest Living Relative.

487 A revision of the CENOFITA collection carried out for this paper showed that *E. europaea* occurs,  
488 in northern Italy, only in two sites reliably dated to the Zanclean (BG-Sento I, PO-Pocapaglia in  
489 Fig. 1) and in three sites of supposed Zanclean age (BG-Sento II, CV-Ca' Viettone, RB-Ronco  
490 Biellese in Fig. 1).

491

492 Family Eucommiaceae

493 *Eucommia europaea* Mädlér

494 Plate I, 5

495 *Occurrence:* East area: AD3 (Gelasian); reported by Gregor (1986) in layers corresponding to LA3  
496 (late Calabrian).

497 *General distribution and characteristics:* this species had been reported in Italy only for the  
498 Castelletto Cervo II site, possibly Gelasian (Cavallo and Martinetto, 2001). The occurrence of  
499 fruits in the Gelasian of N Italy is remarkable because they have not been found in Pliocene  
500 sediments, and there is only a preliminary report in the Calabrian (Gregor, 1986). This pattern of  
501 fossil record would suggest either an immigration or an increase of *Eucommia europaea* in N  
502 Italy, possibly in correspondence of the cooling at the Piacenzian/Gelasian transition (Bertini,  
503 2010). Unfortunately, the scarce woodiness in addition to the large size of the fruits (Plate I, 5)  
504 hampers their preservation in the fossil record (Vassio and Martinetto, 2012), and limits the use of  
505 this species in biochronological analyses.

506

507 Family Hypericaceae

508 *Hypericum tertiarum* P. Nikitin

509 Plate I, 6

510 *Occurrence:* West area: RDB1, RDB6, RT1 (Piacenzian); East area: CQ1 (Piacenzian).

511

512 *General distribution and characteristics:* An overview about the fossil record of this species has  
513 been recently reported by Martinetto et al. (2012). We add here its occurrence in the Piacenzian  
514 sample of Monte Falcone (CQ1). The occurrence of *H. tertiarum* in marine sediments appears to  
515 be extremely scanty, and not suitable for the assessment of the presence/absence of this species in  
516 the contemporary continental environments. For this reason the hypothetical extinction of this



517 species from northern Italy in correspondence to the Gelasian/Calabrian transition (Martinetto et  
518 al., 2012) needs to be confirmed by further data from continental sections.

519 The phylogeny of *Hypericum* has recently been investigated by Meseguer et al. (2013) and Nürk et  
520 al. (2013). The modern relatives of *H. tertiaerum*, at species level, are not yet well assessed  
521 because *Hypericum* is a large genus of ca. 500 species of shrubs, trees and herbs distributed  
522 mainly in temperate regions of the northern Hemisphere, but also in high-altitude tropical and  
523 subtropical areas (Meseguer et al., 2013). Despite some recent work dealing with seed characters  
524 (Meseguer and Sanmartin, 2012), these are still unknown for most of the modern species.  
525 According to Velichkevich and Zastawniak (2003), *H. japonicum* Thunberg in Murray, *H.*  
526 *tubulosum* Walt. and *H. virginicum* are the three extant species of *Hypericum* whose seeds most  
527 closely resemble those of *H. tertiaerum*. This would suggest a tentative placement of the fossil  
528 species into the clade (Meseguer et al., 2013) comprising the 3 modern species (“New World clade  
529 B”), whose divergence is estimated in the Oligocene (ca. 30 Ma) on a molecular basis. The  
530 members of this clade are a few species of wetland herbs with present disjunct distribution in East  
531 Asia and North America.

532

533 *Hypericum perforatum* L.

534 Plate I, 7

535 *Occurrence*: East area: LA3 (Calabrian).

536 *General distribution and characteristics*: This species has been reported in numerous interglacial  
537 and interstadial floras of eastern Europe (Velichkevich and Zastawniak, 2008). The seed  
538 morphology is very characteristic (checked in MCC1183) and allows an accurate identification of  
539 the species (Meseguer and Sanmartin, 2012). Molecular data (Meseguer et al., 2013) suggest a late  
540 Miocene divergence of *H. perforatum*, and Mai and Walther (1988) reported the oldest fossil

541 records from “Pliocene” localities of Germany and Russia. Our fossil record of layer LA3 (ca 1.0  
542 to 0.8 Ma) is the oldest for Italy, and is immediately followed by the early Middle Pleistocene one  
543 of Pianico (Martinetto, 2009). So the present evidence would suggest a late immigration of *H.*  
544 *perforatum* in Italy (around 1.0 Ma), much later than the diversification of the species.

545

546 Family Magnoliaceae

547 *Liriodendron geminata* Kirchheimer

548 Plate I, 8

549 1996 *Liriodendron geminata* Kirchheimer - Cavallo and Martinetto: pl. 1, fig. 4 (Pocapaglia, layer  
550 PO1).

551 *Occurrence*: West area: PO1 (Zanclean); RDB1 (Piacenzian); East area: CVE2 (Piacenzian);  
552 SNT1 (early Calabrian); EZ38 (late Calabrian).

553 *General distribution and characteristics*: *Liriodendron geminata* is a seed-morphospecies possibly  
554 linked to the leaves described as *L. procaccinii* in the Messinian (Massalongo and Scarabelli,  
555 1859). However the oldest records of seeds in Italy are available only from the Zanclean (Cavallo  
556 and Martinetto, 1996). The CENOFITA database (Martinetto and Vassio, 2010) indicates that it is  
557 abundant in a few coastal-continental sites attributed to the late Zanclean-late Piacenzian (Aulla,  
558 Fossano, Sento II, Terzoglio III), but also occurs in many other sites with a few seeds. Its modern  
559 relatives are tall deciduous trees that produce wind-dispersed fruits, which are poorly lignified, so  
560 that only the hard seeds are often recorded as fossils.

561 By combining the reconstructed habit, dispersal syndrome and the rather dense fossil record of this  
562 species (Fig. 5) we can conclude that *L. geminata* was most probably a deciduous tall tree, broadly  
563 distributed in the Zanclean and Piacenzian of northern Italy, but possibly uncommon or only

locally common. In marine sediments its occurrence is sporadic, so that the absence in CQ1 and MR3 is not surprising. The post-Piacenzian record was so-far poorly known, and the occurrence in two marine carpological assemblages of the Calabrian (EZ38, SNT1 in Fig. 4) points to the late survival of this plant in Italy. Its rarity in Calabrian sediments (absent from the LA1 and LA3 samples, as well as from Lefte, San Pietro di Ragona, Steglio: Ravazzi and Rossignol-Strick, 1995; Ghiotto, 2010; Martinetto et al., 2012) could mean that, after the end of the Piacenzian, *L. geminata* became a rare tree in northern Italy. Furthermore, the record of the long, but poorly dated, section of the Cervo River in the western Po Plain (BU in Fig. 1) (Cavallo and Martinetto, 2001; Martinetto and Festa, 2013), where *L. geminata* occurs only in two out of ca. twenty fruit-bearing layers of possible Gelasian and/or Calabrian age, could indicate that this plant has left a detectable fossil record only in limited Early Pleistocene time slices, which could be better pinpointed by future research. Unfortunately the pollen record of *Liriodendron* is scanty (Bertini and Martinetto, 2011) and cannot help us to clear this situation.

#### *Magnolia allasoniae* Martinetto

Plate I, 9

1996 *Magnolia allasoniae* Martinetto - Cavallo and Martinetto: pl. 4, figs. 5, 6 (Pocapaglia, layer PO1).

*Occurrence:* West area: PO1, BR3 (Zanclean); East area: LU1 (Zanclean), MR3 (Piacenzian).

*General distribution and characteristics:* This is a seed-morphospecies firstly established in the Pliocene of NW Italy (Martinetto, 1995), and it has been recorded in several sites of northern and central Italy in the time span Messinian-Piacenzian. Curiously, it is common in several Italian sites, but it has not yet been reported in the adjoining countries, which could suggest a plant with a limited geographic distribution in the late Cenozoic (endemic?). The leaves, pollen and wood of

the ancient whole plant are unknown. The seeds of *M. allasoniae* have been so-far compared with just a few species within the genus *Magnolia*, so that its modern relatives are not yet precisely indicated, even if an affinity of the fossil seeds to those of the living species *M. tripetala* has been pointed out. This last species is a deciduous tree which can grow in waterlogged environments.

*M. allasoniae* is broadly distributed and often frequent in the Zanclean of northern Italy, and the single occurrence in the Piacenzian is reported here (MR3 in Fig. 4). The fossil record and the ecology of the living *M. tripetala* would suggest that this plant was probably a common river-side or wetland tree in the Zanclean, while it became very rare in the Piacenzian, possibly with a relic distribution just in the southernmost areas (see MR3 in Fig. 2), as well as in central Italy (Martinetto, 2001a). The importance of the *M. allasoniae*-plant in the Zanclean is confirmed even by considering that *Magnolia* seeds have a fleshy tegument adaptive for animal dispersal, which may cause an overrepresentation in the sedimentary deposits (Vassio and Martinetto, 2012).

*Magnolia cor* Ludwig

Plate I, 10

*Occurrence:* West area: RDB1 (Piacenzian); East area: AD3 (Gelasian).

*General distribution and characteristics:* This seed-morphospecies has been recorded in the late Cenozoic of several parts of Europe (Mai, 1975). In Italy it occurs, sometimes abundantly (Cavallo and Martinetto, 2001), only from the Piacenzian to the Calabrian (Fig. 5). The smooth, heart-shaped seeds of *M. cor* (Plate I, 10) are very variable and morphologically similar to those of several modern East-Asian species (*M. kobus*, *M. stellata*, etc.). These modern plants are both deciduous trees and shrubs, growing in mesic forests. For this reason the seeds of *M. cor* are considered to represent an ancient deciduous tree or shrub with animal-dispersed seeds.

611

612 *Magnolia ludwigii* Ettingshausen

613 Plate I, 11

614 *Occurrence:* East area: MR3, CQ1 (Piacenzian).

615 *General distribution and characteristics:* The characteristic morphology of this seed  
616 morphospecies, often reported in the literature under the invalid synonym *M. lignita*, has been  
617 described by Mai (1975), who suggested an affinity with the living *M. ashei*. The habit and  
618 climatic requirements (see Utescher and Mosbrugger, 2013) of this modern species (actually  
619 considered a subspecies or variety of *M. macrophylla* by Azuma et al., 1999) were used to  
620 interpret *M. ludwigii* as an evergreen tree which required a very warm climate, so that its  
621 occurrence in fossil assemblages is often determinant for their climatic characterisation by means  
622 of the coexistence approach (Mosbrugger and Utescher, 1997). For the thermal characterisation of  
623 *Magnolia ludwigii* (MAT > 9°C in Fig. 4) we used *M. macrophylla* as a NLR and data from the  
624 GrimClim database (Grimm and Denk, 2012).

625 *M. ludwigii* is always rare in the Zanclean of northern Italy, and the only two occurrences in the  
626 Piacenzian are those reported here (CQ1, MR3). So this plant was probably uncommon in the  
627 Pliocene vegetation of northern Italy.

628

629 Family Menispermaceae

630 *Sinomenium cantalense* (E.M. Reid) Dorofeev

631 Plate I, 12

632 1996 *Sinomenium cantalense* (E.M. Reid) Dorofeev - Cavallo and Martinetto: pl. 1, figs. 5, 6  
633 (Pocapaglia, layer PO1).

634 *Occurrence*: West area: PO1, BR3 (Zanclean); RDB1 (Piacenzian); East area: LU1 (Zanclean);  
635 CVE2, MR3, CQ1 (Piacenzian).

636 *General distribution and characteristics*: *S. cantalense* is reported, never abundant, in most of the  
637 Zanclean-Piacenzian sites of northern Italy (Martinetto, 1999; Martinetto et al., 2007). Its single  
638 modern relative, *S. acutum*, is an evergreen or semi-evergreen climber, which produces drupes  
639 dispersed by animals (endozoochory). This feature can be hypothesized also for *S. cantalense*. So,  
640 the habit of the single living relative, the dispersal syndrome and the rather dense fossil record of  
641 this species (Fig. 5) suggest that the *S. cantalense* fossil endocarps were produced by an  
642 evergreen-semievergreen climber, broadly distributed in the mesic and river-side forests of  
643 northern Italy, during Zanclean and Piacenzian. In northern Italy there are no post-Piacenzian  
644 records of this species, whereas in central Italy it has been abundantly recorded in the Ponte Naja  
645 Formation of the Cava Toppetti II site, assigned to the Gelasian (Leone et al., 2000; Martinetto,  
646 2001a).

647

648 Family Rutaceae

649 *Toddalia naviculaeformis* (E.M. Reid) Gregor

650 2001a *Toddalia naviculaeformis* (E.M. Reid) Gregor - Martinetto: pl. 1, figs. 14, 15 (Marecchia,  
651 layer MR3).

652 *Occurrence*: West area: RDB1 (Piacenzian); East area: MR3 (Piacenzian).

653 *General distribution and characteristics*: two seeds of this species from the Piacenzian sediments  
654 of the Marecchia section (MR3) had been figured in Martinetto (2001a), and other few records

were published for sites of Zanclean and/or Piacenzian age in northern Italy (Martinetto and Mai, 1996; Basilici et al., 1997; Ferrero et al., 2005). This species is less common than its relatives *T. latisiliquata* and *T. rhenana*, which are more useful for biochronological analyses (Fig. 5). In general, for the species of *Toddalia* there are no records younger than late Piacenzian in northern Italy, whereas in central Italy *T. latisiliquata* is suspected to persist in the Gelasian (see Martinetto, 2001a).

*Zanthoxylum ailanthiforme* (Gregor) Gregor

1996 *Zanthoxylum ailanthiforme* (Gregor) Gregor - Cavallo and Martinetto: pl. 6, fig. 1 (Pocapaglia, layer PO1).

2001a *Zanthoxylum ailanthiforme* (Gregor) Gregor- Martinetto, pl. 1, fig. 21 (Marecchia, layer MR3).

*Occurrence:* West area: PO1 (Zanclean); East area: MR3 (Piacenzian).

*General distribution and characteristics:* The taxonomic treatment and occurrences of this species in central as well as in northern Italy have already been discussed by Martinetto (2001a), who also figured one specimen from the Piacenzian sediments of the Marecchia section (MR3). This species seems to be important for biochronological analyses because in northern Italy is well represented only in the Messinian (Kovar-Eder et al., 2006) and Zanclean, with a single Piacenzian record in the layer MR3 (Marecchia section). Also in central Italy *Z. ailanthiforme* never occurred in sites of well-assessed Gelasian and/or Calabrian age (Martinetto, 2001a; Girotti et al., 2003).

677 Family Sparganiaceae

678 *Sparganium nanum* Dorofeev

679 Plate I, 13

680 1996 *Sparganium nanum* Dorofeev - Cavallo and Martinetto: pl. 3, fig. 7 (Pocapaglia, layer PO1).

681 *Occurrence:* West area: PO1, BR3 (Zanclean); RDB1 (Piacenzian); East area: CQ1 (Piacenzian);

682 AD3 (Gelasian).

683 *General distribution and characteristics:* The record of AD3 represents the last occurrence of this  
684 species in Europe. The single fruitlet is small, globose, and very resistant (Plate I, 13), so that  
685 reworking cannot be excluded and further findings in other layers of the same age would be  
686 necessary to confirm the extension of the range of *S. nanum* to the early Gelasian.

687 The habit of *S. nanum* can be reconstructed as that of an herb living in freshwater wetlands,  
688 possibly with submerged roots and floating leaves as the modern *S. angustifolium* (Pignatti, 1982).

689

690 Family Stemonaceae

691 *Stemona germanica* (Mai) Mai

692 Plate I, 14

693 1996 *Spirellea* aff. *bohemica* Knobloch et Mai - Cavallo and Martinetto: pl. 3, fig. 9 (Pocapaglia,  
694 layer PO1).

695 *Occurrence:* West area: BR5 (layer ca. 5 m above BR3), PO1 (Zanclean); East area: CQ1  
696 (Piacenzian).



697 *General distribution and characteristics:* This carpological species was assigned to the modern  
698 genus *Stemona* by Mai (2008), who reported occurrences from the Late Cretaceous to the Late  
699 Miocene in Central Europe. In Italy, material of this species was often reported as “*Spirellea* aff.  
700 *bohémica* Knobloch et Mai” (Cavallo and Martinetto, 1996; Basilici et al., 1997; Martinetto and  
701 Ravazzi, 1997) or “*Spirellea* sp.” (Martinetto, 2001a) and was assigned to *S. germanica* for the  
702 first time in Martinetto and Vassio (2010).

703 A revision of the CENOFITA collection carried out for this paper confirmed that *S. germanica*  
704 occurs, in northern Italy, in four sites reliably dated to the Zanclean (BG-Sento I, BR-Breolungi,  
705 MZ-Morozzo and PO-Pocapaglia in Fig. 1), in four sites of supposed Zanclean age (BG-Sento II,  
706 CV-Ca’ Viettone, RB-Ronco Biellese, CGL-Cortiglione d’Asti), and in a single site of supposed  
707 Piacenzian age (MO-Momello/Lanzo). So this species disappeared from Italy later than in central  
708 Europe and seems to be useful for biochronological analyses (Fig. 5), since there are no records  
709 younger than late Piacenzian in northern Italy. We revise here as *S. germanica* also the record of  
710 “*Spirellea* sp.”, reported by Martinetto (2001a) from the Dunarobba site (Piacenzian or Gelasian)  
711 in central Italy, which consists in a single seed.

712

713 Family Symplocaceae

714 *Symplocos casparyi* Ludwig

715 1996 *Symplocos lignitarum* (Quenstedt) Kirchheimer - Cavallo and Martinetto, pl. 5, fig. 9  
716 (Pocapaglia, layer PO1).

717 *Occurrence:* West area: BR3, PO1 (Zanclean); RT1 (Piacenzian); East area: LU1 (Zanclean);  
718 CVE2, MR3, CQ1 (Piacenzian).

*General distribution and characteristics:* *S. casparyi*, as circumscribed by Mai and Martinetto (2006), is a morphospecies with largely variable endocarps. In Italy it occurred in several coastal-continental sites attributed to the late Zanclean (Ca' Viettone, Castellengo, Fossano, Sento II). As its modern relatives, which are small evergreen trees or shrubs, produce drupes which favour animal dispersal (endozoochory), this feature can be hypothesized also for *S. casparyi*. By combining the reconstructed habit, dispersal syndrome and the rather dense fossil record of this species (Fig. 5), we can conclude that the *S. casparyi* fruits were most probably produced by an evergreen small tree or shrub, broadly distributed in the Zanclean and Piacenzian mesic forests of northern Italy. This is in agreement with its occurrence in all the marine carpological assemblages of the Zanclean and Piacenzian in the East area (Fig. 4).

#### Family Theaceae

*Eurya stigmosa* (Ludwig) Mai

Plate I, 15

1996 *Eurya stigmosa* (Ludwig) Mai - Cavallo and Martinetto: pl. 2, fig. 1 (Pocapaglia, layer PO1).

*Occurrence:* West area: PO1, BR3 (Zanclean); East area: LU1 (Zanclean); CQ1 (Piacenzian); AD3 (Gelasian).

*General distribution and characteristics:* One of the commonest species in the Zanclean of northern Italy and already reported in sites dated back to a possible Piacenzian age (Martinetto, 1995). The record of 3 seeds in the CQ1 layer, treated in this paper, is a further clue to the survival of this plant till the late Piacenzian, not only in central Italy (Martinetto, 2001a), but also in the northern part of this country. Finally, the interpretation of a single seed found in the Gelasian

741 sample of the AD3 layer is not easy: the good preservation (Plate 1, 15) does not provide any  
742 evidence of reworking, anyway we suggest to take in consideration this possibility.

743

744 Family Thymelaeaceae

745 *Thymelaea* cf. *passerina* L.

746 Plate I, 16

747 *Occurrence*: East area: AD3 (Gelasian); LA3 (Calabrian).

748 *General distribution and characteristics*: The phylogeny of *Thymelaea*, whose present distribution  
749 is limited to the peri-Mediterranean area, has been studied by Galicia-Herbada (2006). The genus  
750 includes 31 species, most of which show a distribution restricted to the westernmost  
751 Mediterranean (Iberia plus Northwest Africa) and, according to the phylogenetic analysis of  
752 Galicia-Herbada (2006), have a recent origin (not older than 2 Ma) and should not be crucial for  
753 the comparison with our fossils. In Italy (Pignatti, 1982) only 4 species of *Thymelaea* occur at  
754 present: *T. dioica*, *T. hirsuta*, *T. passerina* and *T. tartonraira*. The narrow drop-shaped seeds of  
755 samples AD3 and LA3, with a thick and shining wall, are respectively 2.0 and 1.7-mm long and  
756 both have a length/width ratio of 1.8. The 1.6-2.2 mm-long (mean specimens' length around 1.9  
757 mm) seeds of *T. passerina* from NW Italy (MCC1516), with a 1.8-2.2 length/width ratio, agree in  
758 all the morphological characters with those of fossil samples. The seeds of *T. hirsuta* (MCC1767)  
759 and *T. tartonraira* (MCC1768) are easily distinguishable for the larger dimensions, lower  
760 length/width ratio, and coarser cell pattern.

761 The perfect morphological agreement of the fossil seeds with the modern ones of *T. passerina*  
762 would suggest an assignment of the *T. passerina* clade (Galicia-Herbada, 2006). Furthermore, it  
763 does not seem possible the occurrence of species other than *T. passerina* in northern Italy (also in

the past) because the differentiation of other species occurred because of geographic isolation in the western (e.g. *T. salsa*) or eastern (e.g. *T. mesopotamica*) Mediterranean (Urbani, 1992). However, due to strong phylogenetic link between *T. passerina* and *T. salsa*, pointed out by Galicia-Herbada (2006: fig. 7), it would be desirable to study in more detail the seed morphology of these two species in order to evaluate the presence of differential morphological characters which could be checked in the Early Pleistocene fossils.

The record of AD3 most probably represents the oldest known for *T. passerina* in the fossil record. Its occurrence in the Middle Pleistocene of central Europe was reported by Mai (2010). This species was also reported from archaeological sites of the East area dating back from the 2<sup>nd</sup> century BC to the 6<sup>th</sup> century AD (Rinaldi et al., 2013).

Family *incertae sedis*

*Carpolithes minimus* (Szafer) Mai

Plate I, 17

*Occurrence:* East area: LA1, LA3 (Calabrian); EZ38 (Calabrian).

*General distribution and characteristics:* Despite the good morphological and anatomical agreement (Martinetto, 2001c), assignment of *C. minimus* to *Thalictrum* (Ranunculaceae) has to be rejected because the indehiscent achenes of *Thalictrum* do not correspond to the apparently dehiscent condition of *C. minimus* (Plate I, 17). The new efforts in searching the same structure as in *C. minimus* in fruits and seeds of modern plant taxa, just brought to the detection of superficial resemblances (Lamiaceae: *Horminum*; Cannabaceae: *Humulus*). Therefore, this common and characteristic fossil taxon is still to be considered *incertae sedis*, and we have no information

about the habit and ecology of the plant which produced these remains. It was certainly not linked to coastal areas, since these fossils are also found in ancient freshwater intermontane basins (Steggio: Ghiotto, 2010).

## 7. Discussion

Our analyses, aimed at the reconstruction of the distribution in time and space of plant species, are based on the interpretation of the presence of carpological remains in well-dated layers, whereas the absence of fossils is considered less important. In fact, the analyses of fruit and seed assemblages in modern contexts showed that the fruit/seeds of one species may frequently be absent in the sediments formed close to its growth place, due to several biological and taphonomical factors. Absence of the fruits/seeds with hard parts in the fossil record can be used as a clue to the absence of one plant in an ancient environment, but only after the analysis of several samples and localities of the same age. Conversely, the presence of the fruits/seeds of one species in a sedimentary deposit is a strong evidence for the presence of the mother plant in the environment surrounding the place of deposition (at the catchment basin scale: Vassio and Martinetto, 2012). An exception is represented by the reworking of fruits and seeds from older sediments, which seems to be a rare event at the light of the evidence discussed above; nevertheless, should be kept in consideration.

The fossil fruit and seed record of selected plant taxa in Zanclean, Piacenzian, Gelasian and Calabrian layers showed that the distribution of several species was limited to one or a few chronostratigraphic stages (Fig. 4). The climatic characterisation of these species indicates that the major floristic changes are linked to the disappearance or retreat of thermophile elements in the late Zanclean, late Piacenzian, and during the Calabrian. Additionally, most of the species which appear in the Early Pleistocene fossil record are cool-tolerant. In the Zanclean we observe the

highest number of thermophile species, and some of them (Group 1 in Fig. 4) occur only in this stage. However, the independently dated (Barbieri, 1967) Lugagnano assemblage (Fig. 4) shows that a poor Zanclean flora may lack Group 1 elements (excepted *Visnea germanica*, reported by Gregor, 1990), and fit the criteria used to suggest a Piacenzian affinity (simultaneous occurrence of several species of groups 2 and 3, without species of groups 1 and 5). Unfortunately, we have no information on carpological assemblages of the earliest Piacenzian (3.6 to 3.3 Ma), and the oldest known Piacenzian assemblages come from the West area sites re-examined in this work (RDB1, RDB6, RT1), which have been interpreted by Martinetto et al. (2007) as falling within the Mammoth reversed magnetic polarity event (ca. 3.3 to 3.2 Ma). The record of these Piacenzian western sites (Fig. 4) appears consistently different from that of the eastern ones, mainly because of the occurrence of Group 2 species in the last ones (Fig. 4). In our opinion this difference is linked to the slightly younger age (3.0-2.8 Ma) of the eastern sites CQ1 and MR3 (Tab. 1), since the thermophilous Group 2 elements (MAT often  $>8^{\circ}\text{C}$  in Fig. 4) could have profited from the climate warming reported around 3.0 Ma (Bertini, 2010). So, we can argue that the carpoflora of Chiavenna (CVE2 in Fig. 4) lacks of Group 2 elements because it is dated to the reversed magnetic polarity event Kaena (3.1 to 3.0 Ma), which predates the warming. This is also confirmed, in the northern part of the western Po Plain, by the lack of Group 2 elements in the rich palaeoflora of the Stura di Lanzo Fossil Forest, also dated to the Kaena (Martinetto et al., 2007). On the other hand, the successive (ca. 3.0-2.8 Ma) Piacenzian records of Marecchia (MR3) and Monte Falcone (CQ1) point out a thermophilous palaeoflora (MAT $>8^{\circ}\text{C}$ ) in which even the species characterizing the Zanclean palaeoflora (Group 2 in Fig. 4) are still abundant, therefore they seem to correspond to the warming phase. Therefore, in the interpretation of the age of Pliocene floras, we must consider that the thermophilous species of Group 2 did not disappear at the Zanclean/Piacenzian transition (see Martinetto, 1999), but just experienced a temporary retreat from 3.2 to 3.0 Ma, and assumed again greater importance into the late Piacenzian (ca. 3.0 to 2.8 Ma).

The newly obtained carpological assemblage dated to the early Gelasian (AD3), is particularly important because in the whole N-Italy there was only another carpo flora, the Casnigo one (CG in Fig. 1, already mentioned in Martinetto, 1999), which was reliably dated to the Gelasian, in particular to the final part of this age (ca. 2.1 to 1.8 Ma). Another assemblage interpreted as Gelasian has been described at San Pietro di Ragogna (SPR in Fig. 1, NE Italy), but mainly represents an ancient freshwater plant community and provides a very poor record of the terrestrial palaeoflora (Martinetto et al., 2012). Richer assemblages of possible Gelasian age have been described from the Cervo River section (Fig. 1; Castelletto Cervo II and Buronzo floras: Cavallo and Martinetto, 2001; Martinetto and Festa, 2013) for which a confirmation of the age with methods independent from palaeobotany would be needed. Also the ongoing studies in the southern Piedmont Basin, along the Bormida valley (Fig. 2; Irace et al., 2012) are promising for the gathering of data on Gelasian carpo floras. Nevertheless, the Gelasian is still largely "*terra incognita*" as for the distribution of carpological taxa, and this is particularly critical for the better definition of the last appearance in northern Italy of a large number of Zanclean-Piacenzian species, never detected in the Gelasian/Calabrian. The last known occurrences of such "old" taxa (Groups 2-3 in Fig. 5) mostly fall within the late Piacenzian, but in 2 cases (*Eurya stigmosa*, *Sparganium nanum*) Gelasian occurrences were detected in this work (Fig. 4). However, in consideration of the possibility of reworking (see above), we suggest much caution before placing into the Gelasian the local extinction of these taxa.

The Calabrian carpological record of the East area sections starts with the Santerno-SNT1 assemblage, dated at the very base of the Calabrian (ca. 1.8-1.7 Ma), which is followed by a definitely younger Calabrian assemblages (LA1, ca. 1.1 Ma after Gunderson et al., 2012). Due to gap of ca. 600 ka between the SNT1 and LA1 assemblages, at present it is not possible to point out eventual changes of the carpo floras within the lower Calabrian. About the EZ38 and LA3 assemblages, new magnetostratigraphic data (Gunderson et al., 2012, 2014) point to a late

862 Calabrian age, just after the Jaramillo normal subchron. This is also supported by the occurrence  
863 of *Tsuga*, that is unknown in the Middle Pleistocene, even in pollen records (Ravazzi et al., 2005;  
864 Martinetto, 2009; Bertini, 2010). The four Calabrian fruit and seed assemblages studied are  
865 typically formed by an admixture of species inherited from the Pliocene floras (*Carpolithes*  
866 *minimus*, *Juglans bergomensis*, *Liriodendron geminata*, *Magnolia cor*, *Selaginella* gr. *pliocenica*)  
867 and species which appear in the North Italian record only in the Gelasian (*Ajuga* cf. *iva*, *Picea*  
868 *florschuetzii*, *Taxus baccata*, *Thymelaea* cf. *passerina*, *Tsuga* cf. *chiarugii*) or Calabrian  
869 (*Hypericum perforatum*). None of these new occurrences is interpretable as an evolutionary  
870 appearance. Rather, the effect of the environmental factors seems having prevailed, favouring the  
871 immigration of these species from adjoining areas; for the Gymnosperms, this could simply  
872 consist in descending from the mountains (where no fossil record was preserved).

873 The record of those plant taxa with a proven diversification in the last 5 Ma (e.g., Thymeleaceae:  
874 Galicia-Herbada, 2006; Cyperaceae: Jiménez-Mejías and Martinetto, 2013) is much scattered in  
875 the sections studied, but offers intriguing opportunities for a future detection of the time of  
876 evolutionary appearance of a few species. As for the Cyperaceae the 2.5-2.4 Ma record of  
877 *Schoenoplectus* cf. *litoralis* (AD3) falls within an interesting time range to calibrate the phylogeny  
878 of the group. As for *Thymelaea*, the 2.5-2.4 Ma record of AD3 is the oldest known for the *T.*  
879 *passerina* clade (Galicia-Herbada, 2006), and could predate the diversification of *T. salsa*. The  
880 supposed Late Miocene origin of the clade would suggests that plants producing seeds of the *T.*  
881 *passerina*-type were present since that time somewhere in the Mediterranean. So, the appearance  
882 of such seeds in the fossil record of N-Italy, as late as the Gelasian, could be explained by the  
883 expansion of the plant's habitat at that time, due to slightly drier climatic conditions, which were  
884 possibly absent in the Pliocene (more persistent humidity). In fact the plants of the *T. passerina*  
885 group are small herbs linked to open areas with temporarily dry soil (Pignatti, 1982).

886



## 887 7. Conclusions

888 Our analysis of carpological assemblages in Pliocene and Early Pleistocene successions cropping  
889 out along the southern margin of the Po Plain showed that these deposits contain an interesting  
890 record of the terrestrial palaeoflora, which integrates the information provided by the extensive  
891 pollen record available in this area. We could fix in the geological time scale, often with a  
892 precision of 100 ka (Tab. 1), 14 carpo-flora-bearing layers spanning from 5.1 to 0.9 Ma, thus  
893 obtaining a framework which is not available in other parts of Europe. This enabled us to provide  
894 precise ages for the occurrence of fruits and seeds of a few plant groups with a proven late  
895 Cenozoic differentiation, as a possible contribution to future phylogenetic analyses. For the other  
896 species, with a phylogenetic origin dating back to the Miocene, Palaeogene or Cretaceous, we  
897 obtained an updated synthesis of the distribution in time of the fossil record in the studied area.

898 Our dataset was revealed useful for biochronological analyses, since the fossil record of several  
899 species showed an interesting distribution, often limited to one to three chronostratigraphic stages  
900 (Zanclean-Piacenzian-Gelasian-Calabrian). Nevertheless, several relevant differences could be  
901 pointed out by comparing the record of the West and the East area. In particular, several species  
902 occurring only in the Zanclean of the West area were detected in the late Piacenzian of the East  
903 area (Fig. 4), thus suggesting that both datasets, separately, were not reliable for the reconstruction  
904 of the distribution in time of each plant species.

905 Our elementary palaeoclimatic characterisation of the individual species, and of the plant  
906 assemblages in which they occur (Fig. 4), suggests that temperature decrease is one of the causes  
907 of the disappearance from the fossil record of several species, in particular at the  
908 Piacenzian/Gelasian transition. Also the disappearance (or retreat) of some species at the  
909 Zanclean/Piacenzian transition would seem to be due to climate deterioration, since it affects the  
910 thermophilous Group 1 (NLRs' MAT boundaries often  $>15^{\circ}\text{C}$  and mostly  $>9^{\circ}\text{C}$ ).

911 As for the appearance of new species in the studied fossil record, the analysis of the habit,  
912 ecological-climatic requirements and phylogeny of the recent relatives indicates that it is mostly  
913 due to the establishment of favourable local environmental conditions, rather than to evolutionary  
914 events.

915 The input of the new data obtained into the CENOFITA database led to an apparent improvement  
916 of the distribution chart of carpological taxa in the Pliocene and Early Pleistocene. This new chart  
917 shows the carpological species clustered in groups with a similar biochronologic distribution  
918 (Groups 1-6 in Fig. 5) and we suggest that the analysis of the representation of such groups in  
919 undated or poorly dated carpological assemblages would be useful for their biochronological  
920 interpretation. In this sense, the possibility to characterise the Zanclean floras against the  
921 Piacenzian ones is provided by the occurrence of Group 1 species in the first ones. In addition, the  
922 simultaneous occurrence of several species of groups 2 and 3 (Fig. 5), without species of groups 1  
923 and 5, could be used to suggest a Piacenzian affinity for undated assemblages.

924 The characterisation of Gelasian floras is a major problem, because of the lack of record in the  
925 crucial time span between 2.8 and 2.5 Ma, so that we cannot really say whether the Gelasian  
926 carpological floras can be distinguished from the latest Piacenzian ones. Fruit and seed  
927 assemblages with several species of groups 5 and 6, without species of groups 1-3 (Fig. 5), have  
928 not been detected in any Zanclean or Piacenzian carpooflora, and would suggest a Gelasian or  
929 Calabrian affinity.

930 In conclusion, the new data presented here determined several changes in the biochronological  
931 distributions published in preceeding works and, despite the good chronostratigraphic framing  
932 available in the studied sections, we suggest that the biochronologic chart presented in this paper  
933 (Fig. 5) still requires an integration of data from a broader geographic area, before being  
934 considered as definitely stable. The implementation of the N-Italian Plio-Pleistocene carpological  
935 dataset, especially with data from other calibrated continental sections, would be the right

direction for completing our knowledge of the chronologic distribution of single carpological species as well as species-groups.

#### ACKNOWLEDGEMENTS

We thank H.-J. Gregor for fundamental indications to locate the plant-bearing layers of the Marecchia and Stirone sections and S. Dominici for the help provided in the location and stratigraphic framing of the Arda and Stirone samples, as well as for useful suggestions as to improve the whole manuscript, which also profited from the valuable revision carried out by two anonymous referees that we wish to thank. Likewise, we acknowledge the help provided by A. Fornaciari and S. Iaccarino for the Enza section, by S. Mariani and M. Sami for the Santerno one, and by P. Monegatti and S. Raffi for the Castell'Arquato-Monte Falcone section. We also thank M. Roveri for useful information and discussions about the stratigraphy of the Castell'Arquato basin, and V. Picotti and K. Gunderson for discussion on Stirone and Enza sections. Finally, we wish to thank P. Ghiotto for her help in the analysis of the Stirone samples, M. Escudero and P. Jiménez-Mejías for providing information on the taxonomy of Cyperaceae, and A. S. Meseguer for *Hypericum*. We thank also A. Bruch, T. Denk, G. Grimm, T. Utescher for the useful climatic resignements, and the Vertebrate Palaeontology group of the Turin University for the access to microphotographic equipment.

#### REFERENCES

Azuma, H., Thien, L.B., Kawano, S., 1999. Molecular phylogeny of *Magnolia* (Magnoliaceae) inferred from cpDNA sequences and evolutionary divergence of the floral scents. Journal of Plant Research 112, 291-306.

959 Barbieri, F., 1967. The Foraminifera in the Pliocene section Vernasca-Castell'Arquato including  
 960 the «Piacenzian stratotype» (Piacenza Province). *Memorie della Società Italiana di Scienze*  
 961 *Naturali e del Museo Civico di Storia Naturale di Milano* 15, 145-163.

962 Basilici, G., Martinetto, E., Pavia, G., Violanti, D., 1997. Paleoenvironmental evolution in the  
 963 Pliocene marine-coastal succession of Val Chiusella (Ivrea, NW Italy). *Bollettino della Società*  
 964 *Paleontologica Italiana* 36 (1-2), 23-52.

965 Bertini, A., 2001. Pliocene climatic cycles and altitudinal forest development from 2.7 Ma in the  
 966 northern Apennines (Italy): evidences from the pollen record of the Stirone section (5.1 to ca. 2.2  
 967 Ma). *Geobios* 34, 253-265.

968 Bertini, A., 2010. Pliocene to Pleistocene palynoflora and vegetation in Italy: State of the art.  
 969 *Quaternary International* 225, 5-24.

970 Bertini, A., Martinetto, E., 2011. Reconstruction of vegetation transects for the Messinian /  
 971 Piacenzian of Italy by means of comparative analysis of pollen, leaf and carpological records.  
 972 *Palaeogeography, Palaeoclimatology, Palaeoecology* 304, 230-246.

973 Bertoldi, R., Martinetto, E., 1996. Ricerche paleobotaniche (palinologiche e paleocarpologiche)  
 974 sulla successione villafranchiana del Rio Ca' Viettone. *Il Quaternario (Italian Journal of*  
 975 *Quaternary Sciences)* 8 (2), 403-422.

976 Bertolotto, G., Martinetto, E., Vassio, E., 2012. Applicazioni paleobotaniche dello studio di resti  
 977 carpologici in suoli e depositi fluviali attuali del Piemonte, con particolare riferimento alle  
 978 *Cyperaceae*. *Informatore Botanico Italiano* 44 (2), 72-74.

979 Blanc-Vernet, L., 1969. Contribution à l'étude des foraminifères de Méditerranée. - Travaux de la  
 980 Station Marine d'Endoume, Marseille, France.

- 981 Boano, P., Bormioli, D., Carraro, F., Lanza, R., 1999. Magnetic stratigraphy of the Villafranchian  
982 type-section:(Villafranca d'Asti, Italy). *Annales of Geophysics* 42, 9-19.
- 983 Boufford, D. E. 1997. Urticaceae Jussieu, Nettle family. *Flora of North America North of*  
984 *Mexico*. Volume 3, 400-413.
- 985 Calabrese, L., Di Dio, G., 2009. Note Illustrative della Carta Geologica D'Italia alla scala 1:50.000  
986 foglio 180 Salsomaggiore Terme. Ente realizzatore: Regione Emilia-Romagna, Servizio  
987 Geologico, Sismico e dei Suoli. Editore: ISPRA - Servizio Geologico d'Italia.
- 988 Cande, S.C., Kent, D.V., 1995. Revised calibration of the geomagnetic polarity time scale for the  
989 Late Cretaceous and Cenozoic. *Journal of Geophysical Research* 100 B4, 6093-6095.
- 990 Cappers, R.T.J., 1993. Seed dispersal by water: a contribution to the interpretation of seed  
991 assemblages. *Vegetation History and Archaeobotany* 2, 173-186.
- 992 Carraro, F. (Ed.). 1996. The Villafranchian in the Villafranca d'Asti type-area: a revision. II.  
993 *Quaternario (Italian Journal of Quaternary Sciences)* 9, 5-120.
- 994 Casnedi, R., 1971. Contributo alle conoscenze del Pliocene del Bacino Terziario Piemontese. *Atti*  
995 *Istituto Geologico Università di Pavia* 22, 110-120.
- 996 Cavallo, O., Martinetto, E., 1996. Flore plioceniche del bacino del Tanaro. *Alba Pompeia* 17 (1),  
997 5-31.
- 998 Cavallo, P., Martinetto, E., 2001. Flore carpologiche del Pliocene di Castelletto Cervo (Biella).  
999 *Bollettino del Museo regionale di Scienze naturali di Torino* 18 (2), 277-343.
- 1000 Cavallo, O., Macagno, M., Pavia, G., 1986. Fossili dell'Albese. *Famija Albèisa*, Alba.

- 1001 Cita, M.B., 1975. Studi sul Pliocene e sugli strati di passaggio dal Miocene al Pliocene. VIII.  
1002 Planktonic foraminiferal biozonation of the Mediterranean Pliocene deep sea record. A revision.  
1003 *Rivista Italiana di Paleontologia e Stratigrafia* 81, 527-544.
- 1004 Cita, M.B., Capraro, L., Ciaranfi, N., Di Stefano, E., Marino, M., Rio, D., Sprovieri, R., Vai, G.B.,  
1005 2006. Calabrian and Ionian: A proposal for the definition of Mediterranean stages for the Lower  
1006 and Middle Pleistocene. *Episodes* 29, 107-114.
- 1007 Colalongo, M.L., 1969. Cenozone a foraminiferi ed ostracodi del Pliocene e basso Pleistocene  
1008 della serie del Santerno e dell'Appennino romagnolo. *Giornale Geologia* 35, 29-61.
- 1009 Colalongo, M.L., Ricci Lucchi, F., Berardi, F., Nanni, L., 1982a. Il Pliocene neoautoctono di  
1010 Poggio Berni in Val Marecchia (Appennino romagnolo). In: Cremonini, G., Ricci Lucchi, F.  
1011 (Eds.), Guida alla geologia del margine appenninico-padano. Società Geologica Italiana, Guide  
1012 Geologiche Regionali, Bologna, pp. 177-180.
- 1013 Colalongo, M.L., Ricci Lucchi, F., Guarnieri, P., Mancini, E., 1982b. Il Plio-Pleistocene del  
1014 Santerno (Appennino Romagnolo). In: Cremonini, G., Ricci Lucchi, F. (Eds.), Guida alla geologia  
1015 del margine appenninico-padano. Società Geologica Italiana, Guide Geologiche Regionali,  
1016 Bologna, pp. 161-166.
- 1017 Cremaschi, M., 1982. La formazione fluvio-lacustre del Pleistocene inferiore-medio nel  
1018 pedeappennino emiliano. In: Cremonini, M., Ricci Lucchi, F. (Eds.), Guida alla Geologia del  
1019 margine appenninico-padano, Società Geologica Italiana, Guide Geologiche Regionali, Bologna,  
1020 pp. 145-149.
- 1021 Cremonini, G., Elmi, C., Monesi, A., 1969. Osservazioni geologiche e sedimentologiche su alcune  
1022 sezioni plio-pleistoceniche dell'Appennino romagnolo. *Giornale di Geologia* 35, 85-96.

1023 Denk, T., Grimm, G.W., Grímsson, F., Zetter, R., 2013. Evidence from “Köppen signatures” of  
 1024 fossil plant assemblages for effective heat transport of Gulf Stream to subarctic North Atlantic  
 1025 during Miocene cooling. *Biogeosciences* 10, 7927–7942.

1026 Dominici, S., 2001. Taphonomy and paleoecology of shallow marine macrofossil assemblages in a  
 1027 collisional setting (Late Pliocene-Early Pleistocene, Western Emilia, Italy). *Palaios* 16, 336-353.

1028 Ercole, E., Pistarino, A.L., Martinetto, E., Soldano, A., Siniscalco, C., 2012. Atlante fotografico  
 1029 dei frutti e dei semi della flora del Piemonte e della Valle d’Aosta: Cyperaceae. *Bollettino del*  
 1030 *Museo regionale di Scienze Naturali di Torino* 29 (1), 5-28.

1031 Escudero, M., Hipp, A., Waterway, M., Valente, L., 2012. Diversification rates and chromosome  
 1032 evolution in the most diverse angiosperm genus of the temperate zone (*Carex*, Cyperaceae).  
 1033 *Molecular Phylogenetics and Evolution* 63, 650- 655.

1034 Escudero, M., Hipp, A., 2013. Shifts in diversification rates and clade ages explain species  
 1035 richness in higher-level sedge taxa (Cyperaceae). *American Journal of Botany* 100 (12), 2403-  
 1036 2411.

1037 Fang, J., Wang, Z., Tang, Z., 2011. Atlas of Woody Plants in China, distribution and climate,  
 1038 volume 1. Springer, 1972 pp.

1039 Ferrero, E., Merlino, B., Provera, A., Martinetto, E., 2005. Associazione a molluschi marini e  
 1040 vegetali terrestri del Pliocene di Castellengo (Biella, Italia NW). *Rendiconti Società*  
 1041 *Paleontologica Italiana* 2, 87-108.

1042 Galicia-Herbada, D., 2006. Origin and diversification of *Thymelaea* (Thymelaeaceae): inferences  
 1043 from a phylogenetic study based on ITS (rDNA) sequence. *Plant Systematics and Evolution* 257,  
 1044 159-187.

1045 Gastaldo, R., 1994. The genesis and sedimentation of phytoclasts with examples from coastal  
1046 environments. In: Traverse, A. (Ed.), Sedimentation of organic particles. Cambridge University  
1047 Press, Cambridge, UK, 103-127.

1048 Gee, G.T., 2005. The genesis of mass carpological deposits (bedload carpodeposits) in the tertiary  
1049 of the lower Rhine basin, Germany. *Palaios* 20, 463-478.

1050 Ghielmi, M., Minervini, M., Nini, C., Rogledi, S., Rossi, M., 2010. Sedimentary and tectonic  
1051 evolution in the eastern Po Plain and northern Adriatic Sea area from Messinian to Middle  
1052 Pleistocene (Italy). In: Sassi, F.P. (Ed.). Nature and Geodynamics of the Lithostere in Northern  
1053 Adriatic. *Rendiconti Lincei Scienze Fisiche e Naturali* 21 (Suppl. 1), 131-166.

1054 Ghiotto, P., 2010. La carpoflora del bacino lacustre villafranchiano di Steggio (Treviso, Prealpi  
1055 orientali). *Bollettino del Museo Regionale di Scienze Naturali di Torino* 27, 3-99.

1056 Gibbard, P.L., Head, M.J., 2010. The newly-ratified definition of the Quaternary System/Period  
1057 and redefinition of the Pleistocene Series/Epoch, and comparison of proposals advanced prior to  
1058 formal ratification. *Episodes* 33, 152-158.

1059 Girotti, O., Capasso Barbato, L., Esu, D., Gliozzi, E., Kotsakis, A., Martinetto, E., Petronio, C.,  
1060 Sardella, R., Squazzini, E., 2003. The section of Torre Picchio (Terni, Umbria, Central Italy): A  
1061 Villafranchian site rich in Mammals, Molluscs, Ostracods and Plants. *Rivista Italiana di*  
1062 *Paleontologia e Stratigrafia* 109 (1), 77-98.

1063 Gottschling, M., Mai D.H., Hilger, H.H., 2002. The systematic position of *Ehretia* fossils  
1064 (Ehretiaceae, Boraginales) from the European Tertiary and implications for character evolution.  
1065 *Review of Palaeobotany and Palynology* 121, 149-156.

1066 Gradstein, F., Ogg, J., Smith, A., 2004. A Geologic Time Scale 2004. Cambridge University Press,  
1067 Cambridge.



1068 Gregor, H.J., 1986. Preliminary results of the Plio-Pleistocene flora from Stirone river (Parma,  
1069 Fidenza). *Memorie della Società Geologica Italiana* 31, 219-227.

1070 Gregor, H.J., 1990. Contribution to the Late Neogene and Early Quaternary floral history of the  
1071 Mediterranean. *Review of Palaeobotany and Palynology* 62, 309-338.

1072 Grimm, G.W., Denk, T., 2012. Reliability and resolution of the coexistence approach – a  
1073 revalidation using modern-day data. *Review of Palaeobotany and Palynology* 172, 33-47.

1074 Gunderson, K.L., Kodama, K.P., Anastasio, D.J., Pazzaglia, F.J., 2012. Rock-magnetic  
1075 cyclostratigraphy for the Late Pliocene-Early Pleistocene Stirone section, Northern Apennine  
1076 mountain front, Italy. Italy. In: Jovane, L., Herrero-Bervera, E., Hinnov, L.A., Housen, B. (Eds.),  
1077 Magnetic Methods and the Timing of Geological Processes. Geological Society of London Special  
1078 Publication 373, London, pp. 309–323.

1079 Gunderson, K.L., Pazzaglia, F.J., Picotti, V., Anastasio, D.A., Kodama, K.P., Rittenour, T.,  
1080 Frankel, K.F., Ponza, A., Berti, C., Negri, A., Sabbatini, A., 2014. Unraveling tectonic and  
1081 climatic controls on synorogenic growth strata (northern Apennines, Italy). *Geological Society of*  
1082 *America Bulletin*. doi:10.1130/B30902.1

1083 Hansen A., Sunding P., 1993. Flora of Macaronesia: checklist of vascular plants, 4th edition.  
1084 *Sommerfeltia* 17, 1-295.

1085 Iaccarino, S., 1985. Mediterranean Miocene and Pliocene planktic foraminifera. In: Bolli, H.M.,  
1086 Saunders, J.B., Perch-Nielsen, K. (Eds.). *Plankton Stratigraphy*. Cambridge University Press,  
1087 Cambridge, UK, pp. 283-314.

1088 Iaccarino, S., Monegatti, P., 1996. Stop 6 - La successione pleistocenica del T. Enza. XIII  
1089 Convegno Società Paleontologica Italiana Guida alle Escursioni. University of Parma, Parma,  
1090 Italy, pp. 63-67.

1091 Irace, A., Monegato, G., Tema, E., Martinetto, E., Pini, R., Gianolla, D., Bellino, L., 2012. The  
1092 Pliocene-Pleistocene transition in Southern Piedmont: new data from the Alessandria Basin.  
1093 INQUA-SEQS 2012: Sassari, Italy, pp. 41-42.

1094 Jacques, F.M.B., Liu, C.Y.-S., Martinetto, E., Zhou, Z.-K., 2011. Revised taxonomy of selected  
1095 fossil endocarp species in the Menispermaceae using a morphometric approach. *Geodiversitas* 33  
1096 (1), 177-197.

1097 Jiménez-Mejías, P., Martinetto, E., 2013. Towards an accurate taxonomic interpretation of *Carex*  
1098 fossil fruits (Cyperaceae): a case study in section *Phacocystis* in the Western Palearctic. *American*  
1099 *Journal of Botany* 100, 1580-1603.

1100 Jiménez-Mejías, P., Luceño, M., Castroviejo, S., 2007. *Schoenoplectus corymbosus*: a tropical  
1101 Old-World sedge (Cyperaceae) discovered in Spain and Morocco. *Nordic Journal of Botany* 25,  
1102 70-74.

1103 Jung, J., Choi, H.-K., 2011. Taxonomic Study of Korean *Scirpus* L. s.l. (Cyperaceae) II: Pattern of  
1104 Phenotypic Evolution Inferred from Molecular Phylogeny. *Journal of Plant Biology* 54, 409-424.

1105 Kovar-Eder, J., Kvaček, Z., Martinetto, E., Roiron, P., 2006. Late Miocene to Early Pliocene  
1106 vegetation of southern Europe (7–4 Ma) as reflected in the megafossil plant record.  
1107 *Palaeogeography, Palaeoclimatology, Palaeoecology* 238, 321–339.

1108 Kovar-Eder, J., Jechorek, H., Kvaček, Z., Parashiv, V., 2008. The integrated plant record: an  
1109 essential tool for reconstructing Neogene zonal vegetation in Europe. *Palaios* 23, 97-111.

1110 Leone, G., Bonadonna, F., Zanchetta, G., 2000. Stable isotope record in mollusca and pedogenetic  
1111 carbonate from Late Pliocene soils of central Italy. *Palaeontology, Palaeoclimatology,*  
1112 *Palaeoecology* 163, 115-131.

- 1113 Lona, F., Bertoldi, R., 1973. La storia del Plio-Pleistocene italiano in alcune sequenze  
1114 vegetazionali lacustri e marine. Atti Accademia Nazionale dei Lincei, Memorie serie 8, 11 (1), 1-  
1115 47.
- 1116 Mai, D.H., 1975. Beiträge zur Bestimmung und Nomenklatur fossiler Magnolien. Feddes  
1117 Repertorium 86, 559-640.
- 1118 Mai, D.H., 1994. Fossile Koniferenreste in der meridionalen Zone Europas. Feddes Repertorium  
1119 105, 207-227.
- 1120 Mai, D.H., 1995. Paleocarpological investigations in the Villafranchian (Pliocene) of Italy.  
1121 Bollettino del Museo Regionale di Scienze Naturali di Torino 13, 407-437.
- 1122 Mai, D.H., 2008. Palaeokarpologische Nachweise von neuen Arten aus Reliktfamilien im  
1123 Europäischen Tertiär. Feddes Repertorium 119, 191-206.
- 1124 Mai, D.H., 2010. Karpologische Untersuchungen in einem Interglazial von Neumark-Nord  
1125 (Geiseltal). Palaeontographica Abt. B 282, 99-187.
- 1126 Mai, D.H., Walther, H., 1988. Die pliozänen Floren von Thüringen. Deutsche Demokratische  
1127 Republik. Quartärpaläontologie 7, 55-297.
- 1128 Mai, D.H., Martinetto, E., 2006. A reconsideration of the diversity of *Symplocos* in the European  
1129 Neogene on the basis of fruit morphology. Review of Palaeobotany and Palynology 140, 1-26.
- 1130 Martinetto, E., 1994. Analisi paleocarpologica dei depositi continentali pliocenici della Stura di  
1131 Lanzo. Bollettino del Museo Regionale di Scienze Naturali di Torino 12, 137-172 .
- 1132 Martinetto, E., 1995. Significato cronologico e paleoambientale dei macrofossili vegetali  
1133 nell'inquadramento stratigrafico del "Villafranchiano" di alcuni settori del Piemonte (Italia NW).  
1134 Ph.D. Thesis, University of Turin, Italy.

- 1135 Martinetto, E., 1999. Chronological framing of Pliocene to Early Pleistocene plant macrofossil  
1136 assemblages from northern Italy. *Acta Palaeobotanica Suppl.* 2, 503-511.
- 1137 Martinetto, E., 2001a. The role of central Italy as a centre of refuge for thermophilous plants in the  
1138 late Cenozoic. *Acta Palaeobotanica* 41 (2), 299-319.
- 1139 Martinetto, E. (Ed.), 2001b. Pliocene Plants, Environment and Climate of Northwestern Italy.  
1140 *Flora Tertiaria Mediterranea* 5 (8), 1-88.
- 1141 Martinetto, E., 2001c. Studies on some exotic elements of the Pliocene floras of Italy.  
1142 *Palaeontographica Abt. B* 259, 149-166.
- 1143 Martinetto, E., 2003. Leaves of terrestrial plants from the shallow marine and transitional Pliocene  
1144 deposits of Asti (Piedmont, NW Italy). *Bollettino della Società Paleontologica Italiana* 42 (1-2),  
1145 75-11.
- 1146 Martinetto, E., 2009. Palaeoenvironmental significance of plant macrofossils from the Piànico  
1147 Formation, Middle Pleistocene of Lombardy, North Italy. *Quaternary International* 204, 20-30.
- 1148 Martinetto, E., Mai, D.H., 1996. Macrofossili vegetali. In: Carraro, F. (Ed.). *Revisione del*  
1149 *Villafranchiano nell'area-tipo di Villafranca d'Asti. Il Quaternario (Italian Journal of Quaternary*  
1150 *Sciences)* 9, 73-79.
- 1151 Martinetto, E., Ravazzi, C., 1997. Plant biochronology of the Valle della Fornace succession  
1152 (Varese) based on the Plio-Pleistocene record in northern Italy. *Geologia Insubrica* 2 (2), 81-98.
- 1153 Martinetto, E., Sami, M., 2001. Paleoflora delle "sabbie gialle" medio-pleistoceniche di Oriolo,  
1154 presso Faenza (Ravenna). *Quaderno di Studi e Notizie di Storia Naturale della Romagna* 14, 1-28.

- 1155 Martinetto, E., Vassio, E., 2010. Reconstructing “Plant Community Scenarios” by means of  
1156 palaeocarpological data from the CENOFITA database, with an example from the Ca’ Viettone  
1157 site (Pliocene, Northern Italy). *Quaternary International* 225, 25-36.
- 1158 Martinetto, E., Festa, C., 2013. Frutti e semi fossili del Biellese. *DocBi Studi e ricerche sul*  
1159 *Biellese* 2013, 169-188.
- 1160 Martinetto, E., Scardia, G., Varrone, D., 2007. Magnetobiostratigraphy of the Stura di Lanzo fossil  
1161 forest succession (Piedmont, Italy). *Rivista Italiana di Paleontologia e Stratigrafia* 113 (1), 109-  
1162 125.
- 1163 Martinetto, E., Monegato, G., Vassio, E., 2012. An Early Pleistocene Plant Assemblage with East  
1164 European Affinity in the Venetian-Friulian Basin (NE Italy). *Alpine and Mediterranean*  
1165 *Quaternary* 25 (2), 91-104.
- 1166 Martinetto, E., Bouvet, D., Vassio, E., Magni, P., Jiménez-Mejías, P., 2014. A standard protocol  
1167 for the collection and cataloguing of reference material for the study of fossil Cyperaceae fruits:  
1168 the Modern Carpological Collection. *Review of Palaeobotany and Palynology* 201, 56-74.
- 1169 Massalongo, A., Scarabelli, G.F., 1859. *Studii sulla flora fossile e geologia stratigrafica del*  
1170 *Senigalliese*. Premiata Litografia Penuti, Verona, Italy.
- 1171 Mayer-Eymar, K., 1858, Versuch einer synchronistischen Tabelle der Tertiär Gebilde Europas.  
1172 *Verhandlungen der schweizerischen naturforschenden Gesellschaft* 1857, 1-32.
- 1173 Meseguer, A.S., Sanmartín, I., 2012. Paleobiology of the genus *Hypericum* (Hypericaceae): A  
1174 survey of the fossil record and its palaeogeographic implications. *Anales del Jardín Botánico de*  
1175 *Madrid* 69, 97-106.

1176 Meseguer, A.S., Aldasoro, J.J., Sanmartín, I., 2013. Bayesian inference of phylogeny, morphology  
 1177 and range evolution reveals a complex evolutionary history in St. John's wort (*Hypericum*).  
 1178 Molecular Phylogenetics and Evolution 67 (2), 379-403.

1179 Monegatti, P., Raffi, S., Raineri, G., 1997. The Monte Falcone-Rio Riorzo Composit section:  
 1180 biostratigraphic and ecobiostratigraphic remarks. Bollettino della Società Paleontologica Italiana  
 1181 36 (1-2), 247-262.

1182 Monegatti, P., Canali, G., Bertoldi, R., Albianelli, A., 2002. The classical Late Piacenzian Monte  
 1183 Falcone - Rio Crevalse section (Northern Italy): palynological evidence and  
 1184 biomagnetostratigraphic constraints for climatic cyclicity and local mollusc extinctions. Geobios  
 1185 Mémoire spécial 24, 219-227.

1186 Mosbrugger, V., Utescher, T., 1997. The coexistence approach - a method for quantitative  
 1187 reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils. Palaeogeography,  
 1188 Palaeoclimatology, Palaeoecology, 134, 61-86.

1189 Muasya, A.M., Simpson, D.A., Verboom, G.A., Goetghebeur, P., Naczi, R.F.C., Chase, M.W.,  
 1190 Smets, E., 2009. Phylogeny of Cyperaceae based on DNA sequence data: current progress and  
 1191 future prospects. Botanical Review 75, 2-21.

1192 Muttoni, G., Scardia, G., Kent, D.V., Morsiani, E., Tremolada, F., Cremaschi, M., Peretto, C.,  
 1193 2011. First dated human occupation of Italy at ~0.85 Ma during the late Early Pleistocene climate  
 1194 transition. Earth and Planetary Science Letters 307, 241–252.

1195 Napoleone, G., Albianelli, A., Azzaroli, A., Bertini, A., Magi, M., Mazzini, M., 2003. Calibration  
 1196 of the Upper Valdarno basin to the Plio-Pleistocene for correlating the Apennine continental  
 1197 sequences. Il Quaternario (Italian Journal of Quaternary Sciences) 16 (1bis), 131-166.

- 1198 Nürk, N.M., Madriñán, S., Carine, M.A., Chase, M.W., Blattner, F.R., 2013. Molecular  
1199 phylogenetics and morphological evolution of St. John's wort (*Hypericum*; Hypericaceae).  
1200 Molecular Phylogenetics and Evolution 66 (1), 1-16.
- 1201 Papani, G., Pelosio, G., 1963. La serie Plio-Pleistocenica del Torrente Stirone (Parmense  
1202 occidentale). Bollettino della Società Geologica Italiana 81 (2), 293-335.
- 1203 Pareto, M., 1865. Sur les subdivisions que l'on pourrait établir dans les terrains Tertiaires de  
1204 l'Apennin septentrional. Bulletin de la Société géologique de France 22, 210-277.
- 1205 Pavia, G., 1970. Resti di *Anancus arvernensis* e flore ad affinità plioceniche nel Villafranchiano  
1206 inferiore della cava Arboschio (Villafranca d'Asti). Memorie della Società Geologica Italiana 9,  
1207 157-176.
- 1208 Pavia, G., 1976. I molluschi del Pliocene inferiore di Monteu Roero (Alba, Italia NW). Bollettino  
1209 della Società Paleontologica Italiana 14, 99-175.
- 1210 Pavia, G., Chiambretto, L., Oreggia, G., 1989. Paleocomunità a Molluschi nel Pliocene inferiore di  
1211 Breolungi (Mondovì, Italia NW). In: Di Geronimo, S.I. (Ed.), Atti del 3° Simposio di Ecologia e  
1212 Paleoecologia delle Comunità Bentoniche. Catania, Italy, pp. 521-569.
- 1213 Pignatti, S., 1982. Flora d'Italia. Edagricole, Bologna.
- 1214 Pignotti, L., 2003. *Scirpus* L. and related genera (Cyperaceae) in Italy. Webbia 58 (2), 281-400.
- 1215 Rasmussen, T.L., 2005. Systematic paleontology and ecology of benthic foraminifera from the  
1216 Plio-Pleistocene Kallithea Bay Section, Rhodes, Greece. Cushman Foundation Special Publication  
1217 39, 53-157.

- 1218 Raffi, S., Rio, D., Sprovieri, R., Valleri, G., Monegatti, P., Raffi, I., Barrier, P., 1989. New  
1219 stratigraphic data on the Piacentian stratotype. *Bollettino della Società Geologica Italiana* 108,  
1220 183-196.
- 1221 Ravazzi, C., Rossignol Strick, M., 1995. Vegetation change in a climatic cycle of Early  
1222 Pleistocene age in the Leffe basin (Northern Italy). *Palaeogeography, Palaeoclimatology,*  
1223 *Palaeoecology* 117, 105-122.
- 1224 Ravazzi, C., Pini, R., Breda, M., Martinetto, E., Mattoni, G., Chiesa, S., Confortini, F., Egli, R.,  
1225 2005. The lacustrine deposits of Fornaci di Ranica (late Early Pleistocene, Italian Pre-Alps):  
1226 stratigraphy, palaeoenvironment and geological evolution. *Quaternary International* 131, 35-58.
- 1227 Reid, E.M., 1923. Nouvelles recherches sur les graines du Pliocène inférieur du Pont-de-Gail  
1228 (Cantal). *Bulletin de la Société Géologique de France* 23, 308-355.
- 1229 Repetto, G., 1997. La malacofauna pliocenica di Pocapaglia. *Alba Pompeia* 18, 57-65.
- 1230 Ricci Lucchi, F., 1986. The Oligocene to recent foreland basins of the northern Apennines. In:  
1231 Allen, P.A., Homewood, P. (Eds.), *Foreland Basins*. International Association of Sedimentologists  
1232 Special Publication 8, Wiley-Blackwell, Hoboken, NJ, pp. 105-139.
- 1233 Rinaldi, R., Bandini Mazzanti, M., Bosi, G., 2013. Archaeobotany in urban sites: the case of  
1234 Mutina. *Annali di Botanica* 2013 (3), 217-230.
- 1235 Rio, D., Channell, J.E.T., Bertoldi, R., Poli, M.S., Vergerio, P.P., Raffi, I., Sprovieri, R., Thunell,  
1236 R.C., 1997. Pliocene sapropels in Northern Adriatic area: chronology and paleoenvironmental  
1237 significance. *Palaeogeography, Palaeoclimatology, Palaeoecology* 135, 1-25.
- 1238 Roveri, M., Taviani, M., 2003. Calcarene and sapropel deposition in the Mediterranean Pliocene:  
1239 shallow- and deep-water record of astronomically driven climatic events. *Terra Nova* 15, 279-286.



1240 Schweizer, M., 2006. Evolution and molecular phylogeny of *Cibicides* and *Uvigerina* (Rotaliida,  
1241 Foraminifera). *Geologica Ultraiectina* 261, 1-167.

1242 Sprovieri, R., 1992. Mediterranean Pliocene biochronology: a high resolution record based on  
1243 quantitative planktonic foraminifera distribution. *Rivista Italiana di Paleontologia e Stratigrafia* 98,  
1244 61-100.

1245 Stevens, P. F., 2014. Angiosperm Phylogeny Website. [www.mobot.org/MOBOT/research/](http://www.mobot.org/MOBOT/research/APweb/)  
1246 APweb/ [accessed 5 March 2014]

1247 Thomasson, J.R., 1991. Sediment-borne “seeds” from Sand Creek, northwestern Kansas:  
1248 taphonomic significance and paleoecological and paleoenvironmental implications.  
1249 *Palaeogeography, Palaeoclimatology, Palaeoecology* 85, 213-225.

1250 Thompson, R.S., Anderson, K.H., Bartlein, P.J., 1999a. Atlas of relations between climatic  
1251 parameters and distributions of important trees and shrubs in North America — Introduction and  
1252 conifers. U.S. Geological Survey Professional Paper 1650–A, 1-269.

1253 Thompson, R.S., Anderson, K.H., Bartlein, P.J., 1999b. Atlas of relations between climatic  
1254 parameters and distributions of important trees and shrubs in North America — Hardwoods. U.S.  
1255 Geological Survey Professional Paper 1650–B, 1-423.

1256 Thompson, R.S., Anderson, K.H., Pelltier, R.T., Strickland, L.E., Bartlein, P.J., Shafer, S.L., 2012.  
1257 Quantitative estimation of climatic parameters from vegetation data in North America by the  
1258 mutual climatic range technique. *Quaternary Science Reviews* 51, 18–39.

1259 Urbani, M., 1992. La Sezione *Ligia* (Fasano) Meisn. del genere *Thymelaea* P. Miller  
1260 (*Thymelaeaceae*) in Italia. *Giornale Botanico Italiano* 126, 391.

1261 Utescher, T., Mosbrugger, V., 2013. The Palaeoflora Database. [http://www.geologie.](http://www.geologie.unibonn.de/Palaeoflora)  
1262 [unibonn.de/Palaeoflora](http://www.geologie.unibonn.de/Palaeoflora) [accessed 4 October 2013]

1263

1264 Vaiani, S.C., 2000. Testing the applicability of Strontium Isotope Stratigraphy in marine to deltaic  
1265 Pleistocene deposits: an example from the Lamone River Valley (northern Italy). *Journal of*  
1266 *Geology* 108, 585-599.

1267 Vassio, E., 2012. Palaeovegetation reconstructions and palaeoclimatic interpretations of  
1268 Quaternary carpological assemblages with an actuopalaeobotanical approach. Ph.D. Thesis,  
1269 University of Turin. Turin, Italy.

1270 Vassio, E., Martinetto, E, 2012. Biases in the frequency of fruits and seeds in modern fluvial  
1271 sediments in northwestern Italy: the key to interpreting analogous fossil assemblages. *Palaaios* 27,  
1272 779-797.

1273 Velichkevich, F.Y., Zastawniak, E., 2003. The Pliocene flora of Kholmech, south-eastern Belarus  
1274 and its correlation with other Pliocene floras of Europe. *Acta Palaeobotanica* 43 (2), 137-259.

1275 Velichkevich, F.Y., Zastawniak, E., 2008. Atlas of the Pleistocene Vascular Plant Macrofossils of  
1276 Central and Eastern Europe - Part 2: Herbaceous Dicotyledones. W. Szafer Institute of Botany,  
1277 Polish Academy of Sciences, Kraków.

1278 Vigna, B., Fiorucci, A., Ghielmi, M., 2010. Relations between stratigraphy, groundwater flow and  
1279 hydrogeochemistry in Poirino Plateau and Roero areas of the Tertiary Piedmont Basin, Italy,  
1280 *Memorie descrittive della Carta Geologica d'Italia*, Istituto Superiore per la Protezione e la  
1281 *Ricerca Ambientale* 90, 267-292.

1282 Violanti, D., 2012. Pliocene Mediterranean Foraminiferal Biostratigraphy: A Synthesis and  
1283 Application to the Paleoenvironmental Evolution of Northwestern Italy. In: Elitok, Ö. (Ed.),  
1284 *Stratigraphic Analysis of Layered Deposits*, pp. 123-160.

1285 Zavala, C., Arcuri, M., Blanco Valiente, L., 2012. The importance of plant remains as diagnostic  
1286 criteria for the recognition of ancient hyperpyncites. *Revue de Paléobiologie* Vol. spéc. 11, 457-  
1287 469.

1288

1289

1290 Figures and tables captions:

1291 Fig. 1. Schematic map of northern-central Italy showing the location of the Zanclean-Calabrian  
1292 localities with published carpological assemblages. AD: Castell'Arquato-Arda; BE Benasso; BG:  
1293 Sento I and II; BL: Candelo; BO: Boca; BR: Breolungi; BU: Buronzo; GA: Castelletto Cervo I  
1294 and II; CGL: Cortiglione d'Asti; CM: Crava di Morozzo; CO: Cossato; CVE: Chiavenna; CG:  
1295 Casnigo; CQ: Castell'Arquato-Monte Falcone - Rio Crevalese; CV: Ca' Viettone; EZ: Enza; FR:  
1296 Front; LA: Stirone-Laurano; LF: Leffe; LU: Lugagnano; MO: Momello-Lanzo; MR: Val  
1297 Marecchia; MZ: Morozzo; PO: Pocapaglia; RBS: Ranica Borgo Sale; RDB: RDB Quarry,  
1298 Villafranca d'Asti; RDL: Rio della Lupa; RT: Roatto; SG: Steggio; SNT: Santerno; SPR: San  
1299 Pietro di Ragnogna; STU: Stura di Lanzo Fossil Forest; TC: Castellengo; TZ: Terzoglio; VF: Valle  
1300 della Fornace. Towns: Tu, Turin; Ve, Venice.

1301

1302 Fig. 2. Schematic map of the southern margin of the Po Plain with the location of the studied  
1303 sections. Red lines indicate the frontal thrust arcs. The star indicates the Villafranca d'Asti type  
1304 area; Br: Breolungi; Po: Pocapaglia; 1: Chiavenna Rocchetta; 2: Lugagnano; 3: Monte Falcone and  
1305 Arda (Castell'Arquato section); 4: Stirone; 5: Enza; 6: Santerno; 7: Marecchia.

1306 Fig. 3. Simplified logs of the studies sections with data useful for the chronostratigraphic  
1307 calibration. Chronostratigraphy after Gradstein et al. (2004), Cita et al. (2006), Gibbard and Head

1308 (2010). Magnetostratigraphy after Cande and Kent (1995). Planktonic foraminifer biostratigraphy  
 1309 after Cita (1975), Iaccarino (1985), and Sprovieri (1992). Stratigraphy of analyzed sections are  
 1310 modified after 1) Roveri and Taviani (2003); 2) Barbieri (1967); 3a) Monegatti et al. (2002); 3b)  
 1311 Dominici (2001), Roveri and Taviani (2003); 4) Papani and Pelosio (1963), Gunderson et al.  
 1312 (2012); 5) Iaccarino and Monegatti (1996), Gunderson et al. (2014); 6) Cremonini et al. (1969),  
 1313 Colalongo (1969); 7) Colalongo et al. (1982a and 1982b), Rio et al. (1997).

1314

1315 Fig. 4. List of selected carpological taxa from the West and East area sections (see Tab. 1). Black  
 1316 boxes indicate occurrence in our samples, grey boxes are referred to the report of *Eucommia*  
 1317 *europaea*, *Juglans bergomensis*, *Tsuga cf. chiarugii* and *Visnea germanica* by Gregor (1990), and  
 1318 *Picea florschuetzii* by Mai (1994). Z = Zanclean; G = Gelasian. Palaeoenvironment: C = coastal;  
 1319 M = marine; P = palustrine. The listed taxa have been gathered in 6 main groups, as explained in  
 1320 the text.

1321 A simple characterisation of the growth forms of the listed taxa is reported in the two columns  
 1322 named respectively “Deciduous/Evergreen” and “Habit”, where “c”, “h”, “s” and “t” stand for  
 1323 climber, herb, shrub and tree. The thermal climatic requirements of each species have been  
 1324 synthetically expressed by the boundary values of Mean Annual Temperature (NLR Clim.  
 1325 boundaries), e.g.: >13 = distribution of Nearest Living Relatives (NLRs) mainly in areas where the  
 1326 MAT is higher than 13°C; <14 = distribution of NLRs mainly in areas where the MAT is lower  
 1327 than 14°C; u = uncertain, b = broad climatic distribution (literature sources: ^Denk and Grimm,  
 1328 2012; \*Utescher and Mosbrugger, 2013; #other sources, reported in Supplementary materials).  
 1329 The column "exotic/native" (ex/n) is referred to the NLRs, and “X” means no-NLR (extinct genus  
 1330 or morphogenus).

1331

Fig. 5. Scheme summarizing the biochronological distribution of the carpological records of selected plant taxa in N-Italy. The order of the species and the groups are the same as in Fig. 4. Mes = Messinian; MP = Middle Pleistocene.

The records from all the localities indicated in Fig. 1, deduced from the whole N-Italian CENOFITA database (Martinetto and Vassio, 2010), are reported. The age range for each species is indicated by black bars, for the records originating from reliably dated localities: Lugagnano (5.1-4.5 Ma), Pocapaglia (4.5-3.8 Ma), Breolungi (5.1-3.8 Ma), Morozzo (5.1-3.8 Ma), Sento I (3.8-3.6 Ma), Candelo (4.5-3.6 Ma), Roatto (3.3-3.2 Ma), RDB Quarry of Villafranca d'Asti (3.3 to 3.0 Ma), Stura di Lanzo Fossil Forest (3.1-3.0 Ma), Casnigo (2.1-1.9 Ma), Santerno (1.8-1.7 Ma), Leffe (1.7-1.5 Ma), Stirone-LA1 (1.1-1.0 Ma), Enza-EZ38 (1.1-1.0 Ma), Ranica (1.1-1.0 Ma), Oriolo (1.0-0.8 Ma), Stirone-LA3 (1.0-0.8 Ma). Hatched bars indicate records from the following poorly dated localities (see Martinetto, 1999): Ca' Viettone (late Zanclean, 3.8 to 3.6 Ma); Fossano (late Zanclean, 4.0 to 3.6 Ma); Sento II (late Zanclean, 3.8 to 3.6 Ma); Front (Piacenzian, 3.3 to 3.0 Ma); Momello-Lanzo (Piacenzian, 3.6 to 3.0 Ma); Castelletto Cervo I (late Piacenzian, 2.8 to 2.6 Ma); Castelletto Cervo II (Gelasian, 2.6 to 2.0 Ma); San Pietro di Ragona (Gelasian, 2.6 to 1.8 Ma); Buronzo (Gelasian, 2.2 to 1.8 Ma); Steggio (Calabrian, 1.8 to 0.9 Ma). The dashed lines indicate the presence of records in central Italy, considered a refuge area in the Piacenzian-Calabrian (Martinetto, 2001a).

Tab. 1. Lithologic and stratigraphic data for the studied palaeocarpological samples.

**Plate I** - Examples of fossil fruits from the studied sections. Figs. 1-5 and 8-17 scale bars = 1mm; Figs. 6-7 scale bars = 0.5 mm. The 3-characters codes (e.g. B3E) assign the fossil specimen to a

1355 precise sediment sample recorded in the CENOFITA database; the label MGPT-PU+number is  
 1356 referred to the collections' catalogue of the Earth Sciences Department of the Turin University.

1357 1) *Trichosanthes fragilis* E.M. Reid: a) outer surface, b) inner surface of the seed, Marecchia  
 1358 section, sample MR3 B3E, Piacenzian; MGPT-PU105884;

1359 2) *Cymodocea* cf. *nodosa* (Ucria) Ascherson: a, b) lateral views of the endocarp, Marecchia  
 1360 section, sample MR3 B3E, Piacenzian, MGPT-PU105885;

1361 3) *Carex* cf. *elata* All.: achene, Stirone section, sample LA1 A5B, Calabrian, MGPT-PU105886;

1362 4) *Schoenoplectus* cf. *litoralis* (Schr.) Palla: a) ventral view and b) dorsal view of the achene,  
 1363 Stirone section, sample LA3 A5C, Calabrian, MGPT-PU105887;

1364 5) *Eucommia europaea* Mädl.: fruit, Arda section, sample AD3 B7E, Gelasian, MGPT-  
 1365 PU105888;

1366 6) *Hypericum tertiaerum* Nikitin: seed, Monte Falcone subsection, sample CQ1 B6D, Piacenzian,  
 1367 MGPT-PU105889;

1368 7) *Hypericum perforatum* L.: seed, Stirone section, sample LA3 A5C, Calabrian, MGPT-  
 1369 PU105890;

1370 8) *Liriodendron geminata* Kirchheimer: seed, Santerno section, sample SNT1 B6L, Calabrian,  
 1371 MGPT-PU105891;

1372 9) *Magnolia allasoniae* Martinetto: seed, Marecchia section, sample MR3 B3E, Piacenzian,  
 1373 MGPT-PU105892;

1374 10) *Magnolia cor* Ludwig: seed fragment, Arda section, sample AD3 B7E, Gelasian, MGPT-  
 1375 PU105893;

- 1376 11) *Magnolia ludwigii* Ettingshausen: seed, Marecchia section, sample MR3 B3E, Piacenzian,  
1377 MGPT-PU105894;
- 1378 12) *Sinomenium cantalense* (E.M. Reid) Dorofeev: endocarp, Monte Falcone subsection, sample  
1379 CQ1 B6D, Piacenzian, MGPT-PU105895;
- 1380 13) *Sparganium nanum* Dorofeev: endocarp, Monte Falcone subsection, CQ1 B6D, Piacenzian,  
1381 MGPT-PU105896;
- 1382 14) *Stemona germanica* (Mai) Mai: a) apical view, b) basal view of one seed, Monte Falcone  
1383 subsection, sample CQ1 B6D, Piacenzian, MGPT-PU105897;
- 1384 15) *Eurya stigmosa* (Ludwig) Mai: seed, Arda section, sample AD3 B7E, Gelasian, MGPT-  
1385 PU105898;
- 1386 16) *Thymelaea* cf. *passerina* L.: seed, Arda section, sample AD3 B7E, Gelasian, MGPT-  
1387 PU105899;
- 1388 17) *Carpolithes minimus* (Szafer) Mai: a) outer surface, b) inner surface of the seed, Stirone  
1389 section, sample LA3 A5C, Calabrian, MGPT-PU105900.

1390