



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Science of atmospheric phenomena with JEM-EUSO

This is a pre print version of the following article:						
Original Citation:						
Availability:						
This version is available http://hdl.handle.net/2318/1578422 since 2016-06-30T15:15:33Z						
Published version:						
DOI:10.1007/s10686-014-9431-0						
Terms of use:						
Open Access						
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.						

(Article begins on next page)

Science of atmospheric phenomena with JEM-EUSO

The JEM-EUSO Collaboration

Received: date / Accepted: date

Abstract The main goal of the JEM-EUSO experiment is the study of Ultra High Energy Cosmic Rays (UHECR, $10^{19} - 10^{21}$ eV), but the method which will be used (detection of the secondary light emissions induced by cosmic rays in the atmosphere) allows to study other luminous phenomena. The UHECRs (ultra high energy cosmic rays) will be detected through the measurement of the emission in the range between 290 and 430 nm, where some part of Transient Luminous Events (TLEs) emission also appears.

This work discusses the possibility of using the JEM-EUSO Telescope to get new scientific results on TLEs. The high time resolution of this instrument allows to observe the evolution of TLEs with great precision just at the moment of their origin.

JEM-EUSO Collaboration for full author list see first paper of this volume

Kinga Slominska Space Research Centre E-mail: kslominska@cbk.waw.pl

Jan Blecki Space Research Centre Tel.: +48 – 22– 4966 –203 E-mail: jblecki@cbk.waw.pl

Jan Slominski Space Research Centre E-mail: abe@cbk.waw.pl

Gali Garipov Lomonosov Moscow State University Tel.: + 7 - 495 - 939 - 44 - 96 E-mail: garipov@eas.sinp.msu.ru

Pavel Klimov Lomonosov Moscow State University Tel.: + 7 - 495 - 939 - 57 - 34 E-mail: pavel.klimov@gmail.com The paper consists of four parts: review of the present knowledge on the TLE, presentation of the results of the simulations of the TLE images in the JEM-EUSO telescope, results of the Russian experiment Tatiana-2 and discussion of the possible progress achievable in this field with JEM-EUSO as well as possible cooperation with other space projects devoted to the study of TLE – TARANIS and ASIM.

In atmospheric physics, the study of TLEs became one of the main physical problems after their discovery in 1989. In the years 1992 - 1994 detection was performed from satellite, aircraft and space shuttle and recently from the International Space Station. These events have short duration (milliseconds) and small scales (km to tens of km) and appear at altitudes 50 - 100 km. Their nature is still not clear and each new experimental data can be useful for a better understanding of these mysterious phenomena.

Keywords Transient Luminous Events \cdot Elves \cdot Sprites \cdot Jets atmospheric phenomena

1 Introduction

The Transient Luminous Events are bright, short-lived, atmospheric phenomena occurring above the clouds level, some of them reaching the ionosphere.

The first suggestion on the existence of that kind of phenomena was given by the Scottish physicists C.T.R. Wilson in 1925 [1], who observed that at a certain altitude above the clouds level the value of the electric field connected to the thunder cloud discharge equals the critical value of the electric field allowing for the conventional breakdown mechanism. This in turn means the production of light.

First confirmed observations of the light phenomena above clouds level date since 1989. The experimental discovery of one type of these events was made accidentally in 1989 by scientists from the University of Minnesota Robert Franz, Robert Nemzek and John Winckler. These first detections were related to a phenomenon, which now is called the red sprite. The observation was performed during night time, and only in 1998 Mark Stanley and his collaborators from New Mexico Tech observed a day time sprite [2]. Since then, thousands of observations of different types of TLEs have been carried out from aircrafts, balloons, space stations and satellites. Apart from red sprites, other types of TLE – blue jets, Elves, halos and trolls - were observed. (Different types of TLE are schematically shown in the Fig. 1).

According to all collected up to date observations, the TLEs are triggered by a significant electric discharge either inside a cloud or between the cloud and the Earth. It is estimated that the energy produced in the atmosphere by all types of TLEs is about 700 MJ per minute [3]. In the case of Sprites and Elves the positive or negative cloud to Earth discharges are the necessary condition for this type of TLE to occur. Therefore, a strong correlation between the frequency of the thunderstorms and the Sprites and Elves is observed but the mechanism responsible for triggering the TLEs is still not clearly explained.



Fig. 1 A schematic overview of different types of the Transient Luminous Events. This picture comes from [4]

The wavelengths that these phenomena may produce depend on the composition of the atmosphere above the clouds - mainly Nitrogen - and on the allowed transitions observed in the molecules.

As JEM-EUSO detects wavelengths in the range 290 to 430 nm, we will consider only the second positive line of Nitrogen with possible wavelengths of 268 - 546 nm. Other lines are either outside the visible spectrum of JEM-EUSO or their intensity is much lower than the second positive line of Nitrogen [5].

2 JEM-EUSO in the TLE context

2.1 General description

The JEM-EUSO mission is devoted to the observation of Extensive Air Showers (EAS) produced by UHECRs traversing the Earth's atmosphere from above. For each event, the detector will make accurate measurements of the energy, arrival direction and nature of the primary particle using a target volume far greater than what is achievable from ground [6]. The corresponding increase in statistics will help to clarify the origin and sources of UHECRs as well as to bring new light onto particle physics mechanisms responsible of production, propagation and acceleration.

$2.2~{\rm The~detector}$

JEM-EUSO will be attached to the Japanese Experiment Module (JEM) of the International Space Station (ISS) at an orbit of 400 km above the surface of the Earth. The JEM-EUSO detecting system consists of a telescope with three Fresnel lenses and a focal surface made of photomultipliers and associated electronics, which converts the photons collected by the lenses into photoelectrons. The focal surface consists of some $3 \cdot 10^5$ pixels, each with an independent read out. The diameter of the telescope aperture is about 2.5 meters. The time unit used for sampling the data is 2.5 microseconds. The field of view (FoV) of the detector is 60° . The wavelengths measured by the detector may range from 290 to 430 nm. In addition to this main telescope part, the detection system is completed by an Atmospheric Monitoring system consisting of an Infrared Camera and a LIDAR to provide information concerning the cloud top height. A calibration system, which measures the efficiencies of the optics, of the focal surface and the data acquisition system is provided as well.

2.3 Triggering

The photomultipliers of the JEM-EUSO detectors operate correctly with the total charge not exceeding 250 pC per KI per time unit, which, in the case of JEM-EUSO, is $2.5\mu s$, which is assumed to be the Gate Time Unit (GTU). A KI is a 4 by 2 pixels unit. Assuming the 10 % efficiency of the optics and the electronics of the detector, this corresponds to $6.24 \cdot 10^{15}$ photoelectrons per KI per second, or $1.56 \cdot 10^{10}$ photoelectrons per KI per GTU. If this value is exceeded the voltage on the photomultipliers will be reduced so that the detection system remains safe.

3 Estimation of TLE frequency for JEM-EUSO based on Tatiana-2 experimental data.

The global occurrence of TLE can be obtained in space-borne experiments. Only few detectors devoted to TLE measurements were launched to the orbit: ISUAL on board FORMOSAT-2 satellite [7], UV detector on board Universitetsky-Tatiana satellite [8], UV and IR detector on board Universitetsky-Tatiana-2 satellite [9] and JEM-GLIMS on board ISS [10]. ISUAL and JEM-GLIMS have spatial resolution and provide spectral measurements in wide wavelength



Fig. 2 Global distribution of all transient events

bands, but Tatiana-1 and Tatiana-2 detectors provide measurements in 240 - 400 nm, in the same wavelength range as EAS fluorescent radiation. It is important to mention that in both Tatiana experiments and in JEM-EUSO PMTs with UV filters are used to measure radiation. Oscillograms of fast luminous events are recorded. Tatiana-1 detector recorded only one flash (transient) event per orbit (because of poor transmission channel between satellite and ground receiving station) and obtained small statistics (≈ 300 TLE events during two years of operation). Tatiana-2 could measure the most powerful event per each minute during night part of orbit, which allowed to collect larger transient statistics (more than 2500 TLE events during three months). More details about detectors and their results are in [11], [12]. Global distribution of all transient events, obtained by Tatiana-2 satellite is presented in the Fig. 2. The region of observations during local night period has distinct borders: 60° of latitude in the Northern hemisphere and 30° in the Southern hemisphere. Most of events are concentrated above continents in equatorial regions of America, Africa and Indo – China, this distribution seems similar to common lightning distribution. But the unexpected effect can be seen if we divide all events into groups with various energy ranges (number of photons). Events with smaller number of photons have more even geographical distribution and do not obviously correlate with huge thunderstorm regions (Fig. 3).

Tatiana's UV detectors have no spatial resolution and it is impossible to definitely distinguish types of TLE, to determine their actual size and spatial light emission distribution. However, the collected statistics allows us to estimate the low limit of transient event rate expected for space-borne detectors like JEM-EUSO and its influence on their duty cycle. In the period October 2009 – January 2010 Tatiana-2 detectors orbited the Earth 797 times with 320 hours of operation time in shadowed "night" part. During these three months 2628 events with $N_{ADC} > 80$ were measured. The average rate of TLE events from this data is 0.13 min⁻¹ or $10^{-4} \text{ hr}^{-1}\text{km}^{-2}$. The instantaneous frequency of TLE in local thunderstorm area could be significantly



Fig. 3 Global map of transients with photon number less than $5 \cdot 10^{21}$ in the atmosphere

higher (~ 10^{-3} hr⁻¹km⁻²). It is worth noticing that this number is order of magnitude higher than TLE frequency, measured by ISUAL [12], [13]. It could be explained by the sensitivity of the detector whose instrumental threshold is ~ 10^{20} in photon number. As it was mentioned above, these faint flashes are distributed more evenly in global map. ISUAL can not measure them.

To estimate the number of TLE events which could be measured by JEM-EUSO or influence its operation, we should take into account the ratio of detectors fields of view. Tatiana-2 observed an area of $7 \cdot 10^4$ km², JEM-EUSO in nadir mode will observe $2 \cdot 10^5$ km², three times larger than Tatiana-2. It means that JEM-EUSO will measure one UV flash from atmospheric phenomena at least every third minute. In the worst case, when the size of TLE is comparable to JEM-EUSO FoV (for example in case of a large Elve) or very intensive TLE which produces a huge amount of light scattered in detectors parts, it can cause a pause in EAS measurements for 70 seconds (it is the most conservative estimation of dead time within which the light source will leave JEM-EUSO FoV). Such dead time is ~ 40 % of the duty cycle. This is the upper bound. To obtain more feasible numbers we should take into account energy (number of photons) distribution of observed events.

The typical event measured by Tatiana-2 satellite has about 10^4 photons per ms in 0.4 cm² aperture of detector. It means that more than 10^6 transient photons per GTU (2.5 μ s) will come in JEM-EUSO aperture (4.5 m²). The JEM-EUSO optics efficiency (0.4) and the difference between orbit altitude of ISS (≈ 400 km) and Tatiana-2 satellite (≈ 830 km) were taken into account. In these calculations spatial light distribution in transients was not considered. Transient photons may be distributed among many pixels of photo receiver and signal in each pixel could be smaller. Actual size of TLE depends on its type. In case of Elves this signal will be distributed among almost all pixels and will give few photoelectrons per GTU. Smaller events will significantly affect the operation of several specific Photo Detector Modules (PDM) and just slightly reduce the FoV. The rest of the photo detector will be illuminated by diffuse light scattered in the detector (for our estimations we will consider diffuse light as 10 % of the total number of photons per event). Diffuse light gives a signal increase of ≈ 0.5 photoelectrons per GTU in one pixel. This is a minor addition to background signal which will not influence detector threshold and therefore a duty cycle. The fraction of flashes which produce more than 2 photoelectrons per GTU of scattered light in each pixel is 20 % of all measured by Tatiana-2 satellite. If we consider 70 seconds pause only for these TLE the dead time becomes near 8 %. This number is consistent with other estimations of duty cycle which take into account clouds and thunderstorm regions.

4 Estimation of the TLE light collected by the JEM-EUSO experiment

4.1 Elves

The most common phenomena are ELVES - Emission of Light and VLF perturbations due to Electromagnetic Pulse Sources [14]. They occur at the lower ionosphere, at altitudes of about 90 km. They are concentric rings of light expanding with speed close to the speed of light. The diameter of the Elves disk may range from 200 to 500 km. The whole process of expansion takes up to a millisecond. The Elves result from the interaction of an electromagnetic pulse produced with the cloud-to-ground discharge with the molecules and particles in the ionosphere. To estimate the image viewed by the JEM-EUSO detector while observing an Elve example, we have estimated the parameters of the average Elve basing our assumptions on the results provided by the ISUAL experiment and presented in [3]. The occurrence rate of the Elves is 35 per minute around the world. The spatially averaged brightness of an elve presented in [3] is: $0.17 \pm 0.08MR = 1.36 \cdot 10^{14}$ ph sr⁻¹m⁻²s⁻¹. The unit R stands for "Rayleigh" (unit of photon flux) typically used for these atmospheric phenomena. We assume an elve with the diameter of 300 km and the timescale of 2 ms = 800 GTU, which should allow for recording of the whole event and the observation directly in the JEM-EUSO detector in the nadir mode. This is the least favorable scenario in the sense that the geometry of the system allows for the maximum amount of light reaching the focal surface of the detector.

With all these assumptions, we get $9.25 \cdot 10^{11}$ photons, which is an estimate for the whole event, in all wavelength channels. As elves are ionospheric phenomena, they are mostly red. The contribution of "blue" wavelengths is marginal. If we take 13 % of the obtained photons fitting our wavelength requirements (second positive line of Nitrogen), we will get the estimated number as: $1.2 \cdot 10^{11}$ photons. This number of photons will be recorded as a total number by all pixels of the focal surface throughout the duration of the event.

The number of pixels in the JEM-EUSO detector is 315 648, it equals to about 40 000 KIs. The distribution of the photons among the separate KIs

has not been considered here. Similarly, the time distribution of the incoming photons has not been taken into account in this work.

Both these factors will reduce the number of photons reaching a KI in one GTU by a few orders of magnitude. It is also important to note that the estimation relates to one such event and not to the total number of photons coming from other sources in the atmosphere. The calculations here presented are based on averaged results, while the most dangerous for the photomultipliers are the extreme cases of events producing light, for which the order of magnitude of the brightness may be a few hundred times greater.

The observed characteristics of different types of TLE can be found in the Table 1.

4.2 Sprites

Sprites are associated with giant storm clouds with dimensions over 1000 km producing strong electric field in the mesosphere in a volume of over 10^4 km³. Sprites are massive but weak luminous flashes, appearing at altitudes of 40-90km. The heads of sprites are predominantly red. The brightest region lies in the altitude range 65 - 75 km, above which there is often a faint red glow structure that extends to about 90 km. Below the bright red region, blue tendril-like filamentary structures often extend downward to 40 km. Sprites rarely appear singly, usually occurring in groups – two, three or more. The duration of sprites is of the order of milliseconds. Currents associated with sprites have intensity of over kA. The pictures of sprites obtained from the ground and from aircraft show complex structures that assume a variety of forms. Early research reports for these events called them "upward lightning", "upward discharges", "cloud - to - stratosphere discharges" and "cloud - to - ionosphere discharges". Now they are called sprites. The tendril – streamer structure under the upper part of the phenomenon – the so called head of a sprite shows extremely interesting and as yet not fully explained phenomenon of merging streamers, which form bright spherical structures – the beads. Measurements of microsecond time resolution may help to explain the mechanism behind the phenomenon. The brightness measured by ISUAL is 1.5 ± 1.1 MR, which translates into some $3.48 \cdot 10^{11}$ ph in total and $1.46 \cdot 10^{11}$ in the wavelength region detected by the JEM-EUSO. This number is the spatially integrated amount of photons recorded throughout the whole event duration.

4.3 Other types of TLE

Blue Jets Blue jets are optical ejections from the top of the electrically most active regions of thunderstorms. Following their emergence from the top of the thundercloud, they typically propagate upward in narrow structure in form of cones of about 15 degrees full width at vertical speeds of roughly 100 km/s, fanning out and disappearing at heights of about 40 - 50 km. At the base their

Type of TLEs	Altitude Regime	Transverse Dimensions	Spatial Characteristics	Apparent Motion	Duration
Sprites	$\sim 50-90~{\rm km}$	$\sim 1-20~{\rm km}$	Top (> 80 km) diffuse Bottom (< 70 km) structured	Top - upward Bottom - downward	few ms
Elves	$\sim 90~{\rm km}$	> 100 km	Diffuse	Lateral expansion	$\sim \mathrm{ms}$
Blue jets	$\sim 18-45~{\rm km}$	few km	Structured	Upward	hundreds ms
Giant Blue jets	$\sim 18-75~{\rm km}$	few km	Structured	Upward	hundreds ms
Hallos	$\sim 75~{\rm km}$	50 km	Diffuse	Downward	$\sim \mathrm{ms}$
Trolls	$\sim 60 - 70 \text{ km}$	kms	Structured	Upward within decaying sprite tendrils	hundreds ms

Table 1 Principal Types of TLE in the Upper Atmosphere [16]

diameter is about 400 m. Blue jets are rather rare events. Their appearance rate is much lower than that of sprites. Blue jets are not aligned with the local magnetic field. They are triggered by the electric field inside the cloud. Their appearance is not connected to a cloud-to-ground discharge. The brightness of blue jets is estimated as 0.5 MR [15].

Gigantic Jets They are much larger than the Blue Jets. Gigantic Jets originate form the cloud top and develop into a tree-like structure reaching the lower ionosphere. Their propagate upward at the speed of about 100 km/s. The ISUAL measured only 8 such events. The brightest one was 0.8 MR.

5 Conclusions and the Future

The frequency of occurrence of the TLE events is large. Their characteristics have been described by different groups working with detectors into orbit and with on ground detecting systems. JEM-EUSO is unique in that its time and spatial resolution is very high and will allow time evolution measurement. Two more experiments dedicated to measuring TLEs will soon be put into orbit: ASIM and Taranis. The set of data produced by all these experiments may create an exceptional chance to further investigate the spatial and time evolution of the TLEs.

The amount of light produced by the two most common types of TLEs does not pose danger to the photomultipliers of the detector. However, these phenomena constitute a significant background to the detection of UHECRs. Many aspects of the detection of the signal from the TLEs remain to be considered in next developments of this work. One interesting path would also be to test the possibility of finding correlations between the occurrence of very energetic particles in the Earth's atmosphere and the occurrence and magnitude of sprites. These goals are in the reach of the JEM - EUSO mission in space and, together with its unique features above mentioned, they may lead to significant steps forward in the field of Transient Luminous Events research.

6 Acknowledgments

This work is supported partially by the grant MNiSW N307 065834 and grant of Russian Foundation for Basic Research No. 12-05-31025-mol-a.

References

- 1. Wilson, C. T. R., The electric field of a thundercloud and some of its effects, Proc. Phys. Soc. London, (1925), 37, 32D
- 2. Schneider, D., Scientific American, (2000), vol. 88, no 4., p. 1
- 3. Kuo et al., Radiative emission and energy deposition in transient luminous events, J. Phys. D: Appl. Phys. 41 (2008) 234014 (14pp)
- 4. http://upload.wikimedia.org/wikipedia/commons/c/c4/Upperatmoslight1.jpg
- Kuo, C.-L., et al., Modeling elves observed by FORMOSAT-2 satellite, J. Geophys. Res., 112, A11312, doi:10.1029/2007JA012407 (2007)
- Adams Jr., J.H. et al. (JEM-EUSO Collaboration), An evaluation of the exposure in nadir observation of the JEM-EUSO mission, Astroparticle Physics, 44, 76-90 (2013)
- Chern, J. L., R. R. Hsu, H. T. Su, S. B. Mende, H. Fukunishi, Y. Takahashi, and L. C. Lee, Global survey of upper atmospheric transient luminous events on the ROCSAT-2 satellite, J. Atmos. Sol. Terr. Phys., 65(5), 647-659, (2003)
- Garipov, G.K., Panasyuk, M.I., Rubinshtein, I.A., et al., UV Detector onboard Research and Education MicroSatellite Universitetsky-Tatiana of Moscow University, Prib. Tekh.Eksp., (2006), no. 1, pp. 135-141
- Sadovnichy V.A., M.I. Panasyuk, I.V. Yashin et al., Investigation of Space Environment Aboard Universitetsky-Tatiana and Universitetsky-Tatiana-2 Microsatellites, Solar System Res., (2011), 45(1), 3-29
- Sato, M., Y. Takahashi, M. Kikuchi, M. Suzuki, A. Yamazaki, and T. Ushio, Lightning and Sprite Imager (LSI) onboard JEM-GLIMS, IEEJ Trans. FM, 131 (12), pp. 994-999, (2011)
- Garipov, G.K., Khrenov, B.A., Panasyuk, M.I., et al., UV Radiation from the Atmosphere: Results of the MSU Tatiana Satellite Measurements, Astropart.Phys., (2005), vol. 24, pp. 400-408
- N. N. Vedenkin, G. K. Garipov, P. A. Klimov, V. V. Klimenko, E. A. Mareev, O. Martinez, V. S. Morozenko, I. H. Park, M. I. Panasyuk, U.Salazar, V. I. Tulupov, B. A. Khrenov, and I. V. Yashin, Atmospheric Ultraviolet and Red Infrared Flashes from Universitetsky Tatiana-2 Satellite Data, JETP, 113, 5, ISSN 1063-7761, (2011)
- A. B. Chen, C. L. Kuo et al. Global distribution and occurrence rate of TLE. (2008) JGR, v.118, A0806, doi101029/029/2008JAO013101
- 14. Fukunishi, H., et al., Elves: Lightning-induced transient luminous events in the lower ionosphere Geophys. Res. Let., 23, 2157-2160, (1996)
- Wescott, Sentman, et al., Sprite 94 campaign, Geophysical Research Letters, (1995), vol. 22, iss. 10, pp. 1205-1208
- Sentman, D. D., Overview of Transient Luminous Events, Presentation at Space Science Laboratory Seminar, University of Berkeley, California, 2005.