

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Chronic unpredictable stress and long-term ovariectomy affect arginine-vasopressin expression in the paraventricular nucleus of adult female mice**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/149901> since 2016-07-01T17:38:50Z

*Published version:*

DOI:10.1016/j.brainres.2014.09.006

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in BRAIN RESEARCH, 1588, 2014, 10.1016/j.brainres.2014.09.006.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.brainres.2014.09.006

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0006899314011937>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/149901>

**BRES-D-14-01077-R1**

**Chronic unpredictable stress and long-term ovariectomy affect arginine-vasopressin expression in the paraventricular nucleus of adult female mice.**

Grassi D.<sup>a,b,d</sup>, Lagunas N.<sup>a,1</sup>, Calmarza-Font I.<sup>a</sup>, Diz-Chaves Y.<sup>a</sup>, Garcia-Segura L.M.<sup>a</sup>,  
Panzica G.C.<sup>b,c</sup>

(a) Cajal Institute, CSIC, Madrid, Spain, (b) University of Torino, Department of Neuroscience "Rita Levi Montalcini", Torino (Italy), and Neuroscience Institute Cavalieri Ottolenghi (NICO), Orbassano (Torino, Italy), (c) National Institute of Neuroscience, Torino, Italy (INN), (d) Institute of Anatomy and Cell Biology, Department of Molecular Embryology, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany

<sup>1</sup>**Present address:** Université François Rabelais · INSERM U 930 Imagerie et Cerveau, Tours, France

**Running title:** Stress, ovariectomy and arginine-vasopressin in PVN

**Corresponding author:** Professor GianCarlo Panzica, PhD, Department of Neuroscience, Laboratory of Neuroendocrinology, Neuroscience Institute Cavalieri-Ottolenghi (NICO), Regione Gonzole 10, Orbassano (Torino, Italy). Voice: +39 011 6706607. Fax: +39 011 2366607. e-mail: giancarlo.panzica@unito.it

**ABSTRACT**

Arginine-Vasopressin (AVP) may regulate the hypothalamic-pituitary-adrenal axis (HPA) and its effects on depressive responses. In a recent study, we demonstrated that Chronic Unpredictable Stress (CUS) depressive effects are enhanced by long-term ovariectomy (a model of post-menopause). In the present study, we investigated the effects of long-term ovariectomy and CUS on AVP expression in different subdivision of the paraventricular nucleus (PVN) of female mice. Both long-term ovariectomy and CUS affect AVP immunoreactivity in some of the PVN subnuclei of adult female mice. In particular, significant changes on AVP immunoreactivity were observed in magnocellular subdivisions, the paraventricular lateral magnocellular (PaLM) and the paraventricular medial magnocellular (PaMM), the 2 subnuclei projecting to the neurohypophysis for the hormonal regulation of body homeostasis. AVP immunoreactivity was decreased in the PaLM by both the long-term deprivation of ovarian hormones and the CUS. In contrast, AVP immunoreactivity was increased in the PaMM by CUS, whereas it was decreased by ovariectomy. Therefore, present results suggest morphological and functional differences among the PVN's subnuclei and complex interactions among CUS, gonadal hormones and AVP immunoreactivity.

**Key words:** Vasopressin; depression; menopause; stress; hypothalamus.

## 1. INTRODUCTION

Evolutionary success depends on our ability to adapt to changing circumstances. The neuroendocrine response to stress is an excellent example of a plastic system that responds to threats to homeostasis and alters its output to meet current and expected future demands.

In recent years, magnocellular neurons of the paraventricular nucleus of the hypothalamus (PVN) have been reported to be involved in the regulation of hypothalamic-pituitary-adrenal axis (HPA) and its behavioral and emotional effects (see for extensive reviews, Landgraf, 2001, Engelmann et al., 2004, Landgraf et al., 2007). In particular, HPA-related stress responses involve the action of several factors including arginine-vasopressin (AVP) in magnocellular nuclei (for a review see Rotzinger et al., 2010). The contribution of centrally released vasopressin in the regulation of blood pressure and cardiovascular functions (Pyner, 2009), water intake (Bolignano and Zoccali, 2010, Moeller and Fenton, 2012) and in the emotional and social behavior is well established (Insel, 2010, Albers, 2012, Bosch and Neumann, 2012). On the other hand, it has been postulated that enhanced action of vasopressin in the brain may contribute to generation of pain, anxiety, panic attacks, aggression and depression (Coccaro et al., 1998, Siegel et al., 2007, Caldwell et al., 2008, Surget and Belzung, 2008). Among the different animal models to study the response to stress, Chronic Unpredictable Stress (CUS) has long been used as a model of emotional stress that can induce depressive and anxiety like-behaviors in rodents (Mineur et al., 2006).

The response to stress depends on gonadal hormones and is sexually dimorphic (Patchev et al., 1995, Viau, 2002). Sex differences exist in the response of the HPA axis to stress: i) women react more robustly than men (Kudielka and Kirschbaum, 2005) and ii) in both sexes, activation of the HPA axis may inhibit reproductive functions

(Cameron, 1997). Conversely, the sex differences in HPA function are in part due to differences in the circulating gonadal steroid hormone milieu (Handa et al., 1994). A recent magnetic resonance imaging study demonstrated sex differences in the activation of brain circuitries sensible to stress. In women the activation varies upon the hormonal cycle, whereas in man activated circuitries are similar to those observed in the early follicular menstrual phase of women (when estrogen and progesterone are low, Goldstein et al., 2010).

Recently, we have focused our attention on the inter-regulation between the stress response and gonadal hormones. We used an animal model of post-menopause (long-term ovariectomy), field in which we could study the emotional behavioral effect of the CUS. In a recent study (Lagunas et al., 2010), we demonstrated that CUS emotional effects on depressive and anxiety like-behaviors are enhanced by long-term gonadal hormones deprivation.

Based on these findings, the aim of the present study was to investigate the response of the different PVN subnuclei to the CUS in long-term ovariectomized female mice using corticosterone plasma levels and AVP immunoreactivity as parameter of stress response.

## **2. RESULTS**

### **2.1 Serum corticosterone assay**

Serum corticosterone levels were not significantly different between control animals and ovariectomized (OVX) animals (not submitted to CUS). However, plasma corticosterone levels were significantly increased in OVX-CUS animals compared to controls (Fig. 1). Data were analyzed by a two-way ANOVA with CUS and the

ovariectomy as the two independent variables. The results of two-way ANOVA showed no effect of CUS on serum corticosterone levels [ $F_{(1,12)}= 3.52$ ,  $p= 0.09$ ], a significant effect of ovariectomy [ $F_{(1,12)}= 8.71$ ,  $p= 0.016$ ] and a significant effect for the interaction of the two variables [ $F_{(1,12)}= 8.64$ ,  $p= 0.016$ ]. The Bonferroni's test detected a significant increase of corticosterone levels in OVX females exposed to CUS in comparison to the other groups ( $p<0.05$ ).

## 2.2 AVP expression in the PVN.

The rodents' PVN consists of several distinct subnuclei with different cytological, neurochemical and circuitry features: the paraventricular dorsal cap (PaDC), the paraventricular lateral magnocellular (PaLM), the paraventricular medial magnocellular (PaMM) and the paraventricular medial parvocellular (PaMP) subnuclei (Benarroch, 2005, Ferguson et al., 2008). We decided, therefore, to measure AVP immunoreactivity in the four different subnuclei, separately.

Qualitative observation of AVP immunopositive staining revealed differences in the immunoreactivity pattern among the different experimental groups (Fig. 2). These qualitative differences were confirmed by the morphometric analysis (Fig. 3).

For each different PVN subnucleus, we analyzed the data by two-way ANOVA with CUS and ovariectomy as independent variables and the volume occupied by AVP immunoreactivity as dependent variable.

In the PaDC, the two-way ANOVA showed a significant effect of stress [ $F_{(1,12)}= 13.62$ ,  $p= 0.003$ ], no significant effect of ovariectomy [ $F_{(1,12)}= 3.06$ ,  $p= 0.105$ ] and a significant effect of the interaction of the two variables [ $F_{(1,12)}= 19.72$ ,  $p= 0,0008$ ]. The post-hoc Bonferroni's test showed that the volume occupied by AVP immunoreactive material

was significantly decreased in ovariectomized females compared to all the other groups ( $p < 0.01$  vs Control,  $p < 0.05$  vs CUS,  $p < 0.001$  vs CUS+OVX).

In the PaLM, the two-way ANOVA showed a significant effect of stress [ $F_{(1,12)} = 6.04$ ,  $p = 0.03$ ], of ovariectomy [ $F_{(1,12)} = 42.25$ ,  $p < 0.0001$ ] and of interaction of the two variables [ $F_{(1,12)} = 11.69$ ,  $p = 0.005$ ]. The Bonferroni's test showed a significant decrease of AVP immunoreactivity in all groups compared to control female mice [ $p < 0.01$  vs CUS,  $p < 0.001$  vs OVX and CUS+OVX].

In the PaMM, the two-way ANOVA showed a significant effect of stress [ $F_{(1,12)} = 52$ ,  $p < 0.0001$ ] and ovariectomy [ $F_{(1,12)} = 16.96$ ,  $p = 0.001$ ]. No significant effect of interaction was detected [ $F_{(1,12)} = 2.93$ ,  $p = 0.112$ ]. The Bonferroni's test showed a significant decrease of AVP immunoreactivity in OVX group compared to all the other groups ( $p < 0.01$  vs Control,  $p < 0.001$  vs CUS, and  $p < 0.001$  vs OVX+CUS). AVP immunoreactivity significantly increased in CUS group compared to controls ( $p < 0.01$ ) and OVX ( $p < 0.001$ ).

In the PaMP, the two-way ANOVA showed a significant effect of stress [ $F_{(1,12)} = 5.47$ ,  $p = 0.037$ ], but no effect of ovariectomy [ $F_{(1,12)} = 0.12$ ,  $p = 0.74$ ] and of the interaction [ $F_{(1,12)} = 2.12$ ,  $p = 0.17$ ]. The post hoc Bonferroni's test reported no significant differences among the four experimental groups.

### 3. DISCUSSION

Our results show that CUS did not affect plasma corticosterone levels in gonadally intact females, but it increased plasma corticosterone levels in OVX females submitted to CUS compared to control females (Fig. 1). These findings are in concordance with our previous results demonstrating that gonadally intact females submitted to CUS did not show differences in anxiety and depression-like behavior compared to control



group. All these data suggest that the gonads carry out a protective effect against the stress-dependent anxiety and depression-like behavior, as reported also in other studies (Lagunas et al., 2010, Vega-Rivera et al., 2013). Since the corticosterone secretion is normally regulated by the light-dark cycle, and the exposure to constant light disrupts this circadian rhythm (Park et al., 2013), the manipulations of this cycle on the last day of CUS could have altered the corticosterone levels. However, we have not observed differences between the CUS group and controls, therefore it seems unlikely that this created a significant problem. The significant difference between OVX animals submitted to CUS and controls, with a significant increase of plasma corticosterone levels in OVX animals submitted to CUS, could be related with the lack of estrogen inhibition of the HPA axis response on PVN (Lund et al., 2006, Weiser and Handa, 2009), and also with the decreased levels of estrogen receptor  $\beta$  mRNA expression in the PVN of OVX animals (Jin et al., 2005, Barker and Galea, 2009, Liu et al., 2011).

The present findings indicate that both long-term ovariectomy and CUS may affect AVP immunoreactivity in the PVN of adult female mice. In particular, significant changes of the AVP immunoreactive material compared to control animals were observed in magnocellular subdivisions, the PaLM and the PaMM, the 2 subnuclei projecting to the neurohypophysis for the hormonal regulation of body homeostasis. On the other hand, the parvocellular subnuclei, the PaDC and the PaMP, were not influenced.

Long-term gonadal hormones' deprivation induced a significant decrease of AVP immunoreactivity in the PVN and in particular in the PaLM, not influencing the other magnocellular and parvocellular subnuclei. Similarly, De Vries et al. (1984) demonstrated that during the next 2 weeks from the ovariectomy no effect of ovariectomy could be detected upon the AVP innervation of dorsal vagal complex that

is derived from the parvocellular division of the PVN (Sawchenko and Swanson, 1982) in mice. Moreover, it has been shown that administration of estradiol (E<sub>2</sub>) to postmenopausal women increases both circulating levels of AVP (Bossmar et al., 1995) and rapid intrahypothalamic AVP release (Wang et al., 1995, Burbach et al., 2001).

In our females, CUS significantly affects AVP immunoreactivity in the PVN but only in the magnocellular subnuclei, PaLM and PaMM, with no significant effects in the parvocellular subdivisions. Moreover, CUS affects AVP immunoreactivity in a different way within the magnocellular subnuclei: in the PaLM, CUS induced a decrease of AVP immunoreactivity, while in the PaMM, CUS caused an increase in both gonadally intact (CUS) and ovariectomized animals (OVX+CUS).

The difference within magnocellular and parvocellular neuronal subpopulations could be due to the differential expression of estrogen receptors (ER) in the PVN. In the rat and mouse, hypothalamus ER $\beta$  is the predominant ER isoform found in the PVN (Mitra et al., 2003, Merchenthaler et al., 2004). Interestingly, E<sub>2</sub> has been shown to activate the Mitogen-Activated Protein Kinases (MAPK) pathway via Extracellular signal-Regulated Kinases (ERK) 1/2 in the PVN and the SON of mice (Abraham et al., 2004). In the PVN, this is potentially mediated via either ER $\beta$  (Grassi et al., 2013) and/or G protein-coupled receptor 30 (GPR30). In the hypothalamus intense GPR30 expression was found in the PVN. In the rat PVN, GPR30 immunoreactivity is prominent in the magnocellular PVN with low-level labelling in the parvocellular PVN. Dual labelling of hypothalamic sections revealed that GPR30 immunoreactivity was present in AVP magnocellular neurons and 50–70% co-expressed GPR30 and AVP in the PVN (Hazell et al., 2009). In addition, looking in detail into the picture of Hazell's paper (Fig. 4 A-B-C), the higher levels of AVP and GPR30 co-expression are localized in the part of

magnocellular subdivision that we identify as PaLM suggesting an involvement of GPR30 in the regulation of AVP in the PaLM.

Present results, showing that CUS selectively affects AVP immunoreactivity in the PaLM and PaMM subdivisions are in agreement with previous studies indicating that the magnocellular neurosecretory system plays a role in some, but not all, chronic stress paradigms (Albeck et al., 1997, Guccione et al., 2002, Seale et al., 2004). On the contrary, no effects were detectable in AVP immunoreactive levels in the parvocellular PVN subdivision. This is in agreement with (Pinnock and Herbert, 2001) who demonstrated that the AVP response to both acute and repeated stress was unaltered in the low-dose corticosterone-treated rats compared to control animals in the parvocellular PVN, suggesting that a return of AVP to lower levels is necessary to allow this peptide to respond to repeated stress.

In conclusion, our results indicate that magnocellular subdivisions, PaLM and PaMM, known to be the part of the PVN sensible to the ovarian hormone-dependent regulation of AVP system (Wang et al., 1995), are differentially regulated by the exposure to CUS. As previously discussed this is probably related to differences in the distribution of nuclear or membrane-bound estrogen receptors (Merchenthaler et al., 2004, Hazell et al., 2009), or to differences in estrogen-sensitive afferents to PVN (Grassi et al., 2010). It is interesting to underline that the specific stimulation of GPR30 in male and ovariectomized female mice induced anxiogenic effects (Kastenberger et al., 2012). In our previous experiment, we demonstrated an enhancement of anxiety and depressive-like behaviors in OVX mice submitted to CUS (Lagunas et al., 2010) and now, the present findings show a correlation between these altered behaviors and modifications in the levels of AVP immunoreactivity in some PVN's subnuclei, in

agreement with the involvement of this system in depression and anxiety (Landgraf, 2001, Surget and Belzung, 2008, Rotzinger et al., 2010).

#### **4. EXPERIMENTAL PROCEDURE**

Female C57BL6 mice (Harlan, Barcelona, Spain) housed in groups, with unlimited access to food and water, under controlled light schedule (12 hours light:12 hours dark) were used for this experiment. The animals were handled following the European Union guidelines (Directive 86/609/EEC). Our Institutional Animal Use and Care Committee (CSIC) approved the experimental procedures. Special care was taken to minimize animal suffering and to reduce the number of animals used to the minimum.

##### **4.1. *Chronic unpredictable stress protocol (CUS)***

At the age of two months, female mice were assigned to two different groups: intact and ovariectomized. Mice belonging to the second group were bilaterally ovariectomized under isoflurane anesthesia. Four months after ovariectomy, the groups were subdivided into 2 subgroups: stressed and no stressed mice. The stressed groups were submitted to the CUS protocol to define the effect of chronic unpredictable stress in physiological and long-term gonadal hormones' deprivation conditions. The stressed animals were submitted to CUS protocol during 4 weeks. The stress procedure was a modified version of a previously described model (Moreau et al., 1992). The stress consisted of repeated psychological stressors, such as: social isolation, predator proximity, presence of an intruder, continuous room and cage changes, cage tilting, damp sawdust and lights on for a long period (see Table 1). None of the stressors involved food or water deprivation. Control and CUS animals were housed in the same animal facility but in separate rooms.

In summary, we obtained a total of 4 experimental groups: (i) (Control) intact not stressed mice (n=24); (ii) (CUS) intact stressed mice (n=25); (iii) (OVX) ovariectomized not stressed mice (n=24) and (iv) (OVX+ CUS) ovariectomized stressed mice (n=24). At the end of CUS protocol, at the age of 7 months, animals were assessed for forced swimming test and elevated plus-maze test (Lagunas et al., 2010). Some animals (n=4) from all experimental groups were not behaviorally assessed and their brains were used for vasopressin immunohistochemical staining.

#### 4.2. Determination of serum corticosterone.

Blood samples were collected immediately before the perfusion by tail nick and collected into EDTA-coated capillary tubes and kept on ice. The blood was rapidly placed in microfuge tubes and allowed to clot for 1 h. After centrifugation at 3500 rpms serum was stored at -70 °C until assayed.

The serum corticosterone levels were measured using a kit of enzyme-immunoassay (EIA) for the quantitative determination of corticosterone in mouse serum (IDIS OCTEIA) according to the manufacturer's instructions. The sensitivity of the measurement was 0.55 ng/mL. The intra- and inter-assay coefficients of variation were 3.8% and 7.7%, respectively. The results are presented as ng/mL.

#### 4.3. Immunohistochemistry

The animals (seven month-old) were deeply anesthetized with pentobarbital (Normon Veterinary Division, Madrid, Spain, 50 mg/kg) and perfused through the left cardiac ventricle with saline solution (0.9% NaCl) and then with fixative solution (4% paraformaldehyde in 0.1M PBS, pH 7.4). Brains were quickly dissected, immersed for 4–6 h at 4°C in freshly prepared fixative solution and then rinsed with 0.1M PBS. Brains were serially cut in the coronal plane at 50µm thickness with a Vibratome (VT

1000 S, Leica Microsystems, Wetzlar, Germany). The plane of sectioning was oriented to match the drawings corresponding to the transverse sections of the mouse brain atlas (Paxinos and Franklin, 2001). Sections were collected in a cryoprotectant solution (Watson et al., 1986) at  $-20^{\circ}\text{C}$ . Every fourth section (one section every  $200\ \mu\text{m}$ ) was stained for AVP immunohistochemistry according to our standard protocol (Plumari et al., 2002, Grassi et al., 2010). Briefly, free-floating sections were washed overnight at  $4^{\circ}\text{C}$  in PBS 0.1M then in PBS containing 0.2% Triton X-100 and 0.2% BSA, and finally treated with a solution of PBS/methanol (1:1) with 0.3% hydrogen peroxide (to quench endogenous peroxidase activity). The sections were incubated for 48 hours at  $4^{\circ}\text{C}$  with a rabbit polyclonal AVP antibody (Chemicon, Millipore Ibérica, Madrid, Spain) diluted 1:10,000 in PBS-Triton X-100-BSA buffer, containing 3% normal serum goat. A biotinylated goat anti-rabbit antibody (Thermo scientific, Pierce, Rockford, IL, USA) was then used at a dilution of 1:300 for 120 min at room temperature. The antigen-antibody reaction was revealed by incubation with avidin-peroxidase complex (Thermo scientific, Pierce) for 90 min. Peroxidase activity was visualized with a solution containing 0.187mg/ml 3,3-diamino-benzidine (Sigma, Madrid, Spain) in PBS. Sections were mounted on chromallum-coated slides, air-dried, cleared in xylene, and cover slipped.

#### **4.4. Morphometrical analysis**

The morphometric analysis of AVP immunostaining was performed on coded sections without previous knowledge of the experimental group. The quantification of AVP immunoreactivity was achieved within the PVN on 3 coded sections (covering the entire rostrocaudal extension of the right PVN) by estimating the percentage of volume occupied by AVP immunoreactive material with the Weibel's point counting method (Weibel, 1979), using a grid of 20 square boxes of  $8\ \mu\text{m}^2$  each and a x20 objective

(Grassi et al., 2010). Since rodents' PVN consists of several distinct subnuclei with different cytological and neurochemical features (Ferguson et al., 2008), quantification was separately performed on four different subnuclei of the PVN [paraventricular dorsal cap (PaDC), paraventricular lateral magnocellular (PaLM), paraventricular medial magnocellular (PaMM) and paraventricular medial parvocellular (PaMP) subnuclei] (see Fig. 2). The PVN subdivisions were identified following the anatomical description based on Nissl-stained material (Silverman and Pickard, 1983, Armstrong, 1985). Percentage of the volume fraction occupied by immunoreactive material versus the total volume of each PVN subnucleus was considered for statistical analysis. In figure 3, volume fraction values were normalized to control values.

#### **4.5 Data analysis**

Statistical analysis of the experimental data was carried out using SPSS (version 17.0). Two-way analysis of variance (ANOVAs) with Bonferroni post hoc test was conducted to evaluate effects of the ovariectomy, the CUS and the potential interactions. The level for statistical significance was set at  $p < 0.05$ .

**5. Acknowledgements:** This study has been supported by the Ministerio de Ciencia e Innovación, Spain (BFU2008-02950-C03-01) (LMGS), University of Torino (G.C.P.), Fondazione Cavalieri Ottolenghi (Torino) and Fondazione San Paolo (Progetto Neuroscienze, PF-2009.1180 G.C.P.).

**6. Conflict of interest.** The authors declare no conflicts of interest.

## 7. REFERENCES

- Abraham IM, Todman MG, Korach KS, Herbison AE. 2004. Critical *in vivo* roles for classical estrogen receptors in rapid estrogen actions on intracellular signaling in mouse brain. *Endocrinology* 145: 3055-3061.
- Albeck DS, McKittrick CR, Blanchard DC, Blanchard RJ, Nikulina J, McEwen BS, Sakai RR. 1997. Chronic social stress alters levels of corticotropin-releasing factor and arginine vasopressin mRNA in rat brain. *J Neurosci* 17: 4895-4903.
- Albers HE. 2012. The regulation of social recognition, social communication and aggression: vasopressin in the social behavior neural network. *Horm Behav* 61: 283-292.
- Armstrong WE. 1985. Hypothalamic supraoptic and paraventricular nuclei. In: *The rat nervous system*, pp 119-128 Australia: Acad.Press.
- Barker JM, Galea LA. 2009. Sex and regional differences in estradiol content in the prefrontal cortex, amygdala and hippocampus of adult male and female rats. *Gen Comp Endocrinol* 164: 77-84.
- Benarroch EE. 2005. Paraventricular nucleus, stress response, and cardiovascular disease. *Clin Auton Res* 15: 254-263.
- Bolignano D, Zoccali C. 2010. Vasopressin beyond water: implications for renal diseases. *Curr Opin Nephrol Hypertens* 19: 499-504.
- Bosch OJ, Neumann ID. 2012. Both oxytocin and vasopressin are mediators of maternal care and aggression in rodents: from central release to sites of action. *Horm Behav* 61: 293-303.
- Bossmar T, Forsling M, Akerlund M. 1995. Circulating oxytocin and vasopressin is influenced by ovarian steroid replacement in women. *Acta Obstet Gynecol Scand* 74: 544-548.



- Burbach JP, Luckman SM, Murphy D, Gainer H. 2001. Gene regulation in the magnocellular hypothalamo-neurohypophysial system. *Physiol Rev* 81: 1197-1267.
- Caldwell HK, Lee HJ, Macbeth AH, Young WS, 3rd. 2008. Vasopressin: behavioral roles of an "original" neuropeptide. *Prog Neurobiol* 84: 1-24.
- Cameron JL. 1997. Stress and behaviorally induced reproductive dysfunction in primates. *Semin Reprod Endocrinol* 15: 37-45.
- Coccaro EF, Kavoussi RJ, Hauger RL, Cooper TB, Ferris CF. 1998. Cerebrospinal fluid vasopressin levels: correlates with aggression and serotonin function in personality-disordered subjects. *Archives of general psychiatry* 55: 708-714.
- De Vries GJ, Buijs RM, Sluiter AA. 1984. Gonadal hormone actions on the morphology of the vasopressinergic innervation of the adult rat brain. *Brain Res* 298: 141-145.
- Engelmann M, Landgraf R, Wotjak CT. 2004. The hypothalamic-neurohypophysial system regulates the hypothalamic-pituitary-adrenal axis under stress: an old concept revisited. *Front Neuroendocrinol* 25: 132-149.
- Ferguson AV, Latchford KJ, Samson WK. 2008. The paraventricular nucleus of the hypothalamus - a potential target for integrative treatment of autonomic dysfunction. *Expert Opin Ther Targets* 12: 717-727.
- Goldstein JM, Jerram M, Abbs B, Whitfield-Gabrieli S, Makris N. 2010. Sex differences in stress response circuitry activation dependent on female hormonal cycle. *J Neurosci* 30: 431-438.
- Grassi D, Amorim MA, Garcia-Segura LM, Panzica G. 2010. Estrogen receptor alpha is involved in the estrogenic regulation of arginine vasopressin immunoreactivity in the supraoptic and paraventricular nuclei of ovariectomized rats. *Neurosci Lett* 474: 135-139.

- Grassi D, Bellini MJ, Acaz-Fonseca E, Panzica G, Garcia-Segura LM. 2013. Estradiol and testosterone regulate arginine-vasopressin expression in SH-SY5Y human female neuroblastoma cells through estrogen receptors alpha and beta. *Endocrinology* 154:: 2092–2100.
- Guccione M, Silbiger S, Lei J, Neugarten J. 2002. Estradiol upregulates mesangial cell MMP-2 activity via the transcription factor AP-2. *Am J Physiol Renal Physiol* 282: F164-169.
- Handa RJ, Burgess LH, Kerr JE, O'Keefe JA. 1994. Gonadal steroid hormone receptors and sex differences in the hypothalamo-pituitary-adrenal axis. *Horm Behav* 28: 464-476.
- Hazell GG, Yao ST, Roper JA, Prossnitz ER, O'Carroll AM, Lolait SJ. 2009. Localisation of GPR30, a novel G protein-coupled oestrogen receptor, suggests multiple functions in rodent brain and peripheral tissues. *J Endocrinol* 202: 223-236.
- Insel TR. 2010. The challenge of translation in social neuroscience: a review of oxytocin, vasopressin, and affiliative behavior. *Neuron* 65: 768-779.
- Jin M, Jin F, Zhang L, Chen Z, Huang H. 2005. Two estrogen replacement therapies differentially regulate expression of estrogen receptors alpha and beta in the hippocampus and cortex of ovariectomized rat. *Mol Brain Res* 142: 107-114.
- Kastenberger I, Lutsch C, Schwarzer C. 2012. Activation of the G-protein-coupled receptor GPR30 induces anxiogenic effects in mice, similar to oestradiol. *Psychopharmacology (Berl)* 221: 527-535.
- Kudielka BM, Kirschbaum C. 2005. Sex differences in HPA axis responses to stress: a review. *Biol Psychol* 69: 113-132.

- Lagunas N, Calmarza-Font I, Diz-Chaves Y, Garcia-Segura LM. 2010. Long-term ovariectomy enhances anxiety and depressive-like behaviors in mice submitted to chronic unpredictable stress. *Horm Behav* 58: 786-791.
- Landgraf R. 2001. Neuropeptides and anxiety-related behavior. *Endocr J* 48: 517-533.
- Landgraf R, Kessler MS, Bunck M, Murgatroyd C, Spengler D, Zimbelmann M, Nussbaumer M, Czibere L, Turck CW, Singewald N, Rujescu D, Frank E. 2007. Candidate genes of anxiety-related behavior in HAB/LAB rats and mice: focus on vasopressin and glyoxalase-I. *Neurosci Biobehav Rev* 31: 89-102.
- Liu J, Hu P, Qi X, Meng F, Kalsbeek A, Zhou J. 2011. Acute restraint stress increases intrahypothalamic estradiol concentrations in conjunction with increased hypothalamic oestrogen receptor beta and aromatase mRNA expression in female rats. *J Neuroendocrinol* 23: 435-443.
- Lund TD, Hinds LR, Handa RJ. 2006. The androgen 5alpha-dihydrotestosterone and its metabolite 5alpha-androstan-3beta, 17beta-diol inhibit the hypothalamo-pituitary-adrenal response to stress by acting through estrogen receptor beta-expressing neurons in the hypothalamus. *J Neurosci* 26: 1448-1456.
- Merchenthaler I, Lane MV, Numan S, Dellovade TL. 2004. Distribution of estrogen receptor alpha and beta in the mouse central nervous system: in vivo autoradiographic and immunocytochemical analyses. *J Comp Neurol* 473: 270-291.
- Mineur YS, Belzung C, Crusio WE. 2006. Effects of unpredictable chronic mild stress on anxiety and depression-like behavior in mice. *Behav Brain Res* 175: 43-50.
- Mitra SW, Hoskin E, Yudkovitz J, Pear L, Wilkinson HA, Hayashi S, Pfaff DW, Ogawa S, Rohrer SP, Schaeffer JM, McEwen BS, Alves SE. 2003. Immunolocalization of estrogen receptor beta in the mouse brain: comparison with estrogen receptor alpha. *Endocrinology* 144: 2055-2067.

- Moeller HB, Fenton RA. 2012. Cell biology of vasopressin-regulated aquaporin-2 trafficking. *Pflugers Archiv* 464: 133-144.
- Moreau JL, Jenck F, Martin JR, Mortas P, Haefely WE. 1992. Antidepressant treatment prevents chronic unpredictable mild stress-induced anhedonia as assessed by ventral tegmentum self-stimulation behavior in rats. *Eur Neuropsychopharmacol* 2: 43-49.
- Park SY, Walker JJ, Johnson NW, Zhao Z, Lightman SL, Spiga F. 2013. Constant light disrupts the circadian rhythm of steroidogenic proteins in the rat adrenal gland. *Mol Cell Endocrinol* 371: 114-123.
- Patchev VK, Hayashi S, Orikasa C, Almeida OFX. 1995. Implications of Estrogen-Dependent Brain Organization for Gender Differences in Hypothalamo-Pituitary-Adrenal Regulation. *FASEB J* 9: 419-423.
- Paxinos G, Franklin KBJ. 2001. *The Mouse Brain in Stereotaxic Coordinates*. San Diego: Academic Press.
- Pinnock SB, Herbert J. 2001. Corticosterone differentially modulates expression of corticotropin releasing factor and arginine vasopressin mRNA in the hypothalamic paraventricular nucleus following either acute or repeated restraint stress. *Eur J Neurosci* 13: 576-584.
- Plumari L, Viglietti Panzica C, Allieri F, Honda S, Harada N, Absil P, Balthazar J, Panzica GC. 2002. Changes in the Arginine-Vasopressin Immunoreactive Systems in Male Mice Lacking a Functional Aromatase Gene. *J Neuroendocrinol* 14: 971-978.
- Pyner S. 2009. Neurochemistry of the paraventricular nucleus of the hypothalamus: implications for cardiovascular regulation. *J Chem Neuroanat* 38: 197-208.

- Rotzinger S, Lovejoy DA, Tan LA. 2010. Behavioral effects of neuropeptides in rodent models of depression and anxiety. *Peptides* 31: 736-756.
- Sawchenko PE, Swanson LW. 1982. Immunohistochemical identification of neurons in the paraventricular nucleus of the hypothalamus that project to the medulla or to the spinal cord in the rat. *J Comp Neurol* 205: 260-272.
- Seale JV, Wood SA, Atkinson HC, Bate E, Lightman SL, Ingram CD, Jessop DS, Harbuz MS. 2004. Gonadectomy reverses the sexually diergic patterns of circadian and stress-induced hypothalamic-pituitary-adrenal axis activity in male and female rats. *J Neuroendocrinol* 16: 516-524.
- Siegel A, Bhatt S, Bhatt R, Zalcman SS. 2007. The neurobiological bases for development of pharmacological treatments of aggressive disorders. *Curr Neuropharmacol*. 5: 135-147.
- Silverman AJ, Pickard GE. 1983. The hypothalamus. In: *Chemical neuroanatomy* (Emson, PC, ed), New York: Raven Press. pp 295-336.
- Surget A, Belzung C. 2008. Involvement of vasopressin in affective disorders. *Eur J Pharmacol* 583: 340-349.
- Vega-Rivera NM, Fernandez-Guasti A, Ramirez-Rodriguez G, Estrada-Camarena E. 2013. Acute stress further decreases the effect of ovariectomy on immobility behavior and hippocampal cell survival in rats. *Psychoneuroendocrinology* 38: 1407-1417.
- Viau V. 2002. Functional cross-talk between the hypothalamic-pituitary-gonadal and -adrenal axes. *J Neuroendocrinol* 14: 506-513.
- Wang H, Ward AR, Morris JF. 1995. Oestradiol acutely stimulates exocytosis of oxytocin and vasopressin from dendrites and somata of hypothalamic magnocellular neurons. *Neuroscience* 68: 1179-1188.

Watson RE, Wiegand SJ, Clough RW, Hoffman GE. 1986. Use of cryoprotectant to maintain long-term peptide immunoreactivity and tissue morphology. *Peptides* 7: 155-159.

Weibel ER. 1979. *Stereological methods*. Vol. 1. Practical methods for biological morphometry. London: Academic Press.

Weiser MJ, Handa RJ. 2009. Estrogen impairs glucocorticoid dependent negative feedback on the hypothalamic-pituitary-adrenal axis via estrogen receptor alpha within the hypothalamus. *Neuroscience* 159: 883-895.

Accepted manuscript

**FIGURES' CAPTIONS**

**Fig.1.** Corticosterone levels in serum (ng/mL±S.E.) of intact mice (Control), ovariectomized mice (OVX), intact mice submitted to stress (CUS) and ovariectomized mice submitted to stress (OVX+CUS). Different letters denote significant differences ( $p<0.05$ ) between pairs.

**Fig.2.** Representative microphotographs of AVP immunostaining within paraventricular nucleus (PVN) in control mice (control), ovariectomized mice (OVX), control mice submitted to stress (CUS) and ovariectomized mice submitted to stress (OVX+CUS). PaDC - Paraventricular Dorsal Cap; PaLM - Paraventricular Lateral Magnocellular subdivision; PaMM - Paraventricular Medial Magnocellular subdivision; PaMP - Paraventricular Medial Parvocellular subdivision. Scale bar, 100  $\mu\text{m}$ . \* 3<sup>rd</sup> Ventricle.

**Fig.3.** Volume fraction of Paraventricular Dorsal Cap (PaDC), Paraventricular Lateral Magnocellular (PaLM), Paraventricular Medial Magnocellular (PaMM) and Paraventricular Medial Parvocellular (PaMP) subdivisions occupied by AVP immunoreactive material in A) ovariectomized animals (OVX), B) control animals submitted to stress (CUS) and C) ovariectomized animals submitted to stress (OVX+CUS). The results are normalized to control animals' value equal 100. Different letters denote significant differences ( $p<0.05$ ) between pairs.

	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
Day		Continuous room and cage changes (8 h once per hour) 9AM-5PM	Introducing in a cage with rat sawdust (1h) 12PM-1PM				Lights on 9AM-7PM
Night	Social isolation 7PM-9AM	Animals were paired with an unfamiliar mouse 7PM-9AM	Predator proximity: introducing a rata into the cage (2h) 7PM-9PM	Damp sawdust 7PM-9AM	Cage tilting (45°) 7PM-9AM	Lights on 7PM-9AM	Lights on 7PM-9AM


**Table 1.** Schematic representation of chronic unpredictable mild stress (CUS) protocol. Stress protocol was repeated continuously during 4 weeks (Lagunas et al., 2010)





**Highlights**



- Long-term ovariectomy induces a decrease of AVP-ir in some subnuclei of the PVN
- The CUS significantly affects AVP-ir not in the same direction of ovariectomy
- Magnocellular subdivisions of the PVN are significantly affected by CUS
- Long-term ovariectomy and CUS induce changes in AVP-ir with regional specificity

Accepted manuscript









**Ovariectomy and Chronic Unpredictable Stress differentially affect expression of arginine vasopressin in the mouse PVN**