

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Italian Nurses' Experience of Stalking: A Questionnaire Survey

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/154359> since 2017-02-01T10:31:35Z

Published version:

DOI:10.1891/0886-6708.VV-D-12-00078

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Daniela Acquadro Maran; Antonella Varetto; Massimo Zedda. Italian Nurses' Experience of Stalking: A Questionnaire Survey. *VIOLENCE AND VICTIMS*. 29 pp: 109-121.

DOI: 10.1891/0886-6708.VV-D-12-00078

The publisher's version is available at:

<http://www.ingentaconnect.com/content/springer/vav/2014/00000029/00000001/art00007>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/154359>

Violence and Victims

Italian nurses' experience of stalking: a questionnaire survey

--Manuscript Draft--

Manuscript Number:	VV-D-12-00078
Full Title:	Italian nurses' experience of stalking: a questionnaire survey
Article Type:	Original Research
Keywords:	organizational and individual prevention; patient-nurses relationship; stalking; workplace
Corresponding Author:	daniela acquadro maran, ph.d. univeristà degli studi di torino torino, ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	univeristà degli studi di torino
Corresponding Author's Secondary Institution:	
First Author:	daniela acquadro maran, ph.d.
First Author Secondary Information:	
Order of Authors:	daniela acquadro maran, ph.d. antonella varetto massimo zedda
Order of Authors Secondary Information:	
Abstract:	<p>Background: Stalking is a phenomenon characterized by a set of repetitive behaviors, intrusive surveillance and control, communication and search of contact with a victim who is afraid and/or worried and/or annoyed by such unwanted attention. Literature analysis show that nurses are at greater risk of being stalked than the general population, both by ex-partners and acquaintances (such as colleagues and patients) and by unknown stalkers, who may have different motives: relational rejection, an infatuation, an inability to express their own emotions and recognize those of others, a desire for revenge. Objective: the aim of this study was to explore Italian nurses' experience of stalking (incidence, relationship victim-stalker, motive, impact, coping strategy). Design: a copy of the Italian modified version of the Sheridan Questionnaire on Stalking, BDI and Stai Y1-Y2 scales were distributed in four Italian state hospitals after obtaining the permission of the respective institutional managers who had previously signed a formal letter of agreement. Settings: Italian state hospitals located in the North of the Country that agree to participate in the study. Participants: a sample of 2,000 nurses working in Italian state hospitals. Results: a total of 765 nurses returned the questionnaire, 107 had been victims (13.99%). Most part of victims are female stalked by older males (Acquaintance 41.12%), whose behavior included unwanted communications, following, control, assault and threats. With regard to stalkers' motive, the majority was Rejected (32.71%; Incompetent suitors 26.17%, Resentful 22.43% and Intimacy-Seekers 13.08%). The stalking campaign cause in victims both emotional than physical consequences, but level of depression are not higher as expected. The most used coping strategies involve personal protection (offensive tactics) and confronting stalkers, to explain the disruption they are causing (interactional tactics). Conclusion: the incidence of the phenomenon among nurses require to intervene adopting legal, organizational and individual strategies. Hospital managers are seriously committed to addressing legal issues and to organize educational programs. Individual strategies include to refuse to treat certain patients, guarding personal data and not disclosing personal (or other) details to patients personal data and not disclosing personal (or other) details to patients.</p>
Suggested Reviewers:	eloisa Guerrero Barona

	eloisa@unex.es she is an expert of nurses' workplace issues
Author Comments:	

Introduction

1
2
3
4
5
6 Since the eighties social science scholars have been conducting research in an
7
8 attempt to describe and analyze stalking from the legal, social, psychological and
9
10 psychiatric viewpoints. Some authors have described stalking as a form of intimate
11
12 violence (Coleman, 1997; Douglas & Dutton, 2001; Kurt, 1995), others as a
13
14 constellation of behaviors in which a person (stalker) inflicts on another (stalking
15
16 victim) repeated unwanted intrusions and/or communications (Pathè & Mullen, 1997).
17
18 Others, such as Meloy (1998), have defined it as an ongoing course of conduct in which
19
20 a person behaviorally intrudes upon another's life in a manner perceived to be
21
22 threatening. A more comprehensive definition was proposed by Westrup(1998:276),
23
24 stalking is defined as one or more of a constellation of behaviors that "(a) are directed
25
26 repeatedly toward a specific individual (the 'target'), (b) are experienced by the target as
27
28 unwelcome and intrusive, and (c) are reported to trigger fear or concern in the target". It
29
30 is the repetition/the persistent nature of such behaviors that makes the difference
31
32 compared to other forms of violence in the workplace and/or in private life (Sheridan,
33
34 Gillet & Davies, 2000).

35
36 Starting from the nineties, some large-scale studies of stalking have been
37
38 conducted in several countries (USA, Australia, Great Britain, Germany, Austria, Italy).
39
40 The data gathered during these studies differs considerably, depending on the criteria
41
42 used to define the phenomenon (e.g. a certain number of episodes of intrusive behavior
43
44 occurring over a period of more than one month, or the inclusion/exclusion of unwanted
45
46 attention and harassment), the method used (e.g. telephone interviews or questionnaires)
47
48 and the subjects involved (females only or males and females). Despite these
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 limitations, in their analysis of 18 studies on ordinary people, Spitzberg and Cupach
2
3 (2003) reported an average incidence of 13.9%: they found victims to be mostly
4
5 women, aged 18-29 years, stalked by a man, whose behavior consisted of surveillance
6
7 and communications containing threats of injury, in an effort to establish or re-establish
8
9 a relationship (Tjaden & Thonnes, 1998). First, these data confirmed that stalking is an
10
11 issue that could affect everybody, not only celebrities (West & Friedman, 2008) and
12
13 second, they underlined the importance of analyzing and describing the nature and
14
15 evolution of the stalker's behavior in order to recognize the potential risk for the victim
16
17 (author, year). Using preliminary data gathered from studies on the general population,
18
19 some researchers have attempted to analyze the motivations for stalking, the
20
21 relationship between victim and perpetrator, the type of stalking behavior and the
22
23 coping strategies adopted by the victims, the physical and psychological consequences
24
25 for victims, victimization among particular professional groups.
26
27
28
29
30
31

32 *Motivations for stalking*

34
35 Studies aimed at understanding stalkers' motives have described stalking as an
36
37 ongoing process of disjunctive relationship development in which the pursuer endeavors
38
39 to re/establish some form of relationship which the object of pursuit refuses (Emerson *et*
40
41 *al.*, 1998). Mullen *et al* (1999) noted that stalkers have several different motivations,
42
43 such as pursuing a vendetta because of a real or perceived sense of grievance. The
44
45 authors proposed a classification that identifies five types of stalkers, one of the most
46
47 cited in the literature and which is used to evaluate the escalation of aggressive behavior
48
49 (see MacKenzie *et al*, 2008). According to this classification, stalkers may be Rejected
50
51 stalkers (pursue their victims to avoid or avenge a rejection, e.g. divorce or separation),
52
53 Intimacy seekers (seek to establish an intimate or loving relationship with their victims),
54
55
56
57
58
59
60
61
62
63
64
65

1 Incompetent suitors (have a fixation, a sort of entitlement to an intimate relationship
2 with their victim who is the person that has attracted their amorous interest), Resentful
3 stalkers (seek revenge due to a sense of grievance against the victim, perceived as the
4 cause of damage suffered by them or another person), Predatory stalkers (follow and
5 spy on the victim in order to prepare and plan an attack, often sexual).
6
7
8
9
10
11
12

13 The first study to investigate stalker typology involved a sample of 145 subjects
14 (79.31% were male, age range 15-75 years, median 38 years) who had been referred to
15 an Australian forensic psychiatry center for treatment (Mullen *et al*, 1999). The data
16 collected in this study showed that 35.86% of the sample were Rejected stalkers. They
17 had started stalking after being rejected in a personal relationship (by a partner, friend,
18 relative) or work relationship (e.g. dismissal). 33.79% of the sample were Intimacy
19 seekers, for whom the aim of stalking was to seek to establish a romantic relationship
20 with their 'true love'. 15.17% were Incompetent suitors who stalked victims who
21 attracted their romantic interest. Resentful stalkers, who represented 11.04% of the
22 sample, sought a victim to blame for their professional or affective distress. Predatory
23 stalkers accounted for 4.14% of the sample.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 *Relationship between victim and perpetrator*

41
42 In the literature, the relationship between victims and perpetrators has been
43 categorized according to the degree of intimacy: there is stalking by ex-intimates, (e.g.
44 partner, fiancé), acquaintances (neighbors, friends, colleagues, clients, suppliers) and
45 strangers (people with whom the victim has never had any previous contact) (see Zona,
46 Sharma e Lane, 1993; Mullen *et al*, 1999; Kordvani, 2000). This classification has been
47 used in several studies. For example, a Canadian research project conducted by Kong
48 (1996) found that in a sample of 5,023 reported cases of stalking, 53.48% of stalkers
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 were ex-intimates (ex-partners 47.26%, relatives 4.66%, partners 1.56%), 32.81% were
2
3 acquaintances (4.9% were work colleagues), 8.12% were strangers, while 5.59% of the
4
5 victims had no idea who was stalking them. These results have been confirmed by other
6
7 studies, such as that by Sheridan and Davies (2001) at the Suzy Lamplugh Trust (Great
8
9 Britain) involving 95 victims of stalking: 91.58% were women, 49.48% of them had had
10
11 a previous intimate relationship with the stalker (ex-partner), while 37.89% of them
12
13 were stalked by someone they knew and 12.63% by a stranger. Hall (1998) conducted a
14
15 study involving 145 self-declared victims (men 17.24%, women 82.76%) which
16
17 revealed significant differences in terms of gender and relationship with the stalker: 64%
18
19 of men were stalked by an acquaintance, 28% by an ex-intimate, 4% by a stranger, and
20
21 4% had no idea who was stalking them. Among women, 62.5% were stalked by an ex-
22
23 partner, 27.5% by an acquaintance, 6.67% by a stranger and 3.33% had no idea who was
24
25 stalking them. The results revealed that men are more frequently stalked by an
26
27 acquaintance than women, while women are more often victims of stalking by an
28
29 intimate partner than men. Similar findings were also reported by Palarea and colleagues
30
31 (1999). They analyzed 223 cases reported between 1990 and 1996 to the Los Angeles
32
33 Threat Management Unit: 53.19% of the 47 male victims and 62.5% of the 176 female
34
35 victims had been in an intimate relationship with the stalker.

36 37 38 39 40 41 42 43 44 45 *Stalking behaviors and victims' coping strategies*

46
47 Stalking behavior is characterized by repeated intrusions involving unwanted
48
49 contacts (such as following, spying on and approaching the victim) and unwanted
50
51 communications (including telephone calls, e-mails, letters) (Meloy, 2000). Victims
52
53 adopt different coping strategies in relation to the type of stalking behavior. Relatives,
54
55 children and friends may also be harmed by stalking (Sheridan and Davies, 2001), and
56
57
58
59
60
61
62
63
64
65

1 coping strategies can be more or less effective. In an interesting study Nicaastro, Cousins
2 and Spitzberg (2000) analyzed 55 cases of stalking and domestic violence filed with the
3 Domestic Violence Unit in San Diego between 1992 and 1997. Most of the victims were
4 females (92.73%) stalked by males (90.91%), and the case files included information
5 such as police reports, statements by witnesses and the victims, pictures and even some
6 statements by the suspects themselves. All the information was coded. The study
7 revealed the use of several tactics by stalkers: pursuit (e.g. calls, following the victim,
8 messages/letters/notes), threats (e.g. verbal and/or physical, offensive language),
9 violence (e.g. harming with weapons, sexual coercion), violation of privacy (e.g.
10 trespassing). Victims responded to such behaviors by adopting various coping strategies,
11 categorized as below: interactional tactics (e.g. threatening to harm perpetrators, telling
12 perpetrators what they are doing is wrong), offensive avoidance tactics (e.g. calling the
13 police, obtaining a restraining order), defensive avoidance tactics (e.g. ignoring the
14 stalker, changing their phone number), therapy (e.g. seeking professional advice). The
15 findings revealed that stalkers used a variety of tactics, mostly calls (70.91%), verbal
16 threats (67.27%), harassment (61.82%), violation of private property (54.55%),
17 following (50.91%), offensive language (50.91%), vandalism (47.27%), physically
18 hurting (45.45%), physically restraining (38.18%) and driving by work/home/school
19 (36.36%). The coping strategies adopted were equally varied, victims required police
20 intervention (96.36%) and/or threatened to call the police (14.55%), had a physical
21 confrontation (18.18%), hung up when the stalker called (16.36%), asked a friend and/or
22 relative to offer protection (14.55%) and screened phone calls (14.55%). Further studies
23 have confirmed the findings of Nicaastro and colleagues: stalking activities usually
24 involve following around, trying to control, sending flowers and gifts, communicating

1 through letters and e-mails, searching for information about the victim, disseminating
2
3 false and defamatory information, sexual and physical aggression, damaging private
4
5 property, breaking and entering, theft (Spitzberg, 2002). While various studies have
6
7 reported different coping strategies adopted by victims, several found that victims prefer
8
9 to seek the help of friends and relatives rather than law enforcement. These findings
10
11 emerged from an investigation by Buhi, Clayton and Surrency (2009) involving 391
12
13 college women. 19.95% of them were victims of stalking, and more than half (52.56%)
14
15 sought the help of a friend (90.24%), a relative (29.27%), a hall of residence supervisor
16
17 or the police (19.51%). Victims do not usually call the police and would rather minimize
18
19 the problem, ignore the stalker, try to get away from him/her, limit access to information
20
21 (e.g. by changing their e-mail address, using a nickname, blocking e-mails from the
22
23 stalker's address) (Amar & Alexis, 2010) or block access to private data via social
24
25 networks (Kennedy & Taylor, 2010).
26
27
28
29
30
31

32 *Physical and psychological consequences for the victim*

34
35 While the duration of stalking has been found to vary from a few weeks to
36
37 several years, Spitzberg (2002) calculated that the average duration of stalking is
38
39 approximately two years. Stalking has a deep impact on the victim, with physical,
40
41 emotional and psychological consequences. Some authors, such as Kamphuis and
42
43 Emmelkamp (2001), reported that stalking victims show trauma symptoms similar to
44
45 those of the victims of road accidents, people during periods of mourning and those who
46
47 have been repeatedly caught up in bank robberies. Brewster (1998) interviewed 187
48
49 women who were self-declared victims of stalking, 45.97% of whom reported being
50
51 subjected to violence during stalking. 66.28% had been involved one or two accidents,
52
53 33.72% in three or more. The most frequent forms of violence were: punching in the
54
55
56
57
58
59
60
61
62
63
64
65

1 face (59.3%), serious cuts requiring medication (39.53%), nose-bleeds (26.74%),
2
3 internal injuries (13.95%) and bruising (10.47%). The most common reactions were:
4
5 fear (57.22%), anger (23.53%), insomnia (12.83%), frustration (12.3%), depression
6
7 (11.23%), mistrust and sense of insecurity (11.23%) and irritation/anxiety (10.16%).
8
9
10 Victims also said that during stalking they were particularly suspicious (44.38%), scared
11
12 (41.71%) and nervous (31.02%). They also experienced other changes that affected their
13
14 quality of life: 93.58% constantly looked away, 71.67% locked the door (something they
15
16 had not done before), 63.64% took a different route to go to work and avoided certain
17
18 places, 58.82% left the house lights turned on or off, 40.64% carried pepper spray in
19
20 their pocket, 23.53% never left the house on their own and 9.63% took a dog for
21
22 protection. Some victims reported having sustained costs as a result of being stalked.
23
24
25 Such costs were related to attempts to discourage the stalker, purchase security
26
27 measures, legal expenses, lower income (2.14% lost their job), professional help and
28
29 repairs for damage to property. Research conducted in other countries has confirmed
30
31 these results. For example, a study in Pakistan involving 100 female victims of stalking
32
33 (Haider & Haider, 2002) reported that 82 victims suffered from deterioration of mental
34
35 health (70 reported poorer job performance) and 47 victims received a diagnosis of
36
37 PTSD (DSM-IV classification). More than half of the victims (55) suffered from
38
39 symptoms such as flashback and intrusive memories. Physical symptoms included
40
41 chronic sleep disorder (74), headache (55), appetite disorder (48), fatigue (47), weight
42
43 changes (45), sickness and digestive disorders (30), heart palpitations (25).
44
45
46
47
48
49
50
51

52 *Victimization among particular professional groups*

53

54 In an attempt to understand the context in which the phenomena occurred and
55
56 identify any correlations – in addition to gender – including social and professional
57
58
59
60
61
62
63
64
65

1 status, some researchers have investigated whether certain groups are more at risk of
2
3 victimization.
4

5
6 From an analysis of the literature it has been found that health care professionals
7
8 (doctors, nurses, therapists, social workers, psychologists, psychiatrists and so on) are
9
10 more at risk than other professionals (Galeazzi, Elkins & Curci, 2005; Jones &
11
12 Sheridan, 2009; Purcell *et al*, 2005). The reasons for this higher incidence of
13
14 victimization were investigated by Pathè, Mullen and Purcell (2002), who found a
15
16 correlation with the nature of the health care profession, characterized by contact with
17
18 people who are lonely and/or suffer from some disorder: some people may misinterpret
19
20 care and attention as romantic interest, which can provoke aggressive reactions,
21
22 intrusive behavior and harassment (Laskowsky, 2003). Above all their findings showed
23
24 nurses to be exposed to the greatest risk of victimization (see Magnavita & Heponiemi,
25
26 2011; Hahn *et al*, 2012): this appears to be due to the fact that they provide constant and
27
28 daily care to patients. The perception is that nurses are more caring and pay more
29
30 attention to relationships, including intimate relationships, and if this expectation is not
31
32 fulfilled it can trigger resent in partners, friends, relatives and acquaintances. The risk of
33
34 victimization in nurses was confirmed by research carried out in Great Britain by
35
36 Ashmore, Jones, Jackson and Smoyak (2006). The study involved 112 mental health
37
38 nurses, 50% of respondents had been stalked. Most were females (78.57%) who had
39
40 been stalked for less than 12 months by a male stalker (82.14%) aged 17-60 years. In
41
42 some cases (31.43%) the stalker was a patient. Stalking behaviors included unwanted
43
44 communications (calls, letters) and unwanted gifts (41.07%). In 26.79% of cases the
45
46 communications contained threats (21.43% had suffered physical aggression), and/or
47
48 sexual threats (33.93%, with 1.79% of victims sexually assaulted), and/or self-harm
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 threats (19.64%) and/or threats to third parties (19.64%).
2

3 Other stalking behaviors included following around (42.86%), property damage
4 (12.5%), theft (7.14%), trespassing (.79%) and defamation (1.79%). Physical and
5 emotional reactions were similar to those reported in other research, primarily anxiety
6 (46.43%), irritation (37.5%), anger (37.5%), stress (30.36%) and sleep disorders (25%).
7
8 Victims adopted interactional coping strategies, for instance, 50% attempted to reason
9 with the stalker, defensive avoidance tactics (37.5% ignored the stalker), offensive
10 avoidance tactics (12.5% threatened the stalker) and therapy (28.57% sought clinical
11 supervision). Some victims sought help by telling a close friend (46.42%) or a family
12 member (37.5%) about what was happening, while 23.21% called the police. Kaplan
13 (2006) reported that victims took more precautions at work, sought help, shared their
14 experience with a colleague and told a supervisor. Authors such as Sandberg, McNiel
15 and Binder (1998) suggested that under-reporting of episodes of stalking to colleagues,
16 supervisors and the police (including those within the hospital) is related to denial and
17 minimization: it is assumed that intrusions are correlated with the nursing profession,
18 which involves dealing with people with physical, emotional, relational and mental
19 disorders (Smoyak, 2003). The denial strategy might help victims to ignore the situation
20 – which they see as potentially dangerous – and continue with their care work (Brown,
21 Dubin, Lion and Garry, 1996). Another aspect that could influence under-reporting is the
22 possibility of being accused of incompetence, victimization could arouse skepticism
23 among colleagues and supervisors about the relational ability to intervene (McEwan,
24 Mullen and Purcell, 2007).
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 Research involving a sample of nurses in Italy

57 *Aim*

58
59
60
61
62
63
64
65

1 The aim of this study was to explore Italian nurses' experience of stalking.
2

3 Specifically, this study sought answers to the following research questions:
4

- 5 - What is the incidence of stalking among a sample of nurses in Italy?
6
- 7 - What is the prior relationship between victims-stalkers, the motive for
8 stalking?
9
- 10 - What impact does stalking have on Italian nurses, does it have consequences
11 on the quality of life?
12
- 13 - How do Italian nurses cope with being stalked?
14
15
16
17
18
19

20 Method 21

22 In this study we used the modified Italian version of the Sheridan Questionnaire
23 on Stalking (Sheridan & Davies, 2001), to investigate the incidence, relationship
24 between stalkers and victims, motivations and consequences for victims. The
25 questionnaire was adapted for use with an Italian audience, by translating it from British
26 English and then 'back translating' it (White & Elander, 1992), and modified in some
27 parts to adapt it to the research questions. Finally, it was piloted using a sample of 35
28 trainee nurses before use in this study. This process did not result in any further revision
29 of the instrument. The modified questionnaire covered such issues as: demographic
30 details of victims and stalkers, the nature of their relationship, the perceived motivation
31 for stalking, the duration and frequency of stalking and stalking behavior, the reactions
32 of victims and others, physical and emotional consequences. Moreover, as suggested in
33 the literature, we used the Beck Depression Inventory (BDI, Beck & Ward, 1961; Italian
34 version by Scillico, 1978) and the State Trait Inventory (STAI, (Spielberger, 1983,
35 Italian version by Pedrabissi and Santinello, 1989) to investigate the psychological signs
36 in victims and their exasperation during/after stalking. The BDI is a 21-question survey,
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 in which each answer is scored by assigning a value of 0 to 3, designed to determine the
2
3 presence of depression. A total score of 0-13 indicates minimal depression, 14-19
4
5 indicates mild depression, 20-28 moderate depression, and 29-63 severe depression. The
6
7 STAI consists of two forms (Y1 and Y2), each comprising 20 items rated on a 4-point
8
9 scale. It is used for assessing anxiety levels. Form Y1 is used to assess a transitory
10
11 emotional state that varies in intensity, fluctuates over time, and is characterized by
12
13 feelings of tension and apprehension and by heightened activity of the autonomic
14
15 nervous system. It evaluates how stalking victims feel “right now”, at this moment.
16
17 Form Y2 evaluates trait anxiety, a relatively stable predisposition of individuals to
18
19 respond to situations perceived as threatening. This scale evaluates how stalking victims
20
21 feel most of the time. The scores for each of the forms range from 20 to 80, with a value
22
23 of 40 indicating the presence of anxiety symptoms (from 40 to 50: mild anxiety). Scores
24
25 of 50 to 60 indicate moderate anxiety and those of more than 60 indicate severe anxiety
26
27 symptoms. All questionnaires were self-administered.
28
29
30
31
32
33

34 *Sample and procedure*

35
36
37 The research project were send to Italian state hospitals located in the North of
38
39 the Country. Four hospitals agree to participate in the study: the questionnaires were
40
41 distributed to a suitable sample of 2,000 nurses after obtaining the permission of the
42
43 respective institutional managers who had previously signed a formal letter of
44
45 agreement.
46
47

48
49 Full disclosure of the purpose of the investigation was provided in the letter
50
51 accompanying the questionnaire, with a definition of the phenomenon, a synthesis of
52
53 those found in literature as purposed by Galeazzi and Curci (2001): set of repetitive
54
55 behaviors, intrusive surveillance and control, communication and search of contact with
56
57
58
59
60
61
62
63
64
65

1 a victim who is afraid and/or worried and/or annoyed by such unwanted attention. The
 2
 3 accompanying letter also included a privacy statement (with the indication that the
 4
 5 questionnaire was anonymous and the data would be used for research purposes only,
 6
 7 thus treated as confidential), instructions about how to fill-in and return the
 8
 9 questionnaire. The completed questionnaires were placed in a specific box that had been
 10
 11 specially designed to guarantee privacy and anonymity and were collected 15 working
 12
 13 days later. Data were processed using SPSS version 18 to produce mainly descriptive
 14
 15 statistics.
 16
 17
 18
 19

20 Results

21 *Sample and stalkers' characteristics*

22
 23 A total of 765 (38.35%) nurses returned the questionnaire. The majority of
 24
 25 respondents were female (68.89%), aged from 19 to 65 years (mean=40.13, s.d.=10.57);
 26
 27 11.50% were trainees. 107 had been victims of stalking (13.99%), 90 females (84.11%)
 28
 29 and 17 (15.89%) males, aged from 19 to 60 years (mean=37.80, s.d.=10.62). Victims
 30
 31 represented 17.08% of all female respondents and 7.26% of all male respondents.
 32
 33 Before being stalked the nurses had no previous knowledge of the phenomenon
 34
 35 (57.01%).
 36
 37
 38
 39
 40
 41

42 *Relationship between victims and stalkers and motivations for stalking*

43
 44 Stalkers were classified on the basis of their motives (as suggested by Mullen *et*
 45
 46 *al*, 1999: missing item MI=6) and relationship (see table 1).
 47
 48

49 - Table 1 approximately here -

50
 51 None were classified as Predatory stalkers. The majority of the victims knew
 52
 53 their stalker (67.29%), who was generally male (75.7%; MI=7). 17 stalkers stalked
 54
 55 victims of the same sex, 8.64% of male stalkers stalked other men and 52.63% of female
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

1 stalkers stalked other women (MI= 7). Within the acquaintance category 38.64% were
2
3 colleagues, 31.82% friends and 29.55% others (including patients and a chaplain). At
4
5 the time they filled-in the questionnaire, most of the respondents were no longer victims
6
7 (77.57%), while 9.57% were still victims and 13.08% did not know whether they were
8
9 still victims or not. The duration of stalking ranged from 4 to 572 weeks, with a mean
10
11 duration of 67 weeks. The frequency of contact was one or more times a day for most of
12
13 the victims (61.36%). For the 83 subjects who were no longer victims, stalking had
14
15 simply ended (32.53%) or had ended as the result of intimidation by the victim, a
16
17 relative or friend (27.71%), delivery of a police warning (16.87%), the stalker had found
18
19 a new victim (7.23%) or other type of intervention (15.66%, including one case in which
20
21 the stalker died). Stalkers engaged in various stalking activities, mainly unwanted
22
23 communications (in particular by former partners and Rejected stalkers), following and
24
25 control (such as visiting and waiting outside the home and/or workplace, forms of
26
27 behavior used on average by all types of stalkers; following was more frequent among
28
29 Intimacy seekers), assault and threats of assault (this type of behavior was more frequent
30
31 among former partners and Rejected stalkers than others - see table 2).
32
33
34
35
36
37
38
39

40 - Table 2 approximately here -
41

42 Our findings revealed that both unknown and Resentful stalkers (e.g. patients)
43
44 had access to nurses' personal data and could communicate with their victims by
45
46 telephone, text messages, letters and by waiting outside their home. Furthermore these
47
48 types of stalkers were more able than others to find information at the workplace
49
50 (54.17%) and to switch services on/off (such as telephone; 12.5%). Another alarming
51
52 finding regarded threat and assault, including harassment of others (colleagues,
53
54 relatives, friends, current partner, children). This type of aggressive behavior
55
56
57
58
59
60
61
62
63
64
65

1 characterized relationships with former partners (80%), 28.57% of whom suffered
 2
 3 psychological violence (e.g. jealousy) and 32.14% both psychological and physical
 4
 5 violence (e.g. battering,). The ANOVA analysis (Tukey's HSD Post-hoc test) showed
 6
 7 that items related to fear, confusion and increasing distrust ($p < 0.05$) were significantly
 8
 9 different for Ex-partners than for Acquaintances and Unknown stalkers: victims who
 10
 11 had had a previous relationship with their stalker are more exposed than others, since the
 12
 13 stalker knows their routine, friends and relatives. In line with this data, our findings
 14
 15 revealed that Ex-partner stalkers generally involved other individuals (82.14%; e.g.
 16
 17 family members, colleagues, doorkeepers).
 18
 19
 20
 21

22 *Impact on victims' lives*

23
 24
 25 Stalking has both physical and emotional consequences for victims, as shown in
 26
 27 table 3. On average, emotional consequences such as anxiety, anger, confusion and fear
 28
 29 are predominant, while physical consequences include sleep disruption, weight and
 30
 31 appetite disturbances, headaches, tiredness and panic attacks. One interesting finding
 32
 33 was the increasing distrust observed, in particular, among victims of Ex-partners,
 34
 35 Rejected and Resentful stalkers.
 36
 37
 38

39
 40 No victims reached cut-off on the BDI, while anxiety levels were, on average,
 41
 42 mild (< 40), particularly on STAI form Y2 determining how victims felt over time.
 43
 44

45 - Table 3 approximately here -
 46

47
 48 Multivariate ANOVA was used to determine differences in gender, depression
 49
 50 and anxiety symptoms of victims and stalking behavior. The results revealed no
 51
 52 significant associations between scores on the BDI scale for gender or behavior.
 53
 54

55 However the interaction between anxiety and gender was significant (Wilks' Lambda
 56
 57 $= 0.89$, $F(3,17) = 0.11$ $p < 0.05$). This interaction was subsequently analyzed considering
 58
 59
 60
 61
 62
 63
 64
 65

1 the STAI scores (forms Y1 and Y2) and the gender and stalking behavior variables. Post
2
3 hoc ANOVAs revealed a significant difference in trait anxiety scores ($p < 0.001$), and
4
5 forms of behavior such as following ($p < 0.05$) and unwanted communications ($p < 0.05$),
6
7 suggesting a difference in how the possible hazard is perceived by men and women.
8
9 Physical and emotional consequences were not significant.
10

11 *Coping strategies*

12
13 To cope with the phenomenon, victims adopted a range of strategies as shown in
14
15 table 4. From our results it appears that no single strategy successfully stopped the
16
17 stalking, and that stalking behavior did not end with the implementation of coping
18
19 strategies. Victims of Rejected stalkers constituted an exception, since gathering
20
21 evidence proved useful for 14 (40%) of them. The victims of Ex-partner stalkers were
22
23 also an exception, with half of them finding that reducing contact with friends was
24
25 helpful. A common strategy was to talk to somebody about the stalking experience, for
26
27 example, the victim's partner, family members, friends and/or colleagues, but this
28
29 strategy was only considered useful for victims of Rejected (47.20%) and Ex-partner
30
31 (50.00%) stalkers. On average, the most used strategies were offensive tactics (e.g.
32
33 collecting evidence) and interactional tactics (e.g. answering calls, text messages and e-
34
35 mails), while defensive tactics (e.g. having a security plan) and therapy (e.g. seeking
36
37 professional advice) were used less. ANOVA and multivariate analyses were used to
38
39 determine differences in gender, relationship, motives and coping strategies, but the
40
41 results revealed no significant associations.
42
43
44
45
46
47
48
49
50

51 - Table 4 approximately here -
52
53

54 Conclusion

55 The analysis of the literature has shown that health professionals and in
56
57
58
59
60
61
62
63
64
65

1 particular nurses are at greater risk of being stalked than the general population, both by
2
3 ex-partners and acquaintances (such as colleagues and patients, Pathè *et al*, 2002) and
4
5 by unknown stalkers, who may have different motives: relational rejection, an
6
7 infatuation, an inability to express their own emotions and recognize those of others, a
8
9 desire for revenge (Mullen *et al*, 1999). Despite these data, in our sample the percentage
10
11 of nurses' victims (13.99%) was very similar to the average across the general
12
13 population (13.9%; Spitzberg and Cupach, 2003). Another difference regarded the type
14
15 of relationship: the majority of stalkers in our study were Acquaintances (41.12%),
16
17 while in the studies by Kong (1996) and Sheridan and Davies (2001) stalkers were
18
19 generally former partners. With regard to stalkers' motives, according to Mullen *et al*
20
21 (1999) the majority are Rejected stalkers (32.71%), but we found more Incompetent
22
23 suitors than Intimacy-Seekers (respectively 26.17% and 13.08%), and a large proportion
24
25 of Resentful stalkers (22.43%). These data could be explained in relation to the
26
27 particular nature of the work performed by nurses. The fact that they care for others
28
29 could cause patients, colleagues, friends, family members or partners to develop
30
31 disillusioned beliefs about the care they provide: they expect nurses to pay more
32
33 attention to the relationship and failure to fulfill this expectation could trigger reactions
34
35 such as anger and revenge. This causes high levels of anger and anxiety among victims,
36
37 especially women, who perceive themselves to be more exposed, and the emotional
38
39 consequences include fear, professional insecurity, a sense of isolation and in some
40
41 cases adverse effects on performance at work. Our results for depression were
42
43 interesting: we expected to report higher levels, as suggested in the literature (see
44
45 Brewster, 1998; Haider & Haider, 2002), than those actually found. Further research is
46
47 needed to investigate levels of depression and anxiety in nurses – both among victims of
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 stalking and those who are not – also in relation to the organizational context, in order to
2
3 compare results.
4

5
6 In line with other studies (see Meloy, 2000), our findings showed most victims to
7
8 be females stalked (on average) by older males, whose behavior included unwanted
9
10 communications, following, control, assault and threats. Victims who had a closer
11
12 relationship with the stalker (former partner) were more exposed to the phenomenon,
13
14 but findings about access to information on victims was alarming: anybody could obtain
15
16 private information, for instance by asking colleagues or using the internet. Much
17
18 information was actually available on the hospital's official website (e.g. CV, e-mail
19
20 address, work phone). We found the average duration of stalking to be more than one
21
22 year, and that victimization affected the quality of life, for example by limiting social
23
24 life and forcing people to change their routine. The most used coping strategies involve
25
26 personal protection (offensive tactics) and confronting stalkers, to explain the disruption
27
28 they are causing (interactional tactics).
29
30
31
32
33

34
35 Can this phenomenon be prevented? The findings of research into victimization
36
37 among nurses could be used by institutions and individuals to acquire further
38
39 knowledge of stalking, its impact on nurses' lives, and the strategies that can be
40
41 implemented to promote safety, including at the workplace. Researchers have suggested
42
43 at least adopting legal, organizational and individual strategies (both to prevent and to
44
45 intervene). Hospital managers are seriously committed to addressing legal issues (rights
46
47 and duties of each worker): it is important to pay attention to this problem in order to
48
49 ensure the timely intervention of safety officers and avoid any infringement of the law.
50
51 Thus means that it is important know the laws adopted in the country under which
52
53 stalkers and stalking behaviour (the different degrees based on the number of instances
54
55
56
57
58
59
60
61
62
63
64
65

1 and gravity of intent) are punishable (Sheridan & Davies, 2001). Organizational
2
3 strategies should include formal educational programs, especially for nurses in the early
4
5 stages of their career, who should be informed of this professional risk (Kaplan, 2006).
6
7 Educational programs could focus on raising awareness, paying special attention to all
8
9 unwanted behavior (own and other) likely to generate annoyance and physical assault.
10
11 Assessments of risks among nurses should contribute to the development of
12
13 management approaches that reduce the emotional consequences of stalking.
14
15
16
17

18 About individual strategies, researchers have suggested that nurses should
19
20 protect themselves by refusing to treat certain patients that who have experienced an
21
22 attraction or a certain kind of attention, guarding personal data and not disclosing
23
24 personal (or other) details to patients (Hughes, Thom & Dixon, 2007). Finally, in case of
25
26 victimization, the treatment of nurses must be based on a comprehensive approach
27
28 including practical measures, such as recommended behavior, advice and practical
29
30 assistance for accessing police and other legal protection (Mullen *et al*, 2006), education
31
32 and supportive psychotherapy, such as group therapy with stalking survivors, Cognitive
33
34 Behavioral Therapy (CBT), Eye Movement Desensization and Reprocessing (EMDR)
35
36 and other stress management techniques (Abrams & Robinson, 1998; Bisson *et al*,
37
38 2007). This is an expensive phenomenon, from both a personal and organizational point
39
40 of view.
41
42
43
44
45
46
47
48
49

50 References

51
52 Abrams, K.M. & Robinson, G.E. (1998). Stalking. Part II: Victims problems with legal
53
54 system and therapeutic considerations. *Canadian Journal of Psychiatry*, 43(5):
55
56 477-81.
57
58
59
60
61
62
63
64
65

- 1 Ashmore, R., Jones, J., Jackson, A. & Smoyak, S. (2006). A survey of mental health
2
3 nurses' experiences of stalking. *Journal of Psychiatric and Mental Health*
4
5 *Nursing*, 13: 562-569.
6
7
- 8 Bernstein, H.A. (1981). Survey of threats and assaults directed toward psychotherapists.
9
10 *American Journal of Psychotherapy*, 35(4): 542-549.
11
12
- 13 Bisson, J.I., Ehlers, A., Matthews, R., Pilling, S., Richards, D. & Turner, S. (2007).
14
15 Psychological treatments for chronic post traumatic stress disorder. Systematic
16
17 review and meta-analysis. *British Journal of Psychiatry*, 190: 97-104.
18
19
- 20 Budd, T. & Mattinson, J. (2000). *Stalking: Findings from the 1998 British Crime*
21
22 *Survey*. London: Home service.
23
24
- 25 Coleman, F.L. (1997). Stalking Behavior and the Cycle of Domestic Violence. *Journal*
26
27 *of Interpersonal Violence*, 12(3-4): 420-432.
28
29
- 30 Douglas, K.S., & Dutton, D.G. (2001). Assessing the link between stalking and
31
32 domestic violence. *Aggression and Violent Behavior*, 6(6): 519-546.
33
34
- 35 Emerson, R.M., Ferris, K.O. & Gardner, C.B. (1998). On being Stalked. *Social*
36
37 *Problems*, 45(3): 289-314.
38
39
- 40 Flannery, R.B., Hansen, M.A., Penk, W.E. & Flannery, G.J. (1994). Violence against
41
42 women: Psychiatric patient assaults on female staff. *Professional Psychology:*
43
44 *Research and Practice*, 25(2): 182-184.
45
46
- 47 Galeazzi, G.M. & De Fazio, L. (2006). A review on the stalking of mental health
48
49 professionals by patients, prevention and management issues. *Primary Care*
50
51 *and Community Psychiatry*, 11(2):57-66.
52
53
- 54 Galeazzi, G.M., Elkins, K. & Curci, P. (2005). The stalking of mental health
55
56 professionals by patients. *Psychiatric Services*, 56:137-138.
57
58
59
60
61
62
63
64
65

- 1 Gentile, S.R. (2001). The stalking of psychologists by their clients: A descriptive study.
2
3 [Dissertation Abstract] Dissertation Abstracts International: Section B: The
4
5 Sciences and Engineering. Vol 62 (2-B), pp. 1077.
6
7
8 Gentile, S.R., Asaman, J.K., Harmell, P.H. & Weathers, R. (2002). The stalking of
9
10 psychologists by their clients. *Professional Psychology: Research and*
11
12 *Practice*, 33:490–494.
13
14
15 Guy, J.B., Brown, K. & Poelstra, P. (1990). Who gets attacked? A national survey of
16
17 patient violence directed at psychologists in clinical practice. *Professional*
18
19 *Psychology: Research and Practice*, 21(6): 493–495.
20
21
22
23 Guy, J.B., Brown, K. & Poelstra, P. (1992). Safety concerns and protective measures
24
25 used by psychotherapists . *Professional Psychology: Research and Practice*,
26
27 23(5): 421-423.
28
29
30 Hahn, S. et al (2012). Patient and visitor violence in the general hospital, occurrence,
31
32 staff interventions and consequences: a cross-sectional survey. *Journal of*
33
34 *Advanced Nursing*, doi: 10.1111/j.1365-2648.2012.05967.x.
35
36
37 Hudson-Allez, G. (2002). The prevalence of stalking of psychological therapists
38
39 working in primary care by current or former clients. *Counseling and*
40
41 *Psychotherapy Research*, 2:139-146.
42
43
44 Hughes, F.A., Thom, K., & Dixon, R. (2007). Nature & prevalence of stalking among
45
46 New Zealand mental health clinicians. *Journal of Psychosocial Nursing &*
47
48 *Mental Health Services*, 45(4): 32-39.
49
50
51 Kamphuis, J.H., Galeazzi, G.M., De Fazio, L., Emmelkamp, P.M.G., Farnham, F.,
52
53
54 Groenen, A., James, D. & Vervaeke, G. (2005). Stalking perceptions and
55
56 attitudes among helping professions: An EU cross-national comparison.
57
58
59
60
61
62
63
64
65

Clinical Psychology and Psychotherapy, 12:215-225.

Krammer, A., Stepan, A., Baranyi, A., Kapfhammer, H.P. & Rothenhäusler, H.B.

(2007). Auswirkung von Stalking auf Psychiater, Psychotherapeuten und
Psychologen: Häufigkeit von Stalking und dessen emotionale Folgen.

Nervenarzt, 78(7): 809-817.

Kurt, J.L. (1995). Stalking as a variant of domestic violence. *Bulletin of the American*

Academy of Psychiatry and the Law, 23: 219–230.

Lion, J.R. & Herschler, J. A. (1998). The stalking of clinicians by their patients. In *The*

Psychology of Stalking. Clinical and Forensic Perspectives (Meloy JR ed.),

Academic Press, San Diego, CA, pp.165-173.

Luberto, S. (2005). Introduzione in *Donne vittime di Stalking: riconoscimento e*

modelli di intervento in ambito europeo. Franco Angeli, Milano.

Magnavita, N. & Heponiemi, T. (2011). Workplace Violence against Nursing Students

and Nurses: An Italian Experience. *Journal of Nursing Scholarship*, 43(2):203-
211.

McIvor, R.J., Potter, L. & Davies, L. (2006). Stalking Behaviour by Patients Towards

Psychiatrists in a Large Mental Health Organization. *International Journal of*
Social Psychiatry, 54(4): 350-357.

Meloy, J.R. (Ed) (1998). *The Psychology of Stalking: Clinical and Forensic*

Perspectives. Academic Press, San Diego.

Mullen, P.E., Mackenzie, R., Ogloff, J.R., Pathé, M., McEwan, T. & Purcell, R.J.

(2006). Assessing and managing the risks in the stalking situation. *American*
Academy Psychiatry Law, 34(4): 439-50.

Pathé, M. & Mullen, P.E. (1997). The impact of stalkers on their victims. *British*

1 *Journal of Psychiatry*, 170: 12–17.

2
3 Pathè, M., Mullen, P.E. & Purcell, R. (2002). Patients who stalk doctors: their motives
4
5 and management. *Medical Journal of Australia*, 176: 335–338.

6
7
8 Pope, K.S. & Tabachnick, B.G. (1993). Therapists' anger, hate, fear, and sexual feelings:
9
10 National survey of therapist responses, client characteristics, critical events,
11
12 formal complaints, and training. *Professional Psychology: Research &*
13
14 *Practice*, 24(2), 142-145.

15
16
17
18 Purcell, R., Pathè, M. & Mullen, P.E. (2002). The prevalence and nature of stalking in
19
20 the Australian community. *Australian and New Zealand Journal of Psychiatry*,
21
22 36:114–120.

23
24
25 Purcell, R., Powell, M.B. & Mullen, P.E. (2005). Clients who stalk psychologists:
26
27 Prevalence, methods, and motives. *Professional Psychology: Research and*
28
29 *Practice*, 36: 537-543.

30
31
32
33 Romans, J.S., Hays, J.R. & White, T.K. (1996). Stalking and related behaviours
34
35 experienced by counseling centre staff members from current or former
36
37 clients. *Professional Psychology: Research and Practice*, 27:595-599.

38
39
40 Sandberg, D.A., McNiel, D.E. & Binder, R.L. (2002), Stalking, threatening, and
41
42 harassing behavior by psychiatric patients toward clinicians. *Journal of the*
43
44 *American Academy of Psychiatry and Law*, 30:221-229.

45
46
47 Sheridan, L. & Davies, G.M. (2001). Stalking: The elusive crime. *Legal and*
48
49 *Criminological Psychology*, 6, 133-147.

50
51
52 Smoyak, S. (2003). Perspectives of mental health clinicians on stalking continue to
53
54 evolve. *Psychiatric Annals*, 33: 641–648.

55
56
57 Tryon, G.S. (1986). Abuse of therapists by patients: A national survey. *Professional*
58
59
60
61
62
63
64
65

1 *Psychology: Research and Practice*, 17, 357-363.
2

3 Tjaden, P. & Thoennes, N. (1998). *Stalking in America: Findings from the National*
4
5 *Violence Against Women Survey*. National Institute of Justice and Centers for
6
7 Disease Control and Prevention, Washington, DC.
8
9

10 Tjaden, P. & Thoennes, N. (2000). *Extent, Nature, and Consequences of Intimate*
11
12 *Partner Violence: Findings From the National Violence Against Women*
13
14 *Survey*. National Institute of Justice and Centers for Disease Control and
15
16 Prevention, Washington, DC.
17
18
19

20 Zona, M.A., Sharma, K.K. & Lane, J.C. (1993). A comparative study of erotomantic and
21
22 obsessional subjects in a forensic sample. *Journal of Forensic Sciences*, 38:
23
24 894–903.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 – Socio-demographic characteristic, relationship victims-stalkers and motivation of stalking campaign

	Victims						Stalkers				
	f	M%	F%	Age Range	Means	s.d.	M%	F%	Age Range	Means	s.d.
Relationship											
Ex- partner	28	21.43	78.57	19-55	38.25	10.03	75.00	25.00	20-56	39.65	9.32
Acquaintance	44	13.64	86.36	21-56	35.91	10.03	84.09	15.91	20-80	38.84	13.37
Unknown ¹	35	22.86	77.14	20-60	39.88	11.69	65.71	14.29	25-60	38.38	11.24
Motivation											
Rejected	35	22.86	77.14	19-55	36.77	10.10	77.14	22.86	20-57	39.74	10.52
Intimacy Seeker	14	7.14	92.86	21-54	38.86	11.48	78.57	21.43	20-80	42.93	10.31
Incompetent ²	28	21.43	78.57	20-60	38.36	11.67	78.57	14.29	25-60	36.60	10.32
Resentful ³	24	16.67	83.33	22-52	36.04	9.30	79.19	16.67	20-54	35.43	10.59

¹ stalkers gender and age MI= 7

² stalkers gender and age MI=2

³ stalkers gender and age MI=1

Table 2 – Harassing behavior characterizing stalking campaign (values expressed in percentage)

Behaviors	Ex-partner n=28	Acquaintance n=44	Unknown n=35	Rejected n=35	Intimacy S. n=14	Incompetent n=28	Resentful n=24
Calls	60.71	50.00	62.86	57.14	42.86	67.86	58.33
Letters	17.86	15.91	20.00	22.86	14.29	21.43	12.50
Sms	46.43	34.09	8.57	48.57	28.57	17.86	20.83
Gifts	7.14	11.36	14.29	8.57	14.29	7.14	8.33
Other unwanted communication	14.29	29.55	5.71	22.86	42.86	7.14	12.50
Following	46.43	40.91	31.43	42.86	57.14	28.57	33.33
Breaking home	3.57	6.82	0.00	2.86	14.29	0.00	4.17
Visiting							
- Home	10.71	11.36	0.00	11.43	14.29	7.14	4.17
- Workplace	25.00	27.27	20.00	22.86	14.29	32.14	29.17
Waiting							
- Outside Home	50.00	40.91	20.00	42.86	50.00	21.43	45.83
- Outside Workplace	32.14	29.55	17.14	37.14	21.43	14.29	29.17
Vandalizing (home, car, properties...)	28.57	15.91	5.71	31.43	14.29	3.57	12.50
Physical assault	14.29	11.36	0.00	8.57	14.29	3.57	12.50
Sexual assault	3.57	0.00	0.00	2.86	0.00	0.00	0.00
Threat							
- Physical assault	35.71	11.36	5.71	28.57	0.00	3.57	25.00
- Sexual assault	0.00	9.09	2.86	0.00	7.14	7.14	8.33
- Third parties assault	10.71	2.27	5.71	5.71	7.14	7.14	4.17
Harass others (relatives, friends, colleagues...)	14.29	2.27	8.57	11.43	0.00	10.71	4.17
Manipulate others	17.86	9.09	8.57	14.29	14.29	10.71	8.33
Spread rumors	35.71	13.64	11.43	25.71	14.29	7.14	29.17

Table 3 – Consequences of stalking campaign on victims

	Ex-partner n=28	Acquaintance n=44	Unknown n=35	Rejected n=35	Intimacy S. n=14	Incompetent n=28	Resentful n=24
Physical %							
- Digestive disorder	60.71	22.73	22.86	51.43	35.71	7.14	37.50
- Sleep disorder	50.00	34.09	37.14	45.71	28.57	35.71	45.83
- Headache	28.57	18.18	11.43	20.00	21.43	14.29	25.00
- Tiredness	32.14	13.64	20.00	20.00	7.14	17.86	37.50
- Sickness	3.57	6.82	0.00	5.71	14.29	0.00	0.00
- Weakness	7.14	11.36	8.57	8.57	7.14	3.57	20.83
- Self harm	3.57	2.27	0.00	0.00	7.14	0.00	4.17
- Purging	0.00	6.82	0.00	2.86	7.14	0.00	4.17
- Panic attack	25.00	15.91	17.14	14.29	28.57	14.29	29.17
Emotive %							
- Suicidal thoughts	0.00	2.27	0.00	0.00	7.14	0.00	0.00
- Depression	21.43	6.82	8.57	14.29	7.14	0.00	25.00
- Anxiety	64.29	45.45	48.57	62.86	42.86	46.43	54.17
- Anger	42.86	50.00	48.57	48.57	50.00	50.00	54.17
- Confusion	35.71	25.00	8.57	22.86	28.57	10.71	37.50
- Fear	64.29	43.18	45.71	54.29	57.14	46.43	54.17
- Increased distrust	21.43	6.82	2.86	14.29	7.14	0.00	16.67
- Aggression	14.29	13.64	0.00	20.00	7.14	0.00	8.33
- Paranoia	14.29	13.64	11.43	17.14	14.29	14.29	8.33
- Annoyance	17.86	27.27	25.71	22.86	21.43	28.57	29.17
- Agoraphobia	0.00	4.55	0.00	0.00	14.29	0.00	0.00
Psychological Mean (s.d.)							
- BDI	10.06 (8.93)	11.30 (8.39)	11.26 (6.29)	9.45 (8.21)	10.33 (6.04)	10.21 (5.48)	11.53 (6.97)
- STAI Y1	40.17 (12.92)	41.51 (13.56)	44.65 (8.76)	38.49 (11.62)	36.15 (12.77)	42.62 (10.91)	42.50 (11.18)
- STAI Y2	42.73 (13.92)	44.55 (12.87)	44.85 (11.91)	40.50 (12.34)	38.85 (8.93)	43.42 (12.14)	43.68 (11.41)

Table 4 – Coping strategies (values expressed in percentage)

	Ex-partner n=28	Acquaintance n=44	Unknown n=35	Rejected n=35	Intimacy S. n=14	Incompetent n=28	Resentful n=24
Get evidence of stalker in action	57.14	43.18	25.71	60.00	35.71	32.14	37.50
Hold on any evidence	53.57	47.43	20.00	51.43	28.57	21.43	62.50
Have a safety plan	28.57	15.91	2.86	31.43	21.43	0.00	8.33
Tell to relatives, colleagues, friends...	57.14	43.18	22.86	48.57	42.86	28.57	45.83
Assume more alcohol or drugs than usual	10.71	11.36	4.55	7.41	21.43	7.14	8.33
Reduce social activity	28.00	22.73	4.55	28.60	7.14	10.71	20.83
Change routine	28.57	36.36	20.00	34.29	50.00	25.00	16.67
Carry a weapon	14.29	2.27	2.86	11.43	0.00	3.57	8.33
See a counselor or psychologist	21.43	13.64	2.86	14.29	35.71	0.00	12.50
Ignore the stalker	14.29	11.36	28.57	8.57	28.57	32.14	12.50
Answer:							
- Calls	46.43	52.27	42.86	48.57	42.86	53.57	54.17
- E-mail/letters	21.43	4.55	2.86	22.86	0.00	3.57	0.00
- Sms	35.71	20.45	2.86	42.86	21.43	3.57	4.17
Confront the stalker	35.71	34.09	11.43	45.71	28.57	7.14	29.17
Threat the stalker	14.29	22.73	11.43	20.00	28.57	10.71	16.67
Attack the stalker	10.71	13.64	2.86	17.14	0.00	3.57	12.50
Ask to stop	53.57	45.45	17.14	51.43	28.57	14.29	62.50

Title page:

Italian nurses' experience of stalking: a questionnaire survey.