

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Olivine thermal emissivity under extreme temperature ranges: implication for Mercury surface

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/135211> since 2016-07-21T10:03:17Z

Published version:

DOI:10.1016/j.epsl.2013.03.038

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in EARTH AND PLANETARY SCIENCE LETTERS, 371-372, 2013, 10.1016/j.epsl.2013.03.038.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.epsl.2013.03.038

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0012821X13001593>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/135211>

Olivine thermal emissivity under extreme temperature ranges: implication for Mercury surface

Nestola F. ¹, Ferrari S. ¹, Maturilli A. ², Massironi M. ^{1,3}, Helbert J. ², Redhammer G. J. ⁴, Capria M.T. ⁵, Carli C. ⁶, Capaccioni F. ⁶, Bruno M. ⁷

(1) Dipartimento di Geoscienze, Università degli Studi di Padova, Italy, (2) Institute for Planetary Research, DLR, Berlin-Adlershof, Germany, (3) CISAS, Università degli studi di Padova, Italy, (4) Institute for Crystallography for the RWTH, Aachen, Germany, (5) IFSI-INAF Roma, Italy, (6) IFSI-INAF Roma, Italy, (7) Università di Torino, Dipartimento di Scienze della Terra, Italy.

Due to its proximity to the Sun and a weak exosphere, the surface of Mercury is subjected to temperature variation up to 625 K during a hermean day. Minerals elastically expand and contract adjusting their crystal structure through time and from place to place as a function of temperature variations. This significantly influences the interpretation of remote sensing data by instantaneously and reversibly modifying the mineral spectral signatures, which are intimately related to the crystal structure. Here we show, for the first time, how the thermal emissivity spectra of a Mg-rich olivine (i.e. $\text{Mg}_{1.84}\text{Fe}_{0.16}\text{SiO}_4$) significantly changes as a function of temperature. An increase of 420 K modifies the olivine spectrum shifting in wavelength its emissivity features, simulating the spectrum of an olivine strongly enriched in iron (Fe abundance close to 20%). This “chameleon-like behaviour” indicates that the spectroscopic investigations of the surface of Mercury must take into account the local time dependency to properly infer the compositions of planetary materials by remote sensing acquisition of future space missions.

It is well known that the surface of Mercury shows extremely large temperature variations at specific locations¹. This is due to its closeness to the Sun but also to its slow rotation (one hermean day last 88 terrestrial days) which allows the surface temperature of the side of the planet facing the Sun to reach equilibrium with the solar flux while its dark side is much colder; the surface temperature can vary over a hermean day between about 70 and 700 K¹ at different latitudes. These temperature variations significantly affect the crystal structure and density of minerals and rocks present on the surface of a planet².

As a consequence, several kinds of remote information linked to mineral structures, including relative spectral signatures, are strictly dependent on the environmental conditions and vary according to the surface temperature. By “looking at the planet” in remote sensing, Mercury may change its outer face through time and from place to place as a function of solar irradiation. If this is not taken into account it could lead to important misinterpretations and could limit our capability of inferring compositions and rheological properties of materials from remote sensing acquisition.

Thus, we need to investigate the *in situ* behaviour of geological materials, i.e. under extreme temperature environments, and improve the geological interpretation and compositional inferences by applying the knowledge acquired to the analysis of the available remote sensing data.

In this light, here we present a new *in situ* multi-methodological laboratory approach comprehensive of high-temperature X-ray diffraction (single-crystal and powder materials) and high-temperature thermal infrared (TIR) spectroscopy in the range 1400 to 700 cm⁻¹ (7 to 14 μm). Such range is the same that will be investigated by the Mercury Radiometer and Thermal Infrared Spectrometer³ (MERTIS) on board of BepiColombo Mercury Planetary Orbiter spacecraft. In such wavelength range, the spectra can be used very effectively to identify the fine-scale structural properties of silicates (e.g. stretching and bending motions in the silicon-oxygen anions, metal – oxygen and lattice vibrations⁴). In addition, for some important mineral family, like olivine, the emissivity peak positions are a good indicator of the composition^{5,6}. TIR spectroscopy was performed at the Planetary Emissivity Laboratory in Berlin (PEL), which has been built up over the

last 5 years to study systematically the effect of temperature on TIR spectral signatures in the 1400-700 cm^{-1} (7-14 μm) range⁷.

In this work we investigated an olivine sample having composition $\text{Mg}_{1.84}\text{Fe}_{0.16}\text{SiO}_4$ (Fo_{92})^{8,9,10,11}. Olivine has been chosen as one of the major component of Earth's mantle, of many meteorites classes and has been identified on the surfaces of planetary bodies, and in the spectra of astronomical targets³. Moreover, the presence of olivine has been considered in several petrological models of Mercury^{12,13,14,15} and should likely be a primary mineralogical phase in the wide volcanic plains recently discovered on its surface¹⁶, particularly if the new MESSENGER X-ray observations pointing to a basic to ultrabasic compositions for these effusions are taken into account¹⁷. The olivine studied here occurs in high-pressure ultramafic nodules from the Newer Volcanics of Mt. Leura^{8,9} (Victoria, Australia) and was fully characterized by single-crystal X-ray diffraction and powder X-ray diffraction. The chemistry has been assumed to be identical to that published for the same sample by 9, as confirmed by the X-ray structure refinement of our sample and by the excellent agreement between the unit-cell parameters obtained by our characterization and their data (Supplementary Tables 1, 2, 3).

The minerals present on the Mercury surface undergo temperature variations that depend on solar irradiation. Known thermal models^{18,19} of Mercury show that at the equator a point on the surface is subjected to a temperature variation of about 420 K between sunrise days and the midday.

In order to evaluate how olivine would behave on the Mercury surface under such significant temperature variations, we have measured its volume - and density - change as a function of temperature (Fig. 1a) (see Methods Summary). In detail, we have measured the unit-cell volume between room temperature (the minimum temperature presently achievable by our set-up and at the same time the temperature at which all the TIR mineral databases are available^{6,7}) and 773 K, in order to cover most of the possible temperature variations on Mercury surface. The volume change as a function of temperature (Table 1) gave us a measure of the linear volume thermal expansion coefficient $\alpha_V = 3.07(2) \times 10^{-5} \text{ K}^{-1}$ (calculated as: $\alpha_V = [(V-V_0)/V_0]/\Delta T$) in agreement with previous

results²⁰. Assuming a temperature variation of 420 K, as for example the temperature variation between sunrise and midday on Mercury at the equator, we calculate a volume variation of 3.77(8) Å³ (calculating using the volume thermal expansion coefficient). In Fig. 1b the unit-cell volume is plotted against the composition of olivines for compositions ranging between Fo₉₂ and Fo₆₂, the compositional range most representative of the geological environments²¹. It is evident that increasing the iron content the unit-cell volume significantly increases due to the larger cation radius of Fe²⁺ (in octahedral coordination the radius of Fe²⁺ is 0.78 Å against 0.72 Å of Mg⁽²²⁾). Hence, the increase in temperature and the increase in iron have the same effect on olivine, increasing the unit-cell volume. According to the equation $V = 308.6(6) - 0.181(8) \times \text{Fo}\%$ ^{10,11}, an increase in temperature by about 420 K corresponds to an increase in iron from Fo₉₂ to Fo₇₁.

From such crucial experimental observation we decided to determine the effect of temperature on the TIR spectral signature of olivine, performing measurements in vacuum at different temperatures (see Methods Summary). The olivine sample was hand-reduced to a grain size range between 100 µm and 160 µm which is considered close to a most likely end-member of the Mercury regolith size²³. In detail, on the same sample we have collected two spectra at two different temperatures between about 350 K and 770 K (the error in temperature calibration is estimated to be up to 5 K) in order to simulate the typical diurnal equatorial temperature variation of the hermean surface.

In order to define the effect of the thermal annealing during the emissivity data collection, the sample was re-analyzed *ex-situ* by X-ray powder diffraction and single-crystal X-ray diffraction and did not show any significant difference in the unit-cell parameters with respect to the pre-annealing data (Supplementary Table 4). In particular, we can state that the sample did not undergo any iron oxidation due to the temperature effect: in fact, as it is well known, an iron oxidation would have reduced the Fe²⁺ to Fe³⁺ with a consequent reduction of the cation radius from 0.78 Å to 0.645 Å, respectively, and a consequent decrease of the unit-cell parameters.

In Figure 2a the main olivine bands between 1000 and 850 cm⁻¹ are shown. Bands 1, 2 and 3 are the most intense in olivine in this spectral range and their exact position is a strong function of the iron

abundance⁴. In the inset 2b we highlight the shift of 7 cm^{-1} in the band 3 position, between the 352 K-spectrum and the 773 K-spectrum. This is a significant result as the same wavenumber shift corresponds to a change in composition from Fo_{92} to Fo_{71} according to the data of Hamilton⁴ in Table 6.

On the surfaces of planetary bodies we have rocks and soils and we must then consider the simultaneous presence of different minerals, each with its characteristic thermal expansion coefficient. This would result in a more complex behaviour and in an even more difficult interpretation of the spectra: in fact, for example, a further important mineral assumed to be present on Mercury is the plagioclase and plagioclase have very different thermal expansion coefficients with respect to olivine. Data obtained on several plagioclase compositions²⁴, using the same equation used considered for olivine, gave thermal expansion coefficients very variable between $\alpha_V = 2.8 \times 10^{-5} \text{ K}^{-1}$ for pure albite $\text{NaAlSi}_3\text{O}_8$ and $\alpha_V = 1.6 \times 10^{-5} \text{ K}^{-1}$ (for pure anorthite $\text{CaAl}_2\text{Si}_2\text{O}_8$ (data calculated between 298 and 774 K). Clinopyroxene and orthopyroxene also have been assumed to be possible phases for Mercury surface¹⁵: for these minerals different thermal expansion coefficient data have been determined. For compositions between diopside ($\text{CaMgSi}_2\text{O}_6$) and hedenbergite ($\text{CaFeSi}_2\text{O}_6$) literature data show values between $3.3^{(25)}$ and $2.9^{(26)} \times 10^{-5} \text{ K}^{-1}$; for orthopyroxenes (between orthoenstatite $\text{Mg}_2\text{Si}_2\text{O}_6$ and orthoferrosilite $\text{Fe}_2\text{Si}_2\text{O}_6$) values close to $3.0\text{-}3.1 \times 10^{-5} \text{ K}^{-1}$ have been determined²⁷. Thus, considering such different thermal expansion behaviour, we could easily state that spectral signature collected at room temperature could be totally different from that collected at high temperature.

Our cross-related data by in situ X-ray diffraction and TIR spectroscopy demonstrate that the effect of temperature must be taken into account when interpreting spectral features, in order to correctly determine the abundance and composition of specific mineral phases.

Our results should be considered reliable for all those terrestrial bodies and asteroids, which undergo significant surface temperature variations and that could be characterized by what we called “chameleon-like surface”.

METHODS SUMMARY

In situ high-temperature diffraction. In situ X-ray measurements were performed by powder diffraction experiments (5° - 100° in 2θ , continuous scan mode) in the temperature range 295 K - 1273 K to include a possible temperature range encountered on the Hermean surface. Data were collected on a Philips X'Pert diffractometer system (University of Salzburg, Austria), equipped with an Anton PAAR HTK-16 high temperature chamber (Cu Ka radiation, primary and secondary side 0.04 rad Soller slits, secondary graphite monochromator, 4 hours/ measurement). After each temperature increase the sample was allowed to equilibrate for one hour before starting measurement at this temperature. NIST silicon was used as an internal standard; lattice parameters were obtained from whole pattern refinement using the FULLPROF-suite²⁸ and are reported at each temperature in Table 1.

In situ high-temperature emissivity measurements. Emissivity data were measured in the Planetary Emissivity Laboratory²⁸ (PEL), at the Institute for Planetary Research of the German Aerospace Center (DLR) in Berlin. The instrument used is a Bruker Vertex 80V, with a liquid nitrogen cooled HgCdTe detector and a KBr beamsplitter. The spectrometer is coupled to an evacuated planetary simulation chamber, having an automatic sample transport system to maintaining the vacuum while changing the samples. Radiance is collected by an Au-coated parabolic 90° off-axis mirror and reflected to the spectrometer entrance port. The sample is heated via a copper induction pancake coil while staying always under 1 mbar of pressure²⁹. Measurements of a calibration body are taken at 2 temperatures to derive the instrumental response function. The artefact that in a certain point of the spectrum, called Christiansen feature (CF), emissivity goes to unity is used for the calibration procedure³⁰ and to determine the temperature of the sample at the emitting skin.

1. Strom, R. G., Sprague, A. L. *Exploring Mercury: The Iron Planet* (Springer/Praxis, New York, 2003).
2. Hazen, R. M. & Downs, R. T. High-Temperature and High-Pressure Crystal Chemistry. In Hazen, R. M. and Downs, R. T. (eds) *Reviews in Mineralogy and Geochemistry*, Vol. 41 (Mineralogical Society of America, Washington DC, 2000).
3. Hiesinger, H. & Helbert, J. The Mercury Radiometer and Thermal Infrared Spectrometer (MERTIS) for the BepiColombo mission. *Planet. Space Sci.* **58**, 144-165 (2010).
4. Hamilton, V. E. Thermal infrared (vibrational) spectroscopy of Mg-Fe olivines: A review and applications to determining the composition of planetary surfaces. *Chem. Erde* **70**, 7-33 (2010).
5. Koike, C. *et al.* Compositional dependence of infrared absorption spectra of crystalline silicate - II. Natural and synthetic olivines. *Astron. Astrophys.* **399**, 1101-1107 (2003).
6. Christensen, P. *et al.* A thermal emission spectral library of rock-forming minerals. *J. Geophys. Res. Planets* **105**, 9735-9739 (2000).
7. Maturilli, A., Helbert, J. & Moroz, L. The Berlin emissivity database (BED). *Planet. Space Sci.* **56**, 420-425 (2008).
8. Dal Negro, A. *et al.* Crystal chemistry and evolution of the clinopyroxene in a suite of high pressure ultramafic nodules from the Newer Volcanics of Victoria, Australia. *Contrib. Mineral. Petrol.* **86**, 221-229 (1984).
9. Princivalle, F. & Secco, L. Crystal structure refinement of 13 olivines in the forsterite-fayalite series from volcanic rocks and ultramafic nodules. *Mineral. Petrol.* **34**, 105-115 (1985).
10. Nestola, F. *et al.* New accurate elastic parameters for the forsterite-fayalite solid solution. *Am. Miner.* **96**, 1742-1747 (2011).
11. Nestola *et al.* First crystal-structure determination of olivine in diamond: Composition and implications for provenance in the Earth's mantle. *Earth Planet. Sci. Lett.* **305**, 249-255 (2011).
12. Brown, S. M. & Elkins-Tanton, L. T. Compositions of Mercury's earliest crust from magma ocean models. *Earth Planet. Sci. Lett.* **286**, 446-455 (2009).
13. Riner, M. A., McCubbin, F. M., Lucey, P. G., Jeffrey Talyor, G. & Gillis-Davis, J. J. Mercury surface composition: Integrating petrologic modeling and remote sensing data to place constraints on FeO abundance. *Icarus* **209**, 301-313 (2010).
14. Sprague, A. L. *et al.* Spectral emissivity measurements of Mercury's surface indicate Mg- and Ca-rich mineralogy, K-spar, Na-rich plagioclase, rutile, with possible perovskite, and garnet. *Planet. Space Sci.* **57**, 364-383 (2009).
15. Warell, J. *et al.* Constraints on Mercury's surface composition from MESSENGER and ground-based spectroscopy. *Icarus* **209**, 138-163 (2010).

16. Head, J. W. *et al.* Flood volcanism in the northern high latitudes of Mercury revealed by MESSENGER. *Science* **333**, 1853-1856 (2011).
17. Nittler, L. R. *et al.* The major-element composition of Mercury's surface from MESSENGER X-ray spectrometry. *Science* **333**, 1847-1850 (2011).
18. Vasavada, A. R., Paige, D. A. & Wood, S. E. Near surface temperatures on Mercury and the Moon and the stability of polar ice deposits. *Icarus* **141**, 179-193 (1999)
19. Bauch K., Hiesinger, H. & Helbert, J. Insolation and resulting surface temperatures of studyregions on Mercury. *Lunar Planet. Sci. XLII*, abstr. 2257 (Lunar Planet. Inst., 2011).
20. Andrault, D. *et al.* Thermal pressure in the laser-heated diamond-anvil cell: An X-ray diffraction study. *Eur. J. Mineral.* **10**, 931-940 (1998).
21. Morgan, J.W., & Anders, E. Chemical composition of the Earth, Venus and Mercury. *Proc. Natl. Acad. Sci.* **77**, 6973-6977 (1980).
22. Shannon, R. D. Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Crystallogr., Sect. A* **32**, 751-767 (1976).
23. Emery, J. P. *et al.* Mercury: thermal modeling and mid-infrared (5–12 μm) observations. *Icarus* **136**, 104-123 (1998).
24. Tribaudino, M. *et al.* Thermal expansion of plagioclase feldspars. *Contrib. Mineral. Petrol.* **160**, 899-908 (2010).
25. Cameron, M., Sueno, S., Prewitt, C. T. and Papike, J. J. High temperature crystal chemistry of acmite, diopside, hedenbergite, jadeite, spodumene, and ureyite. *Am. Mineral.* **58**, 594-618 (1973).
26. Tribaudino, M., Nestola, F., Bruno, M., Boffa Ballaran, T. & Liebske, C. Thermal expansion along the $\text{NaAlSi}_2\text{O}_6$ - $\text{NaFe}^{3+}\text{Si}_2\text{O}_6$ and $\text{NaAlSi}_2\text{O}_6$ - $\text{CaFe}^{2+}\text{Si}_2\text{O}_6$. *Phys. Chem. Miner.* **35**, 241-248 (2008).
27. Hugh-Jones, D. Thermal expansion of MgSiO_3 and FeSiO_3 ortho- and clinopyroxenes. *Am. Mineral.* **82**, 689-696 (1997).
28. Rodriguez-Carvajal, J. Magnetic structure determination from powder diffraction symmetry analysis and simulated annealing RID C-4362-2008. *Epdic 7: European Powder Diffraction, Pts 1 and 2* **378-3**, 268-273 (2001).
29. Helbert, J. & Maturilli, A. The emissivity of a fine-grained labradorite sample at typical Mercury dayside temperatures. *Earth Planet. Sci. Lett.* **285**, 347-354 (2009).
30. Maturilli, A., Helbert, J., Witzke, A. & Moroz, L. Emissivity measurements of analogue materials for the interpretation of data from PFS on Mars Express and MERTIS on Bepi-Colombo. *Planet. Space Sci.* **54**, 1057-1064 (2006).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgments

Please here....add what must be added!!

Author Contributions N.F. refined X-ray data and fitted them to spectroscopic data; S.F. selected, reduced, prepared the samples and collected spectroscopic data; M.A. collected and processed spectroscopic data; M.M. designed the study; G.R. collected and processed X-ray data; M.T.C. discussed thermal models; M.B. analysed the thermal expansion data; J.H., C.C. and F.C. were involved in study schedule; N.F. and S.F. designed the study, analysed data and wrote the paper. All authors discussed the results and commented on the manuscript.