

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Polyamines and jasmonic acid induce plasma membrane potential variations in Lima bean

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/69558> since 2016-10-11T15:16:10Z

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

R. Ozawa; C.M. Berteà; M. Foti; R. Narayana; G.I. Arimura; A. Muroi; M. Maffei; J. Takabayashi. Polyamines and jasmonic acid induce plasma membrane potential variations in Lima bean. *PLANT SIGNALING & BEHAVIOR*. 5(3) pp: 308-310.

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/69558>

# Polyamines and jasmonic acid induce plasma membrane potential variations in Lima bean

Rika Ozawa, Cinzia M. Berteza, Maria Foti, Ravishankar Narayana, Gen-Ichiro Arimura, Atsushi Muroi, Massimo E. Maffei & Junji Takabayashi  
Page 308-310 | Received 03 Dec 2009, Accepted 04 Dec 2009, Published online: 01 Mar 2010

## Abstract

Exogenous polyamines (PAs) [cadaverine (Cad), putrescine (Put), spermidine (Spd) and spermine (Spm)] elicit the production of volatiles in Lima bean (*Phaseolus lunatus*). Among the tested PAs, Spm induces the production of some volatile terpenoids that are known to be induced by the spider mite *Tetranychus urticae*. Spm treatment elicits the biosynthesis of Jasmonic acid (JA), a phytohormone known to regulate the production of the volatile terpenoids. The treatment with JA together with Spm resulted in the increased volatile emission, and predatory mites *Phytoseiulus persimilis* preferred JA and Spm-treated leaves over those treated with JA alone.<sup>5</sup> JA and Spm treatment has no effects on polyamine oxidase (PAO) and Cu-amine oxidase (CuAO) but has a significant induction of calcium influx, ROS production, enzyme activities for NADPH-oxidase complex, superoxide dismutase, catalase, ascorbate peroxidase, glutathione reductase and glutathione peroxidase, and gene expressions except for NADPH-oxidase complex.<sup>5</sup> Here, we report that a plasma membrane potential ( $V_m$ ) depolarization was observed after polyamine perfusion with an increasing trend: Spm, Cad, Put and Spd. JA perfusion did not alter  $V_m$  but the perfusion of JA and the polyamines significantly increased Cad and Put  $V_m$  depolarization. When JA was perfused with polyamines, a negative correlation was found between  $V_m$  depolarization and the number of amino group of the polyamines tested.

Polyamines are involved in plants' stress responses and growth. By activating biosynthesis of nucleic acids, polyamines concern the plant growth and differentiation.<sup>1</sup> [Walters DR](#). Polyamines in plant-microbe interactions. *Physiol Mol Plant Pathol* 2000; 57:137 - 146 –<sup>3</sup> [Martin-Tanguy J](#). Metabolism and function of polyamines in plants: recent developments (new approaches). *Plant Growth Regul* 2001; 34:135 - 148 Furthermore, it has been reported that polyamines are involved in the response against environmental stress and plant disease.<sup>1</sup> [Walters DR](#). Polyamines in plant-microbe interactions. *Physiol Mol Plant Pathol* 2000; 57:137 - 146 –<sup>4</sup> [Bouchereau A](#), [Aziz A](#), [Larher F](#), [Martin-Tanguy J](#). Polyamines and environmental challenges: recent development. *Plant Sci* 1999; 140:103 - 125 We recently reported that exogenously applied polyamines ~diamines [cadaverine (Cad), putrescine (Put)], triamine [spermidine (Spd)] and tetraamine [spermine (Spm)]~ induce volatile emission in Lima bean leaves.<sup>5</sup> [Ozawa R](#), [Berteza CM](#), [Foti M](#), [Narayana R](#), [Arimura G-I](#), [Muroi A](#), et al. Exogenous polyamines elicit herbivore-induced volatiles in Lima bean leaves: involvement of calcium, H<sub>2</sub>O<sub>2</sub> and jasmonic acid. *Plant Cell Physiol* 2009; 50:2183 - 2199 Membrane potentials ( $V_m$ ) and intracellular calcium variations were also studied in Lima bean leaves after perfusion with the polyamines and with these addition of JA and here we report on these additional results.

The primary candidate for intercellular signaling in higher plants is the stimulus-induced change in  $V_m$ .<sup>6</sup> [Maffei M](#), [Mithöfer A](#), [Boland W](#). Before gene expression: early events in plant-insect interactions. *Trends Plant Sci* 2007; 12:310 - 316 The plasma membrane potential ( $V_m$ ), which lies in the range of -50 to -200 mV in Lima bean leaves,<sup>7</sup> [Maffei M](#), [Bossi S](#), [Spitfeller D](#), [Mithöfer A](#), [Boland W](#). Effects of feeding *Spodoptera littoralis* on Lima beans leaves I. Membrane

potentials, intracellular calcium variations, oral secretions and regurgitate components. *Plant Physiol* 2004; 134:1752 - 1762 may be shifted either to more negative (hyperpolarization) or to more positive values (depolarization) in response to various biotic or abiotic stresses.

Measurement of  $V_m$  were performed and data statistically treated as previously described (ANOVA and Tukey-Kramer's HSD test).<sup>7</sup> [Maffei M](#), [Bossi S](#), [Spitfeller D](#), [Miltho'fer A](#), [Boland W](#). Effects of feeding *Spodoptera littoralis* on Lima beans leaves I. Membrane potentials, intracellular calcium variations, oral secretions and regurgitate components. *Plant Physiol* 2004; 134:1752 - 1762

Perfusion with the polyamines ([Fig. 1](#) single arrow) shows a specific response of the leaf tissues with a different  $V_m$  depolarization, depending on the polyamine. In general, a  $V_m$  depolarization was observed after polyamine perfusion with an increasing trend: Spm, Cad, Put and Spd ([Fig. 1](#)). Spm and Spd  $V_m$  depolarization values were significantly different ( $p < 0.05$ ) from all other polyamines, whereas no significant difference was found between Put and Cad  $V_m$  depolarization ( $p = 0.435$ ). In all cases,  $V_m$  depolarization was reversed by washing polyamine-treated leaves with a fresh buffer solution ([Fig. 1](#) double arrow); however, a full recovery of the  $V_m$  was observed only for Put ([Fig. 1](#)). The linearization of the data from [Figure 1](#) allowed to calculate the rate of  $V_m$  depolarization after perfusion of the polyamines which was higher for Spd ( $6.0 \text{ mV min}^{-1}$ ;  $R = 0.96$ ), equal for Put and Cad ( $4.8 \text{ mV min}^{-1}$ ; Put  $R = 0.95$ ; Cad  $R = 0.97$ ) and lower for Spm ( $3.0 \text{ mV min}^{-1}$ ;  $R = 0.96$ ).

Perfusion with JA caused a slight and not significant ( $p = 0.332$ )  $V_m$  depolarization ([Fig. 2](#)) with respect to control. The addition of JA caused a significant increase ( $p < 0.01$ ) in  $V_m$  depolarization when perfused with Cad, with respect to the sole perfusion with Cad ([Fig. 1](#)). The same was observed when JA was perfused with Put, whereas not significant differences were observed when Spm ( $p = 0.513$ ) and Spd ( $p = 0.107$ ) were perfused with JA ([Fig. 2](#)), with respect to the sole perfusion with Spm and Spd ([Fig. 1](#)). The linearization of the data from [Figure 2](#) allowed to calculate the rate of  $V_m$  depolarization after perfusion of the polyamines + JA, which was higher for Cad ( $24.40 \text{ mV min}^{-1}$ ;  $R = 0.99$ ), almost equal for Put and Spd (Put:  $14.21 \text{ mV min}^{-1}$ ,  $R = 0.99$ ; Spd:  $13.49 \text{ mV min}^{-1}$ ,  $R = 0.99$ ) and lower for Spm ( $1.34 \text{ mV min}^{-1}$ ;  $R = 0.93$ ). For JA the rate of  $V_m$  depolarization was  $0.19 \text{ mV min}^{-1}$  ( $R = 0.96$ ). With the addition of JA, a negative correlation was found between  $V_m$  depolarization and the number of amino group of the polyamines tested.

Since ion fluxes through channels directly influence  $V_m$ , it seems reasonable to assume that molecules able to act on channel activity might be considered as important factors inducing electrical signals. Among the various channels, calcium and potassium channels are predominantly involved in cell signaling.<sup>8</sup> [White PJ](#). Calcium channels in higher plants. *Biochim Biophys Acta* 2000; 1465:171 - 218

In the present study, rapid and reversible  $V_m$  depolarization observed upon perfusion of Lima bean mesophyll cells with polyamines was found to be significantly increased when JA was added to Cad and Put. The reversibility of the  $V_m$  may be linked to the overall physico-chemical amphiphilic properties of polyamines, probably depending on non covalent interaction with plasma membrane molecules, as polyamines occur in plants in free form, bound electrostatically to negatively charged molecules, and conjugated to small molecules and proteins.<sup>9</sup> [Martin-Tanguy J](#). Conjugated polyamines and reproductive development: biochemical, molecular and physiological approaches. *Physiol Plantarum* 1997; 100:675 - 688

Liu et al.<sup>10</sup> [Liu K](#), [Fu H](#), [Bei Q](#), [Luan S](#). Inward potassium channel in guard cells as a target for polyamine regulation of stomatal movements. *Plant Physiol* 2000; 124:1315 - 1326 showed that Spm, Spd, Cad and Put strongly inhibited opening and closing of stomata in *Vicia faba*, suggesting that polyamines target inward potassium channels in guard cells and modulate stomatal movements, so providing a link between abiotic stress, polyamine levels and stomatal regulation. Moreover, the transport of polyamines across the plasma membrane of plant cells is energy-dependent and calcium is involved in the uptake mechanism.<sup>1</sup> [Walters DR](#). Polyamines in plant-microbe interactions. *Physiol Mol Plant*

Pathol 2000; 57:137 - 146 ,[11 Antognoni F, Pistocchi R, Casali P, Bagni N](#). Does calcium regulate polyamine uptake in carrot protoplasts?. Plant Physiol Biochem 1995; 33:701 - 702 Both mechanisms can be correlated to the observed  $V_m$  depolarization, and the positive correlation between intracellular  $Ca^{2+}$  concentration [5 Ozawa R, Berthea CM, Foti M, Narayana R, Arimura G-I, Muroi A](#), et al. Exogenous polyamines elicit herbivore-induced volatiles in Lima bean leaves: involvement of calcium,  $H_2O_2$  and jasmonic acid. Plant Cell Physiol 2009; 50:2183 - 2199 and  $V_m$  depolarizing activity of polyamines confirms the involvement of  $Ca^{2+}$  during polyamine uptake.[11 Antognoni F, Pistocchi R, Casali P, Bagni N](#). Does calcium regulate polyamine uptake in carrot protoplasts?. Plant Physiol Biochem 1995; 33:701 - 702

#### Acknowledgements

This work was supported by: Grant-in-Aid for Scientific Research (S) (No. 19101009 to J.T.); Global COE Program A06 of Kyoto University; CEBIOVEM (Centre of Excellence for Plant and Microbial Biosensing of the University of Turin to M.E.M.).

#### Figures and Tables

Figure 1 Effect of 1 mM polyamines (arrow) on the  $V_m$  of Lima bean palisade cells. Spermine (Spm) caused the lowest  $V_m$  depolarization, whereas spermidine (Spd) showed the highest values of  $V_m$  depolarization. intermediate values were found when putrescine (Put) and cadaverine (cad) were perfused. after washing the tissues with fresh buffer (double arrow)  $V_m$  was always hyperpolarized, however the initial potential was recovered only for Put, while for all other polyamines the  $V_m$  never reached the initial values. Metric bars indicate standard deviation.





Figure 2 Effect of 1 mM polyamines + 0.1 mMJA (arrow) on the  $V_m$  of Lima bean palisade cells. the perfusion with Ja did not cause any variation in the  $V_m$ . addition of JA to Spm and Spd caused the same  $V_m$  depolarization observed in the absence of JA, whereas when JA was added to Put and Cad a stronger and significantly different  $V_m$  depolarization was observed. even in this case washing the tissues with fresh buffer (double arrow) caused a  $V_m$  hyperpolarized, however in this

case Spd reached  $V_m$  values significantly more negative than the initial  $V_m$ . Metric bars indicate standard deviation. For abbreviations see [Figure 1](#).



## References

- Walters DR. Polyamines in plant-microbe interactions. *Physiol Mol Plant Pathol* 2000; 57:137 - 146
- Walters DR. Polyamines and plant disease. *Phytochemistry* 2003; 64:97 - 107
- Martin-Tanguy J. Metabolism and function of polyamines in plants: recent developmentn (new approaches). *Plant Growth Regul* 2001; 34:135 - 148
- Bouchereau A, Aziz A, Larher F, Martin-Tanguy J. Polyamines and environmental challenges: recent development. *Plant Sci* 1999; 140:103 - 125
- Ozawa R, Berteau CM, Foti M, Narayana R, Arimura G-I, Muroi A, et al. Exogenous polyamines elicit herbivore-induced volatiles in Lima bean leaves: involvement of calcium,  $H_2O_2$  and jasmonic acid. *Plant Cell Physiol* 2009; 50:2183 - 2199
- Maffei M, Mithöfer A, Boland W. Before gene expression: early events in plant-insect interactions. *Trends Plant Sci* 2007; 12:310 - 316
- Maffei M, Bossi S, Spitfeller D, Miltho'fer A, Boland W. Effects of feeding *Spodoptera littoralis* on Lima beans leaves I. Membrane potentials, intracellular calcium variations, oral secretions and regurgitate components. *Plant Physiol* 2004; 134:1752 - 1762
- White PJ. Calcium channels in higher plants. *Biochim Biophys Acta* 2000; 1465:171 - 218

Martin-Tanguy J. Conjugated polyamines and reproductive development: biochemical, molecular and physiological approaches. *Physiol Plantarum* 1997; 100:675 - 688

Liu K, Fu H, Bei Q, Luan S. Inward potassium channel in guard cells as a target for polyamine regulation of stomatal movements. *Plant Physiol* 2000; 124:1315 - 1326

Antognoni F, Pistocchi R, Casali P, Bagni N. Does calcium regulate polyamine uptake in carrot protoplasts?. *Plant Physiol Biochem* 1995; 33:701 - 702