

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cytotoxic response in human lung epithelial cells and ion characteristics of urban-air particles from Torino, a northern Italian city.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/144095> since 2015-09-07T13:05:47Z

Published version:

DOI:10.1007/s11356-013-2468-1

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Luca Alessandria; Tiziana Schilirò; Raffaella Degan; Deborah Traversi; Giorgio Gilli.. Cytotoxic response in human lung epithelial cells and ion characteristics of urban-air particles from Torino, a northern Italian city.. ENVIRONMENTAL SCIENCE AND POLLUTION RESEARCH INTERNATIONAL. 21 (8) pp: 5554-5564.
DOI: 10.1007/s11356-013-2468-1

The publisher's version is available at:

<http://link.springer.com/content/pdf/10.1007/s11356-013-2468-1>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/144095>

Cytotoxic response in human lung epithelial cells and ion characteristics of urban-air particles from Torino, a northern Italian city

Luca Alessandria ¹

Tiziana Schilirò ^{1,2,*}

Phone +39-011-6705820

Fax +39-011-6705874

Email tiziana.schiliro@unito.it

Raffaella Degan ¹

Deborah Traversi ¹

Giorgio Gilli ¹

¹ Department of Public Health and Pediatrics, University of Torino, Piazza Polonia, 94, 10126 Torino, Italy

² Torino,

Abstract

Recently, much attention has been devoted to urban air pollution because epidemiological studies have reported health impacts related to particulate matter (PM). PM₁₀ and PM_{2.5} were collected during different seasons in Torino, a northern Italian city, and were characterised by inorganic chemical species (secondary particulates and bio-available iron). The biological effects of aqueous and organic solvent PM extracts on human epithelial lung A549 were evaluated, and the effects on cell proliferation and lactate dehydrogenase (LDH) release were assayed. The average PM₁₀ concentration during the sampling period was $47.9 \pm 18.0 \mu\text{g}/\text{m}^3$; the secondary particles accounted for $49 \% \pm 9 \%$ of the PM₁₀ total mass, and the bio-available iron concentration was $0.067 \pm 0.045 \mu\text{g}/\text{m}^3$. The PM_{2.5}/PM₁₀ ratio in Torino ranged from 0.47 to 0.90 and was higher in cold months than in warm months. The PM₁₀ and PM_{2.5} extracts inhibited cell proliferation and induced LDH release in a dose-dependent manner with a seasonal trend. The PM₁₀ extract had a stronger effect on LDH release, whereas the PM_{2.5} extract more strongly inhibited cell proliferation. No significant differences were observed in the effects induced by the two extracts, and no significant correlations were found between the biological effects and the PM components evaluated in this study, thus emphasising the importance of the entire mixture in inducing a cytotoxic response.

Keywords

PM
A549
LDH
Metals
Secondary particulate
Cytotoxicity

Responsible editor: Philippe Garrigues

Introduction

Particulate matter (PM) is an air-suspended mixture of solid and liquid particles that vary in number, size, shape, surface area, chemical composition, solubility and origin (Pope and Dockery [2006](#)); PM can be classified as PM10 (particles with an aerodynamic diameter \varnothing less than 10 μm), PM2.5 ($\varnothing < 2.5 \mu\text{m}$), and PM1 ($\varnothing < 1 \mu\text{m}$). Currently, major sources of ambient PM include vehicular and industrial emissions, power plants, refuse incineration and geological material. The fine fractions (PM2.5 and PM1) are dominated by combustion-derived particles, whereas the coarse fraction (PM10) contains a major component of mineral compounds and certain adsorbed endotoxins. Anthropogenic contributions, i.e. organic carbon, are greater in urban environments, and specific metals may also be more prevalent in these areas. In an urban setting, approximately 40 % of the particle mass can be attributed to fossil fuel use (Ghio et al. [2012](#)).

The respiratory system is the primary target of airborne particles, which are inhaled and tend to accumulate in the airways. The exposure dose in the tissues depends on the PM atmospheric concentration, the deposition rate in the airways, the clearance mechanisms and the subsequent retention of particles within the respiratory tract. The epidemiological associations are well established between PM mass concentrations and increased human mortality and morbidity (Brook et al. [2010](#), Pope and Dockery [2006](#)). Currently, the hypothesis that long-term exposure to air pollution can produce human morbidity and mortality is unanimously accepted, and epidemiological evidence suggests an increased risk of lung cancer for people living in urban areas (Nerriere et al. [2005](#), Sun et al. [2008](#)). Long-term exposure to air pollution PM increases the risk of lung cancer, respiratory diseases and arteriosclerosis, whereas short-term exposure can exacerbate several forms of respiratory diseases, i.e. bronchitis and asthma, and changes in heart rate variability (Dominici et al. [2007](#); Peacock et al. [2011](#); Pope et al. [2009](#); Raaschou-Nielsen et al. [2011](#); Rusconi et al. [2011](#)). However, fundamental uncertainties still exist with respect to the underlying mechanisms of toxicity that are responsible for the mortality or morbidity increases following exposure to current levels of air pollution PM (Schwarze et al. [2007](#)).

The PM-induced human health effects are still not completely understood, but oxidative stress and inflammatory reaction seem to be of fundamental importance (Baulig et al. [2009](#). Garcon et al. [2006](#). Lodovici and Bigagli [2011](#). Lonkar and Dedon [2011](#)). Experimental and epidemiological data indicate that lung disorders are often associated with a pro-oxidant/antioxidant imbalance and inflammatory reaction, and there is increasing evidence that air pollution via PM induces acute responses and exacerbates existing

inflammatory diseases in the lungs (Maestrelli et al. **2011**. Peacock et al. **2011**. Raaschou-Nielsen et al. **2011**. Rusconi et al. **2011**. Terzano et al. **2010**). Under physiological conditions, the normal production of reactive oxygen species (ROS) is counteracted by antioxidant scavengers and enzymes, which include both enzymes and non-enzymatic scavengers (de Kok et al. **2006**; Garcon et al. **2006**; Sun et al. **2008**). However, under abnormal conditions, excessive levels of ROS exceed the detoxification capacity of the antioxidant defences, thereby causing a change in the redox status of the cell (Jomova and Valko **2011**; Lonkar and Dedon **2011**; Ziech et al. **2011**). Oxidative stress subsequently triggers a cascade of events closely associated with inflammation, which is believed to play a key role in the air pollution/PM-induced development and/or exacerbation of acute and/or chronic lung diseases (Dagher et al. **2005**, **2006**, **2007**; Garcon et al. **2006**; Lodovici and Bigagli **2011**; Oh et al. **2011**; Schwarze et al. **2007**). Transition metals (i.e. Fe) (Carter et al. **1997**; Hutchison et al. **2005**; Roig et al. **2013**; Schins et al. **2004**), organics (i.e. polycyclic aromatic hydrocarbons or PAHs) (Billet et al. **2008**) and endotoxins (Becker et al. **2003**; Longhin et al. **2013**; Oberdorster et al. **2000**; Traversi et al. **2010**) in the PM seem to be involved in this process (Donaldson et al. **2003**; Schwarze et al. **2006**; Sorensen et al. **2003**). Moreover, although secondary inorganic aerosols show less toxic activity when tested under controlled laboratory conditions, epidemiological studies demonstrate significant associations between sulphates and nitrates and various health outcomes. In ambient air, this fraction may act as a carrier for other components or as a surrogate for PM emitted from the combustion of sulphur-containing fuels (Schwarze et al. **2006**).

The PM size is another important parameter that influences biological effects. The PM₁₀ fraction has been associated with pro-inflammatory and cytotoxic effects (Gualtieri et al. **2010**; Hetland et al. **2005**), and the PM_{2.5} fraction has been associated primarily with a higher genotoxic potential (Billet et al. **2008**) also related to higher PAH and metal contents. Moreover, positive relationships were found between the formation of ROS and the induction of DNA damage. Depending on the amount formed, ROS may result in necrosis as well as apoptosis (de Kok et al. **2006**; Gualtieri et al. **2011**).

The aim of the present study was to contribute to a better knowledge of the cytotoxic effects induced by PM extracts. The PM₁₀ and PM_{2.5} materials were sampled in traffic conditions and at an urban background station. The inhalable fraction of ambient particulate matter (PM₁₀) is widely used as an air quality indicator (European Commission (EC) Directive **2008**; WHO 2006a); moreover, PM₁₀ and PM_{2.5} pollution is specifically related to urban environments (Pelucchi et al. **2009**). The PM samples were analysed in terms of the amounts of inorganic species (bio-available iron and secondary particulates of sulphates and nitrates). Additionally, A549 cells were exposed to increasing concentrations of PM₁₀ and PM_{2.5} (aqueous and organic solvent) extracts, and the experiments were performed at concentrations according to previous data (Schilirò et al. **2010**). Next, the effects on cell proliferation and LDH release induced by extracts from cold and warm season samples were evaluated. Another aim of this work was to define whether the cytotoxic effects produced in the A549 cells by extracts of PM₁₀ and PM_{2.5} might be related to the chemical compounds evaluated on the particles.

The chemical and *in vitro* toxicological characterisations of urban-air PM could provide a basis for environmental and sanitary actions, particularly in one of the most industrialised areas of Italy (Hazenkamp-von Arx et al. **2003**; Marcazzan et al. **2003**), with high levels of particulate matter among European nations (WHO **2006b**).

Material and methods

PM sampling

Sampling was performed from January 2007 to December 2008 at two meteorological–chemical stations of the Environmental Protection Regional Agency (Piedmont A.R.P.A.) located in Torino, a city in the northwest of the Padana Plain (Italy). The first sampling site (station 1) was located outdoors in a small green area within an enclosure zone classified as urban background. The second sampling site was located in the centre of the city (station 2) in a traffic-regulated street and was classified as a traffic station (Putaud et al. **2010**). The PM₁₀ (PM passing through a size-selective inlet with a 50 % efficiency cut-off at 10 µm aerodynamic diameter) material was sampled at station 2 on glass micro-fibre filters (Type A/E, 8 × 10 in., Gelman Sciences, MI, USA) with a Sierra Andersen High-Volume Sampler 1200/VFC (Andersen Samplers, Atlanta, Georgia, USA) at a flow rate of 1.16 L/min. The sampling duration was controlled using a timer accurate to ±15 min over a 24-h sampling period. The PM_{2.5} was sampled at station 1 with glass micro-fibre filters (Type A/E 47 mm, Gelman Sciences, MI, USA) using a PM_{2.5} MicroVol 1100 Low-Volume Air Sampler with a flow rate of approximately 32 L/min. This sampler is certified in compliance with norm EN-14907. The exact flow rate was calculated daily and corrected for variations in atmospheric pressure and the actual differential pressure across the filter. The filters were conditioned for 48 h and were weighed with an analytical balance (±10 µg) before and after sampling to calculate the PM mass trapped on the filter. The procedures were conducted according to the European Committee for Standardization (CEN **1998**), as previously described (Gilli et al. **2007c**).

Particles extractions

The PM₁₀ filter strips were pooled to obtain one weekly sample, and particles were extracted as previously described (Schilirò et al. **2010**). In brief, each pool was treated with two different extraction media, acetone or RPMI1640 without foetal calf serum (FCS), which were chosen for different capabilities for the extraction of different compounds. Acetone was chosen as the solvent for organic-extractable compounds, and RPMI1640 without FCS was chosen to extract a water-soluble component theoretically comparable to the extraction in the lung cells (Schilirò et al. **2010**).

Each portion of the filter was cut into thin strips and placed in a 50-mL polypropylene sterile tube with 15 mL of the extraction medium. The tubes were placed in an ultrasonic water bath for 10 min, followed by 1 min of vortexing. This procedure was repeated three times. The samples were centrifuged at 5,000×g for 10 min to remove the filter material, and the supernatant was collected. Acetone extracts were evaporated with a rotary evaporator and re-suspended in culture medium. Aqueous extracts were directly assayed.

The PM_{2.5} filters were pooled to obtain one monthly sample, and the same extraction procedure was followed. Unless otherwise specified, all chemicals were purchased from Sigma, St. Louis, MO, USA.

Iron and bio-available iron

Iron determination was performed according to the procedure of Gilli et al. (2007b). In brief, the metals were extracted from the filter strips (3.18 cm by 20.32 cm) using a nitric acid solution. After cooling, the sample was mixed and centrifuged, and the trace element concentrations in each sample were determined by atomic absorption spectrometry (Varian GTA-96).

The bio-available iron from urban particulates was determined as previously described (Lund and Aust 1990) with certain modifications (Gilli et al. 2007b; Smith and Aust 1997). In brief, filter strips (3.18 cm by 20.32 cm) were suspended in 50 nM NaCl, mixed, and adjusted to pH 7.5. Citrate was added to the samples to a final concentration of 1 mM, and all samples were placed on a wrist-action shaker in the dark for 24 h. Samples of 1 mL were withdrawn and centrifuged to remove the particulates. The amount of iron mobilised as the citrate/Fe complex in the supernatant was determined using a spectrophotometric total non-heme iron assay (Brumby and Massey 1967).

Sulphates and nitrates

Sulphates and nitrates were measured according to the procedure of Gilli et al. (2007a), 2007b). In brief, each filter strip (3.18 cm by 20.32 cm) was extracted in 15 mL of distilled and deionised water via 30-min sonication, 30-min agitation, and overnight refrigeration at 4 °C. Prior to the analysis, the samples were centrifuged to remove particles. Ion chromatography was used to determine the soluble ion content of sulphate (SO_4^{2-}) and nitrate (NO_3^-) using a Dionex DX-100 ion chromatograph with 0.3 mM NaHCO₃ and 2.7 mM Na₂CO₃ 2.7 for the eluent and IonPac analytical column S12A for the anions. The applied standards, i.e. sodium sulphate and sodium nitrate (71959 and 71759 FLUKA, respectively), ranged from 0.1 to 100 g/ml. The ions were identified by their elution/retention times (~8.75 min and 12.5 min for nitrates and sulphates, respectively) and were quantified by the conductivity peak area or peak height (300A method, US EPA 1996).

Cell culture

The human lung epithelial cell line A549 (non-small cell lung cancer) from the Interlab Cell Line Collection (Genova, IT) was used as a model for human epithelial lung cells. Cells were grown as a monolayer and were maintained and treated in RPMI1640 supplemented with 10 % (v/v) FCS, 2 % L-glutamine (200 mM) and 1 % penicillin/streptomycin (10 mg/ml) at 37 °C in a humidified atmosphere containing 5 % CO₂.

Cell viability

The MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay was used to

measure cell viability (Gualtieri et al. 2010). Cells were seeded in 24-well plates at a density of 4×10^4 cells/well and exposed to PM extracts equivalent to particle concentrations of 50, 100 and 200 $\mu\text{g/ml}$ (Schilirò et al. 2010). After treatment with PM₁₀ and PM_{2.5}, the medium was discharged; cells were rinsed with PBS, and MTT (final concentration of 0.5 mg/ml) reagent was added for 4 h. The medium was removed, and the MTT reduction product (formazan crystals) was dissolved in 300 μl of DMSO. Cell proliferation was determined after 72 h of exposure by measuring the absorbance at 570 nm using a micro-plate reader (ELX 800 UV, Bio-Tek Instruments, Inc.). Blank filters were used as negative controls, and all experiments were performed in triplicate. The percentage of cell proliferation was calculated by comparing the absorbance of exposed cultures with the absorbance of non-exposed cultures.

LDH assay

To evaluate PM cytotoxicity, the released LDH activities from damaged cells were measured in cell-free culture supernatants as previously described (Schilirò et al. 2010). In brief, A549 cells were seeded in six-well plates at a density of 1×10^6 cells/well and exposed to PM₁₀ and PM_{2.5} extracts containing an equivalent of 50, 100 and 200 $\mu\text{g/ml}$ of particles. Blank filters were treated in the same manner. At 72 h, LDH activity was measured in the supernatant and cell lysate. The LDH activity, the ratio of extracellular LDH (measured in the supernatant) and total LDH (expressed as the sum of the LDH measured in the supernatant and cell lysate) were measured as consumption of NADH in terms of the absorbance at 340 nm, read using a micro-plate reader (Benchmark Plus Microplate Reader, Biorad). All experiments were performed in triplicate. The LDH activity of the exposed cells is expressed as a percentage of that of non-exposed cells.

Statistical analysis

Statistical analyses were performed using the SPSS Package (version 17.0) for Windows. The mean values were compared with the Wilcoxon–Mann–Whitney *U* test, and the Spearman rank correlation coefficient (*r*_S) was used to assess the relationships between variables. The mean difference and correlation were considered significant at $p < 0.05$.

Results

PM concentration

A total of 117 PM₁₀ filters and 362 PM_{2.5} filters were analysed during 2007 and 2008. The annual means for PM₁₀ and PM_{2.5} were 47.9 ± 18.0 and $35.2 \pm 22.7 \mu\text{g/m}^3$, respectively.

The PM_{2.5}/PM₁₀ ratio ranged from 0.47 to 0.90 and was higher in winter/autumn than in spring/summer. This ratio is affected by the different sampling sites in terms of traffic pressure but is comparable to those values reported in the literature (Querol et al. 2004).

The particulate matters often showed concentrations higher than the daily and annual targets proposed in the Air Quality Directive 2008/50/CE (EU, 2008). The highest values both for PM₁₀ and PM_{2.5} were observed in winter, whereas the lowest were

measured in summer (Figure 1). Moreover, significant differences ($p < 0.05$) were observed between the seasonal (autumn/winter vs. spring/summer) PM10 and PM2.5 concentrations (Table 1).

Table 1

PM10, PM2.5, sulphates, nitrates and Fe concentrations (micrograms per cubic metre), means \pm standard deviation

	PM10	PM2.5	SO ₄ ⁼	NO ₃ ⁻	SO ₄ ⁼	NO ₃ ⁻	Fe total
	($\mu\text{g}/\text{m}^3$)	($\mu\text{g}/\text{m}^3$)	($\mu\text{g}/\text{m}^3$)	($\mu\text{g}/\text{m}^3$)	(% PM10)	(% PM10)	($\mu\text{g}/\text{m}^3$)
Whole period	47.9 \pm 18.0	35.2 \pm 22.7	12.5 \pm 3.5	9.5 \pm 5.5	29.7 \pm 10.4	19.3 \pm 5.4	1.953 \pm 0.680
Winter/autumn	58.4 \pm 18.6*	52.7 \pm 19.8*	11.5 \pm 3.3	12.1 \pm 5.9	22.7 \pm 9.5*	20.5 \pm 5.9	2.085 \pm 0.954
Summer/spring	37.3 \pm 10.5*	17.7 \pm 2.8*	13.5 \pm 3.8	6.7 \pm 3.8	36.7 \pm 5.7*	18.2 \pm 5.1	1.821 \pm 0.254

* $p < 0.05$ level of statistically significant differences (autumn/winter vs. spring/summer)(Mann-Wh

Iron and bio-available iron

The total iron mean concentration was $1.953 \pm 0.680 \mu\text{g}/\text{m}^3$. The highest value was observed in winter ($4.263 \mu\text{g}/\text{m}^3$), whereas the lowest was observed in summer ($0.600 \mu\text{g}/\text{m}^3$). Table 1 also shows the mean concentrations of bio-available iron, and 3.9 ± 3.2 % of the total iron was bio-available. Statistically significant differences were observed for the mean seasonal concentrations ($p < 0.01$). The bio-available iron showed significant positive correlations with the PM10 and PM2.5 concentrations (respectively, $r_S = 0.711$, $p < 0.05$ and $r_S = 0.846$, $p < 0.01$).

Sulphates and nitrates


The mean concentration of sulphates was $12.5 \pm 3.5 \mu\text{g}/\text{m}^3$ relative to PM10, such that sulphates represented 29.7 ± 10.4 % of the total PM10 mass. The mean concentration of nitrates was $9.5 \pm 5.5 \mu\text{g}/\text{m}^3$, such that nitrates represented 19.3 ± 5.4 % of the total PM10 mass. Considering the sum of sulphates and nitrates as the principal components of secondary particulate matter, these materials accounted for 49 ± 9 % of the PM10 total mass (Table 1). No statistically significant differences were found for the seasonal concentrations for both sulphates and nitrates expressed as micrograms per cubic metre. Considering the concentrations of the two species relative to PM10 (percentage), it was observed that the sulphate content (36.7 ± 5.7 %) in the hot season was significantly elevated compared with that in the cold season (22.7 ± 9.5 %, $p < 0.05$) whereas, for the

nitrates, the amounts were similar in the two seasons.

As expected, the sulphates and nitrates showed significant correlations with the particulate matter concentrations. More specifically, the sulphates displayed a negative correlation with both PM10 and PM2.5, whereas the nitrates had a positive correlation ([Fig. 1](#)).

Fig. 1

Monthly trends of the PM10 (*top*) and PM2.5 (*bottom*) concentrations




Cell proliferation

Extracts from blank filters had no significant effects on cell proliferation. Figure 2 shows

the effects on cell proliferation of organic and aqueous extracts. The PM10 and PM2.5 extracts inhibited cell proliferation in a dose-dependent manner. The PM10 extracts induced a significant inhibition of cell proliferation only at the highest exposure dose (200 µg/ml), both for the organic and aqueous extracts. However, the PM2.5 extracts also had strong effects on cell proliferation (especially for the aqueous samples) at doses of 50 and 100 µg/ml. No seasonal (autumn/winter vs. spring/summer) differences were observed in the inhibition of proliferation of A549. Moreover, the chemical parameters were not correlated with the inhibition of cell proliferation.

Fig. 2

Inhibition of proliferation of A549 cells after incubation with PM10 (*top*) and PM2.5 (*bottom*) organic solvent extracts (acetone) and aqueous extracts (RPMI) (50, 100 and 200 µg/ml). Inhibition of cell proliferation was calculated by comparing the absorbance of the exposed cultures with the absorbance of non-exposed cultures. *Significant differences from the control, $p < 0.05$ (Mann–Whitney U test)




LDH release

An increase in extracellular LDH enzyme activity reflects an increase in the amount of membrane-damaged cells (Lobner 2000). The release of the cytoplasmic enzyme LDH into

the culture supernatant was used to measure the cytotoxicity of the PM10 extracts. Extracts from blank filters had no significant effect on LDH release. As shown in **Fig. Figure 3**, incubation of A549 cells with PM10 extracts at a concentration of 50 µg/ml had no effect on LDH release compared with control cells.

Fig. 3

LDH release from A549 cells after their incubation with PM10 (*top*) and PM2.5 (*bottom*) organic solvent extracts (acetone) and aqueous extracts (RPMI) (50, 100 and 200 µg/ml). LDH release was calculated by comparing the absorbance of exposed cultures with the absorbance of non-exposed cultures. *Statistically significant differences from the control, $p < 0.05$ (Mann–Whitney U test)



Statistically significant increases in extracellular LDH activities were observed after cell exposure at 100 and 200 µg/mL of particle concentration for both the organic and aqueous

extracts. Furthermore, the organic extracts showed seasonal differences (autumn/winter vs. spring/summer) in the induction of LDH release for the two highest concentrations of exposure.

The PM_{2.5} extracts induced a percentage release of LDH in exposed cells compared with control cells that always ranged between 15 % and 35 %, and these differences were significant only for three cases. No seasonal (autumn/winter vs. spring/summer) differences were found in the LDH release induced by the PM_{2.5} extracts.

The LDH increase in A549 due to either the organic or aqueous extract was not correlated with the PM₁₀ and PM_{2.5} concentrations or with any other parameters.

Discussion and conclusions

The current study focused on the chemical characterisation and the cytotoxic effects of particulate matter in the city of Torino. Moreover, in this work, we intended to determine whether the cytotoxic effects produced in A549 cells from the extracts of PM₁₀ and PM_{2.5} might be related to specific compounds evaluated on the particles.

The north of Italy, and in particular the Padana Plain where Torino is located, is an area of widespread air pollution (Cadum et al. 2009). The weak dispersion rate observed during winter due to the conformation of the territory represents a relevant factor (Poncino et al. 2009), and different air pollution indicators exceed the WHO guidelines, especially for particulate matter, i.e. PM₁₀ and PM_{2.5}. In addition, the exposure concentrations of 40 and 25 µg/m³ (for PM₁₀ and PM_{2.5}, respectively) defined by the European Directive 2008/50/EC (ECDirective 2008), which must be met by 2015, were clearly exceeded, and it is difficult to imagine better results for the upcoming years.

In contrast with our previous finding (Gilli et al. 2007a, b) but in agreement with the latest data from another city in the Padana Plain (Camatini et al. 2012), no significant seasonal differences were observed in total Fe concentrations. A small amount of iron is bio-available (approximately 4 % of total Fe), and this fraction is involved in oxidant generation after PM exposure by directly supporting electron transport, thus producing ROS (Ghio et al. 2012). Bio-available Fe showed a seasonal trend, with higher content of this metal in samples from the cold season compared with those from the warm seasons and was also correlated with the PM₁₀ and PM_{2.5} concentrations.

The secondary PM components represented approximately 49 % of the total PM₁₀ in term of mass, as confirmed by the findings reported on other urban sites (Lee and Hieu 2013) in the Italian Po valley, in particular in Gilli et al. (2007b) and Marcazzan et al. 2003). The seasonal trend of both sulphates and nitrates is normally due to the photochemical reactions that occur more frequently in the warm season.

Many other studies have investigated the cytotoxicity of PM in airway epithelial cells (Scapellato and Lotti 2007), and PM induces different biologic effects depending on the sampling site (Rosas Perez et al. 2007), size fraction (Alfaro-Moreno et al. 2002; Hetland

et al. 2004; Osornio-Vargas et al. 2003) sampling time (Frampton et al. 1999) and contaminants adsorbed on the particles (Baulig et al. 2003; Billet et al. 2007; Calcabrini et al. 2004; Frampton et al. 1999; Muller et al. 2006). In this study, PM10 and PM2.5 extracts inhibited cell proliferation and induced LDH release in a dose-dependent manner, similar to the results found in other studies on epithelial cells (Alfaro-Moreno et al. 2002; Osornio-Vargas et al. 2003; Schilirò et al. 2010). Significant seasonal trends were observed only with the organic PM10 extracts; otherwise, the effects produced by the samples from cold seasons were similar to those induced by the other samples. As previously observed (Schilirò et al. 2010), aqueous PM extracts seemed to have a greater effect on cell proliferation than organic-solvent (acetone) extracts, whereas organic extracts induced a greater release of LDH. Both types of PM10 extracts have a significant impact on cells, and it is likely that water-soluble components could cross the membranes and induce rapid inflammatory responses; organic-phase compounds still cross the membranes but have a more deleterious effect on the cells (e.g. LDH release). It is important to emphasise that PM10 extracts had a stronger effect on LDH release than PM2.5 extracts (47.38 % vs. 26.77 %, respectively), according to a similar result on the pro-inflammatory potential of PM10 compared with PM2.5 presented by numerous authors (Duvall et al. 2008; Gualtieri et al. 2008; Hetland et al. 2005). As reported in another study (Shang et al. 2013), combined MTT data and LDH data show that PM exposure does not produce cell death by damaging the cell membrane of A549 cells. For the aqueous PM2.5 extract, for example, the discrepancy between LDH release data and the reduction in viability detected by MTT may suggest that the impairment of viability (approximately 25-30 % cell death was observed at concentrations of 200 µg/mL; Fig. 2) may be due to the induction of the apoptotic cell death pathway in which the integrity of the plasma membrane is maintained. In addition, it should be noted that the MTT assay measures cellular metabolic activity, and thus, low MTT values might indicate reduced metabolic activity/mitochondrial damage or reduced cell proliferation.

Finally, no significant correlations were found between the biological effects and PM components (iron, bio-available iron and secondary particulates) evaluated in this study. These results confirm the results of other recent studies in which inconsistent associations were reported between both cell proliferation and LDH release and secondary particulates and metals for in vitro lung cell exposure (Happo et al. 2008, Roig et al. 2013). One possible explanation for the paucity of correlations could be that the biological end-points and the PM components were quantified in chemically separate fractions. Furthermore, the aqueous extracts did not reflect the activity of any particular PM10 chemical component but rather only highlighted the activity of physiologically soluble components. Moreover, the lack of correlations indicates that certain other chemical parameters should be assessed.

The results of the current study supported the assertion that short-term bioassays using human lung cells might be adequate for completing routine chemical analysis and for carrying out preliminary screening of the potential effects of PM-associated airborne pollutants. Our results must be further explored to identify the most dangerous PM fractions, even though the precise components responsible for cytotoxicity following exposure of cells to PM remain to be defined.

Acknowledgements

This study was financed by a Local Research Grant from the Piedmont University of Torino. The authors kindly thank Drs. M. Sacco, M. Maringo, F. Lollobrigida and M. Grosa of the Environmental Protection Regional Agency (Piedmont A.R.P.A.).

References

- Alfaro-Moreno E, Martinez L, Garcia-Cuellar C, Bonner JC, Murray JC, Rosas I, Rosales SP, Osornio-Vargas AR (2002) Biologic effects induced in vitro by PM10 from three different zones of Mexico City. *Environ Health Perspect* 110:715–720
- Baulig A, Sourdeval M, Meyer M, Marano F, Baeza-Squiban A (2003) Biological effects of atmospheric particles on human bronchial epithelial cells. Comparison with diesel exhaust particles. *Toxicol In Vitro* 17:567–573
- Baulig A, Singh S, Marchand A, Schins R, Barouki R, Garlatti M, Marano F, Baeza-Squiban A (2009) Role of Paris PM2.5 components in the pro-inflammatory response induced in airway epithelial cells. *Toxicology* 261:126–135
- Becker S, Soukup JM, Sioutas C, Cassee FR (2003) Response of human alveolar macrophages to ultrafine, fine, and coarse urban air pollution particles. *Exp Lung Res* 29:29–44
- Billet S, Garcon G, Dagher Z, Verdin A, Ledoux F, Cazier F, Courcot D, Aboukais A, Shirali P (2007) Ambient particulate matter (PM(2.5)): physicochemical characterization and metabolic activation of the organic fraction in human lung epithelial cells (A549). *Environ Res* 105:212–223
- Billet S, Abbas I, Goff JL, Verdin A, Andre V, Lafargue PE, Hachimi A, Cazier F, Sichel F, Shirali P, Garcon G (2008): Genotoxic potential of polycyclic aromatic hydrocarbons-coated onto airborne particulate matter (PM(2.5)) in human lung epithelial A549 cells. *Cancer Lett* 270:144-155
- Brook RD, Rajagopalan S, Pope CA, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong YL, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC, Whitsel L, Kaufman JD, Amer Heart Assoc Council E, Council Kidney Cardiovasc D, Council Nutr Phys Activity M (2010) Particulate matter air pollution and cardiovascular disease an update to the scientific statement from the American Heart Association. *Circulation* 121:2331–2378
- Brumby PE, Massey V (1967) Determination of nonheme iron, total iron, and copper. In: Estabrook RW, Pullman ME (eds) *Methods in enzymology*, vol 10. Academic Press, New York, pp 463–474
- Cadum E, Berti G, Biggeri A, Bisanti L, Faustini A, Forastiere F, Grp Collaborativo E

(2009) The results of EpiAir and the national and international literature. *Epidemiologia & Prevenzione* 33:113–119

Calcabrini A, Meschini S, Marra M, Falzano L, Colone M, De Berardis B, Paoletti L, Arancia G, Fiorentini C (2004) Fine environmental particulate engenders alterations in human lung epithelial A549 cells. *Environ Res* 95:82–91

Camatini M, Corvaja V, Pezzolato E, Mantecca P, Gualtieri M (2012) PM10-biogenic fraction drives the seasonal variation of proinflammatory response in A549 cells. *Environ Toxicol* 27:63–73

Carter JD, Ghio AJ, Samet JM, Devlin RB (1997) Cytokine production by human airway epithelial cells after exposure to an air pollution particle is metal-dependent. *Toxicol Appl Pharmacol* 146:180–188

CEN (1998) Air quality—determination of the PM10 fraction of suspended particulate matter – reference method and field test procedure to demonstrate reference equivalence of measurement methods. Brussels: European Committee for Standardization; (European Standard EN 12341)

Dagher Z, Garcon G, Gosset P, Ledoux F, Surpateanu G, Courcot D, Aboukais A, Puskaric E, Shirali P (2005) Pro-inflammatory effects of Dunkerque city air pollution particulate matter 2.5 in human epithelial lung cells (L132) in culture. *J Appl Toxicol* 25:166–175

Dagher Z, Garcon G, Billet S, Gosset P, Ledoux F, Courcot D, Aboukais A, Shirali P (2006) Activation of different pathways of apoptosis by air pollution particulate matter (PM2.5) in human epithelial lung cells (L132) in culture. *Toxicology* 225:12–24

Dagher Z, Garcon G, Billet S, Verdin A, Ledoux F, Courcot D, Aboukais A, Shirali P (2007) Role of nuclear factor-kappa B activation in the adverse effects induced by air pollution particulate matter (PM2.5) in human epithelial lung cells (L132) in culture. *J Appl Toxicol* 27:284–290

de Kok TM, Driee HA, Hogervorst JG, Briede JJ (2006) Toxicological assessment of ambient and traffic-related particulate matter: a review of recent studies. *Mutat Res* 613:103–122

Dominici F, Peng RD, Ebisu K, Zeger SL, Samet JM, Bell ML (2007) Does the effect of PM10 on mortality depend on PM nickel and vanadium content? A reanalysis of the NMMAPS data. *Environ Health Perspect* 115:1701–1703

Donaldson K, Stone V, Borm PJ, Jimenez LA, Gilmour PS, Schins RP, Knaapen AM, Rahman I, Faux SP, Brown DM, MacNee W (2003) Oxidative stress and calcium signaling in the adverse effects of environmental particles (PM10). *Free Radic Biol Med* 34:1369–1382

Duvall RM, Norris GA, Burke JM, Mcgee JK, Gilmour MI, Devlin RB (2008): Source apportionment of particulate matter in the US and associations with in vitro and in vivo lung inflammatory markers. *Air Pollution Modeling and Its Application XIX*, 721-722

European Commission (EC) Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. Technical Report 2008/50/EC, L152Off J Eur Comm; 2008

Frampton MW, Ghio AJ, Samet JM, Carson JL, Carter JD, Devlin RB (1999) Effects of aqueous extracts of PM(10) filters from the Utah valley on human airway epithelial cells. *Am J Physiol* 277:L960–L967

Garcon G, Dagher Z, Zerimech F, Ledoux F, Courcot D, Aboukais A, Puskaric E, Shirali P (2006) Dunkerque city air pollution particulate matter-induced cytotoxicity, oxidative stress and inflammation in human epithelial lung cells (L132) in culture. *Toxicol In Vitro* 20:519–528

[Ghio AJ, Carraway MS, Madden MC \(2012\): Composition of Air Pollution Particles and Oxidative Stress in Cells, Tissues, and Living Systems. *Journal of Toxicology and Environmental Health-Part B-Critical Reviews* 15, 1-21](#)

Gilli G, Traversi D, Rovere R, Pignata C, Schilirò T (2007a) Airborne particulate matter: ionic species role in different Italian sites. *Environ Res* 103:1–8

Gilli G, Pignata C, Schilirò T, Bono R, La Rosa A, Traversi D (2007b) The mutagenic hazards of environmental PM2.5 in Turin. *Environ Res* 103:168–175

Gilli G, Traversi D, Rovere R, Pignata C, Schilirò T (2007c) Chemical characteristics and mutagenic activity of PM10 in Torino, a northern Italian city. *Sci Total Environ* 385:97–107

Gualtieri M, Mantecca P, Cetta F, Camatini M (2008) Organic compounds in tire particle induce reactive oxygen species and heat-shock proteins in the human alveolar cell line A549. *Environ Int* 34:437–442

Gualtieri M, Ovrevik J, Holme JA, Perrone MG, Bolzacchini E, Schwarze PE, Camatini M (2010) Differences in cytotoxicity versus pro-inflammatory potency of different PM fractions in human epithelial lung cells. *Toxicol In Vitro* 24:29–39

Gualtieri M, Ovrevik J, Mollerup S, Asare N, Longhin E, Dahlman HJ, Camatini M, Holme JA (2011) Airborne urban particles (Milan winter-PM2.5) cause mitotic arrest and cell death: effects on DNA, mitochondria, AhR binding and spindle organization. *Mutat Res* 713(1–2):18–31

Happo MS, Hirvonen MR, Halinen AI, Jalava PI, Pennanen AS, Sillanpaa M, Hillamo R, Salonen RO (2008) Chemical compositions responsible for inflammation and tissue damage in the mouse lung by coarse and fine particulate samples from contrasting air

pollution in Europe. *Inhal Toxicol* 20:1215–1231

Hazenkamp-von Arx ME, Gotschi Fellmann T, Oglesby L, Ackermann-Liebrich U, Gislason T, Heinrich J, Jarvis D, Luczynska C, Manzanera AJ, Modig L, Norback D, Pfeifer A, Poll A, Ponzio M, Soon A, Vermeire P, Kunzli N (2003) PM_{2.5} assessment in 21 European study centers of ECRHS II: method and first winter results. *J Air Waste Manag Assoc* 53:617–628

Hetland RB, Cassee FR, Refsnes M, Schwarze PE, Lag M, Boere AJ, Dybing E (2004) Release of inflammatory cytokines, cell toxicity and apoptosis in epithelial lung cells after exposure to ambient air particles of different size fractions. *Toxicol In Vitro* 18:203–212

Hetland RB, Refsnes M, Cassee FR, Lag M, Dybing E, Schwarze PE (2005) Cytokine release from alveolar macrophages exposed to ambient particulate matter: relation to size, city, season and metal content. *Toxicol Lett* 158:S80–S81

Hutchison GR, Brown DM, Hibbs LR, Heal MR, Donaldson K, Maynard RL, Monaghan M, Nicholl A, Stone V (2005) The effect of refurbishing a UK steel plant on PM₁₀ metal composition and ability to induce inflammation. *Respir Res* 6:43

Jomova K, Valko M (2011) Advances in metal-induced oxidative stress and human disease. *Toxicology* 283:65–87

Lee B-K, Hieu NT (2013) Seasonal ion characteristics of fine and coarse particles from an urban residential area in a typical industrial city. *Atmospheric Research* 122:362–377

Lobner D (2000) Comparison of the LDH and MTT assays for quantifying cell death: validity for neuronal apoptosis? *J Neurosci Methods* 96:147–152

Lodovici M, Bigagli E (2011) Oxidative stress and air pollution exposure. *J Toxicol* 2011:487074

Longhin E, Pezzolato E, Mantecca P, Holme JA, Franzetti A, Camatini M, Gualtieri M (2013) Season linked responses to fine and quasi-ultrafine Milan PM in cultured cells. *Toxicol In Vitro* 27:551–559

Lonkar P, Dedon PC (2011) Reactive species and DNA damage in chronic inflammation: reconciling chemical mechanisms and biological fates. *Int J Cancer* 128:1999–2009

Lund LG, Aust AE (1990) Iron mobilization from asbestos by chelators and ascorbic acid. *Arch Biochem Biophys* 278:61–64

Maestrelli P, Canova C, Scapellato ML, Visentin A, Tessari R, Bartolucci GB, Simonato L, Lotti M (2011) Personal exposure to particulate matter is associated with worse health perception in adult asthma. *J Investig Allergol Clin Immunol* 21:120–128

Marcazzan GM, Ceriani M, Valli G, Vecchi R (2003) Source apportionment of PM₁₀ and PM_{2.5} in Milan (Italy) using receptor modelling. *Sci Total Environ* 317:137–147

Muller A, Wichmann G, Massolo L, Rehwagen M, Grabsch C, Loffhagen N, Herbarth O, Ronco A (2006) Cytotoxicity and oxidative stress caused by chemicals adsorbed on particulate matter. *Environ Toxicol* 21:457–463

Nerriere E, Zmirou-Navier D, Desqueyroux P, Leclerc N, Momas I, Czernichow P (2005) Lung cancer risk assessment in relation with personal exposure to airborne particles in four French metropolitan areas. *J Occup Environ Med* 47:1211–1217

Oberdorster G, Finkelstein JN, Johnston C, Gelein R, Cox C, Baggs R, Elder AC (2000): Acute pulmonary effects of ultrafine particles in rats and mice. *Res Rep Health Eff Inst*, 5-74; disc 75-86

Oh SM, Kim HR, Park YJ, Lee SY, Chung KH (2011) Organic extracts of urban air pollution particulate matter (PM_{2.5})-induced genotoxicity and oxidative stress in human lung bronchial epithelial cells (BEAS-2B cells). *Mutat Res* 723:142–151

Osornio-Vargas AR, Bonner JC, Alfaro-Moreno E, Martinez L, Garcia-Cuellar C, Ponce-de-Leon Rosales S, Miranda J, Rosas I (2003) Proinflammatory and cytotoxic effects of Mexico City air pollution particulate matter in vitro are dependent on particle size and composition. *Environ Health Perspect* 111:1289–1293

Peacock JL, Anderson HR, Bremner SA, Marston L, Seemungal TA, Strachan DP, Wedzicha JA (2011): Outdoor air pollution and respiratory health in patients with COPD. *Thorax*

Pelucchi C, Negri E, Gallus S, Boffetta P, Tramacere I, La Vecchia C (2009) Long-term particulate matter exposure and mortality: a review of European epidemiological studies. *BMC Public Health* 9:453

Poncino S, Bande S, Muraro M (2009) Meteo-diffusive analysis: a case-study of Turin. *Epidemiol Prev* 33:27–33

Pope CA 3rd, Dockery DW (2006) Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 56:709–742

Pope CA, Ezzati M, Dockery DW (2009) Fine-particulate air pollution and life expectancy in the United States. *New England J Med* 360:376–386

Putaud JP et al (2010) A European aerosol phenomenology-3: physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe. *Atmos Environ* 44:1308–1320

Querol X, Alastuey A, Ruiz CR, Artinano B, Hansson HC, Harrison RM, Buringh E, ten

Brink HM, Lutz M, Bruckmann P, Straehl P, Schneider J (2004) Speciation and origin of PM10 and PM2.5 in selected European cities. *Atmos Environ* 38:6547–6555

Raaschou-Nielsen O, Andersen ZJ, Hvidberg M, Jensen SS, Ketzel M, Sorensen M, Hansen J, Loft S, Overvad K, Tjonneland A (2011) Air pollution from traffic and cancer incidence: a Danish cohort study. *Environ Health* 10

Roig N, Sierra J, Rovira J, Schuhmacher M, Domingo JL, Nadal M (2013) In vitro tests to assess toxic effects of airborne PM(10) samples. Correlation with metals and chlorinated dioxins and furans. *Sci Total Environ* 443:791–797

Rosas Perez I, Serrano J, Alfaro-Moreno E, Baumgardner D, Garcia-Cuellar C, Martin Del Campo JM, Raga GB, Castillejos M, Colin RD, Osornio Vargas AR (2007) Relations between PM10 composition and cell toxicity: a multivariate and graphical approach. *Chemosphere* 67:1218–1228

Rusconi F, Catelan D, Accetta G, Peluso M, Pistelli R, Barbone F, Di Felice E, Munia A, Murgia P, Paladini L, Serici A, Biggeri A (2011) Asthma symptoms, lung function, and markers of oxidative stress and inflammation in children exposed to oil refinery pollution. *J Asthma* 48:84–90

Scapellato ML, Lotti M (2007) Short-term effects of particulate matter: an inflammatory mechanism? *Crit Rev Toxicol* 37:461–487

Schilirò T, Alessandria L, Degan R, Traversi D, Gilli G (2010) Chemical characterisation and cytotoxic effects in A549 cells of urban-air PM10 collected in Torino, Italy. *Environ Toxicol Pharmacol* 29:150–157

Schins RP, Lightbody JH, Borm PJ, Shi T, Donaldson K, Stone V (2004) Inflammatory effects of coarse and fine particulate matter in relation to chemical and biological constituents. *Toxicol Appl Pharmacol* 195:1–11

Schwarze PE, Ovreivik J, Lag M, Refsnes M, Nafstad P, Hetland RB, Dybing E (2006) Particulate matter properties and health effects: consistency of epidemiological and toxicological studies. *Hum Exp Toxicol* 25:559–579

Schwarze PE, Ovreivik J, Hetland RB, Becher R, Cassee FR, Lag M, Lovik M, Dybing E, Refsnes M (2007) Importance of size and composition of particles for effects on cells in vitro. *Inhal Toxicol* 19(1):17–22

Shang Y, Fan L, Feng J, Lv S, Wu M, Li B, Zang YS (2013) Genotoxic and inflammatory effects of organic extracts from traffic-related particulate matter in human lung epithelial A549 cells: the role of quinones. *Toxicol In Vitro* 27:922–931

Smith KR, Aust AE (1997) Mobilization of iron from urban particulates leads to generation of reactive oxygen species in vitro and induction of ferritin synthesis in

human lung epithelial cells. *Chem Res Toxicol* 10:828–834

Sorensen M, Autrup H, Moller P, Hertel O, Jensen SS, Vinzents P, Knudsen LE, Loft S (2003) Linking exposure to environmental pollutants with biological effects. *Mutat Res* 544:255–271

Sun JJ, Kim HJ, Seo HG, Lee JH, Yun-Choi HS, Chang KC (2008) YS 49, 1-(alpha-naphthylmethyl)-6,7-dihydroxy-1,2,3,4-tetrahydroisoquinoline, regulates angiotensin II-stimulated ROS production, JNK phosphorylation and vascular smooth muscle cell proliferation via the induction of heme oxygenase-1. *Life Sci* 82:600–607

Terzano C, Di Stefano F, Conti V, Graziani E, Petroianni A (2010) Air pollution ultrafine particles: toxicity beyond the lung. *Eur Rev Med Pharmacol Sci* 14:809–821

Traversi D, Alessandria L, Schilirò T, Chiado Piat S, Gilli G (2010) Meteo-climatic conditions influence the contribution of endotoxins to PM10 in an urban polluted environment. *J Environ Monit* 12:484–490

[U.S. EPA \(1996\): Air quality criteria for particulate matter. vol 1-3. EPA/600](#)

[/P-95/001a](#). WHO, World Health Organization (2006) Health risks of particulate matter from long-range transboundary air pollution. Joint WHO/Convention Task Force on the Health Aspects of Air Pollution, Copenhagen, Denmark

WHO, World Health Organization Regional Office for Europe (2006b) Health impact of PM10 and ozone in 13 Italian cities. Copenhagen, Denmark

Ziech D, Franco R, Pappa A, Panayiotidis MI (2011) Reactive oxygen species (ROS)—induced genetic and epigenetic alterations in human carcinogenesis. *Mutat Res* 711:167–173