

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effects of Subchronic Finasteride Treatment and Withdrawal on Neuroactive Steroid Levels and their Receptors in the Male Rat Brain

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1616906> since 2017-05-14T11:42:19Z

Published version:

DOI:10.1159/000442982

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Giatti, Silvia; Foglio, Benedetta; Romano, Simone; Pesaresi, Marzia; Panzica, Giancarlo; Garcia-Segura, Luis Miguel; Caruso, Donatella; Melcangi, Roberto C.. Effects of Subchronic Finasteride Treatment and Withdrawal on Neuroactive Steroid Levels and their Receptors in the Male Rat Brain. NEUROENDOCRINOLOGY. None pp: 1-12.
DOI: 10.1159/000442982

The publisher's version is available at:

<http://www.karger.com/Article/Pdf/442982>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

This full text was downloaded from iris - AperTO: <https://iris.unito.it/>

iris - AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

EFFECTS OF SUBCHRONIC FINASTERIDE TREATMENT AND WITHDRAWAL ON NEUROACTIVE STEROID LEVELS AND THEIR RECEPTORS IN THE MALE RAT BRAIN

Silvia Giatti^{1#}, Benedetta Foglio^{2#}, Simone Romano¹, Marzia Pesaresi¹, Giancarlo Panzica², Luis Miguel Garcia Segura³, Donatella Caruso¹, Roberto C. Melcangi^{1*}

¹Dipartimento di Scienze Farmacologiche e Biomolecolari, Università degli Studi di Milano, Milan, Italy; ²Department of Neuroscience, University of Turin, 10126, Turin, Italy and Neuroscience Institute Cavalieri Ottolenghi (NICO), 10143 Orbassano, Italy; ³Instituto Cajal, C.S.I.C., 28002 Madrid, Spain.

These two authors contributed equally to this work

Short title: Finasteride and nervous system

Keywords: neuroactive steroids, finasteride, 5alpha reductase, androgen receptor, estrogen receptor, GABA A receptor, brain, cerebrospinal fluid, plasma.

Key words: pregnenolone, progesterone, testosterone, metabolites, steroid receptor, GABA A receptor

Corresponding author:

*Roberto Cosimo Melcangi, Dipartimento di Scienze Farmacologiche e Biomolecolari Center of Excellence on Neurodegenerative Diseases, Università degli Studi di Milano, Italy, Via Balzaretti 9, 20133 Milano, Italy. Tel. +39 02 50318238; Fax: +39 02 50318204.

Email: roberto.melcangi@unimi.it



Abstract

The enzymatic conversion of progesterone and testosterone by the enzyme 5 α reductase exerts a crucial role in the control of nervous function. The effects in the brain of finasteride, an inhibitor of this enzyme used for the treatment of human benign prostatic hyperplasia and androgenic alopecia, have been poorly explored. Therefore, the effects of a subchronic treatment of finasteride at low doses (3mg/kg/day) and the consequences of its withdrawal on neuroactive steroid levels in plasma, cerebrospinal fluid and some brain regions as well as on the expression of classical and non classical steroid receptors have been evaluated in male rats. After subchronic treatment (i.e., 20 days) the following effects were detected: i) depending on the compartment considered, alteration in the levels of neuroactive steroids, not only in 5 α reduced metabolites but also in its precursors and in neuroactive steroids from other steroidogenic pathways and ii) upregulation of androgen receptor in the cerebral cortex and beta3 subunit of GABA A receptor in the cerebellum. One month after the last treatment (i.e., withdrawal period), some of these effects persisted (i.e., upregulation of androgen receptor in the cerebral cortex, increase of dihydroprogesterone in cerebellum, decrease of dihydrotestosterone in plasma). Moreover, other changes in neuroactive steroid levels, steroid receptors (i.e., upregulation of estrogen receptor alpha and downregulation of estrogen receptor beta in cerebral cortex) and GABA A receptor subunits (i.e., decrease of alpha 4 and beta 3 mRNA levels in cerebral cortex) were detected. These findings suggest that finasteride treatment may have broad consequences for brain function.



Introduction

Neuroactive steroids, such as progesterone (PROG) and testosterone (T), are converted by the enzyme 5 α reductase (5 α R) into dihydroprogesterone (DHP) and dihydrotestosterone (DHT), respectively. These metabolites are further converted in other steroids, such as tetrahydroprogesterone (THP) and isopregnanolone in the case of DHP and 5 α androstane 3 β ,17 β diol (3 β diol) or 5 α androstane 3 β ,17 β diol (3 β diol) in the case of DHT. These enzymatic conversions occur not only in the peripheral glands but also in the nervous system [1].

Three 5 α R isozymes, defined as type 1, 2 and 3, have been identified in the brain [2, 3], even if it is important to highlight that 5 α R type 3 is still under intensive investigation. At variance to type 1 and 2, type 3 is thought to play a role in protein glycosylation via dolichol phosphate pathway [3, 4]. It is still unclear whether specific isozymes are responsible for the conversion of T or PROG [2, 3]. The 5 α R inhibitor finasteride inhibits both type 1 and type 2 isozymes, although it has higher affinity for the isozyme type 2 [2, 3].

The enzymatic conversion mediated by 5 α R exerts a crucial role in the control of nervous function. Indeed, the metabolites of PROG and T are involved in the regulation of neuroendocrine events, behavior, affection, learning and memory, synaptic and glial plasticity and adult hippocampal neurogenesis as well as in the response of brain tissue to injury and neurodegeneration (i.e., regulating neuronal survival, axonal regeneration and gliosis) [1, 5–7]. Finasteride has been used as a tool to evaluate whether the effects of PROG or T in the nervous tissue could be ascribed to their conversion into 5 α reduced metabolites [2, 8–18]. However, despite of the wide therapeutic use of this inhibitor (e.g., human benign prostatic hyperplasia and androgenic alopecia), the effects of finasteride *per se* in the nervous system have been poorly explored. The effects of short exposures (i.e., few days) to high doses (i.e., 50mg/kg) of finasteride have been assessed in experimental models [2, 8–18]. However, the effects of subchronic treatments with finasteride at low doses, more similar to those used in clinical practice, and the consequences of its withdrawal have yet to be clarified.



Sexual dysfunction and anxious/depressive symptomatology have been reported in a subpopulation of male patients with androgenic alopecia that received finasteride [19–23]. These side effects were also reported in a subset of patients after discontinuation of the therapy [23–28]. Interestingly, these patients also showed altered neuroactive steroid levels in plasma and cerebrospinal fluid (CSF) in comparison to healthy individuals [29, 30]. Our hypothesis was that the treatment with low doses of finasteride may also cause alterations in neuroactive steroid levels in the central nervous system. Therefore, the aim of this study was to determine if a subchronic treatment with low doses of finasteride affects the levels of neuroactive steroids in the brain of male rats. In particular, since finasteride inhibits 5α R activity, we expected that the treatment with this drug would affect the levels of 5α R substrates and products. We also hypothesized that by the alteration of the levels of neuroactive steroids, finasteride would also affect the expression of steroid receptors in the brain. On this basis, we have here evaluated in male rats the effects of a subchronic treatment with low doses of finasteride (i.e., 3 mg/kg/day for 20 days) and the consequences of its withdrawal (i.e., evaluated 1 month after the last treatment) on neuroactive steroid levels in plasma, CSF, cerebral cortex, cerebellum and hippocampus. Neuroactive steroids assessed (Fig. 1) were: pregnenolone (PREG), PROG, DHP, THP, isopregnanolone, dehydroepiandrosterone (DHEA), T, DHT, 3 α -diol, 3 β -diol, 17 β -estradiol (17 β -E) and 17 α -estradiol (17 α -E). On the basis of the changes observed in the levels of neuroactive steroids, we analyzed possible changes of their receptors. To this aim, the expression of steroid receptors was evaluated in those brain structures in which neuroactive steroid levels were affected by finasteride. Thus, depending on the specific binding capability of the neuroactive steroids affected by finasteride, the expression of androgen (AR), estrogen (ER), progesterone (PR) receptor or GABA A receptor subunits (i.e., α 2, α 4, β 3, δ and γ 2) were assessed.



Material and Methods

Animals

Sprague Dawley male rats (150–175 g at the arrival, Charles River, Italy) were housed in the animal care facility of the Neuroscience Institute Cavalieri Ottolenghi (NICO) at the University of Torino. Animals were maintained in standard rat cages (cage size: 59.5 cm x 38.0 cm x 20.0 cm, Eurostandard type IV) with food and tap water available *ad libitum* and under controlled temperature ($21 \pm 4^\circ\text{C}$), humidity (40–60%), room ventilation (12.5 air changes per hour) and light cycle (12 h light / dark cycle; on 8 AM / off 8 PM).

Rats were acclimated to the new environment for ten days before being randomly assigned to one of the experimental groups described below. Animals were maintained according to the National Institutes of Health Guide for the Care and Use of Laboratory Animals. The experimental protocol was approved by the Bioethics and Animal Welfare Committee of the University of Torino and the procedures were performed in accordance with EU Council directive (2010/63/EU) and with the Italian DL 116/1992.

Treatments

Finasteride (3 mg/kg/day, Sigma Aldrich, Milan, Italy) was suspended in a vehicle solution of sesame oil and ethanol (5% v/v). Either this solution, or vehicle, was administered to the animals subcutaneously, at a volume of 100 μl /day for 20 days.

Finasteride and control rats were sacrificed at 24 hours and 1 month after last treatment in order to investigate the effects of the treatment and the effects of the withdrawal. Body weight and relative testis and prostate weights were assessed at the end of the treatment and at the end of the withdrawal period.



Liquid chromatography tandem mass spectrometry analysis

The cerebral cortex, cerebellum, hippocampus, plasma and CSF were collected and stored at 80°C until analyzed. In particular, blood samples were first collected in heparin tubes, then placed in centrifuge tubes and centrifuged at 2500 g for 15 min at 4°C to obtain plasma. CSF was obtained by puncture of *cisterna magna* as previously described [31]. Extraction and purification of the samples were performed according to Caruso and colleagues [32].

Briefly, samples were spiked with $^{13}\text{C}_3$ 17 β E (1 ng/sample), D₉ PROG (0.2 ng/sample) and D₄ PREG (5 ng/sample), as internal standards and homogenized in 2 ml of MeOH/acetic acid (99:1 v/v) using a tissue lyser (Qiagen, Italy). After an overnight extraction at 4°C, samples were centrifuged at 12,000 rpm for 5 min and the pellet was extracted twice with 1 ml of MeOH/acetic acid (99:1 v/v). The organic phases were combined and evaporated to dryness. The organic residues were resuspended with 3 ml of MeOH/H₂O (10:90 v/v) and passed through SPE cartridges, previously activated with MeOH (5ml) and MeOH:H₂O 1:9 v/v (5ml). The steroids were eluted in MeOH, concentrated and transferred in autosampler vials before liquid chromatography tandem mass spectrometry (LC MS/MS) analysis. Quantitative analysis was performed on the basis of calibration curves daily prepared and analyzed as previously described [32]. Briefly, blank samples (6% albumin in PBS) or cortex homogenate (obtained by control rats) were spiked with $^{13}\text{C}_3$ 17 β E (1 ng/sample), D₉ PROG (0.2 ng/sample) and D₄ PREG (5 ng/sample), as internal standards. Increasing amounts (0.05–5 ng/sample) of each steroid were added. Calibration curves were extracted and analyzed as described for experimental samples.

Positive atmospheric pressure chemical ionization (APCI+) experiments were performed with a linear ion trap mass spectrometer (LTQ, ThermoElectron Co, San Jose, CA, USA) using nitrogen as sheath, auxiliary and sweep gas. The instrument was equipped with a Surveyor liquid chromatography (LC) Pump Plus and a Surveyor Autosampler Plus (ThermoElectron Co, San Jose, CA, USA). The mass spectrometer (MS) was employed in tandem mode (MS/MS) using helium as collision gas.



The LC mobile phases were previously described [32]. The Hypersil Gold column (100x3 mm, 3 µm; ThermoElectron Co, San Jose, CA, USA) was maintained at 40°C. Peaks of the LC MS/MS were evaluated using a Dell workstation by means of the software Excalibur® release 2.0 SR2 (ThermoElectron Co, San Jose, CA, USA). Samples were analyzed using the transitions previously reported [32].

Real time polymerase chain reaction (RT PCR)

Total RNA from tissues was extracted using the standard Trizol protocol based on the method developed by Chomczynski and Sacchi [33]. RNA was subsequently extracted in accordance with the manufacturer's protocol (Invitrogen, San Giuliano Milanese, Italy) and prepared by using Direct zol™ RNA MiniPrep kit (Zymo Research, Irvine, CA, USA). RNA was analyzed using a TaqMan quantitative real time PCR instrument (CFX96 real time system; Bio Rad Laboratories, Segrate, Italy) using the iScript™ one step RT PCR kit for probes (Bio Rad Laboratories, Segrate, Italy). Samples were run in 96 well formats in duplicate as multiplexed reactions with a normalizing internal control, 18s rRNA (Applera, Monza, Italy). Specific TaqMan MGB probes (Applera, Monza, Italy) were: 5 R (Rn00567064_m1); GABA A subunit 2 (Rn01413643_m1); GABA A subunit 4 (Rn00589846_m1); GABA A subunit 3 (Rn00567029_m1); GABA A subunit 1 (Rn01517017_m1); GABA A subunit 2 (Rn00788325_m1).

Western blotting

Tissues were homogenized on ice in excess of lysis buffer (phosphate buffered saline (PBS), pH 7.4, added with 1% Nonidet P 40) supplemented with protease cocktail inhibitor (Roche Diagnostic spa, Monza, Italy), with a tissue lyser II (Qiagen; Hilden, Germany). Homogenate was sonicated and centrifuged at 400 g at 4°C to remove particulate matter. The protein concentration of each sample was assayed relative to the bovine serum albumin standard according to the method of Bradford [34]. In the case of steroid receptors, equal amounts of each sample were solubilized in 0.1% sodium



decylsulphate (SDS) sample buffer, left for 5 min at 100°C, resolved on a 10% SDS polyacrylamide gel, and electroblotted overnight to a nitrocellulose membrane (Trans blot; Bio Rad, Milan, Italy). The membranes were blocked at room temperature in PBS with added 0.1% Tween 20 and 5% nonfat dried milk. Successively, the filters were cut and the top was incubated overnight at 4°C with rabbit polyclonal antibody against ER α and ER β (Santa Cruz, Heidelberg, Germany) or over three days at 4°C with either mouse monoclonal antibody against PR (Thermo Scientific; Rockford, IL, USA) or rabbit monoclonal antibody against AR (Abcam; Cambridge, UK). Primary antibodies were diluted in PBS buffer 0.1% Tween 20 – 3% nonfat dried milk. The filters were then washed for 1h and incubated with an anti mouse or an anti rabbit horseradish peroxidase–conjugated secondary antibody. In parallel, the bottom filters were incubated with a primary monoclonal anti tubulin antibody (Sigma Aldrich, St. Louis, MO, USA) diluted in PBS buffer 0.1% Tween 20–3% nonfat dried milk, as an internal control, then washed and incubated with an anti mouse horseradish peroxidase–conjugated secondary antibody. After washing the filters, bound antibodies were detected with the ECL method (Bio Rad, Milan, Italy). Chemiluminescent signals were acquired with a ChemiDoc™ XRS+ system (Bio Rad, Milan, Italy) and analyzed with Image Lab™ software version 3.0 (Bio Rad, Milan, Italy). The mean control value within a single experiment was set to 100 and all the other values were expressed as a percentage. Values of controls from different experiments were all within 10%.

Statistical analysis

The quantitative data obtained by the experiments were analyzed by inferential statistical analysis in accordance with the experimental protocols and the nature of the data (i.e. Student's t test). $P \leq 0.05$ was considered significant. Analyses were performed using GRAPHPAD PRISM, version 4.00 (GraphPad Inc, La Jolla, CA).



Results

Body weight and prostate and testis weights

At the sacrifice body weight, relative prostate and relative testis weights were assessed. Body weight [Control (n=8): 346.3 ± 9.8 g vs. Finasteride (n=8): 351.3 ± 8.3 g] and relative testis weight [Control (n=8): 0.8920 ± 0.0213 g vs. Finasteride (n=8): 0.8560 ± 0.0297 g] were unaffected by finasteride at the end of the treatment. On the contrary, a significant decrease of relative prostate weight was detected [Control (n=8): 0.1111 ± 0.0116 g vs. Finasteride (n=8): 0.0786 ± 0.0072 g; $p \leq 0.05$]. At the end of the withdrawal period, body weight [Control (n=8): 445.6 ± 11.9 g vs. Finasteride (n=8): 443.6 ± 13.6 g], relative prostate weight [Control (n=8): 0.1259 ± 0.0061 g vs. Finasteride (n=8): 0.1113 ± 0.0107 g] and relative testis weight [Control (n=8): 0.7434 ± 0.0148 g vs. Finasteride (n=8): 0.7071 ± 0.0196 g] were not significantly different between the two experimental groups.

Assessment of neuroactive steroid levels after subchronic treatment with finasteride

Table 1 shows neuroactive steroid levels, assessed by LC MS/MS, in the cerebral cortex, cerebellum, hippocampus, CSF and plasma of the animals treated with finasteride or vehicle. PREG levels in the CSF were significantly upregulated by finasteride. In contrast, PREG levels in plasma and in the brain structures analyzed were not significantly affected by the treatment. The levels of the PREG product (i.e., PROG) were increased in the hippocampus after finasteride treatment, but were not affected in the cerebral cortex, cerebellum, plasma and CSF. The levels of the first PROG metabolite (i.e., DHP) were significantly upregulated in the cerebellum by finasteride. In agreement with the upregulation of this 5 α R product, a significant increase of 5 α R gene expression was also detected in the same brain area [Control (n=5): 0.796±0.036 vs. Finasteride (n=5): 0.992±0.063, $p \leq 0.05$]. In the case of the further metabolites of PROG, only isopregnanolone levels were significantly affected by finasteride, with an increase in the cerebellum and plasma (Table 1).

The levels of DHEA and its metabolites (i.e., androgens and estrogens) were also affected by finasteride. As reported in Table 1, DHEA levels in the cerebral cortex were significantly decreased in



finasteride treated rats. DHT levels (i.e., metabolite of T) were reduced in plasma, but not in the CSF and brain tissue after finasteride treatment. In the case of the further metabolites of DHT (i.e., 3 α -diol and 3 β -diol), their levels were significantly decreased and enhanced, respectively, in the cerebellum and CSF of finasteride treated animals. In the case of estrogens, only the levels of 17 β -E were significantly affected by finasteride. Indeed, the levels of this T metabolite were significantly increased in the cerebellum and plasma of finasteride treated rats.

Assessment of neuroactive steroid levels at the end of the withdrawal period

Table 2 shows neuroactive steroid levels, assessed by LC MS/MS, in the cerebral cortex, cerebellum, hippocampus, CSF and plasma of the animals treated for 20 days with finasteride or vehicle and sacrificed 1 month after the last treatment. Finasteride withdrawal induced a different regulation of PREG levels in the cerebellum and the hippocampus. Thus, compared to control animals an upregulation of PREG levels was observed in the cerebellum, while in the hippocampus this neuroactive steroid was decreased at the end of the withdrawal period. PROG levels were significantly decreased in the cerebral cortex, hippocampus and plasma after withdrawal. Finasteride withdrawal also affected the levels of the PROG metabolite DHP, which were decreased in the cerebral cortex and upregulated in the cerebellum and hippocampus compared to control animals (Table 2). In agreement with these changes, 5 α -R gene expression was significantly decreased in the cerebral cortex [Control (n=5): 1.428 \pm 0.105 vs. Finasteride (n=5): 1.038 \pm 0.093, $p \leq 0.05$] and increased in the cerebellum [Control (n=5): 1.722 \pm 0.076 vs. Finasteride (n=5): 2.192 \pm 0.145, $p \leq 0.05$]. The levels of the further metabolites of DHP were also affected (Table 2). In particular, the levels of THP in the cerebral cortex and plasma and the levels of isopregnanolone in the cerebral cortex were decreased after finasteride withdrawal.

The levels of DHEA and its metabolites (i.e., androgens and estrogens) were also affected at the end of the withdrawal period. Thus, as reported in Table 2, DHEA levels were significantly decreased in the cerebral cortex and upregulated in CSF. In the hippocampus, a tendency to decrease, which



however did not reach statistical significance ($p=0.07$), was observed. T levels were significantly increased in the cerebellum as well as in CSF, while the levels of its first metabolite (i.e., DHT) were significantly decreased not only in the cerebellum and CSF but also in plasma. The levels of the further metabolites of DHT were also affected by finasteride withdrawal. In particular, the levels of 3 α -diol and 3 β -diol were significantly decreased in plasma and the cerebellum, respectively. In case of estrogens, only the levels of 17 β -E were significantly affected by finasteride withdrawal. The levels of this neuroactive steroid were significantly reduced in the cerebral cortex and upregulated in the CSF. In the hippocampus, a tendency to decrease, which however did not reach the statistical significance ($p=0.063$), was observed.

Effects of the subchronic treatment with finasteride and its withdrawal on the expression of classical and non classical steroid receptors

On the basis of the observed changes in the levels of neuroactive steroids, we evaluated whether finasteride treatment and its withdrawal did also affect the expression of classical (i.e., AR, PR, ER α , ER β) and non classical (GABA A receptor $\alpha 2$, $\alpha 4$, $\beta 3$, δ and $\gamma 2$ subunits) steroid receptors. These receptors were analyzed only in the brain regions in which changes in corresponding neuroactive steroid levels were detected.

As reported in Figure 2 panel A, after finasteride treatment we observed a significant upregulation of AR protein levels in the cerebral cortex. On the contrary, the expression levels of PR (panel B), ER α (panel C), and ER β (panel D) in cerebellum as well as PR (panel E) in the hippocampus were unaffected. As reported in Figure 3, at the end of the withdrawal period the protein levels of AR (panel A) and ER α (panel C) were upregulated in the cerebral cortex while a downregulation of ER β (panel D) was observed. The expression levels of PR in the cerebral cortex (panel B), the cerebellum (panel E) and the hippocampus (panel G) as well as of AR in the cerebellum (panel F) were unaffected.



After finasteride treatment, an increase in the gene expression of the $\alpha 3$ subunit of the GABA A receptor was observed in the cerebellum (Figure 4). At the end of the withdrawal period the mRNA levels of the $\alpha 4$ and $\alpha 3$ subunits of the GABA A receptor in the cerebral cortex were decreased (Figure 5).

Discussion

In this study we have assessed whether a subchronic treatment with a low dose of finasteride was able to affect neuroactive steroid levels and their receptors in the brain of male rats. Indeed, to our knowledge the effects of finasteride have been generally explored so far after the administration of very high doses [2, 8 18]. We demonstrated that, in agreement with the well known effect of finasteride in the prostate [35], a significant decrease in its relative weight was observed at the end of the subchronic treatment with the drug. In addition, subchronic treatment with finasteride did also affect neuroactive steroid levels and expression of classical and non classical steroid receptors in the brain.

Since finasteride inhibits 5α R activity, we hypothesized that the subchronic treatment with this drug would affect the levels of 5α R substrates, PROG and T and the levels of the 5α R products, DHP and DHT. As mentioned above, T and PROG may be converted by different 5α R isozymes in several brain areas [2, 3]. For instance, type 1 is expressed in olfactory bulb [36, 37], cerebellum [37, 38], prefrontal and cerebral cortex [37, 39, 40], hippocampus, striatum, thalamus and amygdala [37]. As recently observed, enzyme type 2 shows a very similar brain distribution [41]. However, a possible difference may be the cellular localization of these two isozymes. Indeed, type 1 is typically localized in glial cells [1] while type 2 is consistently expressed in neurons [41]. Therefore, this dichotomy may reflect differential roles in the modulation of brain function. Finasteride did not significantly affect the levels of PROG and T in plasma, CSF and brain tissues, with the exception of the hippocampus, in which, as expected, the 5α R inhibitor caused a significant increase in the levels of PROG, the substrate of the enzyme. However, the increase in the levels of PROG was not accompanied by a



decrease in DHP levels. Concerning the 5α R products, we detected a decrease in DHT levels in plasma after finasteride treatment, in agreement to what is observed in patients treated with this drug [42]. However, DHT levels in the CSF and brain tissue were not affected by the subchronic finasteride treatment. Surprisingly, in the cerebellum finasteride treatment resulted in a significant increase in DHP levels. This was unexpected since DHP is a product of 5α R. The cause for this increase in DHP levels remains unexplained. Finasteride treatment did not significantly affect DHP levels in the other brain areas considered. In general, we may conclude that the subchronic treatment with a low dose of finasteride do not have major effects on the levels of the 5α R substrates and products in the brain of male rats but it is only able to affect peripheral DHT production. Indeed, in this context it is important to recall that in periphery 5α R type 1 gene is predominantly expressed in the skin (e.g., pubic skin and scalp) while 5α R type 2 is mainly expressed in the prostate and gonads [1]. Whether the effect we observed might be due to an impairment of a specific isozyme is still unclear. Indeed, even if with a different affinity, both isozymes are able to convert T into DHT [1].

However, finasteride caused important changes in the levels of other neuroactive steroids, such as isopregnanolone, DHT and 17β E in plasma, PREG, 3α diol and 3β diol in the CSF, DHEA in the cerebral cortex and isopregnanolone, 3α diol, 3β diol and 17β E in the cerebellum. These changes, involving secondary metabolites of PROG and T, as well as PROG and T precursors, show regional specificity and suggest that 5α R inhibition impact on other steroidogenic steps. Additional studies should determine whether subchronic finasteride treatments affect the expression or activity of other steroidogenic enzymes and alternative steroidogenic pathways. For instance, it has been reported that 20α reduction of PROG is enhanced in the brain when 5α reduction is inhibited by finasteride [43]. Furthermore, changes in neurotransmission caused by modifications in steroids with activity on GABA A receptor may affect general steroidogenic activity. For instance, aromatase activity and therefore 17β E synthesis is regulated by excitatory neurotransmission [44]. Interestingly, 17β E levels are increased in the cerebellum in parallel to increased levels of



isopregnanolone. Since isopregnanolone antagonizes the effects of THP on the GABA A receptor [1, 45, 46], the altered isopregnanolone levels may cause a modification in the activity of GABA A and consequently in the balance in excitation/inhibition.

The analysis of neuroactive steroids one month after finasteride withdrawal revealed long term consequences. In plasma, we detected a significant decrease in the levels of PROG, THP, DHT and 3 α -diol compared to control animals. In the CSF, the levels of T and 17 β -E were increased, while the levels of DHT were decreased, suggesting an increased accumulation of T and an increased conversion of T in 17 β -E, due to the inhibition of the conversion of T in DHT. Interestingly, post finasteride patients with persistent symptomatology showed significant decreased THP levels in plasma, as well as increased T levels in CSF that were associated with a decrease in DHT levels [30]. Stronger effects at the end of the withdrawal period were detected in the cerebral cortex, in which the levels of PROG, DHP, isopregnanolone, THP, DHEA and 17 β -E were significantly decreased. In the cerebellum, the levels of PREG, DHP and T were increased while the levels of DHT and 3 α -diol were decreased. In the hippocampus, the levels of PREG and PROG were decreased while the levels of DHP were increased. These findings indicate that finasteride treatment has long term effects in the levels of neuroactive steroids, even after a long period of withdrawal. In fact, the effects of finasteride withdrawal were stronger than the effects of finasteride treatment *per se*, in particular in brain regions like for instance the cerebral cortex and the hippocampus. This suggests that the steroidogenic system is able to partially adapt and compensate for the chronic inhibition of 5 α -R activity. However, the system, after this adaptation, may be less flexible to compensate for a new alteration and could be further disrupted when 5 α -R activity is recovered.

Changes in the levels of neuroactive steroids could imply changes in the expression of their receptors. Therefore, we hypothesized that finasteride treatment and its withdrawal could also affect the expression of AR (i.e., the receptor able to bind DHEA, T and DHT), PR (i.e., the receptor able to bind PROG and DHP) ER α and ER β (i.e., the receptor isoforms able to bind estrogens) and/or GABA A receptor α 2, α 4, β 3, δ and γ 2 subunits (i.e., the receptor able to bind THP and 3 α -diol). This



hypothesis has been confirmed, but only in some cases. For instance, in the cerebral cortex, in which we detected decreased levels of DHEA immediately after finasteride treatment and at the end of the withdrawal period, we detected an upregulation of AR expression. Even if the mechanism of action of DHEA in the nervous system has not yet been fully characterized, some observations suggest an interaction with AR [47–49]. It is interesting to note that the upregulation of AR has been also reported in the prostate of patients treated with finasteride for benign prostatic hyperplasia [50] as well as in the prepuce of patients treated with the drug for male androgenetic alopecia and showing persistent side effects [51]. In addition, in the cerebral cortex, the decrease in the levels of 17β -E after finasteride withdrawal was associated with the upregulation of ER α . In contrast, the expression of ER β was decreased. This finding suggests that the expression of ER α and ER β is regulated by different mechanisms in the cerebral cortex. Indeed, a differential regulation of ER α and ER β in the cerebral cortex has been previously observed in aged male rats [52]. In addition, low sex steroid levels were associated with high expression of classical steroid receptors (i.e., AR and ER) in the brain of tropical male passerine birds [53]. In the brain of rodents, modifications in 17β -E levels are known to affect the expression and localization of ERs with regional specificity [54–56]. Moreover, it is interesting to note that finasteride treatment induces intersex animals in Western clawed frogs showing high expression in the brain of AR and ER [57].

On the contrary, as we here reported for the cerebellum at the end of the withdrawal period, changes in the levels of other ligands of AR, such as T or DHT, were not associated with a significant change in the expression of this steroid receptor, suggesting a different regional regulation of AR by the neuroactive steroid environment. Regional regulation of AR is also suggested by a previous study showing that aging induced a decreased expression of brain androgen synthesis enzymes in both the cerebral cortex and cerebellum, but the expression of AR was decreased only in the cerebral cortex and was not affected in the cerebellum [52].

Similarly, changes in the levels of other neuroactive steroids, like for instance the increase in the levels of 17β -E in the cerebellum at the end of the withdrawal period, were not associated with



changes in the expression of ER . Another example is provided by the changes of neuroactive steroids able to interact with PR, like for instance DHP [1]. Indeed, its upregulation in the cerebellum at the end of the withdrawal period was not associated with significant changes in the expression of PR.

The GABA A receptor has a pentameric structure formed by multiple subunits. Up to now, 19 subunits have been identified and among these, $\alpha 2$, $\alpha 4$, $\beta 3$, δ and $\gamma 2$ subunits are more widely expressed and are targets for neuroactive steroids [58, 59]. THP is one of the neuroactive steroids able to interact directly with GABA A receptors. The decrease in its levels in the cerebral cortex at the end of the withdrawal period was associated with the decrease in the gene expression of $\alpha 4$ and $\beta 3$ subunits. The levels of isopregnanolone, which in contrast to THP does not bind directly to the GABA A receptor, but, as mentioned before can antagonize the effects of THP on the GABA A receptor [1, 45, 46], were also decreased in the cerebral cortex at the end of the withdrawal period and this was also associated with the decrease in the gene expression of $\alpha 4$ and $\beta 3$ subunits of GABA receptors. The mRNA levels of $\beta 3$ subunit were also affected in the cerebellum after the subchronic treatment with finasteride. The increased gene expression of this subunit was associated with increased isopregnanolone levels and with decreased levels of 3β diol (i.e., a T metabolite that interacts with GABA A receptors). Data available in the literature indicate that THP treatment as well as its withdrawal (i.e., 24 hours) transiently increases the expression of $\alpha 4$ subunit in the hippocampus [60]. A decrease of this GABA A subunit has been reported after neonatal THP treatment [61] or following the development of acute THP tolerance [62]. Usually, the effect of THP on the expression of $\alpha 4$ subunit is associated with changes in the expression of δ subunit [60], which was not affected in our experimental model. This indicates that THP treatment and THP withdrawal affect the expression of different GABA A subunits than those affected by finasteride treatment and withdrawal, suggesting that the effect of finasteride on GABA A subunits is mediated by other mechanisms in addition to the modification of THP levels.



In summary, our findings indicate that finasteride treatment causes modifications in the levels of neuroactive steroids and their receptors in brain tissue. These changes are even more prominent after a long time of drug withdrawal. Given the pleiotropic regulatory functions exerted by neuroactive steroids in the nervous system, these persistent changes in their levels may have multiple consequences for brain function. Indeed, alterations of neuroactive steroid levels have been reported to be associated with neurodegenerative disorders, such as Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple sclerosis, traumatic brain injury and stroke as well as with psychiatric disorders, like for instance autism, depression, anxiety, posttraumatic stress and schizophrenia [63]. Moreover, as recently reviewed [64], emerging evidence in the literature suggests that 5 α R inhibitors, such as finasteride and dutasteride, may have important and serious adverse effects, like for instance on sexual function, insulin resistance and diabetes, high grade prostate cancer, cardiovascular disease, bone metabolism, neurobehavioral functions and neuropsychiatric disturbances. Further studies should explore the possible behavioral consequences of the modifications in neuroactive steroid levels caused by finasteride treatment and its withdrawal.

Acknowledgements

We acknowledge financial support from the Fondazione Cariplo to R.C.M. (grant number 2012 0547) and the Ministerio de Economía y Competitividad, Spain to L.M.G.S. (BFU2011 30217 C03 01). We also thank the Post Finasteride Foundation for the financial support to R.C.M.

References

- 1 Melcangi RC, Garcia Segura LM, Mensah Nyagan AG: Neuroactive steroids: state of the art and new perspectives. *Cell Mol Life Sci* 2008; 65:777 797.
- 2 Finn DA, Beadles Bohling AS, Beckley EH, Ford MM, Gililand KR, Gorin Meyer RE, Wiren KM: A new look at the 5alpha reductase inhibitor finasteride. *CNS Drug Rev* 2006; 12:53 76.
- 3 Traish AM: 5alpha reductases in human physiology: an unfolding story. *Endocr Pract* 2012; 18:965 975.
- 4 Cantagrel V, Lefeber DJ, Ng BG, Guan Z, Silhavy JL, Bielas SL, Lehle L, Hombauer H, Adamowicz M, Swiezewska E, De Brouwer AP, Blumel P, Sykut Cegielska J, Houliston S, Swistun D, Ali BR, Dobyns WB, Babovic Vuksanovic D, van Bokhoven H, Wevers RA, Raetz CR, Freeze HH, Morava E, Al Gazali L, Gleeson JG: SRD5A3 is required for converting polyprenol to dolichol and is mutated in a congenital glycosylation disorder. *Cell* 2010; 142: 203 217.
- 5 Melcangi RC, Panzica G, Garcia Segura LM: Neuroactive steroids: focus on human brain. *Neuroscience* 2011; 191: 1 5.
- 6 Panzica GC, Balthazart J, Frye CA, Garcia Segura LM, Herbison AE, Mensah Nyagan AG, McCarthy MM, Melcangi RC: Milestones on Steroids and the Nervous System: 10 years of basic and translational research. *J Neuroendocrinol* 2012; 24: 1 15.
- 7 Melcangi RC, Panzica GC: Allopregnanolone: state of the art. *Prog Neurobiol* 2014; 113: 1 5.
- 8 Ciriza I, Carrero P, Frye CA, Garcia Segura LM: Reduced metabolites mediate neuroprotective effects of progesterone in the adult rat hippocampus. The synthetic progestin medroxyprogesterone acetate (Provera) is not neuroprotective. *J Neurobiol* 2006; 66: 916 28.
- 9 Ford MM, Yoneyama N, Strong MN, Fretwell A, Tanchuck M, Finn DA: Inhibition of 5alpha reduced steroid biosynthesis impedes acquisition of ethanol drinking in male C57BL/6J mice. *Alcohol Clin Exp Res* 2008; 32: 1408 1416.
- 10 Biagini G, Longo D, Baldelli E, Zoli M, Rogawski MA, Bertazzoni G, Avoli M: Neurosteroids and epileptogenesis in the pilocarpine model: evidence for a relationship between P450 α induction and length of the latent period. *Epilepsia* 2009; 50 Suppl 1:53 58.
- 11 Atif F, Sayeed I, Yousuf S, Ishrat T, Hua F, Wang J, Brat DJ, Stein DG: Progesterone inhibits the growth of human neuroblastoma: in vitro and in vivo evidence. *Mol Med* 2011; 17: 1084 1094.
- 12 Bortolato M, Devoto P, Roncada P, Frau R, Flore G, Saba P, Pistritto G, Soggiu A, Pisanu S, Zappala A, Ristaldi MS, Tattoli M, Cuomo V, Marrosu F, Barbaccia ML: Isolation rearing induced reduction of brain 5alpha reductase expression: relevance to dopaminergic impairments. *Neuropharmacology* 2011; 60: 1301 1308.



- 13 Paris JJ, Brunton PJ, Russell JA, Walf AA, Frye CA: Inhibition of 5 α reductase activity in late pregnancy decreases gestational length and fecundity and impairs object memory and central progesterone milieu of juvenile rat offspring. *J Neuroendocrinol* 2011; 23: 1079 1090.
- 14 Darbra S, Modol L, Vallee M, Pallares M: Neonatal neurosteroid levels are determinant in shaping adult prepulse inhibition response to hippocampal allopregnanolone in rats. *Psychoneuroendocrinology* 2013; 38: 1397 1406.
- 15 Handa RJ, Kudwa AE, Donner NC, McGivern RF, Brown R: Central 5 α reduction of testosterone is required for testosterone's inhibition of the hypothalamo pituitary adrenal axis response to restraint stress in adult male rats. *Brain Res* 2013; 1529: 74 82.
- 16 Izumi Y, Svrakic N, O'Dell K, Zorumski CF: Ammonia inhibits long term potentiation via neurosteroid synthesis in hippocampal pyramidal neurons. *Neuroscience* 2013; 233: 166 173.
- 17 Modol L, Darbra S, Vallee M, Pallares M: Alteration of neonatal Allopregnanolone levels affects exploration, anxiety, aversive learning and adult behavioural response to intrahippocampal neurosteroids. *Behav. Brain Res* 2013; 241: 96 104.
- 18 Jeffrey M, Lang M, Gane J, Chow E, Wu C, Zhang L: Novel anticonvulsive effects of progesterone in a mouse model of hippocampal electrical kindling. *Neuroscience* 2014; 257: 65 75.
- 19 Kaufman KD, Olsen EA, Whiting D, Savin R, DeVillez R, Bergfeld W, Price VH, Van Neste D, Roberts JL, Hordinsky M, Shapiro J, Binkowitz B, Gormley GJ: Finasteride in the treatment of men with androgenetic alopecia. Finasteride Male Pattern Hair Loss Study Group. *J Am Acad Dermatol* 1998; 39: 578 589.
- 20 McClellan KJ, Markham A: Finasteride: a review of its use in male pattern hair loss. *Drugs* 1999; 57: 111 126.
- 21 Olsen EA, Hordinsky M, Whiting D, Stough D, Hobbs S, Ellis ML, Wilson T, Rittmaster RS: The importance of dual 5 α reductase inhibition in the treatment of male pattern hair loss: results of a randomized placebo controlled study of dutasteride versus finasteride. *J Am Acad Dermatol* 2006; 55: 1014 1023.
- 22 Traish AM, Hassani J, Guay AT, Zitzmann M, Hansen ML: Adverse side effects of 5 α reductase inhibitors therapy: persistent diminished libido and erectile dysfunction and depression in a subset of patients. *J Sex Med* 2011; 8: 872 884.
- 23 Irwig MS: Safety concerns regarding 5 α reductase inhibitors for the treatment of androgenetic alopecia. *Curr Opin Endocrinol Diabetes Obes* 2015; 22: 248 253
- 24 Altomare G, Capella GL: Depression circumstantially related to the administration of finasteride for androgenetic alopecia. *J Dermatol* 2002; 29: 665 669.



- 25 Rahimi Ardabili B, Pourandarjani R, Habibollahi P, Mualeki A: Finasteride induced depression: a prospective study. *BMC Clin Pharmacol* 2006; 6: 7.
- 26 Irwig MS, Kolukula S: Persistent sexual side effects of finasteride for male pattern hair loss. *J Sex Med* 2011; 8: 1747 1753.
- 27 Irwig MS: Depressive symptoms and suicidal thoughts among former users of finasteride with persistent sexual side effects. *J Clin Psychiatry* 2012; 73: 1220 1223.
- 28 Irwig MS: Persistent sexual side effects of finasteride: could they be permanent? *J Sex Med* 2012; 9: 2927 2932.
- 29 Melcangi RC, Caruso D, Abbiati F, Giatti S, Calabrese D, Piazza F, Cavaletti G: Neuroactive Steroid Levels are Modified in Cerebrospinal Fluid and Plasma of Post Finasteride Patients Showing Persistent Sexual Side Effects and Anxious/Depressive Symptomatology. *J Sex Med* 2013; 10: 2598 2603.
- 30 Caruso D, Abbiati F, Giatti S, Romano S, Fusco L, Cavaletti G, Melcangi RC: Patients treated for male pattern hair with finasteride show, after discontinuation of the drug, altered levels of neuroactive steroids in cerebrospinal fluid and plasma. *J Steroid Biochem Mol Biol* 2015; 146: 74 79.
- 31 Lebedev SV, Blinov DV, Petrov SV: Spatial characteristics of cisterna magna in rats and novel technique for puncture with a stereotactic manipulator. *Bull Exp Biol Med* 2004; 137: 635 638.
- 32 Caruso D, Pesaresi M, Abbiati F, Calabrese D, Giatti S, Garcia Segura LM, Melcangi RC: Comparison of plasma and cerebrospinal fluid levels of neuroactive steroids with their brain, spinal cord and peripheral nerve levels in male and female rats. *Psychoneuroendocrinology* 2013; 38: 2278 2290.
- 33 Chomczynski P, Sacchi N: Single step method of RNA isolation by acid guanidinium thiocyanate phenol chloroform extraction. *Anal Biochem* 1987; 162: 156 159.
- 34 Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal Biochem* 1976; 72: 248 254.
- 35 Zhang MG, Wu W, Zhang CM, Wang XJ, Gao PJ, Lu YL, Shen ZJ: Effects of oral finasteride on erectile function in a rat model. *J Sex Med* 2012; 9: 1328 1336.
- 36 Kiyokage E, Toida K, Suzuki Yamamoto T, Ishimura K: Localization of 5alpha reductase in the rat main olfactory bulb. *J Comp Neurol* 2005; 493:381 395.
- 37 Agis Balboa RC, Pinna G, Zhubi A, Maloku E, Veldic M, Costa E, Guidotti A: Characterization of brain neurons that express enzymes mediating neurosteroid biosynthesis. *Proc Natl Acad Sci U S A* 2006; 103:14602 14607.



- 38 Kiyokage E, Toida K, Suzuki Yamamoto T, Ishimura K: Cellular localization of 5 α reductase in the rat cerebellum. *J Chem Neuroanat* 2014; 59 60:8 16.
- 39 Sanchez P, Torres JM, Gavete P, Ortega E: Effects of swim stress on mRNA and protein levels of steroid 5 α reductase isozymes in prefrontal cortex of adult male rats. *Neurochem Int* 2008; 52:426 431.
- 40 Agis Balboa RC, Guidotti A, Pinna G: 5 α reductase type I expression is downregulated in the prefrontal cortex/Brodman's area 9 (BA9) of depressed patients. *Psychopharmacology (Berl)*. 2014; 231:3569 3580.
- 41 Castelli MP, Casti A, Casu A, Frau R, Bortolato M, Spiga S, Ennas MG: Regional distribution of 5 α reductase type 2 in the adult rat brain: an immunohistochemical analysis. *Psychoneuroendocrinology* 2013; 38:281 293.
- 42 Duskova M, Hill M, Starka L: Changes of metabolic profile in men treated for androgenetic alopecia with 1 mg finasteride. *Endocr Regul* 2010; 44: 3 8.
- 43 Mukai Y, Higashi T, Nagura Y, Shimada K: Studies on neurosteroids XXV. Influence of a 5 α reductase inhibitor, finasteride, on rat brain neurosteroid levels and metabolism. *Biol Pharm Bull* 2008; 31: 1646 1650.
- 44 Charlier TD, Cornil CA, Patte Mensah C, Meyer L, Mensah Nyagan AG, Balthazart J: Local modulation of steroid action: rapid control of enzymatic activity. *Front Neurosci* 2015; 9:83.
- 45 Hedstrom H, Bixo M, Nyberg S, Spigset O, Zingmark E, Backstrom T: Studies of pharmacokinetic and pharmacodynamic properties of isoallopregnanolone in healthy women. *Psychopharmacology (Berl)* 2009; 203: 85 98.
- 46 Bengtsson SK, Nyberg S, Hedstrom H, Zingmark E, Jonsson B, Backstrom T, Bixo M: Isoallopregnanolone antagonize allopregnanolone induced effects on saccadic eye velocity and self reported sedation in humans. *Psychoneuroendocrinology* 2015; 52: 22 31.
- 47 Lu SF, Mo Q, Hu S, Garippa C, Simon NG: Dehydroepiandrosterone upregulates neural androgen receptor level and transcriptional activity. *J Neurobiol* 2003; 57: 163 171.
- 48 Mo Q, Lu SF, Simon NG: Dehydroepiandrosterone and its metabolites: differential effects on androgen receptor trafficking and transcriptional activity. *J Steroid Biochem Mol Biol* 2006; 99: 50 58.
- 49 Mo Q, Lu S, Garippa C, Brownstein MJ, Simon NG: Genome wide analysis of DHEA and DHT induced gene expression in mouse hypothalamus and hippocampus. *J Steroid Biochem Mol Biol* 2009; 114: 135 143.
- 50 Hsieh JT, Chen SC, Yu HJ, Chang HC: Finasteride upregulates expression of androgen receptor in hyperplastic prostate and LNCaP cells: implications for chemoprevention of prostate cancer. *Prostate* 2011; 71: 1115 1121.



- 51 Di Loreto C, La Marra F, Mazzon G, Belgrano E, Trombetta C, Cauci S: Immunohistochemical evaluation of androgen receptor and nerve structure density in human prepuce from patients with persistent sexual side effects after finasteride use for androgenetic alopecia. *PLoS One* 2014; 9: e100237.
- 52 Munetomo A, Hojo Y, Higo S, Kato A, Yoshida K, Shirasawa T, Shimizu T, Barron A, Kimoto T, Kawato S: Aging induced changes in sex steroidogenic enzymes and sex steroid receptors in the cortex, hypothalamus and cerebellum. *J Physiol Sci* 2015; 65:253 263.
- 53 Canoine V, Fusani L, Schlinger B, Hau M: Low sex steroids, high steroid receptors: Increasing the sensitivity of the nonreproductive brain. *Dev Neurobiol* 2007; 67: 57 67.
- 54 Yamada S, Noguchi D, Ito H, Yamanouchi K: Sex and regional differences in decrease of estrogen receptor alpha immunoreactive cells by estrogen in rat hypothalamus and midbrain. *Neurosci Lett* 2009; 463: 135 139.
- 55 Mitterling KL, Spencer JL, Dziedzic N, Shenoy S, McCarthy K, Waters EM, McEwen BS, Milner TA: Cellular and subcellular localization of estrogen and progestin receptor immunoreactivities in the mouse hippocampus. *J Comp Neurol* 2010; 518: 2729 2743.
- 56 Reyes Guerrero G, Guzman C, Garcia DE, Camacho Arroyo I, Vazquez Garcia M: Extremely low frequency electromagnetic fields differentially regulate estrogen receptor alpha and beta expression in the rat olfactory bulb. *Neurosci Lett* 2010; 471: 109 113.
- 57 Langlois VS, Duarte Guterman P, Ing S, Pauli BD, Cooke GM, Trudeau VL: Fadrozole and finasteride exposures modulate sex steroid and thyroid hormone related gene expression in *Silurana (Xenopus) tropicalis* early larval development. *Gen Comp Endocrinol* 2010; 166: 417 427.
- 58 Belelli D, Lambert JJ: Neurosteroids: endogenous regulators of the GABA(A) receptor. *Nat Rev Neurosci* 2005; 6: 565 575.
- 59 Gunn BG, Cunningham L, Mitchell SG, Swinny JD, Lambert JJ, Belelli D: GABA receptor acting neurosteroids: A role in the development and regulation of the stress response. *Front. Neuroendocrinol.* 2014; 36: 28 48.
- 60 Smith SS, Shen H, Gong QH, Zhou X: Neurosteroid regulation of GABA(A) receptors: Focus on the alpha4 and delta subunits. *Pharmacol Ther* 2007; 116: 58 76.
- 61 Darbra S, Modol L, Llido A, Casas C, Vallee M, Pallares M: Neonatal allopregnanolone levels alteration: effects on behavior and role of the hippocampus. *Prog Neurobiol* 2014; 113: 95 105.
- 62 Turkmen S, Backstrom T, Wahlstrom G, Andreen L, Johansson IM: Tolerance to allopregnanolone with focus on the GABA A receptor. *Br J Pharmacol* 2011; 162: 311 327.



- 63 Giatti S, Garcia Segura LM, Melcangi RC. New steps forward in the neuroactive steroid field. *J Steroid Biochem Mol Biol* 2015; doi: 10.1016/j.jsbmb.2015.03.002.
- 64 Traish AM, Melcangi RC, Bortolato M, Garcia Segura LM, Zitzmann M. Adverse effects of 5alpha reductase inhibitors: What do we know, don't know, and need to know? *Rev Endocr Metab Disord*. 2015. DOI 10.1007/s11154 015 9319 y

Manuscript



Figure 1

Schematic representation of neurosteroidogenesis. Framed neuroactive steroids reported have been assessed by LC MS/MS.

Manuscript



Figure 2

Protein levels of classical steroid receptors detected by Western blotting in control (n=5) and subchronically treated finasteride animals (n=5). Expression of androgen receptor (AR; panel A) in cerebral cortex, progesterone receptor (PR; panel B), estrogen receptor α (ER α ; panel C) and estrogen receptor β (ER β ; panel D) in cerebellum, and PR (panel E) in hippocampus. In each panel a



representative blot is reported. The columns represent the mean \pm SEM after normalization with α tubulin. Data were analyzed by Student's t test. *p 0.05 vs. control.

Manuscript



Manuscript

Figure 3

Protein levels of classical steroid receptors detected by Western blotting at the end of the withdrawal period in control (n=5) and finasteride treated animals (n=5). Expression of androgen



receptor (AR; panel A), progesterone receptor (PR; panel B), estrogen receptor α (ER α ; panel C) and estrogen receptor β (ER β ; panel D) in cerebral cortex, PR (panel E) and AR (panel F) in cerebellum, and PR (panel G) in hippocampus. In each panel a representative blot is reported. The columns represent the mean \pm SEM after normalization with α tubulin. Data were analyzed by Student's t test.

*p 0.05 vs. control.

Manuscript



Figure 4

Gene expression of $\alpha 2$, $\alpha 4$, $\beta 3$, $\beta 2$ GABA A receptor subunits detected by Real Time PCR in cerebellum after chronic treatment. The columns represent the mean \pm SEM after normalization with 18s rRNA in control (n=5) and chronically treated finasteride animals (n=5). Data were analyzed by Student's t test. **p 0.01 vs. control.



Figure 5

Gene expression of $\alpha 2$, $\alpha 4$, $\beta 3$, $\beta 2$ GABA A receptor subunits detected by Real Time PCR in cerebral cortex at the end of the withdrawal period. The columns represent the mean \pm SEM after normalization with 18s rRNA in control (n=5) and chronically treated finasteride animals (n=5). Data were analyzed by Student's t test. *p < 0.05 and **p < 0.01 vs. control.



Table 1. Levels of neuroactive steroids in plasma, cerebrospinal fluid and brain areas of control and finasteride treated male rats: effect after subchronic treatment

	Plasma		CSF		Cerebral Cortex		Cerebellum		Hippocampus	
	control	Finasteride	control	Finasteride	control	Finasteride	control	Finasteride	control	Finasteride
PREG	0.20±0.046	0.19±0.03	0.31±0.091	0.53±0.07*	1.70±0.24	2.01±0.23	1.60±0.45	1.08±0.29	5.61±1.25	8.68±1.63
PROG	1.31±0.218	0.94±0.12	0.62±0.07	0.53±0.035	2.61±0.68	2.94±0.31	0.58±0.16	0.98±0.24	6.26±0.44	9.64±1.45*
DHP	0.56±0.07	0.52±0.14	0.49±0.12	0.31±0.06	5.18±0.81	5.49±0.76	2.94±1.06	5.56±0.89*	9.40±2.53	10.3±1.48
ISOPREG	u.d.l.	0.14±0.01*	0.13±0.022	0.11±0.007	2.54±0.62	3.48±1.03	0.51±0.06	1.16±0.20**	0.84±0.11	0.84±0.27
THP	0.25±0.07	0.14±0.04	2.31±0.15	2.23±0.15	15.6±1.75	19.0±1.95	5.34±0.48	9.34±2.33	u.d.l.	u.d.l.
DHEA	0.11±0.03	0.11±0.04	0.096±0.02	0.060±0.004	0.81±0.11	0.48±0.06**	0.51±0.08	0.75±0.19	0.29±0.09	0.30±0.08
T	4.23±1.19	3.16±0.91	0.10±0.01	0.12±0.037	2.38±0.37	1.71±0.32	1.86±0.26	1.89±0.26	3.21±0.56	1.97±0.78
DHT	0.79±0.09	0.30±0.02***	1.54±0.12	1.65±0.20	17.5±1.89	16.9±1.55	0.83±0.50	1.08±0.11	3.87±0.63	4.00±0.41
3 α -diol	0.84±0.10	1.04±0.28	0.076±0.014	u.d.l.*	0.57±0.06	0.60±0.11	0.54±0.07	0.34±0.07*	0.26±0.07	0.30±0.05
3 β -diol	u.d.l.	u.d.l.	0.056±0.004	0.10±0.02*	0.79±0.17	0.60±0.08	0.25±0.034	0.72±0.21*	u.d.l.	u.d.l.
17 α -E	u.d.l.	u.d.l.	u.d.l.	u.d.l.	0.076±0.02	0.053±0.01	0.054±0.02	0.077±0.02	u.d.l.	u.d.l.
17 β -E	0.022±0.001	0.048±0.01*	0.06±0.02	0.04±0.012	0.15±0.03	0.10±0.01	0.032±0.006	0.066±0.02*	0.03±0.007	0.05±0.008

Levels of pregnenolone (PREG), progesterone (PROG), dihydroprogesterone (DHP), tetrahydroprogesterone (THP), isopregnanolone (ISOPREG), dehydroepiandrosterone (DHEA), testosterone (T), dihydrotestosterone (DHT), 5 α -androstane-3 α , 17 β -diol (3 α -diol), 5 α -androstane-3 β , 17 β -diol (3 β -diol), 17 α -estradiol (17 α -E) and 17 β -estradiol (17 β -E) in plasma, cerebrospinal fluid (CSF), cerebral cortex, cerebellum and hippocampus of male rats. Data (n = 8 for each group) are expressed as pg/ μ l \pm SEM in case of plasma and CSF and pg/mg \pm SEM in case of brain areas. Student's t test analysis: * p < 0.05, ** p < 0.01 and *** p < 0.001 versus control. Under detection limit (u.d.l.). Detection limit were 0.05 pg/ μ l or pg/mg for 3 β -diol; 0.02 pg/ μ l or pg/mg for 17 α -E; 0.1 pg/ μ l or pg/mg for ISOPREG and THP.

Table 2. Levels of neuroactive steroids in plasma, cerebrospinal fluid and brain areas of control and finasteride treated male rats: effect at the end of the withdrawal period

	Plasma		CSF		Cerebral Cortex		Cerebellum		Hippocampus	
	control	Finasteride	control	Finasteride	control	Finasteride	control	Finasteride	control	Finasteride
PREG	0.19±0.03	0.18±0.04	0.20±0.051	0.27±0.04	1.07±0.48	1.24±0.18	0.66±0.03	1.06±0.12**	7.44±1.07	4.37±1.35*
PROG	2.12±0.35	1.32±0.21*	0.64±0.06	0.57±0.05	3.08±0.44	1.83±0.31*	1.48±0.30	1.52±0.57	8.44±1.07	5.27±0.56*
DHP	0.32±0.03	0.34±0.04	0.45±0.12	0.56±0.07	12.03±1.57	3.52±0.21***	2.59±0.61	4.38±0.70*	6.67±0.72	9.88±1.27*
ISOPREG	u.d.l.	0.13±0.04	0.11±0.004	0.11±0.004	2.93±0.35	1.89±0.32*	0.65±0.08	0.88±0.16	0.47±0.06	0.59±0.10
THP	0.50±0.09	0.14±0.03**	2.49±0.19	2.29±0.19	17.8±1.75	7.5±0.47***	4.12±0.37	4.58±0.63	u.d.l.	u.d.l.
DHEA	u.d.l.	u.d.l.	0.064±0.007	0.14±0.03*	1.26±0.31	0.30±0.04**	0.53±0.10	0.40±0.09	0.95±0.43	0.29±0.08
T	3.30±0.7	3.31±0.78	0.15±0.03	0.30±0.07*	3.89±0.61	3.65±0.63	1.22±0.27	2.26±0.25**	7.56±2.70	4.84±1.14
DHT	0.44±0.04	0.33±0.04*	1.35±0.04	0.98±0.10**	9.73±1.38	9.00±0.59	0.60±0.16	0.23±0.12*	3.03±0.43	4.00±0.37
3 -diol	1.63±0.29	1.00±0.19*	u.d.l.	u.d.l.	0.64±0.18	0.52±0.041	0.49±0.14	0.31±0.08	0.26±0.03	0.23±0.04
3 -diol	u.d.l.	u.d.l.	u.d.l.	u.d.l.	0.52±0.21	0.53±0.09	0.63±0.18	0.19±0.03*	u.d.l.	u.d.l.
17 -E	u.d.l.	u.d.l.	u.d.l.	u.d.l.	0.046±0.009	0.045±0.009	0.064±0.03	0.041±0.03	u.d.l.	u.d.l.
17 -E	0.028±0.004	0.024±0.002	u.d.l.	0.04±0.009**	0.12±0.02	0.06±0.01*	0.037±0.01	0.055±0.01	0.074±0.02	0.034±0.01

Levels of pregnenolone (PREG), progesterone (PROG), dihydroprogesterone (DHP), tetrahydroprogesterone (THP), isopregnanolone (ISOPREG), dehydroepiandrosterone (DHEA), testosterone (T), dihydrotestosterone (DHT), 5 α -androstane-3 α , 17 β -diol (3 α -diol), 5 α -androstane-3 β , 17 β -diol (3 β -diol), 17 α -estradiol (17 α -E) and 17 β -estradiol (17 β -E) in plasma, cerebrospinal fluid (CSF), cerebral cortex, cerebellum and hippocampus of male rats. Data (n = 8 for each group) are expressed as pg/ μ l \pm SEM in case of plasma and CSF and pg/mg \pm SEM in case of brain areas. Student's t test analysis: * p < 0.05, ** p < 0.01 and *** p < 0.001 versus control. Under detection limit (u.d.l.). Detection limit were 0.05 pg/ μ l or pg/mg for DHEA, 3 α -diol and 3 β -diol; 0.02 pg/ μ l or pg/mg for 17 α -E and 17 β -E; 0.1 pg/ μ l or pg/mg for ISOPREG and THP.

Neuroendocrinology (*International Journal for Basic and Clinical Studies on Neuroendocrine Relationships*)

Journal Editor: Millar R.P. (Edinburgh)

ISSN: 0028-3835 (Print), eISSN: 1423-0194 (Online)

www.karger.com/NEN

Disclaimer: Accepted, unedited article not yet assigned to an issue. The statements, opinions and data contained in this publication are solely those of the individual authors and contributors and not of the publisher and the editor(s). The publisher and the editor(s) disclaim responsibility for any injury to persons or property resulting from any ideas, methods, instructions or products referred to in the content. Copyright: All rights reserved. No part of this publication may be translated into other languages, reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, recording, microcopying, or by any information storage and retrieval system, without permission in writing from the publisher or, in the case of photocopying, direct payment of a specified fee to the Copyright Clearance Center.

© 2015 S. Karger AG, Basel

