

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Advances in microbiology, infectious diseases and public health: Refractory *Trichophyton rubrum* infections in Turin, Italy: a problem still present

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1617533> since 2018-01-19T13:19:46Z

Publisher:

Springer International Publishing

Published version:

DOI:DOI 10.1007/5584_2015_5012


Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Metadata of the chapter that will be visualized online

Chapter Title	Advances in Microbiology, Infectious Diseases and Public Health Short Data Report/Revised Version
Chapter Sub Title	 Refractory <i>Trichophyton rubrum</i> Infections in Turin, Italy: A Problem Still Present
Copyright Year	2015
Copyright Holder	Springer International Publishing Switzerland
Author	<div>Family Name Tullio</div> <div>Particle</div> <div>Given Name Vivian</div> <div>Suffix</div> <div>Division Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics</div> <div>Organization University of Torino</div> <div>Address via Santena 9, 10126 Turin, Italy</div>
Author	<div>Family Name Cervetti</div> <div>Particle</div> <div>Given Name Ornella</div> <div>Suffix</div> <div>Division Department of Medical Sciences</div> <div>Organization University of Turin</div> <div>Address Turin, Italy</div>
Author	<div>Family Name Roana</div> <div>Particle</div> <div>Given Name Janira</div> <div>Suffix</div> <div>Division Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics</div> <div>Organization University of Torino</div> <div>Address via Santena 9, 10126 Turin, Italy</div>
Author	<div>Family Name Panzone</div> <div>Particle</div> <div>Given Name Michele</div> <div>Suffix</div> <div>Division A.O.U. Città della Salute e della Scienza</div> <div>Organization San Lazzaro Hospital</div> <div>Address Turin, Italy</div>
Author	<div>Family Name Scalas</div>

	Particle	
	Given Name	Daniela
	Suffix	
	Division	Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics
	Organization	University of Torino
	Address	via Santena 9, 10126 Turin, Italy
Author	Family Name	Merlino
	Particle	
	Given Name	Chiara
	Suffix	
	Division	Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics
	Organization	University of Torino
	Address	via Santena 9, 10126 Turin, Italy
Author	Family Name	Allizond
	Particle	
	Given Name	Valeria
	Suffix	
	Division	Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics
	Organization	University of Torino
	Address	via Santena 9, 10126 Turin, Italy
Author	Family Name	Banche
	Particle	
	Given Name	Giuliana
	Suffix	
	Division	Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics
	Organization	University of Torino
	Address	via Santena 9, 10126 Turin, Italy
Author	Family Name	Mandras
	Particle	
	Given Name	Narcisa
	Suffix	
	Division	Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics
	Organization	University of Torino
	Address	via Santena 9, 10126 Turin, Italy

Corresponding Author	Family Name	Cuffini
	Particle	
	Given Name	Anna Maria
	Suffix	
	Division	Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics
	Organization	University of Torino
	Address	via Santena 9, 10126 Turin, Italy
	Email	annamaria.cuffini@unito.it

Abstract	<p>Dermatophytosis caused by <i>Trichophyton rubrum</i> is the most common cutaneous fungal infection in industrialized countries and worldwide with high recurrence and lack of treatment response. In addition, patients with cutaneous and concurrent toenail lesions are often misdiagnosed and therefore treated with an inappropriate therapy. In this study, we evaluated five previously misdiagnosed cases of <i>T.rubrum</i> chronic dermatophytosis sustained by two variants at sites distant from the primary lesion. Our patients were successfully treated by systemic and topical therapy, and 1 year after the end of therapy follow-up did not show any recurrence of infection.</p> <p>Our data indicate that the localization of all lesions, the isolation and the identification of the causative fungus are essential to establish the diagnosis and the setting of a correct therapeutic treatment to avoid recurrences.</p>
----------	--

Keywords (separated by '-')	<i>Trichophyton rubrum</i> - Chronic dermatophytosis - Misdiagnosis
-----------------------------	---

AU1

Advances in Microbiology, Infectious Diseases and Public Health ~~Short Data Report/Revised Version~~

Refractory *Trichophyton rubrum* Infections in Turin, Italy: A Problem Still Present

Vivian Tullio, Ornella Cervetti, Janira Roana, Michele Panzone, Daniela Scalas, Chiara Merlino, Valeria Allizond, Giuliana Banche, Narcisa Mandras, and Anna Maria Cuffini

Abstract

Dermatophytosis caused by *Trichophyton rubrum* is the most common cutaneous fungal infection in industrialized countries and worldwide with high recurrence and lack of treatment response. In addition, patients with cutaneous and concurrent toenail lesions are often misdiagnosed and therefore treated with an inappropriate therapy. In this study, we evaluated five previously misdiagnosed cases of *T. rubrum* chronic dermatophytosis sustained by two variants at sites distant from the primary lesion. Our patients were successfully treated by systemic and topical therapy, and 1 year after the end of therapy follow-up did not show any recurrence of infection.

Our data indicate that the localization of all lesions, the isolation and the identification of the causative fungus are essential to establish the diagnosis and the setting of a correct therapeutic treatment to avoid recurrences.

Keywords

Trichophyton rubrum • Chronic dermatophytosis • Misdiagnosis

V. Tullio, J. Roana, D. Scalas, C. Merlino, V. Allizond, G. Banche, N. Mandras, and A.M. Cuffini (✉)
Bacteriology and Mycology Laboratory, Department of Public Health and Pediatrics, University of Torino, via Santena 9, 10126 Turin, Italy
e-mail: annamaria.cuffini@unito.it

O. Cervetti
Department of Medical Sciences, University of Turin, Turin, Italy

M. Panzone
A.O.U. Città della Salute e della Scienza, San Lazzaro Hospital, Turin, Italy

AU2

Chronic dermatophytosis is a condition in which the clinical symptoms persist for more than 1 year with episodes of exacerbation and remission (Hay 1982; Zaia and Rebell 2003; Prasad et al. 2005). The main etiologic agent is *Trichophyton rubrum* responsible for 90 % of chronic infections (Di Chiacchio et al. 2014; Nenoff et al. 2014). Chronicity is probably related both to fungal cell wall components, such as mannan, that play an important role in the process of down-modulation of cell-mediated immune response of the host and to a lack of treatment response (Blake et al. 1991; Sato and Tagami 2003; Waldman et al. 2010). Patients with cutaneous and concurrent toenail lesions are often misdiagnosed and, therefore, treated with an inappropriate therapy (Larruskain et al. 2005).

In this study, we evaluated previously misdiagnosed cases of *T. rubrum* chronic dermatophytosis in five patients admitted to the Medical Sciences Department, University of Torino (Italy), through an investigation of clinical and mycological infection aspects.

Case 1 A 42-year old male, born in Ecuador, reported a 7-year history of itchy and squamous lesions on the soles, toenails, palms and the nail plates, before arriving in Italy (Fig. 1a–d). Despite therapies with topical antibacterial agents in his native country, the patient had extensive erythema with painful papules, pustules and crusts in the chin and beard (Fig. 1e, f). Incomplete alopecia, associated with follicular nodules most prevalent above the upper lip was seen. Hands and fingernails examination revealed hyperkeratosis and distal onycholysis.

Case 2 A Caucasian male of 48 years presented erythematous and squamous lesions on the feet and toenails. A closer examination revealed scaling lesions on the inguinal area and buttocks, hands and fingernails plate hyperkeratosis and distal onycholysis.

Case 3 A Caucasian female of 78 years reported a 2-week history of extensive erythema with papules and fine pustules appearing at the

opening of hair follicles in the inguinal region (Fig. 2a, b). An intense erythema involved both buttocks and thighs (Fig. 2c). Examination of the left foot revealed sole and toenail/fingernail hyperkeratosis, with nail plate thickened, friable and yellowish (Fig. 2e, f). The left knee (Fig. 2d) and the right leg were also involved with flaking in net margins.

Case 4 A Caucasian female of 69 years, with rheumatoid arthritis, treated for 20 years with therapeutic cycles of methotrexate (7.5 mg/week) and prednisone (5 mg/day), presented a chronic erythematous scaly dermatitis extended to the lower back and rear thigh area, diagnosed as psoriasis (Fig. 3e). Since 2006, she was treated with emollient cream and topical steroids without benefit. On physical examination, the patient revealed *tinea pedis* and *tinea unguium* with sole and toenails plate hyperkeratosis (Fig. 3a, b), squamous lesions on the elbow, on the back and left palm (Fig. 3c, d, g). Involvement of the scalp with flaking dandruff and thinning hair was observed (Fig. 3f).

Case 5 A Caucasian female of 68 years, with rheumatoid arthritis, treated for several years with prednisone (25 mg/day), presented a history of chronic erythematous scaly dermatitis diagnosed as psoriasis and treated with emollient cream without benefit. A closer examination revealed an intense lamellar desquamation of the toenails and fingernails, hyperkeratosis of the soles and the palms, scaling lesions with sharp margins in the breast, abdomen, inguinal area, buttocks and thighs, neck and chin.

Mycological analysis of all patient lesions was performed. Skin and nail samples were collected, examined under a light microscope (20 % KOH + 40 % DMSO preparation) and inoculated into Mycobiotic agar (Merck, KGAA, Germany) to detect dermatophytes. Molds identification was based on macroscopic and microscopic characters of the colonies after 15 days of incubation at 25 °C.

All patients had dermatophytosis and concurrent lesions caused by two variants of *T. rubrum*:



Fig. 1 Case 1. A 42-year old, male, born in Ecuador. Squamous lesions on the soles, toenails, palms and nail plates (a–d); extensive erythema in the chin and beard with follicular nodules above the upper lip (e, f)

119 downy white-colored colonies with reverse pig-
 120 ment brownish-yellow (Cases 1, 2, and 3) or deep
 121 wine-red (Cases 4, and 5). Scant teardrop-shaped
 122 microconidia along septate hyphae were
 123 observed on microscopic colonies examination.
 124 The primary lesion was localized always in
 125 the foot (*tinea pedis*), in agreement with other

studies (Larruskain et al. 2005). Secondary 126
 lesions distributed in other sites were the main 127
 demand for medical consultation: in all five 128
 cases, the anatomical sites mainly interested 129
 were the inguinal area, buttocks, palms and 130
 fingernails (*tinea unguium*). In only one case, 131
tinea capitis was observed (Case 4). Patient 132



Fig. 2 Case 3. A 78-year old, female, Caucasian. Extensive erythema with papules at the opening of hair follicles in the inguinal region (**a, b**), buttocks and thighs (**c**); left

knee with flaking in net margins (**d**); toenail and fingernail hyperkeratosis (**e, f**)

133 4 under methotrexate therapy and patient
134 5, under corticosteroid therapy had risk factors
135 predisposing them to fungal spread. *Tinea* in
136 such cases tends to be chronic and extended,
137 mimicking various skin diseases, such as psoria-
138 sis, eczema, etc., as in Patients 4 and 5 (Atzori
139 et al. 2012; Tan et al. 2014).

For all patients a successful treatment with 140
topical (azoles) and systemic (terbinafine hydro- 141
chloride 250 mg/day) antimycotics was carried 142
out. In details, in patient 1, after 4 weeks of 143
treatment, all skin lesions were completely 144
healed and culture results were negative; both 145
direct mycological and culture were negative 146



Fig. 3 Case 4. A 69-year old, female, Caucasian, with rheumatoid arthritis. Sole and toenails hyperkeratosis (a, b); back and left palm squamous lesions (c, d); extensive

erythema on lower back and rear thigh area diagnosed as psoriasis (e); scalp with flaking dandruff and thinning hair (f); squamous lesions on the elbow (g)

also for nails after 3 months. In patient 2, all lesions were completely healed and culture results were negative after 12-weeks of treatment. In patient 3, all skin lesions were completely healed after 6 weeks of treatment; both direct mycological and culture were negative for nails after 4 months. In patient 4, after 4-weeks of treatment, all skin lesions were completely healed; both direct mycological and culture were negative also for nails and scalp after 5 months. In patient 5, after 6-weeks of treatment, all skin lesions were completely healed and culture results were negative; the nail lesions were alleviated after 5-months therapy.

The five clinical cases reported in this study are considered dermatophytosis, affecting both immunocompetent and immunodeficient patients, and fulfilled the diagnostic criteria of *T. rubrum* chronic dermatophytosis, as indicated by the literature (Zaias and Rebell 1996; Böhmer and Korting 1999; Kick and Korting 2001; Balci and Cetin 2008; Piñeiro et al. 2010; Kong et al. 2015). Since in our group of patients from the beginning a correct therapeutic treatment was not carried out or misapplied, a gradual spread of the infection occurred to the toenails, as secondary site involved, constituting the reservoir of infection that spread later to other sites, such as legs, groin, hands, face and scalp. On the other hand, it has to be underlined that *tinea unguium* is an infection usually more resistant to treatment, whose eradication is difficult even with appropriate therapy (Gupta and Cooper 2008).

For fungal infection eradication, diagnosis must be based on both a correct patient history and an adequate microbiological study that includes the identification of the species isolated. Therefore, it is essential a careful examination of the patient *in toto* to avoid inappropriate or wrong therapeutic treatment. In fact, as in the first patient, the antibiotic treatment was established solely on the observation of highly inflammatory facial injuries that did not present the typical clinical features of *T. rubrum* infection (Yin et al. 2011); hence, the treatment being wrong was ineffective.

In conclusion, our data indicate that in all cases of suspected syndrome or when skin involvement is extended to multiple sites, the localization of all lesions, the isolation and the identification of the causative fungus are essential to establish the diagnosis, prognosis and the setting of a correct antifungal therapy to avoid recurrences.

References


- Atzori L, Pau M, Aste N et al (2012) Dermatophyte infections mimicking other skin diseases: a 154-person case survey of tinea atypica in the district of Cagliari (Italy). *Int J Dermatol* 51:410–415
- Balci DD, Cetin M (2008) Widespread, chronic, and fluconazole-resistant *Trichophyton rubrum* infection in an immunocompetent patient. *Mycoses* 51:546–548
- Blake JS, Dahl MV, Herron MJ et al (1991) An immunoinhibitory cell wall glycoprotein (mannan) from *Trichophyton rubrum*. *J Invest Dermatol* 96:657–661
- Böhmer U, Korting HC (1999) *Trichophyton rubrum* syndrome with axillary tinea infection. *Hautarzt* 50:292–294
- Di Chiacchio N, Madeira CL, Humaire CR et al (2014) Superficial mycoses at the Hospital do Servidor Público Municipal de São Paulo between 2005 and 2011. *An Bras Dermatol* 89:67–71
- Gupta AK, Cooper EA (2008) Update in antifungal therapy of dermatophytosis. *Mycopathologia* 166:353–367
- Hay RJ (1982) Chronic dermatophyte infections. Clinical and mycological features. *Br J Dermatol* 106:1–7
- Kick G, Korting HC (2001) The definition of *Trichophyton rubrum* syndrome. *Mycoses* 44:167–171
- Kong QT, Du X, Yang R et al (2015) Chronically recurrent and widespread *tinea corporis* due to *Trichophyton rubrum* in an immunocompetent patient. *Mycopathologia* 179:293–297
- Larruskain J, Piñeiro L, Idigoras P et al (2005) Dermatophytosis with concurrent lesions in distant locations. Prognostic and therapeutic significance. *Enferm Infecc Microbiol Clin* 23:191–193
- Nenoff P, Krüger C, Schaller J et al (2014) Mycology – an update. Part 1: dermatomycoses: causative agents, epidemiology and pathogenesis. *J Dtsch Dermatol Ges* 12:749–777
- Piñeiro L, Larruskain J, Idigoras P et al (2010) *Trichophyton rubrum* syndrome: the tip of the iceberg and a preventable outcome. *Mycoses* 53:186
- Prasad PVS, Priya K, Kaviarasan PK et al (2005) A study of chronic dermatophyte infection in a rural hospital. *Indian J Dermatol Venereol Leprol* 71:129–130

- 248 Sato N, Tagami H (2003) Severe measles in a young 260
- 249 female patient with chronic, generalized *Trichophyton* 261
- 250 *rubrum* infection showing type 2 helper T cell- 262
- 251 dominant immunologic reactivity. J Am Acad 263
- 252 Dermatol 48(5 Suppl):S43–S46 264
- 253 Tan Y, Lin L, Feng P et al (2014) Dermatophytosis caused 265
- 254 by *Trichophyton rubrum* mimicking syphilid: a case 266
- 255 report and review of literature. Mycoses 57:312–315 267
- 256 Waldman A, Segal R, Berdicevsky I et al (2010) CD4+ 268
- 257 and CD8+ T cells mediated direct cytotoxic effect 269
- 258 against *Trichophyton rubrum* and *Trichophyton*
- 259 *mentagrophytes*. Int J Dermatol 49:149–157
- Yin X, Du X, Zhang H (2011) A case of tinea barbae due 260
- to *Trichophyton rubrum* infection by autoinoculation 261
- from the infected fingernails. Mycoses 54:e864–e866 262
- Zaias N, Rebell G (1996) Chronic dermatophytosis 263
- caused by *Trichophyton rubrum*. J Am Acad Dermatol 264
- 35:S17–S20 265
- Zaias N, Rebell G (2003) Clinical and mycological status 266
- of the *Trichophyton mentagrophytes* (interdigitale) 267
- syndrome of chronic dermatophytosis of the skin and 268
- nails. Int J Dermatol 42:779–788 269

Uncorrected Proof

Author Queries

Chapter No.: 5012 0002619843

Queries	Details Required	Author's response
AU1	Please check the article title and article sub title.	
AU2	The citations Di Chiacchio et al. (2012), Tan et al. (2013), Piñeiro et al. (2009), Kong et al. (2014) have been changed to Di Chiacchio et al. (2014), Tan et al. (2014), Piñeiro et al. (2010), Kong et al. (2015). Please check if okay.	OK

Uncorrected Proof