

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Recommendations for the Management of CML in the Era of Second-Generation TKIs

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1614572> since 2016-11-20T17:26:38Z

Publisher:

Springer Japan

Published version:

DOI:10.1007/978-4-431-55714-2_8

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Molecular Pathogenesis and Treatment of Chronic Myelogenous Leukemia

Masahiro Kizaki

Editor

Molecular Pathogenesis and Treatment of Chronic Myelogenous Leukemia



Springer

Editor

Masahiro Kizaki
Department of Hematology, Saitama Medical Center
Saitama Medical University
Kawagoe, Saitama
Japan

ISBN 978-4-431-55713-5 ISBN 978-4-431-55714-2 (eBook)
DOI 10.1007/978-4-431-55714-2

Library of Congress Control Number: 2015954659

Springer Tokyo Heidelberg New York Dordrecht London

© Springer Japan 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer Japan KK is part of Springer Science+Business Media (www.springer.com)

Preface

Advances in treatment of chronic myeloid leukemia (CML) have been made over the past two decades thanks to research that has furthered our understanding of its molecular pathogenesis. CML is a clonal hematopoietic stem cell disorder characterized by the abnormal proliferation of myeloid cell lineages, which progresses through the chronic phase (CP) to the accelerated and blastic phases. CML is caused by the presence of the Philadelphia (Ph) chromosome in hematopoietic stem cells, which arises from the reciprocal translocation of chromosomes 9 and 22, t(9;22)(q34;q11), resulting in the development of the *bcr-abl* chimeric gene. This chimeric gene produces the BCR-ABL fusion protein that has oncogenic activity. The BCR-ABL fusion protein has constitutive tyrosine kinase (TK) activity that is stronger than that of the naïve ABL protein, conferring a proliferative advantage and aberrant differentiation capacity to affected hematopoietic stem cells, resulting in the oncogenic event of leukemia development. Therefore, formation of the *bcr-abl* chimeric gene and its encoded protein is a primary and central event in the molecular pathogenesis of CML.

Until 2000, drug therapy for CML was limited to non-specific cytotoxic drugs such as busulfan and hydroxyurea, and then interferon (IFN)- α was introduced to regress disease activity, which had a survival benefit. Allogeneic hematopoietic stem cell transplantation (allo-SCT) for CML-CP frequently was a curative therapeutic approach for patients with good performance status and an appropriate stem cell donor, but it also was associated with a high incidence of early morbidity and mortality. Understanding of the molecular pathogenesis of CML resulted in rapid development of new therapeutic agents, including various BCR-ABL specific tyrosine kinase inhibitors (TKIs) such as imatinib, nilotinib, and dasatinib. Current clinical guidelines endorse use of any of these three TKIs for initial management of CML-CP. Molecular-targeted therapy with these TKIs was shown to dramatically improve clinical outcomes of CML patients, increasing the 10-year overall survival (OS) from 20 to 80–90 %. As shown in many clinical studies, CML patients treated with TKIs are expected to live for a long period of time. Thus, identification of appropriate surrogate markers for clinical outcome has become important.

Achieving a more complete and faster molecular response is correlated with good clinical outcome; therefore, improving molecular monitoring techniques for minimal residual disease is crucial. Furthermore, management of TKI-resistant CML and development of new TKIs are also important issues. We now have available multiple TKIs for clinical use, including second- and third-generation agents. In this decade, our goals for the treatment of CML are to optimize the quality of life for patients, to establish the most cost-effective treatment, and to deliver the best treatment and monitoring to each patient anywhere in the world. The ultimate goal for any patient is to discontinue use of TKIs—to achieve treatment-free remission and subsequent cure. In other words, future research into treatment of CML will focus on achieving and maintaining complete molecular remission after discontinuation of TKIs.

This book begins with a discussion of recent advances in basic CML research regarding stem cells and the signaling pathways of leukemic cells; continues by describing various clinical aspects of the use of TKIs in daily clinical practice; and concludes with a discussion of future trials aimed at a cure for this disease. I would like to acknowledge the many excellent colleagues who have contributed to each chapter. In addition, I would like to express my appreciation to the staff of Springer Japan, for all of their efforts in bringing this treatise to publication. It is hoped that this book will encourage implementation of further basic and clinical research projects with the goal of solving the remaining intriguing and important clinical problems of CML treatment.

Kawagoe, Japan

Masahiro Kizaki, MD, PhD

Contents

1	Identification and Biology of CML Stem Cells	1
	Hiromi Iwasaki and Koichi Akashi	
2	Molecular Mechanisms of CML Stem Cell Maintenance	11
	Atsushi Hirao, Yuko Tadokoro, and Masaya Ueno	
3	Roles for Signaling Molecules in the Growth and Survival of CML Cells	29
	Itaru Matsumura	
4	Goals of CML Treatment in the Tyrosine Kinase Inhibitor Era . . .	53
	Jerald Radich and Daniel Egan	
5	Biomarkers for Determining the Prognosis of CML	69
	Naoto Takahashi	
6	Updated European LeukemiaNet Recommendations for the Management of CML	81
	Noriko Usui	
7	Optimal Monitoring of CML Treatment: Molecular and Mutation Analysis	101
	David T. Yeung and Susan Branford	
8	Recommendations for the Management of CML in the Era of Second-Generation TKIs	131
	Alessandro Morotti, Carmen Fava, and Giuseppe Saglio	
9	The Role of New TKIs and Combinations with Interferon-α for the Treatment of CML	147
	Franck E. Nicolini, Marie Balsat, Hélène Labussière-Wallet, Mohamad Sobh, Arthur Bert, and Maël Heiblig	

10	Safety Profiles of First-Line TKIs and Managing Adverse Effects	161
	Gianantonio Rosti, Fausto Castagnetti, Gabriele Gugliotta, and Michele Baccarani	
11	Molecular Mechanism of TKI Resistance and Potential Approaches to Overcome Resistance	167
	Hein Than, Charles Chuah, and S. Tiong Ong	
12	Discontinuation of Therapy and Treatment-Free Remission in CML	183
	David M. Ross and Timothy P. Hughes	