

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

TRACE AND RARE EARTH ELEMENTS IN PM10COLLECTED AT NY-ÅLESUND (SVALBARD ISLANDS, ARCTICA)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1615598> since 2016-11-23T10:28:01Z

Publisher:

Università degli Studi di Messina

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



**Università
degli Studi di Messina**



**Atti del XXVI Congresso
della Divisione di Chimica Analitica
della Società Chimica Italiana**

Giardini Naxos (Messina)

18-22 Settembre 2016

www.analitica2016.it



TRACE AND RARE EARTH ELEMENTS IN PM₁₀ COLLECTED AT NY-ÅLESUND (SVALBARD ISLANDS, ARCTICA)

M. Malandrino¹, O. Abollino¹, A. Giacomino², S. Buoso¹, E. Conca¹, R. Udisti³

¹Dipartimento di Chimica, Università degli Studi di Torino, Via Pietro Giuria, 5 – 10125 Torino

²Dipartimento di Scienze e Tecnologia del Farmaco, Università degli Studi di Torino, Via Pietro Giuria, 9 – 10125 Torino

³Dipartimento di Chimica, Università degli Studi di Firenze, Via della Lastruccia, 3 – 50019, Sesto Fiorentino

The chemical composition of atmospheric aerosol is responsible for the impact of particulate matter on human health and, at a larger scale, on the ongoing climate changes.

The Arctic regions are showing to be the first areas affected by the present climatic variations. Consequently, the study of the chemical composition of atmospheric aerosol in the polar areas is important to understand the feedback processes between the climate forcing and the environmental responses [1, 2].

In this study, the concentrations of main and trace metals in the PM₁₀ collected at Ny-Ålesund (Svalbard Islands) during the spring and summer 2010, 2011 and 2012 campaigns were determined. The results obtained reveal an evident seasonal pattern in the temporal profiles of the majority of the chemical components of the PM₁₀, that show higher atmospheric concentrations in March-April. The most likely explanation for this trend is the influence of continental sources on the composition of the Arctic PM₁₀. Indeed, Svalbard Islands are affected by aerosols coming from anthropized continental areas by long range transport processes, especially occurring in early spring.

The enrichment factors, calculated considering Al as a crustal reference element, are higher than 100 for Zn, Mo, As, Cd, Pb and Na, indicating their non-geogenic origin (anthropogenic sources and sea spray).

The final dataset was treated by chemometric techniques. Principal Component Analysis showed an evident separation between spring and summer Arctic PM₁₀ samples. Factor Analysis identified four factors: F1 – geogenic source (Al, Fe, Mn, Ba, Ti and REEs with exception of Ce); F2 – sea spray source (K, Na and Mg); F3 – combustion processes source (As, Cd, Co, Ni, V, Pb and Zn); F4 – wear-related source (Cu, Zn and Ce).

This work was supported by MIUR and CNR funds and facilities.

[1] M.E. Gagne, N.P. Gillett, J.C. Fyfe, *Geophys. Res. Lett.* 42 (2015) 8481-8488.