

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A peptide from the extracellular region of the synaptic protein α Neurexin stimulates angiogenesis and the vascular specific tyrosine kinase Tie2

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/127094> since 2017-01-18T15:54:03Z

Published version:

DOI:10.1016/j.bbrc.2013.02.045

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

A peptide from the extracellular region of the synaptic protein α Neurexin stimulates angiogenesis and the vascular specific tyrosine kinase Tie2

Stefania Graziano¹, Serena Marchiò², Federico Bussolino^{3*} Marco Arese^{1*}

Department of Oncology, University of Torino Medical School, and laboratory of Neurovascular Biology¹, Tumor Microenvironment², Vascular Oncology³, Institute for Cancer Research @ Candiolo, Candiolo (TO), Italy

Corresponding author: Marco Arese, marco.arese@unito.it

*F.B. and M.A. equally contributed to this report

Fax 011-9933524

Ph 011 -9933508

Abstract

Neurexin (NRXN) and Neuroligin (NLGN) are trans-synaptic proteins involved in vascular biology. NRXN is encoded in long (α) and short (β) isoforms. We have shown that β NRXN modulates blood vessel development in synergy with VEGFA and associates with NLGN. On the other hand α NRXN is also expressed in blood vessels but does not interact with NLGN or act in synergy with VEGFA, thus demonstrating a differential role.

To find clues into the vascular functions of α NRXN, we chose a 7 aa motif that is part of its extracellular region and was formerly selected through a proteomic search for interactors of the vascular receptor Tie2. Next we a) synthesized and modeled such peptide in order to determine its biological activity towards Tie2 in *in vitro* and *in vivo* angiogenesis assays and b) evaluated if α NRXN and Tie2 physically associate in situ during vascular development.

We used biochemical and cellular assays to prove that the synthetic α NRXN peptide a) modulates the angiogenic phenotype of cultured endothelial cells and angiogenesis in vivo and b) efficiently stimulates Tie2 phosphorylation and downstream mediators in endothelial cells. Moreover, we show that α NRXN and Tie2 can be reciprocally immunoprecipitated from chicken blood vessels at late stages of vascular development.

These data have a double significance, i.e. provide a novel tool to modulate Tie2 and further suggest the involvement of the NRXN family of synaptic protein in the vascular system through their interaction with a fundamental vascular player.

keywords: Tie2, Neurexin, peptide, structure

1. INTRODUCTION

Neurexins (NRXNs)¹ are a family of pre-synaptic transmembrane proteins involved in the modulation of synaptic activity in partnership with the post synaptic Neuroligins (NLGNs). In mammals, there are 3 different NRXN genes, each presenting 2 alternative promoters which drive the synthesis of 2 major isoforms per gene (a long α NRXN, and a short C-terminal β NRXN). The extracellular region of α NRXN is constituted by three LNS-EGF-LNS repeats while the short β NRXN is made only of the last LNS (6th) domain (Figure S1).

We recently discovered that a) α and β NRXN as well as NLGN are expressed by blood vessels b) β NRXN and NLGN modulate angiogenesis in the chick embryo chorioallantoic membrane assay; c) while both α NRXN and β NRXN co-precipitate with NLGN in brain, only β NRXN does so in blood vessels [1]d) in zebrafish β NRXN but not α NRXN knock-down causes vascular defect in synergy with VEGFA [2]. All these findings raise questions about α NRXN function and protein partners in the vascular system. In a parallel project we screened a phage-displayed random peptide library on the vascular specific receptor Tie2 in order to find either new small modulators of its activity or discover new physiological partners that could participate in the regulation of the complex family of Angiopoietin/Ties. Among the candidate Tie2 ligands there was a peptide corresponding to a portion of α NRXN. This report analyzes the possibility that Tie2 is linked to the vascular α NRXN and demonstrates that a) a cyclic peptide that mimics a region of the second EGF like domain of α NRXN has angiogenesis-promoting activities and specifically displays agonistic activity towards Tie2 and its downstream signaling and b) that Tie-2 and α NRXN co-precipitate from blood vessels in a developmentally regulated fashion.

2. Methods

2.1 Synthesis and validation of Peptides

Peptide A2 (GTGYLGR) and its cyclic form BbcA2 as well as the control peptide (ADSILRS) were synthesized by New England Peptide (Gardner, MA, USA). The binding of A2 and BbcA2 to Tie2 was verified through ELISA on recombinant Tie2 or by immunofluorescence /FACS on Tie2 overexpressing cells using FITC-conjugated peptides. The control peptide did not bind to Tie-2 in any of these assays (data not shown).

2.2 Proliferation assay

96 well plates were coated with 1% gelatin in PBS. HUVEC cells, pre-incubated for 24 hours with 15 μ M peptides or control medium, were seeded in the presence or absence of peptides. After 24 or 48 hours cells were fixed and colored with crystal violet. Absorbance was read at 595 nm with BioAssay Reader HTS 7000 Perkin Elmer.

2.3 Apoptosis-TUNEL assay

HUVEC cells were incubated with 15 μ M peptides. TUNEL assay (In Situ Cell Death Detection Kit; Roche, #12-156-792-910) was performed on the slides followed by image analysis using fluorescent microscopy (DM IRB; Leica). DAPI stain (Molecular probes, Invitrogen, Carlsbad, CA) was used to assess total cell number.

2.4 Protein immunoprecipitation from cultured cells and tissues

Confluent HeLa cells were washed three times with cold PBS containing 1mM Na orthovanadate and lysed in lysis buffer (Tris HCl pH 7,5 50mM, NaCl 150mM, SDS 0,1%, 1% TRITON X-100) plus protease and phosphatase inhibitors (50 μ g/ml pepstatin, 50 μ g/ml leupeptin, 10 μ g/ml aprotinin, 1mM PMSF, 100 μ M

1) **Abbreviations:** α NRXN (α Neurexin); β NRXN (β Neurexin); A2: 7 aa peptide from the second EGF like domain of alpha Nrnx; BbcA2: backbone cyclic form of A2; Bbc control: control peptide in the cyclic form; EC: endothelial cells; LNS domain: laminin A/ neurexin/ sex hormone-binding globulin repeats; NLGN: Neuroligin

ZnCl₂, 1mM Na orthovanadate, and 10mM NaF) for 30min. After centrifugation (30min at 11,000g), supernatant was quantified with the BCA Protein Assay Reagent Kit (Pierce Chemical Co., Rockford, IL).

1mg of proteins were pre-cleared by incubation for 90 minutes with protein A-Sepharose (Amersham Biosciences, Piscataway, NJ) and then incubated with rabbit anti-NRXN (1,5µg/mg) or rabbit anti-Tie2 C-20 (2,5µg/mg) for 1 h. The immune complexes were recovered on protein A-Sepharose for 90 minutes, beads were washed four times and detected by immunoblot. Proteins were separated by 8% SDS-PAGE electrophoresis gel, transferred to PVDF membrane (Millipore, Billerica, MA), incubated with mouse monoclonal anti-NRXN antibody (BD, Franklin Lakes, NJ), or anti-Tie2 C-20 antibody (Sigma-Aldrich, St. Louis, MO). Horseradish-peroxidase conjugated secondary antibody (Jackson ImmunoResearch, West Grove, PA) was used and visualized by ECL system (Amersham Biosciences, Piscataway, NJ).

Five Chicken Chorioallantoic Membranes were used for each experimental point. Frozen tissues were disaggregated and lysed 40min in ice with a lysis buffer previously set-up (Tris HCl pH 7,5 20mM; aCl 100mM, NaHCO₃ 20mM, KCl 4mM, MgCl₂ 2,5mM; 10% glycerol; 1% triton X-100; 1% CHAPS and protease and phosphatase inhibitors. After centrifugation (30min at 11,000g), supernatant was quantified. 5mg of proteins were pre-cleared by incubation for 90 min with protein A-Sepharose (Amersham Biosciences, Piscataway, NJ) and then incubated with rabbit anti-NRXN (1,5µg/mg) or rabbit anti-Tie2 C-20 (2,5µg/mg) overnight. The immune complexes were recovered on protein A-Sepharose for 90 minutes, beads were washed four times and detected by immunoblot. Proteins were separated by 4-15% gradient SDS-PAGE mini-protean TGX (Biorad, Hercules, CA), transferred to PVDF membrane (Millipore, Billerica, MA), incubated with mouse monoclonal anti-NRXN antibody (BD, Franklin Lakes, NJ), or with rabbit anti-Tie2 C-20 antibody (Sigma-Aldrich, St. Louis, MO).

2.5 Migration assay

Subconfluent HUVEC were starved overnight in M199 2%FBS and 1%BSA. The assay was done with Boyden chamber with a membrane soaked in 1% gelatin. The lower chamber was filled with serum free medium+ stimuli (10 µg/µl VEGF or 15 µM peptide), while the upper was filled with a HUVEC cells suspension (125.000 CELLS/WELL). After 5 hours the chamber was opened and the cells on the membrane were fixed in methanol and colored with Eosin/Hematoxylin.

2.6 Morphogenesis assay in matrigel

48 wells microtiter plates were covered with 150 µl of matrigel (Sigma-Aldrich, St. Louis, MO). 13000 Huvec/well were prepared in M199 10%FBS with or without 15µM of peptides. Photographs were taken after 5 hours. Total length of network of tubuli was calculated through an analysis performed by WinRHIZO program (Regent Instruments).

2.7 Sprouting assay

HUVEC cells suspended at density of 4cells/µl in M199 20% methocel and 200µl for each spheroid was seeded into non adhesive round-bottom 96-well plate. After 12h at 37°C (5% CO₂, 100% humidity) spheroids were harvested, centrifuged (15min at 300g) and resuspended in collagen solution (Roche, Indianapolis, IN) with 15 µM of peptide. Photographs were taken after 24 hours. Total length of the sprouts was calculated through an analysis performed by WinRHIZO program (Regent Instruments).

2.8 CAM assay

Fertilized chick embryos were incubated for 3 days at 37°C at 70% humidity. A small hole was made over the air sac at the end of the egg and a second hole was made directly over the embryonic blood vessels. After 7 days, cortisone acetate-treated filter disks (5 mm) were saturated with 5µl of 100 ng/ml Ang1 (R&D Systems) and peptides at concentration of 15 µM. After 3 days CAMs were fixed with PBS-4%

paraformaldehyde for 10min at room temperature, filter disks were excised and pictures were taken with a QIcam FAST1394 digital color camera (QImaging) connected to the stereomicroscope (model SZX9; Olympus).

2.9 Analysis of the Tie2 phosphorylation pathway

HUVEC cells were incubated with M199 10%FCS with 15 μ M peptides for one hour. The cells were then lysed with hot Laemli Buffer and after centrifugation the protein were quantified. 30 μ g proteins were separated by 7.5% SDS-PAGE mini-protein TGX (Biorad, Hercules, CA), transferred to PVDF membrane (Millipore, Billerica, MA), incubated with mouse monoclonal Anti-Phosphotyrosine, 4G10[®] Platinum (Millipore, Billerica, MA), with rabbit anti-pAkt (S473)(D9E) antibody, with mouse monoclonal p-p44/42 MAPK (T202/Y204)(E10) antibody (both Cells Signalling, Danvers, MA). For total protein normalization purposes we used rabbit anti-Tie2 C-20 antibody (Sigma-Aldrich, St. Louis, MO), rabbit anti-Akt (C67E7) and rabbit p44/42 MAPK (both Cells Signalling, Danvers, MA). Horseradish-peroxidase conjugated secondary antibody (Jackson Immunoresearch, West Grove, PA) was used and visualized by ECL system (Amersham Biosciences, Piscataway, NJ).

3. RESULTS

3.1 A NRXN-EGF domain peptide affects the angiogenic properties of endothelial cells and in vivo angiogenesis.

A phage display screening on Tie2 provided a large set of 7 aa target peptides of which one (A2- NH₂-GTGYPGR-COOH) was highly homologous to a sequence (NH₂-GTGYLGR-COOH) within the second EGF-like domain of α NRXN (Figure S1A). Analysis of this sequence on the α NRXN crystallography (PDB 3POY [3], PDB 3QCW [4]) showed that the peptide is exposed on the external surface of the protein where it is found in a coiled conformation (Figure S1B). Studies were pursued further with the A2 homologous sequence synthesized in a linear (A2) and cyclic (backbone cyclic, named BbcA2) forms and specific assays were performed.

3.1.1 In vitro assays for proliferation, apoptosis and migration

To verify a possible effect of the peptides on proliferation and cell death, EC were treated with A2 and BbcA2 in proliferation and TUNEL assays. Neither had a significant effect even after a 24 hours incubation (Figure S2). On the other hand, migration assays with Boyden chamber showed that BbcA2 promotes the migration of EC cells compared to all controls (Figure 1A).

3.1.2 In vitro morphogenesis

As two of the most relevant assays for angiogenic potential, we measured the ability of EC to form networks on matrigel or to sprout from collagen spheroids upon A2 and BbcA2 treatment. In the case of network formation on matrigel (Figure 1B), as with the migration assay, only BbcA2 induced a statistically significant increase in the tube network length. In the case of spheroid sprouting both A2 and BbcA2 induced positive responses, with BbcA2 inducing a response similar in extent to Ang1 (Figure 2A).

3.1.3 *In vivo* angiogenesis

We finally tested angiogenic activities of the peptides *in vivo*. We chose the chick chorioallantoic membrane (CAM) assay, which is used as an *in vivo* model both to study physiological angiogenesis and to test pro- and anti-angiogenic compounds. The peptides A2 and BbcA2 applied at day 10, induced the formation of capillary bifurcations, with BbcA2 being significantly more effective than its linear counterpart A2 (Figure 2B). Globally, these results indicate that the α NRXN EGF II peptide induces angiogenesis and that mimicking its natural conformation through cyclization positively influences its biological activities.

3.1.4 *BbcA2* induces *Tie2* phosphorylation

As a complement to the phenotypic data shown above we next tested whether A2 could modulate Tie2 signal transduction in EC. Ang1 has been demonstrated to induce activation via phosphorylation of pathways downstream to Tie2 [5]. To determine the activity of our peptides, they were incubated with EC and the phosphorylation of Tie2, Akt and Erk was measured. Figure 3 shows that BbcA2 and A2 enhance the phosphorylation of Tie-2 and of the downstream mediators AKT and ERK over their respective controls. This attests that the effect on EC migration and morphogenesis is complemented by a specific activation of Tie2 and of its main downstream mediators.

3.2 α NRXN and *Tie2* physically associate *in vivo* and their interaction is modulated during development

As an extension to the above data, the physical interaction between Tie2 and α NRXN was tested. Preliminary sets of experiments were dedicated to identifying complexes of these two proteins in overexpressing Hela Cells. α NRXN was immunoprecipitated and the resulting precipitate was immunoblotted with the antibody against Tie2. Figure S3 shows that Neurexin and Tie-2 can indeed be co-immunoprecipitated in these conditions.

To evaluate Nrnx-Tie2 interaction in a physiologically significant environment, we used the chicken chorioallantoic membrane. Figure 4 shows that at the stage E10 α NRXN and Tie2 can be efficiently co-immunoprecipitated from the CAMs but not from the brain, where the two proteins are also present. To evaluate a possible modulation of the interaction between α NRXN and Tie2 during vascular development, we performed co-immunoprecipitation experiments using CAMs at days 5, 10 or 15 days. Interestingly, even though the two proteins are expressed at significant levels at all stages, they interact only from day 10 onward and with increasing intensity (Figure 4B), when blood vessels are mature and well structured.

4. DISCUSSION

Prior to this report, we discovered that the synaptic protein α NRXN was expressed in the vascular system, but unlike the related β NRXN and NLGN, does not interact with VEGFA. Here, starting from the data collected through a parallel project, the α NRXN link to another pivotal family of vascular mediators, the Ang/Tie, was considered.

The data describe above data have two implications:

First, they provide the basis for the construction of a novel targeting tools for Tie2 starting from BbcA2 as lead compound. This receptor and its ligands are involved in disparate activities and its pharmacological modulation may have an impact in wide array of clinical settings, ranging from angiogenesis to inflammation to cancer [6]. BbcA2 is a small peptide activator of the Tie2. Although other instances of peptide activators of signaling pathways are known, in the case of Tie2, another 7aa ligand has been identified with inhibitory effects [7]. One hypothesis is that the latter peptide may function by inhibiting Tie2 dimerization [8] while BbcA2 could relieve the receptor from some inhibitory interactions, of which the most plausible is the one with the strictly related, orphan receptor Tie1[9], or that with α NRXN, shown above. The idea that the interaction between Tie2 and α NRXN is inhibitory (i.e. antiangiogenic) is supported by the fact that they associate only when blood vessels are mature and stabilized. Indeed, the CAM is a well characterized developmental model in which different stages of vascular remodeling, including endothelial “coverage” by developing smooth muscle cells, have been recognized at late stages of development [10]. In addition, it is

well known that Ang1 and Tie2 null mice display defects in the maturation and maintenance of the integrity of blood vessels [11].

Second, our data emphasize the involvement of NRXNs in the vascular system by linking them to a key family of vascular modulators such as the Ang/Tie system. In particular, along with our previous data, they suggest that Tie2 may be a ligand for α NRXN alternative to NLGN, and that α NRXN and β NRXN may have differential activities in blood vessels, i.e. pro-angiogenic for β NRXN and vessel stabilizing for α NRXN.

5 Acknowledgements

This work was supported by grants of: Italian Association for Cancer Research (AIRC) (IG 11503, IG10133; AIRC 5x1000:12182); Regione Piemonte (Finalized Health Research 2008, and 2009; Industrial Research and Precompetitive Development 2006: grants PRESTO and SPLASERBA; Technological Platforms for Biotechnology: grant DRUIDI; Converging Technologies: grant PHOENICS; Industrial Research 2009: grant BANP); Ministry of the University (FIRB, contract: RBAP11BYNP, project: Newton); European Community (contract: 318035, project: Biloba) Cassa di Risparmio di Torino Foundation (to Dr Bussolino); Italian Ministry of Health (Oncological Research Program 2006; Finalized Research 2006) (to Dr Bussolino); supported by the Italian Federation for Cancer Research (FIRC), Italian Association for Cancer Research - My First AIRC Grant (AIRC-MFAG), Banca d'Alba, Piedmont Region (Finalized Health Research Under 40) and Piedmont Foundation for Cancer Research (FPRC) Intramural Grant 5x1000 2008 (to S.M.)

6. References

- [1] A. Bottos, A. Rissone, F. Bussolino, and M. Arese, Neurexins and neuroligins: synapses look out of the nervous system. *Cell Mol Life Sci* 68 (2009) 2655-66.
- [2] A. Rissone, E. Foglia, L. Sangiorgio, S. Cermenati, S. Nicoli, S. Cimbro, M. Beltrame, F. Bussolino, F. Cotelli, and M. Arese, The synaptic proteins beta-neurexin and neuroligin synergize with extracellular matrix-binding vascular endothelial growth factor during zebrafish vascular development. *Arterioscler Thromb Vasc Biol* 32 (2012) 1563-72.
- [3] M.T. Miller, M. Mileni, D. Comoletti, R.C. Stevens, M. Harel, and P. Taylor, The crystal structure of the alpha-neurexin-1 extracellular region reveals a hinge point for mediating synaptic adhesion and function. *Structure* 19 (2011) 767-78.
- [4] F. Chen, V. Venugopal, B. Murray, and G. Rudenko, The structure of neurexin 1alpha reveals features promoting a role as synaptic organizer. *Structure* 19 (2011) 779-89.
- [5] M. Thomas, and H.G. Augustin, The role of the Angiopoietins in vascular morphogenesis. *Angiogenesis* 12 (2009) 125-37.
- [6] S. Kanda, Y. Miyata, Y. Mochizuki, T. Matsuyama, and H. Kanetake, Angiopoietin 1 is mitogenic for cultured endothelial cells. *Cancer Res* 65 (2005) 6820-7.
- [7] R. Tournaire, M.P. Simon, F. le Noble, A. Eichmann, P. England, and J. Pouyssegur, A short synthetic peptide inhibits signal transduction, migration and angiogenesis mediated by Tie2 receptor. *EMBO Rep* 5 (2004) 262-7.
- [8] K.T. Kim, H.H. Choi, M.O. Steinmetz, B. Maco, R.A. Kammerer, S.Y. Ahn, H.Z. Kim, G.M. Lee, and G.Y. Koh, Oligomerization and multimerization are critical for angiopoietin-1 to bind and phosphorylate Tie2. *J Biol Chem* 280 (2005) 20126-31.
- [9] T.C. Seegar, B. Eller, D. Tzvetkova-Robev, M.V. Kolev, S.C. Henderson, D.B. Nikolov, and W.A. Barton, Tie1-Tie2 interactions mediate functional differences between angiopoietin ligands. *Mol Cell* 37 (2010) 643-55.

- [10] D. Ribatti, B. Nico, A. Vacca, L. Roncali, P.H. Burri, and V. Djonov, Chorioallantoic membrane capillary bed: a useful target for studying angiogenesis and anti-angiogenesis in vivo. *Anat Rec* 264 (2001) 317-24.
- [11] C. Suri, P.F. Jones, S. Patan, S. Bartunkova, P.C. Maisonpierre, S. Davis, T.N. Sato, and G.D. Yancopoulos, Requisite role of angiopoietin-1, a ligand for the TIE2 receptor, during embryonic angiogenesis. *Cell* 87 (1996) 1171-80.

LEGEND TO THE FIGURES

Fig.1: BbcA2 induces migration and tube formation of HUVEC cells.

A) Boyden chamber assay shows that only BbcA2 (15 uM) displays a statistically significant effect on migration.

The graph displays the number of migrated cells from 3 different experiments (n=9 for each experimental point).

Data are expressed as mean \pm SEM. ANOVA gave $F= 8.771$.** $P < 0.01$ for BbcA2 vs Bbc Control by Bonferroni posttest.

B) Upper panel: tube morphogenesis assay: representative images showing tube formation on matrigel 5 hours after plating. Lower panel: quantification of total tube length from 3 different experiments (n=9 for each experimental point). Data are expressed as mean \pm SEM. ANOVA gave $F= 30.88$.** $P < 0.01$ for BbcA2 vs Bbc Control by Bonferroni postTest

Control: linear control peptide; Bbc control: Backbone cyclic control peptide

Fig. 2: A2 and BbcA2 induce HUVEC sprouting and in vivo angiogenesis.

A) Sprouting assays. Upper panel : representative images of collagen sprouting upon different treatments.

Lower panel: quantification of total sprout length from 4 different experiments (n=15 for each experimental point). Data are expressed as mean \pm SEM. ANOVA gave $F= 14.53$.*** $P < 0.001$ for BbcA2 vs Bbc Control ** $P < 0.01$ for A2 vs Control by Bonferroni posttest.

B) CAM angiogenesis assay: upper panel: representative images of CAMs upon different treatments. Lower panel: quantification graph from 5 different experrints. (n=25 per each experimental point). Bars represent the number of vessel bifurcations per disc counted after the different treatments Data are expressed as mean \pm SEM. ANOVA gave $F= 45.73$.** * $P < 0.001$ for BbcA2 vs Bbc ** $P < 0.01$ for A2 vs Control by Bonferroni posttest.

Control: linear control peptide; Bbc control: Backbone cyclic control peptide

Fig 3: A2 and BbcA2 induce the the activation of the Tie2 downstream pathway.

A) HUVEC cells were incubated with the different peptides for 1 hour and the phosphorylation status of Tie2, AKT and ERK was revealed by western blot.

B) Quantification analysis: bars represent the ratio between the intensity of the phosphorylated / total protein bands as quantified by image analysis.

Data are expressed as mean \pm SEM from four different experiments




Control: linear control peptide; Bbc control: Backbone cyclic control peptide

Fig 4: Tie2 and alpha Nrnx can be reciprocally immunoprecipitated from the chicken Cam.



A) Reciprocal immunoprecipitation of Tie 2 and aNrnx from chicken CAMs.

Tie2 or Nrnx were immunoprecipitated from E10 chicken embryo CAMs or brain and the precipitates were blotted for the respective partner. Nrnx and Tie2 are expressed by both CAMs and brain, but are co-immunoprecipitated with high efficiency only from the CAMs.



B) Developmental modulation of the Tie2/alphaNrnx interaction. Co-immunoprecipitation experiments performed as in panel A on CAMs at 5, 10 or 15 days of development show that the Tie2/alpha Nrnx interaction occurs only at day 10 and 15 when blood vessels are mature and well structured.





A




B









A



B



A



FigS1:

A) schematic representation of the Neuexins (alpha and beta) and Neuroligins arrangements and structures at the synapse and the position of the A2 peptide.

B) Display of the three dimensional position of the A2 peptide (sequence : GTGYLGR) on the second EGF domain of aNRXN (modeled on PDB 3POY with the Pymol software, The PyMOL Molecular Graphics System, Schrödinger, LLC.).
Green : EGF domain, red: A2 peptide.





Fig S2: Assays for proliferation and apoptosis of HUVEC cells upon A2 and BbCA2 treatment. A) treated or untreated HUVEC cells (15 μ M for 36 hours) were fixed and stained with crystal violet. Proliferation was evaluated by absorbance reading at 595 nm. B) HUVEC cells were treated as above and apoptosis was measured by a TUNEL assay. Quantification of apoptotic cells was manually performed using fluorescence microscopy




Fig S3: in vitro co-immunoprecipitation of Tie2 and alphaNRXN. alphaNRXN was immunoprecipitated from HeLa cells overexpressing alphaNRXN and Tie2 and the resulting precipitate was blotted against Tie2 (upper gel) or Nrxn (lower gel). Lysate obtained from E18 Chick embryo brain was used as a positive control for Neurexin blotting while lysate from Tie2 overexpressing porcine aortic endothelial cells was used as a positive control for Tie2 blotting.