



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

# A case of Feingold type 2 syndrome associated with keratoconus refines keratoconus type 7 locus on chromosome 13q

# This is the author's manuscript Original Citation: Availability: This version is available http://hdl.handle.net/2318/1627472 since 2017-03-08T17:26:24Z Published version: DOI:10.1016/j.ejmg.2017.01.010 Terms of use: Open Access Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright

(Article begins on next page)

protection by the applicable law.





This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in EUROPEAN JOURNAL OF MEDICAL GENETICS, 60 (4), 2017, 10.1016/j.ejmg.2017.01.010.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

(1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

(2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.

(3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), 10.1016/j.ejmg.2017.01.010

The publisher's version is available at: http://linkinghub.elsevier.com/retrieve/pii/S1769721216302555

When citing, please refer to the published version.

Link to this full text: http://hdl.handle.net/

This full text was downloaded from iris - AperTO: https://iris.unito.it/

# A case of Feingold type 2 syndrome associated with keratoconus refines keratoconus type 7 locus on chromosome 13q.

Fabio Sirchia<sup>1\*</sup>, Eleonora Di Gregorio<sup>1,2\*,</sup>Gabriella Restagno<sup>3</sup>, Enrico Grosso<sup>1</sup>, Patrizia Pappi<sup>1</sup>, Flavia Talarico<sup>1</sup>, Elisa Savin<sup>1</sup>, Simona Cavalieri<sup>3</sup>, Elisa Giorgio<sup>2</sup>, Cecilia Mancini<sup>2</sup>, Barbara Pasini<sup>1,2</sup>, Jodhbir S. Mehta<sup>4</sup>, Alfredo Brusco<sup>1,2</sup>.

\*These authors equally contributed to the work.

<sup>1</sup> Città della Salute e della Scienza University Hospital, Medical Genetics Unit, Turin, 10126 Italy
<sup>2</sup> University of Torino, Department of Medical Sciences, Turin, 10126, Italy
<sup>3</sup> Città della Salute e della Scienza University Hospital, Medical Genetics Unit, Turin, 10126, Italy
<sup>4</sup> Singapore National Eye Centre & Duke-NUS Graduate Medical School, Singapore, Singapore.

Corresponding author: Alfredo Brusco, University of Torino, Department of Medical Sciences, via Santena 19, 10126, Torino, Italy. Fax +390116706582; e-mail: <u>alfredo.brusco@unito.it</u>

Keywords: Feingold syndrome; keratoconus; array-CGH; SLITRK1; MBNL2; IPO5; MIR17HG

### Abstract

We report on a 58-year old woman with microcephaly, mild dysmorphic features, bilateral keratoconus, digital abnormalities, short stature and mild cognitive delay. Except for keratoconus, the phenotype was suggestive for Feingold syndrome type 2 (FGLDS2, MIM 614326), a rare autosomal dominant disorder described in six patients worldwide, due to the haploinsufficiency of *MIR17HG*, a micro RNA encoding gene. Karyotype showed a *de novo* deletion on chromosome 13q, further defined by array-Comparative Genomic Hybridization (a-CGH) to a 17.2-Mb region. The deletion included *MIR17HG*, as expected by the FGLDS2 phenotype, and twelve genes from the keratoconus type 7 locus. Because our patient presented with keratoconus, we propose she further refines disease genes at this locus. Among previously suggested candidates, we exclude *DOCK9* and *STK24*, and propose as best candidates *IPO5*, *DNAJC3*, *MBNL2* and *RAP2A*.

In conclusion, we report a novel phenotypic association of Feingold syndrome type 2 and keratoconus, a likely contiguous gene syndrome due to a large genomic deletion on 13q spanning *MIR17HG* and a still to be identified gene for keratoconus.

### 1. Introduction

Feingold syndromes are two very rare genetic entities characterized by digital anomalies, microcephaly, short stature and mild intellectual disability. Both are autosomal dominant, and due to the haploinsufficiency of *MYCN* (MIM 164840)(van Bokhoven, et al., 2005), encoding a transcription factor belonging to the MYC family, and Micro RNA 17 Host Gene (*MIR17HG*)(de Pontual, et al., 2011), a miRNA encoding gene.

Feingold type 1 (FGLDS1) is also associated with gastrointestinal atresia and short palpebral fissures, whereas type 2 (FGLDS2) features include brachimesophalangy, fifth finger clinodactyly and/or other digital anomalies, and dysmorphic features. The patients may also have associated cardiac anomalies (three cases) and hearing loss (two cases). Feingold type 2 is described in six patients worldwide. All have a genomic deletion spanning *MIR17HG* (MIM 609415)(Grote, et al., 2015).

No Feingold type 2 case is reported with keratoconus (MIM 148300, KTCN), a typically bilateral, non-inflammatory, progressive corneal disorder, with no gender predominance and incidence of 1 in 2,000 individuals for the isolated form (ranging from 0.0003% in Russia to 2.3% in Central India)(Gokhale, 2013) Keratoconus is associated with stromal thinning and protrusion, which causes an increase in astigmatism and in advanced cases loss of vision. Positive family history is reported in 6-23% of cases (Bechara, et al., 1996, Nowak and Gajecka, 2011). Eight KTCN loci have been mapped (see MIM 148300), including a 5.6-Mb region on 13q32 from 95,351,409 to 100,940,866 (KTCN7, MIM 614629), containing twenty-five genes.

### 2. Methods

# 2.1 Cytogenetics and array-CGH analyses

G-banding karyotype was performed on a peripheral blood sample following standard procedures. Genomic DNA was extracted from peripheral blood using a standard protocol (Qiagen, Hilden, Germany), and verified using Nanodrop spectrophotometer (Thermo Scientific). Array-CGH was

performed using a 60K whole-genome oligonucleotide microarray following the manufacturer's protocol (Agilent Technologies, Santa Clara, California, USA). Slides were scanned using a G2565BA scanner, and analyzed using Agilent CGH Analytics software ver. 4.0.81 (Agilent Technologies Inc.) with the statistical algorithm ADM-2 and a sensitivity threshold of 6.0. At least three consecutive aberrant probes identified significant copy-number changes. Reference human genomic DNA was GRCh37/hg19. We compared our findings to known CNVs listed in the Database of Genomic Variants (DGV, http://projects.tcag.ca/variation) and in the Database of Chromosomal Imbalance Phenotype Humans Ensembl Resources (DECIPHER, and in using https://decipher.sanger.ac.uk/application/). IRB approved the study.

# 2.2 Real-time PCR

TaqMan real-time quantitative PCR analysis was used to measure copy number variants at 5q35.3 in genomic DNA as follows: (a) *ZNF45* (NM\_1825594.2) exons 2, primers 5'- ggacccttctagcctgagga; 5'- ttagggacaggcaaagtgct; #16 UPL probe (Roche Diagnostics); (b) RNaseP reference gene, VIC-labeled pre-designed TaqMan gene expression assays (P/N 4316844, Applied Biosystems). Reactions were carried out on an ABI 7500 Fast real-time PCR machine using the ABI TaqMan Universal PCR master mix according to the manufacturer's instructions (Applied Biosystems). Efficiencies of the assays were similar and in a range of 90-110%. Samples from affected individuals and unrelated healthy controls were run in triplicate; the mean Ct value was used for calculations using the  $\Delta\Delta$ Ct method (Livak and Schmittgen, 2001).

### 3. Results

# 3.1 Clinical report

Our proband (II-2, Fig. 1) was born by vaginal delivery at 38 weeks of gestation. No history of prenatal exposure to teratogens or maternal illness was recorded. As a child, she suffered from

psychomotor retardation and growth delay. She underwent growth hormone replacement therapy without any improvement.

At 58-yrs., she presented with short stature (height = 134 cm,  $<2^{nd}$  centile), microcephaly (head circumference = 49 cm;  $<2^{nd}$  centile). Parents' height was 160 (father) and 150 cm (mother). The proband showed mild dysmorphic features consisting in bulbous nose and wide philtrum. She suffered from bilateral keratoconus: diagnosed at 40 yrs, with curvature of the left cornea Sim K 52.8D (Max), 46.6 (Min) astigmatism 6.2D and right cornea of Sim K 50.8 D (Max), 46.4 (Min) astigmatism of 4.4D. Both corneas showed increased anterior and poster curvature elevation with thin corneas (RE 443, LE 440) and typical asymmetric 'bow-tie' pattern on orbscan topography. Endothelial cell count on specular microscopy was within normal range for her age. She also presented digital abnormalities (brachymesophalangy, clinodactyly of the fifth fingers) and mild thoracic scoliosis (Figure 1 B-D).

She also had mild mental impairment (IQ = 75, Wechsler Adult Intelligence Scale-Revised scale). She had not been suffering from any notable psychiatric illness during her life. Abdominal ultrasound, echocardiography and thyroid profile showed no abnormalities. She had normal female external genitalia. Hearing was normal. She had only one pregnancy that was interrupted because her fetus inherited her 13q deletion. The patient went through menopause at 47 years old.

Her parents (I-1 and I-2), older sister (II-2) and nephews (III-1 and 2) were healthy (Figure 1A).

### 3.2 Laboratory findings

Because of her psychomotor and growth retardation, she underwent a karyotype analysis, which identified an interstitial deletion on the long arm of chromosome 13. Both parents showed a normal karyotype, proving the deletion was *de novo*. Array-CGH, performed to assess the extension of the deletion, identified two rearrangements: 1) a 400-kb duplication on chromosome 5q35.3, maternally inherited and likely unrelated to the phenotype; 2) a *de novo* 17.2-Mb deletion at 13q31.1q31.2 [arr 5q35.3 (178,078,984-178,457,126)x3,13q31.1q31.2(81,484,523-98,726,969), hg19] (Figure 2A).

This copy number variant (CNV) encompassed *MIR17HG* gene, known to be associated with Feingold syndrome type 2 (Figure 2B). Remarkably, the deletion partially overlapped the keratoconus type 7 *locus* (KTCN7), identified by linkage analysis on Ecuadorian families (MIM 614629) (Figure 2B)(Bechara, et al., 1996, Karolak, et al., 2011).

# 4. Discussion

We describe a never reported association of Feingold type 2 with keratoconus in a sporadic female patient. We demonstrated the patient carried a *de novo* 17.2-Mb deletion on chromosome 13q, including the *MIR17HG* gene whose haploinsufficiency causes FGLDS2. The deletion extends telomerically into the type 7 keratoconus *locus*, thus suggesting that the haploinsufficiency of a still to be identified gene at this region is causing keratoconus in our patient. The gene should map within the smallest region of overlap (SRO) between our deletion and *KTCN7 locus* (chr13:95,052,254-98,726,969 bp, hg19) (Figure 2B).

Eight genes were screened in an original Ecuadorian family where *KTCN7 locus* was mapped (*MBNL2, IPO5, FARP1, RNF113B, STK24, DOCK9, ZIC5 ZIC2* and *RNF113B*) without finding the causative mutation. Three possible candidates were suspected because of the variants found: *DOCK9, STK24*, and *IPO5*. In particular, the c.2262A>C substitution (p. Gln753His; rs191047852) in *DOCK9* segregated with the disease, and it was further shown to alter *DOCK9* splicing pattern using an *in vitro* assay (Karolak, et al., 2015). This variant is reported three times in the ExAC database (~60,000 cases), and was not found in a group of 42 Polish patients with keratoconus (Karolak, et al., 2016). Furthermore, *DOCK9* and *STK24* lay outside the SRO, and are excluded by our patient.

Using a combination of bioinformatics analyses evaluating haploinsufficiency constraint [(ExAC) database (http://exac.broadinstitute.org/)(http://dx.doi.org/10.1101/030338)], expression analysis of genes in the cornea and literature data (Chng, et al., 2013), we tried to prioritize the best candidates for *KTCN* among the 16 genes spanning our SRO (Table 1). We suggest four genes, namely importin 5 (*IPO5*), Muscleblind-Like Splicing Regulator 2 (*MBNL2*), DnaJ heat shock protein family (Hsp40)

member C (*DNAJC3*) and Ras-related protein 2A (*RAP2A*), can be considered as candidates and should be extensively screened in the original Ecuadorian family, because they are expressed in the target tissue and with a high haploinsufficiency constraint (Czugala, et al., 2012). Among them, IPO5 –a nuclear protein import as nuclear transport receptor - and *MBNL2* - a member of the muscle blind protein family that regulates alternative splicing is required for terminal differentiation of muscle and photoreceptor tissues- have already been screened for point mutations in the coding region (Czugala, et al., 2012), but small deletions/duplications, mutations in regulatory regions, deep intronic splice-site mutations or even position effects may have been missed.

Seven cases reported in Decipher database and six described in the literature have a deletion encompassing *MIR17HG*. None of them was reported with keratoconus (see figure 2 for a scheme). Only three of them have a rearrangement spanning *KTCN7 locus;* however, because these patients are infants, a genotype-phenotype correlation is not possible, considering keratoconus generally occurs during puberty (cases A, J and K, fig. 2B). One deletion (Sharaidin, et al., 2013) overlaps ~500 kb with the *KTCN7 locus* without including *MBNL2* and *IPO5* (Tassano, et al., 2013), and two extend ~4-5 Mb with the *KTCN7 locus* (cases A and J, fig.2B).

Among the deleted genes in our patient, we also noted *SLITRK1*, which has been previously associated with psychiatric diseases such as trichotillomania, Gilles de la Tourette syndrome and obsessive-compulsive disorder (Abelson, et al., 2005). No such psychiatric illnesses were present in our proband, likely due to the incomplete penetrance and variable expressivity associated to *SLITRK1* haploinsufficiency (Proenca, et al., 2011).

In conclusion, we describe a possible contiguous gene syndrome phenotypically characterized by Feingold syndrome type 2 and keratoconus, due to a large genomic deletion on 13q overlapping *MIR17HG* and a still to be identified gene for keratoconus.

### Figure legends

# Figure 1. Pedigree and clinical phenotype

A) Family pedigree: filled circles - affected subject; empty symbols - unaffected subjects; arrow indicates the proband; asterisk on the symbol indicates available genomic DNA; dup: duplication; del: deletion; wt: wild type. B) Mild facial dysmorphisms consisting in bulbous nose and wide philtrum. C, D) Brachydactyly of hands and feet.

### Figure 2. Array-CGH analysis and overview of the deleted genomic region

A) Array-CGH results using the 60K Agilent platform showing the 17.2-Mb deletion on chromosome 13q. B) Genes located in the 13q31.1q31.2 genomic region between 81-101 Mb (hg19). In black Keratoconus type 7 locus as described in (Czugala, et al., 2012) and boxed in light grey the SRO defined by our patient's deletion. In red, our patient deletion. Below in grey, cases with a *MIR17HG* gene deletion, reported in the Decipher database and literature. Boxed in light purple the SRO of Feingold 2 disease. The figure 2B was modified from UCSC genome browser, using custom-generated tracks (Raney, et al., 2014).

# Acknowledgments

The authors declare no conflicts of interest related to the publication of this article. The authors are grateful to the patient and her family for agreeing to take part in this study. This work was supported by MURST 60% to AB. This study makes use of data generated by the DECIPHER Consortium. A full list of centers who contributed to the generation of the data is available from http://decipher.sanger.ac.uk/ and via email from decipher@sanger.ac.uk. Funding for the project was provided by the Wellcome Trust. The case reported here has been entered in the "DECIPHER" database (http://decipher.sanger.ac.uk/) with the code number 324479.

### References

Abelson, J.F., Kwan, K.Y., O'Roak, B.J., Baek, D.Y., Stillman, A.A., Morgan, T.M., Mathews, C.A., Pauls, D.L., Rasin, M.R., Gunel, M., Davis, N.R., Ercan-Sencicek, A.G., Guez, D.H., Spertus, J.A., Leckman, J.F., Dure, L.S.t., Kurlan, R., Singer, H.S., Gilbert, D.L., Farhi, A., Louvi, A., Lifton, R.P., Sestan, N., State, M.W., 2005. Sequence variants in SLITRK1 are associated with Tourette's syndrome. Science, 310, 317-320. Bechara, S.J., Waring, G.O., Insler, M.S., 1996. Keratoconus in two pairs of identical twins. Cornea, 15, 90-93.

Chng, Z., Peh, G.S., Herath, W.B., Cheng, T.Y., Ang, H.P., Toh, K.P., Robson, P., Mehta, J.S., Colman, A., 2013. High throughput gene expression analysis identifies reliable expression markers of human corneal endothelial cells. PLoS One, 8, e67546.

Czugala, M., Karolak, J.A., Nowak, D.M., Polakowski, P., Pitarque, J., Molinari, A., Rydzanicz, M., Bejjani, B.A., Yue, B.Y.J.T., Szaflik, J.P., Gajecka, M., 2012. Novel mutation and three other sequence variants segregating with phenotype at keratoconus 13q32 susceptibility locus. European Journal of Human Genetics, 20, 389-397.

de Pontual, L., Yao, E., Callier, P., Faivre, L., Drouin, V., Cariou, S., Van Haeringen, A., Genevieve, D., Goldenberg, A., Oufadem, M., Manouvrier, S., Munnich, A., Vidigal, J.A., Vekemans, M., Lyonnet, S., Henrion-Caude, A., Ventura, A., Amiel, J., 2011. Germline deletion of the miR-17 approximately 92 cluster causes skeletal and growth defects in humans. Nat Genet, 43, 1026-1030.

Gokhale, N.S., 2013. Epidemiology of keratoconus. Indian journal of ophthalmology, 61, 382-383. Grote, L.E., Repnikova, E.A., Amudhavalli, S.M., 2015. Expanding the phenotype of feingold syndrome-2. Am J Med Genet A, 167A, 3219-3225.

Karolak, J.A., Polakowski, P., Szaflik, J., Szaflik, J.P., Gajecka, M., 2016. Molecular Screening of Keratoconus Susceptibility Sequence Variants in VSX1, TGFBI, DOCK9, STK24, and IPO5 Genes in Polish Patients and Novel TGFBI Variant Identification. Ophthalmic Genet, 37, 37-43.

Karolak, J.A., Rydzanicz, M., Ginter-Matuszewska, B., Pitarque, J.A., Molinari, A., Bejjani, B.A., Gajecka, M., 2015. Variant c.2262A>C in DOCK9 Leads to Exon Skipping in Keratoconus Family. Investigative ophthalmology & visual science, 56, 7687-7690.

Karolak, J.A., Kulinska, K., Nowak, D.M., Pitarque, J.A., Molinari, A., Rydzanicz, M., Bejjani, B.A., Gajecka, M., 2011. Sequence variants in COL4A1 and COL4A2 genes in Ecuadorian families with keratoconus. Molecular vision, 17, 827-843.

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods, 25, 402-408.

Nowak, D.M., Gajecka, M., 2011. The genetics of keratoconus. Middle East Afr J Ophthalmol, 18, 2-6. Proenca, C.C., Gao, K.P., Shmelkov, S.V., Rafii, S., Lee, F.S., 2011. Slitrks as emerging candidate genes involved in neuropsychiatric disorders. Trends Neurosci, 34, 143-153. Raney, B.J., Dreszer, T.R., Barber, G.P., Clawson, H., Fujita, P.A., Wang, T., Nguyen, N., Paten, B., Zweig, A.S., Karolchik, D., Kent, W.J., 2014. Track data hubs enable visualization of user-defined genome-wide annotations on the UCSC Genome Browser. Bioinformatics, 30, 1003-1005.

Sharaidin, H.S., Knipe, S., Bain, N., Goel, H., 2013. Clinical features associated with a 15.41 Mb deletion of chromosome 13q encompassing the MIR17HG locus. Clin Dysmorphol, 22, 68-70.

Tassano, E., Di Rocco, M., Signa, S., Gimelli, G., 2013. De novo 13q31.1-q32.1 interstitial deletion encompassing the miR-17-92 cluster in a patient with Feingold syndrome-2. Am J Med Genet A, 161A, 894-896.

van Bokhoven, H., Celli, J., van Reeuwijk, J., Rinne, T., Glaudemans, B., van Beusekom, E., Rieu, P., Newbury-Ecob, R.A., Chiang, C., Brunner, H.G., 2005. MYCN haploinsufficiency is associated with reduced brain size and intestinal atresias in Feingold syndrome. Nat Genet, *37*, 465-467.

Table 1. Constraint metrics (ExAC database) and expression in cornea for genes within the SRO.

Gene	Ζ	pLI	Cornea expression <sup>a</sup>
ABCC4	0.79	0	No expression
CLDN10	0.92	0.05	Young CECs
DNAJC3	1.14	0.95	Relatively low in all
DZIP1	0.35	0	Cultured Cells
FARP1	0.77	0.05	Cultured and Young
HS6ST3	2.16	0.75	No expression
IPO5	2.64	1	Cultured and Young
MBNL2	3.03	0.9	Relatively low in all
OXGR1	1.07	0	No expression
RAP2A	3.63	0.72	Relatively low in all
SOX21	n.d.	n.d.	No expression
UGGT2	1.77	0	Old CECs
N1 /			

Note:

CEC - Corneal Endothelial Cells, Low expression <15 Read a) see supplementary table for details

Figure1











