

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Pinus nigra anthropogenic treelines in the central Apennines show common pattern of tree recruitment

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1622728> since 2017-01-20T11:14:12Z

Published version:

DOI:10.1007/s10342-016-0999-y

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

[Piermattei A., Lingua E., Urbinati C., Garbarino M. 2016. Pinus nigra anthropogenic treelines in the central Apennines show common pattern of tree recruitment. European Journal of Forest Research 135: 1119-1130. doi: 10.1007/s10342-016-0999-y.]

The publisher's version is available at:

[<http://link.springer.com/article/10.1007%2Fs10342-016-0999-y>]

When citing, please refer to the published version.

Link to this full text:

[<http://hdl.handle.net/2318/1622728>]

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

[Click here to view linked References](#)

***Pinus nigra* anthropogenic treelines in the central Apennines show common pattern of tree recruitment**

Alma Piermattei^{1,3}, Emanuele Lingua², Carlo Urbinati¹, Matteo Garbarino^{1*}

1. Department of Agricultural, Food and Environmental Sciences, Marche Polytechnic University, Via delle Breccie Bianche 10, I-60131, Ancona, AN, Italy

2. Department of Land, Environment, Agriculture and Forestry, University of Padova, Viale dell'Università 16, I-35020, Legnaro, PD, Italy

3. Swiss Federal Research Institute WSL, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

* Correspondence: Matteo Garbarino, Department of Agricultural, Food and Environmental Sciences Marche Polytechnic University, Via delle Breccie Bianche 10, I-60131, Ancona, Italy
Tel.: +39 071-2204599
E-mail: m.garbarino@univpm.it

Acknowledgements

This research was partially financed by the Marche Polytechnic University through the project No. 4524/2009 “Spatio-temporal dynamics of forest vegetation and climate change in the Marche region”. We wish to thank the following institutions and people for providing sampling authorization, logistic support, field and laboratory assistance: Monti Sibillini National Park, Gran Sasso-Monti della Laga National Park, Sirente-Velino Regional Park, and Bruno Petriccione, Francesco Renzaglia, Valeria Gallucci, Emidia Santini, Luca Bagnara, Matteo Giove, Andrea Cola, Simone Cingolani, Marco Altieri, Alessandro Vitali, Alan Crivellaro and Edoardo Piermattei.

Abstract

1
2 Treeline position, structure and composition are shaped by natural and anthropogenic factors. At human disturbed
3
4 treelines it is particularly difficult to disentangle the specific role of natural and anthropogenic drivers controlling tree
5
6 recruitment dynamics following land abandonment. We tested for a possible common regeneration pattern of *Pinus nigra*
7
8 Arn. at four upper treeline ecotones in the central Apennines (Italy). The sites were selected based on i) the occurrence of
9
10 natural encroachment of *P. nigra* above 1600 m a.s.l., and ii) the mountain top elevation higher than 2000 m a.s.l. We
11
12 assessed structure and spatio-temporal patterns of *P. nigra* advancing regeneration using point and surface pattern
13
14 analyses. We mapped, measured and dated 845 trees sampled on a total surface area of 336 ha. *P. nigra* is the only tree
15
16 species expanding at high altitude and features a scattered process that started 35-40 years ago, with a maximum
17
18 recruitment frequency between 1995 and 2003. *Pinus* regeneration appeared over-dispersed along the slope at a scale
19
20 range of 12-18 m, but saplings were clumped at short distances (1 m). We found spatial segregation between saplings and
21
22 young trees at intermediate distances (8-17 m) and small patches of young trees distributed along the treeline ecotone.
23
24 The spatial pattern of *P. nigra* encroachment in the central Apennines revealed a replicable model independent of treeline
25
26 topography and local disturbance histories.
27
28
29
30
31
32
33

34 **Keywords** European black pine; pine regeneration; point pattern analysis; secondary succession; surface pattern analysis
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

Upper treelines are typical examples of margin vegetation and important ecological boundaries of dynamic landscapes (Bolli et al. 2007). Margins are of particular interest in vegetation science since they demonstrate a limit to the adaptation of a changing environment. Plant migrations are expected when these changes occur, however it is necessary to distinguish between climate and land use induced changes (Crawford 2008). The distinction between natural and cultural (human dominated) treelines should be very clear when assessing their structures and dynamics (Troll 1973; Miede and Miede 2000). Treelines can be classified as: i) climatic, ii) orographic/edaphic, and iii) anthropogenic depending on their main controlling factors (Holtmeier and Broll 2005). Natural constraints such as climate, soil and topography prevail on sites less disturbed by humans, whereas anthropogenic treelines develop under more severe human impacts (e.g. forest cover reduction or clearing, fire, cattle grazing) that alter site conditions (Holtmeier and Broll 2005). Climate and land use changes appear to be the two main and often synergic drivers that trigger tree establishment at upper treeline ecotones, causing either gap infilling or treeline upward shifts (Vittoz et al. 2008). A recent meta-analysis on 166 treeline sites around the world showed that 52% are advancing, 47% are stable and only 1% in recession (Harsch et al. 2009). The same authors use the treeline form as an indicator for controlling mechanisms and response to climate change. Diffuse and *krummholz* treelines indicate an advancement trend, whereas abrupt and island treelines appear to be more stable (Harsch et al. 2009; Harsch and Bader 2011). Climatic treelines, their position and advancement have been widely and extensively studied worldwide in the last decades (e.g. Troll 1973; Tranquillini 1979; Kullman 1998; Theurillat and Guisan 2001; Holtmeier and Broll 2005; Körner 2012). More recently, studies on anthropogenic treelines developed significantly, given that large parts of European upper treelines have been lowered by human induced disturbances. Their intensity and frequency decrease have caused widespread tree recruitment processes, with or without a treeline upward shift, on several mountain ranges (e.g. Carrer and Urbinati 2001; Dalen and Hofgaard 2005; Motta et al. 2006; Bolli et al. 2007; Rössler et al. 2008; Vittoz et al. 2008; Potthoff 2009).

In areas where man has had a significant impact on mountain ecosystems, especially at regional scale, land use change is an important driver of treeline ecotone dynamics in relation to shifting disturbance intensity (Gehrig-Fasel et al. 2007; Garbarino et al. 2013). In the Italian Alps the climatic upper treelines are limited to rocky, steeper and less accessible slopes and their current altitudes are largely influenced by anthropogenic and topographic factors (Leonelli et al. 2009). Structure and dynamics of treeline ecotones are strictly connected and influenced by species composition (Compostella and Caccianiga 2016). A transition from intensive to extensive grazing may favour tree recruitment during the initial stages, since grass species competition is reduced and soil is more exposed (Motta et al. 2006). Tree recruitment patterns then become largely dependent on the availability of microsites that provide suitable conditions for seedling

1 establishment and growth. The microsite availability can decrease over time reducing the rate of tree encroachment (Bolli
2 et al. 2007).

3
4 Treeline species composition is also a crucial aspect for tree recruitment in Mediterranean mountains such as the
5 Apennines, the Dinaric Alps and the Balkans. Natural treelines were most likely formed by *Pinus heldreichii* (or *P.*
6 *leucodermis*), *P. peuce*, and *Picea abies* which can still be found at some sites up to and above 2000 m a.s.l. (Barbero et
7 al. 1995). *P. mugo* in the form of shrub is often located above treeline up to 2500-2600 m a.s.l., but today several
8 anthropogenic treeline ecotones in the Apennines and less frequently in the Dinaric Mountains feature the dominance of
9 broadleaved montane tree species such as *Fagus sylvatica* and locally *Acer heldreichii*. The reduction of cold-adapted
10 needle-leaved taxa such as *Pinus* is probably due to a combination of past biogeographic and human induced events
11 (Körner 2012). In the Apennines, a NW-SE 1000 km mountain range in Italy with several peaks higher than 2000 m a.s.l.,
12 high elevation forests since post-würmian times were first largely cleared for hunting, and later transformed into wood
13 pastures or grasslands (Pedrotti 1969; Dibari et al. 2015). These ecotones, after previously severe pressure and increasing
14 abandonment, are now marginal land and often included in protected areas. As in other European mountain regions, land
15 abandonment has been a complex and non-linear process. In Italy, the most recent major change in land use occurred in
16 the mid-20th century with the migration of mountain and rural populations downhill or to urban areas. In the Apennines
17 this mass movement led to a dramatic decline in traditional grazing (Santilocchi and D'Ottavio 2005) and agricultural
18 practices, but favoured diffused secondary ecological succession (Torta 2004; Caballero et al. 2009; Pelorosso et al. 2009).
19 Between 1967 and 2007 permanent grassland cover in Italy decreased by 37%, and the dairy cow population declined by
20 35% in 1975-2007 with a negative peak after 1990 (Huyghe et al. 2014).

21
22 The Apennines treelines are on average located between 1600-1800 m a.s.l. and largely formed by European
23 beech (*Fagus sylvatica*); *Pinus* species (*P. mugo*, *P. nigra* var. *laricio*, *P. heldreichii*) only prevail at a few sites, especially
24 in central and southern sectors. The beech treelines can be defined as 'abrupt' and relatively stable due to the species
25 intolerance to soil water deficit at higher altitude and to anthropic disturbances (Stanisci et al. 2000; Pezzi et al. 2008;
26 Catorci et al. 2012). The pine treelines are considered 'diffuse' and more prone to altitude shifts. *Pinus nigra* treelines are
27 an exception since this species is commonly found at lower elevation between 250–1600 m a.s.l. (Isajev et al. 2004). In
28 the Balkans and Iberian Peninsula, they can reach 2000 m a.s.l. and form natural treelines (Richardson 2000). Due to its
29 being a pioneer species *P. nigra* was planted extensively in the Apennines throughout the 20th century on steep, rocky
30 and overgrazed slopes between 1000-1600 m a.s.l., to reduce erosion and landslides. At these sites, we observed diffuse
31 *P. nigra* regeneration, as the only encroaching tree species above the closed forest margin even above 2000 m a.s.l.
32 (Piermattei et al. 2012; Piermattei et al. 2014). This expansion process at high altitude occurred due to a favourable
33 coincidence of several facilitating factors such as grazing pressure decline, climate warming and seed supply from
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 neighbouring pine plantations and also thanks to the ecological traits of this cold-adapted, drought resistant and
2 competitive species (Richardson and Rejmanek 2004; Cousens et al. 2008). Previous papers described the progressive
3 enlargement of the study area and focused on the chronological pattern, tree growth performance and climate sensitivity
4 of *P. nigra* at high elevation in the central Apennines (Piermattei et al. 2012; Piermattei et al. 2014).
5

6
7 In this study, we compared spatial pattern and structure of *Pinus* regeneration at four anthropogenic treeline
8 ecotones of the central Apennines exhibiting different topographic features and land use histories. Tree recruitment
9 processes above the current treeline were studied at fine spatial scales in order to assess the influence of local
10 environmental changes (Veblen 1992). We tested for a possible common spatial pattern suggesting the presence of a
11 predictable underlying process in the central limestone Apennines. Our main hypotheses were: 1) *P. nigra* scattered
12 encroachment at high elevation is a widespread process with a typical spatial pattern independent of microsite conditions;
13 2) trees of different age classes are spatially segregated due to intra-species competition; 3) density, tree size and age
14 structure of *Pinus* regeneration is constrained by elevation.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2. Materials and methods

2.1 Study area and sampling design

We sampled four upper treeline ecotones in the central Apennines (Italy) across the Marche (Mt. Vettore - VET) and Abruzzo (Mt. San Franco - SFR, Mt. Sirente - SIR, and Mt. Ocre - OCR) regions (Fig. 1). The four ecotones were selected by field visits and aerial photographs interpretation according to two requirements: presence of *Pinus* regeneration over 1600 m a.s.l., and a mountain top elevation higher than 2000 m a.s.l. For the purpose of this study we considered as forestline the upper limit of a closed canopy forest (either *F. sylvatica* natural forest or *P. nigra* plantation), and as treeline ecotone the area between the forestline and the encroached trees at the highest elevation reached. All study sites are included in different types of protected areas: VET is in the Sibillini National Park, SFR in the Gran Sasso-Monti della Laga National Park, SIR in the Sirente-Velino Regional Park, and OCR in the Monte Ocre-Acquazzese State Forest. All sites are on calcareous bedrock, and the peak elevation ranges from 2132 m a.s.l. at SFR to 2478 m a.s.l. at VET. The forestline elevation ranges from 1350 m a.s.l. at OCR to 1650 m a.s.l. at SIR and the mean slope from 32° to 36° at all sites, except for SIR (19°) that features a gentler topography (Table 1). All sites share a temperate oceanic macrobioclimate (*sensu* Rivas-Martinez and Rivas-Saenz 2009) with recurrent droughts in July and August and precipitation peaks in autumn and early spring (Table 1). Snowfalls are more common and abundant in late winter (February and March), but snow permanence on the ground is limited and discontinuous (De Bellis et al. 2010).

The four sites share some general common land use traits but have different local histories (Table 2). Livestock grazing at high elevation usually from May to mid-October is a tradition, but is now significantly decreasing at all sites. These grasslands experienced different grazing pressure: sheep, cows and more recently horses have contributed to the pastoral histories of these sites. Local records are lacking but in general grazing pressure was high (Peroni et al. 2000) and sheep largely decreased at all sites in the last 20-40 years. More specifically, sheep decreased by 50% at VET over the last 40 years (Piermattei et al. 2012), and from 5000 to 3000 units in the 1980-2000 period at SFR, in the Gran Sasso area (Coppinger et al. 1983; Rolando et al. 2001). Grazing pressure was the most important human induced disturbance during the 1950s at OCR, but there was a dramatic decline between 1954 and 1995 (Peroni et al. 2000). There is still extensive livestock grazing (cows and then horses) at SIR, due to its gentler topography of a karst plateau and less strict nature conservation constraints (Piermattei et al. 2014). Uncontrolled intensive grazing in the past caused widespread slope erosion processes that required extensive *Pinus* plantations in the studied region between the 1950s and 1970s. Their ages are respectively: 45 (SIR), 48 (SFR), 51 (OCR) and 63 (VET) years and their upper limit is the current forestline. Natural *Pinus nigra* forests were not found in the proximity of the study sites.

2.2. Data collection

Between 2009 and 2013 we surveyed an overall treeline ecotone surface area of 336 ha. At VET and SFR all *Pinus* trees present from the forestline upwards were sampled. At SIR and OCR, due to the larger extension of the ecotone area, we sampled in a transect (length 1700 m and width 218 m at SIR; length 440 m and width 150 m at OCR) established along the maximum slope from 1600 m a.s.l. upwards. A total of 845 encroached *P. nigra* trees were mapped above the forestline with a Trimble GeoXT GPS receiver and their positions corrected with post-processing differential correction (0.5-1 m estimated accuracy). Basal stem diameter, cambial age, total height and number of bearing cones were measured for each *Pinus* individual. Cambial age of all trees with a basal stem diameter ≥ 4 cm was determined after the extraction of one basal increment core. For smaller trees we used the number of internodes (terminal bud scars) as a tree age proxy. Percentage cover of microhabitat types (bare soil, herbs and shrubs) was also visually estimated on 1 m radius subplots centred on each *Pinus* individual position.

2.3. Data analysis

Treeline structure was described by computing classic stand structure attributes such as tree density and size (stem diameter and total height). The relationships between tree recruitment years, tree size and topography were analysed through linear regression analyses.

Point Pattern Analysis (PPA) methods were applied to assess *Pinus* tree spatial patterns within the study area and the association between saplings (age < 10 years) and young trees (age > 15 years) at different spatial scales. *Pinus* regenerating trees were classified as saplings and young trees in order to obtain balanced classes of more than 100 individuals (Wiegand and Moloney 2014) within the narrow age interval (< 40 years) of the observed process. The spatially explicit dataset used for the analyses comprised only 787 living trees (171 VET, 241 SFR, 193 SIR and 182 OCR), 226 saplings, 250 intermediate and 311 young trees. We used pair-correlation functions, both univariate ($g(r)$) and bivariate ($g_{12}(r)$) (Stoyan and Stoyan 1994). As low between-site variability was found from the preliminary ordination analyses, supported by the similar overall univariate spatial pattern found in all sites, we conducted the spatial analyses considering the four sites as pseudo-replications to assess the global average spatial pattern of *P. nigra* treeline ecotone expansion in the central Apennines. The results from each site were thus combined in one average graphic function, using the 'combine replicates tool' included in the Programita software (Wiegand and Moloney 2004; Petritan et al. 2014). The univariate pattern of *Pinus* trees was contrasted against the heterogeneous Poisson null model to account for first-order effects. This null model differs from the complete spatial randomness (CSR) one in that a function $\lambda(x, y)$ varies with location (x, y) but maintains the independence of the occurrence of any point in respect to any other (Wiegand and Moloney 2004). We computed the intensity function λ non-parametrically, directly from the data, using the Epanechnikov

1 kernel estimators (Wiegand and Moloney 2014). Only points of the same category of interest were considered. We used
2 a radius of 20 m for the moving window estimator, considering that the influence of small trees does not exceed this
3 distance in our high elevation sites. For the bivariate analyses between saplings and young trees, we adopted the
4 antecedent condition null model applying the heterogeneous Poisson to only one category (saplings) of trees, maintaining
5 the other fixed (young trees) in order to assess if pioneer individuals act as attracting microsite for younger seedlings,
6 accelerating the encroachment process. The 95% confidence intervals for both univariate and bivariate analyses were
7 computed from 999 Monte Carlo simulations (Stoyan and Stoyan 1994; Wiegand and Moloney 2004) and the goodness-
8 of-fit (GoF) test for null hypothesis was performed (Diggle 2003). All analyses were done only for classes with more than
9 15 trees (Camarero et al. 2000), applying a 1 m lag distance and a maximum distance of 100 m with the grid-based
10 software Programita (Wiegand and Moloney 2004), adopting a grid size of 1 m² and a ring width of 5 m.
11
12
13
14
15
16
17
18
19

20 In order to obtain a spatial localization of tree groups with similar age we used a local index of spatial association
21 (LISA). In particular, the spatio-temporal patterns of tree recruitment were described with the Getis-Ord $G_i^*(d)$ (Getis
22 and Ord 1992) index, using tree age as a quantitative variable. The LISA statistics can detect the local clustering around
23 the individual location and add depth to inferences where a single measure of global association (e.g. Moran's I) can
24 provide little meaningful information (Lamedica et al. 2011; Carrer et al. 2013). Based on previous PPA results, we
25 selected two distance classes (15 and 50 m) to assess fine- and coarse-scale structure potentially related to the processes
26 at stand (Carrer et al. 2013; Petritan et al. 2014) and treeline scale. All local $G_i^*(d)$ analyses were computed with the
27 Rookcase Excel add-in (Sawada 1999).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. Results

3.1. Treeline structure

Pinus nigra recolonization in the central Apennines is a scattered process of isolated trees (average of 5-6 individuals per hectare). Tree size is small at all sites (mean height ranges between 119-211 cm and mean diameter between 7-13 cm) with higher values at SIR (Table 3). The mean tree age ranged between 15 and 19 years, confirming the recent process at all sites. High standard deviation values show high within-site variability. Oldest trees age suggests that the first encroachment events started at least 35-40 years ago (1975-1980). Seeds were coming from neighbouring *P. nigra* plantations located at lower altitude. Recruitment frequencies are normally distributed and exhibit synchronized peaks within the period 1995-2003 at all sites, with a decrease in the last 5-10 years.

Altitude has a negative effect on regeneration density **only at VET and SFR treelines** but not on tree age. Trees of all ages were found at various elevations, revealing that the tree establishment occurred all along the available altitudinal gradient (Fig. 2). Maximum elevation reached by *P. nigra* is 2106 m a.s.l. but most trees are located at 1750-1800 m a.s.l. at VET, SFR and OCR, and 1850-1900 m a.s.l. at SIR. Tree height was positively associated with cambial age and negatively with elevation at all sites (Fig. 3). However, a decrease in tree height and diameter with increasing elevation was evident at VET, weak at SFR, and absent at OCR and SIR. 22.6% of surveyed trees were bearing cones, but only 13% had more than 20 cones/tree. Mean age of these trees was around 20 years, but some individuals reached maturity at 10 years old, whereas the **average maturity age** is 40 years (Farjon 2013).

Microhabitat type around the sampled trees varies at the four sites: grass cover was predominant at OCR (57%) and VET (50%), whereas bare soil (e.g. outcrops, rocky debris) dominated at SIR (69%) and SFR (55%). Presence of shrubs **(mostly *Juniperus communis*)** was less common, ranging from 4% to 14% (SIR and VET).

3.2. Spatial patterns

The overall spatial pattern of *P. nigra* recruitment (univariate PPA under heterogeneous Poisson null-model) showed a significant (GoF: $p \leq 0.01$) tendency towards over-dispersion between 12 and 18 m (Fig. 4). This average pattern is the result of the combined replicates procedure obtained from the random pattern observed at OCR, SFR and SIR, and from the weak departure from randomness observed at VET. Stratifying the dataset according to age groups, young trees showed no significant clustered distribution at any spatial scale and were within the expected distribution up to 40 m (**Online Resource 1 and 2**). According to the overall population structure, both age groups revealed a significant departure from randomness to regularity between 12-17 m and 13-17 m respectively (**Online Resource 1**). Considering the bivariate g_{12} (r) under antecedent condition null model, negative spatial interaction (spatial segregation) existed between the two

age classes at 8-17 m (Fig. 5, GoF: $p \leq 0.01$). The only exception was SFR, where we observed a random spatial relationship between saplings and young trees, without any departure from the confidence envelopes.

The local $G_i^*(d)$ statistic, rasterized on the treeline ecotone maps (Fig. 6), showed groups of even-aged trees older (hot spots) or younger (cold spots) than the mean age and described the spatiotemporal pattern of tree recruitment. Local $G_i^*(d)$ statistic allowed the groups of trees with similar age to be identified and located along the studied slopes. A patchy surface pattern was common to all four sites, where encroaching trees established in different age cohorts in groups of variable size (5-50 m) randomly distributed along the treeline ecotone. The smaller (5 m) significant groups were present at SIR and the pattern was consistent at both 15 and 50 m spatial scales.

4. Discussion

1
2 About 10 years ago, we first observed an upward expansion process of *Pinus nigra* at high altitude in several sites of the
3 central Apennines: this is clearly a natural encroachment process with an anthropic origin. *Pinus* plantations forming the
4 current forestline are important seed sources for supplying the observed process. However, at SIR and at a different site
5 (ACU), not included in this analysis (Piermattei et al. 2012), the planted pines are highly scattered and very distant from
6 the encroachment sites, demonstrating the long-distance dissemination provided by this tree species. Nonetheless given
7 these conditions it is incorrect to define the ongoing process as a typical treeline upward shift driven mainly by natural
8 constraints such as climate and topography.
9

10
11 This research highlighted a common pattern that underlies the tree recruitment process at all sites, regardless of
12 some existing differences of local features such as topography and past land use. At VET and SFR ecotones the density
13 of newly established trees decreased along the elevation gradient, but tree size not very much (height and diameter). A
14 negative correlation between tree growth and elevation is well documented for other pine species such as *P. uncinata* in
15 the Pyrenees (Camarero and Gutierrez 2004; Wiegand et al. 2006) and *P. cembra* in the Alps (Vittoz et al. 2008). A
16 similar pattern could be expected for tree age and altitude but we did not observe it in our sites. The absence of a clear
17 relationship between tree age and elevation was also observed in the Alps (Carrer et al. 2013) and Pyrenees (Camarero et
18 al. 2000, Batllori et al. 2010). All this seems to prove that *Pinus* recruitment is not a progressive process along the slope,
19 enhancing the role of the occurrence of safe sites that are favourable microsites for the establishment and growth of
20 regeneration. In general, encroached trees are sparser at VET and OCR (2-3/ha), compared to SFR and SIR (6-7/ha),
21 probably due to the prevailing grass cover where intraspecific competition between regenerating trees is higher (Table 3).
22 For these reasons, the shrubs cover and the presence of rocky surfaces can be considered safe sites for the establishment
23 of *Pinus* trees.
24

25
26 Despite the differences in topographic features and land use histories a common spatial pattern of *Pinus*
27 regeneration seems to prevail at the four studied sites. This significant overall pattern appears clearly when combining
28 together the observed pair-correlation functions of the four sites. There was no evidence of a clustered structure either for
29 young trees or for the overall population (Fig. 4, 5b). These patterns were consistent in all the sites and suggest that the
30 regeneration is sparsely distributed and not organized in even-aged tree cohorts. Significant negative spatial relationships
31 between saplings and young trees were found in all sites, indicating segregation between the two age classes (Fig. 5).
32 Spatial segregation between younger and older trees may be due to competition mechanisms and this pattern is commonly
33 found in different forest ecosystems where regeneration mainly establishes within canopy gaps. This suggests that in our
34 study sites, older trees do not provide a more favourable microhabitat for younger ones. Anyway, older trees were rare
35 and may have affected younger individuals. A two-fold interpretation follows: the first one implies independence or
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 competition rather than facilitation among trees, since site conditions are not extreme at these treeline ecotones. According
2 to the stress gradient hypothesis, facilitation is more evident in harsher environments (Callaway et al. 2002), as occurring
3 in higher elevations treeline ecotones of the Alps (Lingua et al. 2008; Carrer et al. 2013). The second interpretation is that
4 the pioneer trees, given their current size, are too small to significantly influence the surroundings and improve the
5 microsite conditions for seed germination and seedling growth. The surface pattern analysis revealed that in all the study
6 sites hot spots of older trees were distributed along the slope, without any significant influence of elevation (Fig. 6). Our
7 results confirm that *Pinus* regeneration is only weakly clustered and seems not to be influenced by any particular linear
8 gradient. The absence of such gradient makes it very difficult, at this stage of the process, to predict the further expansion
9 of *P. nigra* regeneration within the studied sites. The common spatial pattern observed may be explained by the favourable
10 contingent factors acting at regional scale, such as land use change and global warming that may overcome the local
11 differences occurring in the studied ecotones.
12

13 **Expansion of *P. nigra* at high altitude in the central Apennines is a spontaneous process** occurring within an
14 anthropogenic ecotone where the forestline pine plantations act as seed sources and pasture abandonment is the
15 predisposing factor. The anthropogenic nature of these upslope secondary successions and the local relevance of land use
16 changes hamper the natural treeline ecological theories being applied to these study cases. However, given the extended
17 distribution of *P. nigra* (especially in plantations) in the central Apennines and it being a pioneer species, important
18 landscape changes are likely to occur at higher elevation forest-grassland ecotones. Studies in southern France (Debain
19 et al. 2005) showed how *P. nigra*, often considered an exogenous and invasive species, recruited more efficiently and
20 grew faster than local *P. sylvestris* as grazing declined and a grassland community took over. Extensive grazing can favour
21 the establishment of *Pinus* seedlings by reducing or eliminating competitors within the resident biota (Hulme 1996),
22 whereas intensive grazing can hamper regeneration due to animal trampling or burrow formation. Herbivores may also
23 indirectly influence *Pinus* recruitment by redistributing nutrients and affecting shrubs density, hence modifying the local
24 microclimate. Indeed, *Pinus* needles contain terpenes, which make the foliage distasteful and not particularly attractive
25 to livestock like cows, sheep and horses that are all selective feeders (Andrews et al. 2000). **However, some pines (e.g.**
26 ***Pinus cembra* and *Pinus peuce*) are grazed when the availability of grasses is limited.** The presence of shrubs, acting as
27 shelter against grazers, may facilitate the establishment of seedlings and influence the spatial distribution pattern
28 (Callaway and Walker 1997).
29

30 *P. nigra* encroachment at high altitude is still an active process because the conditions for an expected further
31 expansion of this species are assured by the increase in propagule supply. Seed sources have increased for two reasons:
32 i) neighbouring artificial plantations remain the most significant supplier given that the mean age of these stands is greater
33 than the *P. nigra* sexual maturity range (15-40 years) (Isajev et al. 2004); ii) several encroached trees (> 20%) have also
34

reached maturity and can act as seed suppliers modifying the future spatial pattern by new cohorts' formation. In this regard, future research should focus on the potential role of these new seed sources as well as long-distance dispersal (LDD) by wind from the neighbouring artificial plantations. Wind direction, tree crown size, cone production and tree height (starting point for seed dispersal) could affect the colonization process (Richardson 2000). Nonetheless, other requirements need to be met to guarantee further recolonization at the treeline ecotone, such as the presence of suitable microsites and microclimate conditions for seed germination and seedling growth. Apart from topography, these factors are dependent on broader vegetation dynamics within a context of climate (especially extreme events) and land use changes. This research is a further contribution to the understanding of *P. nigra* dynamic behaviour at high altitude in the central Apennines, but further investigation with a multi-scale approach combining the appropriate methods at regional, local, population and individual levels is needed to implement a suitable modelling and to include the ongoing process as a real treeline advancement.

References

- 1
2 Andrews J, Miller GR, Armstrong H (2000) The effects of mammalian herbivores on natural regeneration of upland,
3 native woodland. Scottish Natural Heritage, Edinburgh
4
- 5
6 Barbero M, Loisel R, Quezél P, Richardson DM, Romane F (1998) Pines of the Mediterranean Basin. In: Richardson DM
7
8 (ed) Ecology and biogeography of *Pinus*. Cambridge University Press, Cambridge, pp 153-170
9
- 10 Batllori E, Camarero JJ, Gutiérrez E (2010) Current regeneration patterns at the tree line in the Pyrenees indicate similar
11 recruitment processes irrespective of the past disturbance regime. *J Biogeogr* 37:1938-1950
12
- 13
14 Bisci C, Cellini M, Farabollini P, Pittori C (1989) Le gradient thermique dans les Marches meridionales (Italie Centrale).
15
16 Proc. Actes du Colloque de Pavia 2:27-33
17
- 18 Bolli JC., Rigling A, Bugmann H (2007). The influence of changes in climate and land-use on regeneration dynamics of
19 Norway spruce at the treeline in the Swiss Alps. *Silva Fennica* 41(1):55-70
20
- 21
22 Caballero R, Fernandez-Gonzalez F, Badia RP, Molle G, Roggero PP, Bagella S, D'Ottavio P, Papanastasis VP, Fotiadis
23
24 G, Sidiropoulou A, Ispikoudis I (2009). Grazing systems and biodiversity in Mediterranean areas: Spain, Italy and
25 Greece. *Pastos*, 39:9-154
26
- 27
28 Callaway RM, Walker LR (1997) Competition and facilitation: a synthetic approach to interactions in plant communities.
29
30 Ecology 78:1958-65
31
- 32
33 Callaway RM, Brooker RW, Choler P, Kikvidze Z, Lortie CJ, Michalet R, Cook BJ (2002) Positive interactions among
34 alpine plants increase with stress. *Nature* 417:844-848
35
- 36
37 Camarero JJ, Gutiérrez E (2004) Pace and pattern of recent treeline dynamics: response of ecotones to climatic variability
38 in the Spanish Pyrenees. *Climatic Change* 63:181-200
39
- 40
41 Camarero JJ, Gutiérrez E, Fortin MJ (2000) Spatial pattern of subalpine forest-alpine grassland ecotones in the Spanish
42 Central Pyrenees. *Forest Ecol Manag* 134:1-16
43
- 44
45 Carrer M, Urbinati C (2001) Spatial analysis of structural and tree-ring related parameters in a timberline forest in the
46 Italian Alps. *Journal of Vegetation Science* 12:643-652
47
- 48
49 Carrer M, Soraruf L, Lingua E (2013) Convergent space-time tree regeneration patterns along an elevation gradient at
50 high altitude in the Alps. *Forest Ecol Manag* 304:1-9
51
- 52
53 Catorci A, Scapin W, Tardella FM, Vitanzi A (2012) Seedling survival and dynamics of upper timberline in Central
54 Apennines. *Pol J Ecol* 60:79-94
55
- 56
57 Chiarugi A (1950) Le epoche glaciali dal punto di vista botanico. *Accademia Nazionale dei Lincei* 347:55-110
58
- 59
60 Compostella C, Caccianiga M (2016) A comparison between different treeline types shows contrasting responses to
61 climate fluctuations. *Plant Biosystems* doi: 10.1080/11263504.1179695
62
63
64
65

- 1
2 Coppinger R, Lorenz J, Glendinning J, Pinardi P (1983) Attentiveness of guarding dogs for reducing predation on
3 domestic sheep. *J Range Manage* 36:275-279
- 4 Cousens R, Dytham C, Law R (2008) Dispersal in plants: a population perspective. Oxford University Press, Oxford
- 5
6 Crawford RMM (2008) Plants at the margin. Ecological limits and climate change. Cambridge University Press,
7
8 Cambridge.
- 9
10 Dalen L, Hofgaard A (2005): Differential regional treeline dynamics in the Scandes Mountains. In: Arctic, Antarctic, and
11
12 Alpine Research 37:284–296
- 13
14 Debain S, Curt T, Lepart J (2005) Indirect effects of grazing on the establishment of *Pinus sylvestris* and *Pinus nigra*
15
16 seedlings in calcareous grasslands in relation to resource level. *Ecoscience* 12:192-201
- 17
18 De Bellis A, Pavan V, Levizzani V (2010) Climatologia e variabilità interannuale della neve sull'Appennino Emiliano
19
20 Romagnolo. Quaderno Tecnico ARPA-SIMC n. 19
- 21
22 Dibari C, Argenti G, Catolfi F, Moriondo M, Staglianò N, Bindi M (2015) Pastoral suitability driven by future climate
23
24 change along the Apennines. *Italian Journal of Agronomy* 10(659):109-116
- 25
26 Diggle PJ (2003) Statistical analysis of spatial point patterns. Edward Arnold, London
- 27
28 Farjon A (2013) *Pinus nigra* ssp. *nigra*. The IUCN Red List of Threatened Species 2013: e.T20453795A20453836.
29
30 <http://www.iucnredlist.org/details/20453795/0>. Accessed 12 February 2016
- 31
32 Garbarino M, Lingua E, Weisberg PJ, Bottero A, Meloni F, Motta R (2013) Land-use history and topographic gradients
33
34 as driving factors of subalpine *Larix decidua* forests. *Landscape Ecol* 28:805-817
- 35
36 Gehrig-Fasel J, Guisan A, Zimmermann NE (2007) Tree line shifts in the Swiss Alps: climate change or land
37
38 abandonment? *J Veg Sci* 18:571-582
- 39
40 Getis A, Ord JK (1992) The analysis of spatial association by use of distance statistics. *Geogr anal* 24:189-206
- 41
42 Harsch MA, Hulme, PE, McGlone MS, Duncan RP (2009) Are treelines advancing? A global meta-analysis of treeline
43
44 response to climate warming, *Ecol Lett* 12:1040–1049
- 45
46 Harsch MA, Bader MY (2011) Treeline form – a potential key to understanding treeline dynamics, *Global Ecol Biogeogr*
47
48 20:582–596
- 49
50 Harsch MA, McGlone MS, Wilmshurst JM (2014) Winter climate limits subantarctic low forest growth and
51
52 establishment. *PloS one* 9: e93241 doi:10.1371/journal.pone.0093241
- 53
54 Holtmeier KF (2005) Mountain Timberlines. Ecology, Patchiness and Dynamics. Dordrecht, The Netherlands: Kluwer.
- 55
56 Holtmeier F, Broll G (2005) Sensitivity and response of northern hemisphere altitudinal and polar treelines to
57
58 environmental change at landscape and local scales. *Global Ecol Biogeogr* 14:395-410
- 59
60 Hulme PE (1996) Herbivory, plant regeneration, and species coexistence. *J Ecol* 84:609–615
- 61
62
63
64
65

- 1 Huyghe C, De Vlieghe A, van Gils B, Peeters A (2014) Grasslands and herbivore production in Europe and effects of
2 common policies. Editions Quae. Versailles
- 3
4 Isajev V, Fady B, Semerci H, Andonovski V (2004) EUFORGEN Technical Guidelines for genetic conservation and use
5 for European black pine (*Pinus nigra*). International Plant Genetic Resources, Rome
- 6
7 Körner C (1998) A re-assessment of high elevation treeline positions and their explanation, *Oecologia* 115:445–459
- 8
9 Körner C (2012) Alpine Treelines. Functional Ecology of the Global High Elevation Tree Limits. Springer, Basel
- 10
11 Kullman L (1998): Tree-limits and montane forests in the Swedish Scandes: Sensitive biomonitors of climate change and
12 variability, *Ambio* 27:312–321
- 13
14
15 Lamedica S, Lingua E, Popa I, Motta R, Carrer M (2011) Spatial structure in four Norway spruce stands with different
16 management history in the Alps and Carpathians. *Silva Fenn* 45:865-873
- 17
18 Leonelli G, Pelfini M, Morra di Cella U (2009) Detecting climatic treelines in the Italian Alps: the influence of
19 geomorphological factors and human impacts. *Physical Geography* 30: 338-352
- 20
21
22 Lingua E, Cherubini P, Nola P, Motta R (2008) Spatial structure along an altitudinal gradient in the Italian central Alps
23 suggests competition and facilitation among coniferous species. *J Veg Sci* 19:425-436
- 24
25
26
27
28 Lowe JJ, Davite C, Moreno D, Maggi R (1994) Holocene pollen stratigraphy and human interference in the woodlands
29 of the northern Apennines, Italy. *The Holocene* 4:153-164
- 30
31
32 Marchesoni V (1957) Storia climatico-forestale dell'Appennino umbro-marchigiano *Ann Botanica* 25:1-39
- 33
34
35 Miehe, G, Miehe, S (2000): Comparative high mountain research on the treeline ecotone under human impact, *Erdkunde*
36 54:34–50
- 37
38
39 Motta R, Morales M, Nola P (2006) Human land-use, forest dynamics and tree growth at the treeline in the Western
40 Italian Alps. *Ann For Sci* 63:739-747
- 41
42
43 Pedrotti F (1969) Introduzione alla vegetazione dell'Appennino centrale. *Mitt. ostalp-din pflanzensoz.* 9:21-57
- 44
45
46 Pelorosso R, Leone A, Boccia L (2009) Land cover and land use change in the Italian central Apennines: a comparison
47 of assessment methods. *Appl Geogr* 29:35-48
- 48
49
50 Peroni P, Ferri F, Avena GC (2000) Temporal and spatial changes in a mountainous area of central Italy. *J Veg Sci*
51 11:505-514
- 52
53
54 Petritan IC, Marzano R, Petritan AM, Lingua E (2014) Overstory succession in a mixed *Quercus petraea–Fagus sylvatica*
55 old growth forest revealed through the spatial pattern of competition and mortality. *Forest Ecol Manag* 326:9-17
- 56
57
58 Pezzi G, Ferrari C, Corazza M (2008) The altitudinal limit of beech woods in the northern Apennines (Italy). Its spatial
59 pattern and some thermal inferences. *Folia Geobot* 43:447-459
- 60
61
62
63
64
65

- 1
2 Piermattei A, Renzaglia F, Urbinati C (2012) Recent expansion of *Pinus nigra* Arn. above the timberline in the central
3 Apennines, Italy. *Ann For Sci* 69:509-517
- 4 Piermattei A, Garbarino M, Urbinati C (2014) Structural attributes, tree-ring growth and climate sensitivity of *Pinus nigra*
5 Arn. at high altitude: common patterns of a possible treeline shift in the central Apennines (Italy). *Dendrochronologia*
6 32:210-219
- 7
8
9
10 Potthoff K (2009) Grazing history affects the tree-line ecotone: a case study from Hardanger, Western Norway. *Fennia*
11 187 (2):81–98
- 12
13
14 Richardson DM (2000) *Ecology and Biogeography of Pinus*. Cambridge University Press, Cambridge
- 15
16 Richardson DM, Rejmanek M (2004) Conifers as invasive aliens: a global survey and predictive framework. *Divers*
17 *Distrib* 10:321-331
- 18
19
20 Rivas-Martinez S, Rivas-Saenz S (2009) *Worldwide Bioclimatic Classification System, 1996-2015*. Phytosociological
21 Research Center, Spain. <http://www.globalbioclimatics.org>. Accessed 28 July 2015
- 22
23
24 Rössler O, Bräuning A, Löffler J (2008) Dynamics and driving forces of treeline fluctuation and regeneration in central
25 Norway during the past decades. *Erdkunde* 62:117–128
- 26
27
28 Rolando A, Caldoni R, Sanctis A, Laiolo P (2001) Vigilance and neighbour distance in foraging flocks of red-billed
29 choughs, *Pyrrhocorax pyrrhocorax*. *J Zool* 253:225-232
- 30
31
32 Santilocchi R, D'Ottavio P (2005) The evolution of cattle and sheep breeding systems in Central Italy over the past two
33 centuries. In: Georgoudis A, Rosati A, Mosconi C (eds) *Animal production and natural resources utilisation in the*
34 *Mediterranean mountain areas*. Wageningen Academic Pub, Wageningen, pp 15-18
- 35
36
37
38 Sawada M (1999) Rookcase: an Excel 97/2000 Visual Basic (VB) add-in for exploring global and local spatial
39 autocorrelation. *Bull Ecol Soc Am* 80:231-234
- 40
41
42 Slatyer RO, Noble IR (1992) Dynamics of montane treelines. In: di Castri F, Hansen AJ (eds) *Landscape Boundaries.*
43 *Consequences for Biotic Diversity and Ecological Flows*. Springer, New York, pp 346–359
- 44
45
46 Stanisci A, Lavieri D, Acosta A, Blasi C (2000) Structure and diversity trends at *Fagus* timberline in central Italy.
47 *Community Ecol* 1:133-138
- 48
49
50 Stoyan D, Stoyan H (1994) *Fractals, Random Shapes, and Point Fields: Methods of Geometrical Statistics*. Wiley,
51 Chichester
- 52
53
54 Theurillat JP, Guisan A (2001) Potential impacts of climate change on vegetation in the European Alps: A review. *Clim*
55 *Change* 50:77–109
- 56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Torta G (2004) Consequences of rural abandonment in a Northern Apennines landscape (Tuscany, Italy). In: Mazzoleni S, Di Pasquale G, Mulligan M, Di Martino P, Rego F (eds) Recent Dynamics of the Mediterranean Vegetation and Landscape. John Wiley & Sons, Chichester, pp 157-165
- Tranquillini W (1979) Physiological ecology of the alpine timberline. Ecological Series 31. Springer
- Troll C (1973) The upper timberlines in different climatic zones. Arctic Alpine Res 5:3-18
- Veblen TT (1992) Regeneration dynamics. In: Glenn-Lewin DC, Peet RK, Veblen TT (eds) Plant succession: Theory and prediction. Chapman & Hall, London, pp 152-187
- Vittoz P, Rulence B, Largey T, Freléchoux F (2008) Effects of climate and land-use change on the establishment and growth of cembra pine (*Pinus cembra* L.) over the altitudinal treeline ecotone in the Central Swiss Alps. Arct Antarct Alp Res 40:225-232
- Wiegand T, Moloney KA (2004) Rings, circles, and null-models for point pattern analysis in ecology. Oikos 104:209-229
- Wiegand T, Moloney KA (2014) Handbook of Spatial Point-Pattern Analysis in Ecology. Chapman and Hall/CRC, Boca Ranton
- Wiegand T, Camarero JJ, Rüger N, Gutiérrez E (2006) Abrupt population changes in treeline ecotones along smooth gradients. J Ecol 94:880-892

1
2
3
4 **Tables**

5 **Table 1.** Physiographic and climatic features of the surveyed *P. nigra* treeline ecotone sites in the central Apennines.

6 VET: Mt. Vettore, SFR: Mt. San Franco, SIR: Mt. Sirente, OCR: Mt. Ocre. 1961-1990 meteorological records are from
7 the nearest weather stations: Mt. Monaco, 987 m a.s.l. (VET); Campotosto, 1430 m a.s.l. (SFR); L'Aquila, 685 m a.s.l.
8 (SIR and OCR). The temperature and precipitation records of Campotosto and L'Aquila meteorological stations were
9 provided by the Hydrographic and Mareographic Service of the Abruzzo Region, Italy. The temperature and precipitation
10 records of Monte Monaco meteorological station were provided by the Centre of Ecology and Climatology - Geophysical
11 Experimental Observatory at Macerata, Italy. **The mean temperatures at the forestline elevation were extrapolated from
12 the meteorological stations by using a temperature lapse rate of 0.47°/100 m (Bisci et al. 1989).**

Site	Treeline ecotone surface area (ha)	Surveyed surface area (ha)	Slope aspect	Mean slope angle (°)	Forestline elevation (m a.s.l.)	Mountain top elevation (m a.s.l.)	Latitude (N°)	Longitude (E°)	Annual Precip. (mm)	Mean Temp. at Forestline elevation (°C)
VET	181.3	87.6	S - SE	36°	1600	2478	42°81'	13°26'	1209.8	8.2
SFR	53.5	35.6	W - SW	32°	1600	2132	42°45'	13°38'	994.0	7.0
SIR	87.0	37.1	W - SW	19°	1650	2349	42°15'	13°60'	732.4	7.2
OCR	14.0	6.0	N - NE	33°	1350	2204	42°15'	13°27'	732.4	8.6

Table 2. Anthropogenic disturbance histories at VET, SFR, SIR and OCR sites.

Site	Grazing History	Grazing period	Forestline history
VET	Sheep decreased by 50% during 1970-2000	May to mid-October	<i>P. nigra</i> plantation 63 yrs old
SFR	Sheep decreased by 40% during 1980-2000	May to mid-October	<i>P. nigra</i> plantation 48 yrs old
SIR	Grazing of cows and horses still active	May to mid-October	<i>P. nigra</i> plantation 45 yrs old
OCR	Dramatic grazing decline during 1954-1995	May to mid-October	<i>P. nigra</i> plantation 51 yrs old

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 3. Number and main structural variables of sampled *Pinus nigra* encroached trees at the four study sites. Diameter (at root collar), height and age values are means \pm standard deviation (SD).

Site	Living/Dead trees (N)	Density (n/ha)	Diameter (cm)	Height (cm)	Age (years)
VET	177/4	2	9 \pm 7.3	150 \pm 138	15 \pm 7
SFR	241/13	7	7 \pm 6.9	119 \pm 117	16 \pm 7
SIR	194/16	6	13 \pm 8.7	211 \pm 157	19 \pm 8
OCR	182/18	3	8 \pm 5.4	133 \pm 97	17 \pm 6

Figure captions

Fig. 1 Oblique ground-based photographs of four *P. nigra* treelines of the central Apennines: a) OCR, b) SFR, c) SIR, and d) VET.

Fig. 2 Relative frequency (a) and maximum age (b) of regenerating *P. nigra* along the elevation gradient, divided in 20 m elevation classes at four treeline ecotones (VET: Mt. Vettore, SFR: Mt. San Franco, SIR: Mt. Sirente, OCR: Mt. Ocre).

Fig. 3 Scatterplot of tree height and age divided by elevation classes as indicated by grey scale colours and regression lines. All regressions were significant ($p < 0.001$), except for VET and SFR 1900-2000, and R^2 ranged from 0.28 (VET 1800-1900) and 0.85 (SIR > 2000).

Fig. 4 Univariate spatial pattern analysis of 537 *Pinus* trees in four central Apennine treelines adopting combined replicates. Bold line indicates the pair-correlation function $g(r)$ and the dotted line the expected value under the null model (heterogeneous Poisson); shaded areas encompass non-significant, i.e. random, distribution and represent points within the 2.5th and 97.5th percentile $g(r)$ values of the 999 Monte Carlo permutations. The inset graphs show the results of each site separately.

Fig. 5 Bivariate spatial pattern analysis of *Pinus* trees, considering two age classes (saplings < 10 years and young trees > 15 years), using the pair-correlation function $g_{12}(r)$. Black lines indicate the pair-correlation functions $g_{12}(r)$, the dotted line represents the expected value under the null model (antecedent condition and heterogeneous Poisson), and shaded areas encompass non-significant, i.e. random, distribution and represent points within the 2.5th and 97.5th percentile $g(r)$ values of the 999 simulations Monte Carlo permutations. The inset graphs show the results of each site separately.

Fig. 6 Interpolated (Natural Neighbour) and z-transformed local $G_i^*(d)$ values computed for tree age for 15 m and 50 m distance intervals. Red and blue areas are hot spots (i.e. groups of trees with age higher than the mean within the treeline) and cold spots (i.e. groups of trees with age lower than the mean within the treeline) respectively. Colour scale is proportional to the value of the local $G_i^*(d)$ index. Bold contour lines indicate significant ($p < 0.005$) clusters.

Online Resource 1 Univariate spatial pattern analysis of *Pinus* trees, considering two age classes (A- saplings < 10 years; B- young trees > 15 years), using the pair-correlation function $g(r)$. Black lines indicate the pair-correlation functions $g(r)$ and the dotted line represents the expected value under the null model (heterogeneous Poisson); shaded areas encompass non-significant, i.e. random, distribution and represent points within the 2.5th and 97.5th percentile $g(r)$ values of the 999 simulations Monte Carlo permutations. The inset graphs show the results of each site separately.

Online Resource 2 Map of spatial distribution of *Pinus* individuals at four treelines of the central Apennines: VET, SIR, SFR and OCR. Size of points is proportional to the age class of trees.















