

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Gametic embryogenesis through isolated microspore culture in *Corylus avellana* L

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1623005> since 2017-01-25T10:50:36Z

*Published version:*

DOI:10.1007/s11240-015-0921-1

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

Gniech Karasawa, M.M., Chiancone, B., Gianguzzi, V., Abdelgalel A.M., Botta R., Sartor C.,  
Germanà M.A. 2016. Gametic embryogenesis through isolated microspore culture in *Corylus*  
*avellana* L. *Plant Cell Tiss Organ Cult* (2016) 124: 635. doi:10.1007/s11240-015-0921-1

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

<http://link.springer.com/article/10.1007/s11240-015-0921-1>

**Title page**

Marines Marli Gniech Karasawa<sup>1</sup>, Benedetta Chiancone<sup>2</sup>, Valeria Gianguzzi<sup>1</sup>, Ahmed Mohamed Abdelgalel<sup>2</sup>, Roberto Botta<sup>3</sup>, Chiara Sartor<sup>3</sup>, Maria Antonietta Germanà<sup>1</sup>

**Gametic embryogenesis through isolated microspore culture in *Corylus avellana* L.**

<sup>1</sup>Dipartimento di Scienze Agrarie e Forestali. Università degli Studi di Palermo. Viale Delle Scienze, 11. 90128 Palermo –Italy

<sup>2</sup>Dipartimento di Scienze degli Alimenti. Università degli Studi di Parma, Parma, Italy

<sup>3</sup>Dipartimento di Colture Arboree, Università degli Studi di Torino. L.go P. Braccini 2, 10095, Grugliasco (TO) mariaantonietta.germana@unipa.it

**Abstract**

Haploid technology is a valuable tool to obtain homozygosity particularly in woody plants, in order to support plant breeding. Hazelnut, the fourth worldwide nut tree, is a monoecious, anemophilous species and it is characterized by a sporophytic incompatibility system (Germain and Dimoulas, 1978), not allowing to obtain homozygous plants through conventional methods, involving several self-pollination. In this study, gametic embryogenesis was applied to hazelnut and, particularly, isolated microspore culture was performed on four different cultivars. Two culture media were tested and four temperature stress treatments were applied to the isolated microspores, cultivated at the vacuolated stage of development. For the first time in our knowledge, early embryos were recovered in hazelnut *via* isolated microspore culture in all cultivars. The assessment of the zygosity condition of embryos carried out using SSR markers showed that analysed embryos were homozygous indicating that they developed from haploid microspores. The response to the culture treatments, was, however, genotype-dependent, as previously reported in experiments regarding male gamete embryogenesis in other fruit crops.

**Keywords:** Haploid, hazelnut, gametic embryogenesis, temperature stress.

## Introduction

Hazelnut (*Corylus avellana* L.), belonging to the Betulaceae family of the order Fagales and native to European regions, Turkey and the Caucasus mountains (Kafkas and Dogan, 2009), is the sixth worldwide tree nut, with a production of more about 858,000 tons (FAOSTAT, 2013), behind cashew, Persian walnut, chestnut, almond and pistachio. All cultivated forms of hazelnut are diploid with a monoploid number of chromosomes  $n=x=11$ . It is a monoecious, dichogamous, self-incompatible and wind pollinated plant (Rovira et al. 1993). The incompatibility system is sporophytically determined and controlled by a single locus with multiple alleles (Martins 2010). Incompatibility is a main factor limiting genetic improvement, and it prevents to reach homozygosity by conventional methods, involving several self-pollination cycles.

Gametic embryogenesis is recognized as an important tool for plant breeding, making possible to develop, in only one step, completely homozygous lines, from heterozygous parents (Bueno et al. 2006; Seguí-Simaro and Nuez 2008; Islam and Tuteja 2012; Datta 2005; Solis et al. 2008; Germanà 2011a). This opportunity is particularly useful for breeding in woody species, in which it is not possible to obtain homozygosity through conventional methods (Germanà 2011b), because of the long juvenile phase, the high degree of heterozygosity, the large size and, often, of the self-incompatibility (Germanà 2006; 2009).

Haploid (Hs) and doubled haploid (DHs) plants, are very useful to fix parental lines to produce F1 hybrids, for rapid introgression of new characteristics through backcrossing, to develop population molecular maps, to fix characteristics through transformation and mutagenesis (Datta 2005; Szarejko and Foster 2006; Soriano et al. 2013) and to do reverse breeding (Dirks et al. 2009; Ferrie and Möllers 2011). In addition, haploids and double-haploids are important for genomic studies and genome sequencing (Aleza et al. 2009; Dirks et al. 2009; Ferrie and Möllers 2011; Foster et al. 2007; Germanà et al. 2013; Talón and Gmitter 2008), for physical mapping (Leeuwen et al. 2003; Kunzel et al. 2000; Wang et al. 2001), genetic mapping (Houssain et al. 2007; Chu et al. 2008; Zhang et al. 2008), and for the integration between physical and genetic mapping (Kunzel et al. 2000; Wang et al. 2001).

Gametic embryogenesis represents a unique system where, thanks to the plant totipotency, a single immature gamete switches from the gametophytic pathway toward the sporophyte formation (Prem et al 2012). Gametic embryogenesis, that can be achieved from female (gynogenesis) or male (microspore embryogenesis) gametes, rarely occurs in nature (Silva 2012), and it is usually induced subjecting immature gametes to some kinds of stress. Particularly, temperature stresses, hot and cold, are the most effective and commonly used ways to promote microspore embryogenesis in several species (Ferrie and Caswell 2010; Germanà 2011 a; Shariatpanahi et al. 2006; Silva 2012). Specifically, the involvement of heat shock proteins (HSPs), when temperature treatments are applied in the microspore embryogenesis, has been described, but the molecular mechanism of the induction and the relationship between different stresses remain unclear. However, it is supposed that the increased levels of HSPs may be associated with the embryogenic potential acquisition (Karami and Saidi 2010; Silva 2012).

In many crops, anther culture is often the method of choice for H and DH production because of its simplicity (Sopory and Munshi 1996). Nevertheless, the technique of isolated microspore culture, performed by removing somatic anther tissue and requiring better equipment and more skills compared to the first technique, provides a better system for investigating cellular, physiological, biochemical and molecular processes involved in pollen embryogenesis (Nitsch 1977; Reinert and Bajaj 1977; Germanà, 2011b).

In this study, the technique of isolated microspore culture was applied to five hazelnut genotypes, testing two basal media and four temperature stresses, with the aim to obtain micropore-derived embryo regeneration.

## Materials and Methods

### *Plant material and pollen developmental stage*

Immature catkins (Fig. 1a) were harvested in February from trees of the following cultivars of *Corylus avellana* L.: ‘Tonda Gentile Romana’ (2 clones: **TR** and **GR**), ‘Imperatrice Eugenia’ (**IE**), ‘Minnulara’ (**M**) and ‘Carrello’ (**C**). Catkins with anthers (Fig. 1a) mostly containing microspores at the vacuolated/uninucleated stage (Fig. 1b) were selected, assuming that, like in many other fruit crops (Germanà 2011 a; 2011b), this stage of development is the most responsive for gametic embryogenesis also for hazelnut. For this reason, a preliminary investigation was carried out to select the appropriate catkin size staining anthers with few drops of 4',6-diamidino-2-phenylindole (DAPI) solution (1 mg/mL) and observing them under fluorescence microscope (Zeiss, Axiophot, Germany). Catkins with anthers containing microspores at vacuolated/uninucleate stage were selected and, as common pre-treatment, were placed in darkness at 4 °C for two weeks.

### *Catkins sterilization and microspore isolation and culture*

After two weeks of 4 °C pretreatment, 5 catkins for each genotype were surface sterilized by immersion for 3 min in 70% (v/v) ethyl alcohol, followed by immersion for 20 min in 25% (v/v) sodium hypochlorite solution (about 1.5% active chlorine in water) with few drops of Tween® 20, and then rinsed three times with sterile distilled water. After that, the material was maintained in immersion in gentamicin antibiotic solution (0.2%) for 30 minutes. The antibiotic solution was removed and the microspore isolation was performed following the procedure reported by Kumlehn et al., (2006) with limited modifications. Isolated microspores were cultured at the concentration of 100,000 microspores per mL and a final volume of 1.0 mL per each Petri dish (3001-type Petri dishes, BD Biosciences) was placed.

Petri dishes, sealed with Parafilm®, were placed at 26±1 °C, in the dark for the first 30 days and later under cool white fluorescent lamp (Philips TLM 30W/84, France) with a photosynthetic photon flux density of 35 µmol m<sup>-1</sup> s<sup>-1</sup> and a photoperiod of 16 light hours.

### *Experimental design*

In order to evaluate the influence of medium composition on the embryogenic induction of the five genotypes, isolated microspores were cultured on two different media: **P** (Germanà et al. 1996) and **N6** (Germanà and Chiancone 2003). Moreover, because it is well known that usually stress enhances microspore response, isolated microspores cultured in both media, were subjected to the following thermal stress, just after isolation:

- 1) 35 °C for 30 minutes (**H**),
- 2) 40 °C for 60 minutes (**H+**),
- 3) -20 °C for 30 minutes (**F**),
- 4) -20 °C for 60 minutes (**F+**).

Height Petri dishes (repetitions) were prepared per each combination (genotype, medium, thermal stress).

### ***Data collection and statistical analysis***

Isolated microspores in culture were weekly observed by a stereo-microscope (Leica MZ 125) and by an inverted microscope (Zeiss West Germany). In addition, examinations were performed by a fluorescence microscope (Zeiss, Axiophot, Germany), after staining with 4',6-diamidino-2-phenylindole (DAPI). Because changes in the microspore development distribution were observed for a long time, it was decided to report the final monitoring of the different features, registered after 20 months of culture. At this time, per each genotype and per each combination medium/thermal stress, 450 microspores/structures (three repetitions with 150 elements each) were counted, in order to individuate the different structural features: uninucleated microspores, rarely binucleated with two different nucleus sizes beginning their normal gametophytic pathway, binucleated with two equal-size vegetative-type nuclei just starting their sporophytic pathway, trinucleated, and also multinucleated microspores. Moreover, the number of embryos and calli produced per each Petri dish were registered. These values were used to calculate means and to perform the statistical analysis. Differences among genotypes were tested by one-way analysis of variance (ANOVA) with Conover-Inman's test ( $p \leq 0.05$ ). To individuate the influence of the factors "Medium" and "Thermal treatment", means were analysed by two-way ANOVA, and mean separation was performed using Tukey's multiple comparison tests at ( $p \leq 0.05$ ).

### ***Detection of homozygosity***

Simple Sequence Repeat (SSR) markers were used to assess the genetic condition of embryos obtained from pollen microspore culture and to determine their origin (gametic or somatic).

#### **DNA extraction**

Embryos were collected from the culture medium by an insulin syringe, frozen in liquid nitrogen and ground using steel beads in a TissueLyser (QIAGEN®, Germany). DNA extraction for genotyping was performed as described in Doyle and Doyle (1987) with minor modifications.

#### **PCR amplification and SSR allele sizing**

Five fully characterized SSR loci were used for assessing the origin of the embryos: CaT-B107, CaT-B503, CaT-B504, CaT-B505, CaT-B507 (Boccacci et al., 2005). PCR was performed, as described by Boccacci et al. (2008), in two steps in a total volume of 10 µl containing 20 ng of DNA, 0.25 U of KAPA Taq DNA polymerase (KAPABIOSYSTEMS, Wilmington Massachusetts, USA) 1 µl of 10X PCR buffer, 200 µM nucleotide mix and 0.5 mM of each primer. PCR conditions were as follows: an initial denaturation step at 95°C for 3 min followed by 32 cycles of denaturation (30 s at 95°C), annealing (45 s at 52°C), and extension (90 s at 72°C). The final elongation step was at 72°C for 30 min. Two µl of the amplification product were used as template for a second PCR, carried out for 28 cycles with the same conditions of the first one.

PCR products were analyzed on a 3130 Genetic Analyzer (Applied Biosystems, Foster City, California, USA). Data were processed using GeneMapper Software (ver. 4.0; Applied Biosystems) and alleles were defined by their size in base pairs, by comparison with the size standard (GeneScan-500 LIZ, Applied Biosystems).

Data obtained were compared with the hazelnut database of DISAFA.

### **Results and Discussion**

The microspore culture monitoring carried out by fluorescence microscope after 20 months of culture and after DAPI staining, showed uninucleated (Fig. 1b), binucleated presenting nuclei with asymmetrical division (Fig. 2a), binucleated revealing nuclei with symmetrical division (Fig. 2b), trinucleated (Fig. 2c), and multinucleated stage (Fig. 2d); the performances of the different genotypes are reported in table 1-6. Actually, the presence of binucleated microspores with asymmetrical division of the nucleus, indicating of the gametophytic pathway, was very sporadic: never more than 3% and in three genotypes, (C, IE and M), less than 1% of the total (Table 1).

The statistical analysis carried out to study the *in vitro* isolated microspore development in the five hazelnut genotypes tested, showed different responses among the genotype (Table 1). For this reason, the influence of the two parameters “Medium” and “Thermal treatment” on hazelnut microspore development was carried out separately per each genotype. Particularly, regarding the “Medium” influence on the multinucleated microspores production, the P medium was the most valuable in the genotypes C, M and TR (Tables 2, 5 and 6). In the other genotypes, differences between the two media were not statistically significant. Two way ANOVA evidenced also that in the cultivars C and M, the formation of multinucleated microspores was mainly influenced by the factor “Medium”, while different results were recorded for the genotypes GR, IE and TR, where a significant interaction was observed between the two factors. IE, GR and C, were the genotypes with the highest percentages of multinucleated microspores.

For GR, the factor “Treatment” was the main source of variability in the multinucleated microspore induction (Table 3). Particularly, statistically significant differences were observed between the treatment F+ and the treatment H, while the other treatments induced intermediate responses. The analysis of interaction evidenced a statistically significant difference in the thermal treatment H, where N6 induced better response than P (32.0% vs 19.8%) (data not shown).

Diverse response was observed in the cultivar IE concerning the same parameter, in which none of the factors had a predominant influence (Table 4), but differences were observed in the response to thermal treatments between the two media: P better than N6 in H+ (+34.7% vs 24.7%) and in F (40.0 % vs 23.3%), whereas N6 better than P in F+ (30.0% vs 22.7%) (data not shown).

On the contrary, for TR, both factors (“Medium” and “Treatment”) had a significant influence on the multinucleated microspore induction. Analyzing separately the factors, P medium induced a statistically significant higher percentage of multinucleated microspores than N6; while, regarding the thermal treatments, two groups were individuated: Co and H induced statistically higher response than H+, F and F+ (Table 6). Furthermore, regarding the interaction, main differences were observed in the Control where the P medium induced a statistically higher response than N6 (37.3% vs 22.0%)

After three-four months of culture, stereo- microscope observations revealed new structures: light brown calli (Fig. 3) that increased in quantity and volume and, also, the formation of globular embryos was detected (Fig. 4 a). Moreover, different shapes of embryos were observed: elongated (Fig. 4 b) and also often embryos with a suspensor-like structure (Fig. 4c). This kind of structure was observed also in Citrus in the embryos obtained through isolated microspore culture, but not by anther culture.

Previously, in fruit tree crops, Hofer et al (1999) reported embryo regeneration, through isolated microspore culture, in apple and, recently (Chiancone et al., 2015), the production of gamete-derived embryos obtained by in the same way in Citrus was reported.

Because of the data distribution, it was not possible to apply the two-way ANOVA to the average number of calli and embryos registered per Petri dish, but for the novelty and relevance of these results, it is interesting to describe the performance of the different cultivars. The statistical analysis of the average calli number per Petri dish confirmed the morphogenic potential of TR, that, together with GR, produced the highest number of calli (Fig. 5a). Actually, the highest average callus number per Petri dish was recorded in the genotype 'Tonda Gentile Romana' TR, in the treatment H<sup>+</sup> in combination with the N6 medium (13.7) (Fig. 5f). As reported for the microspore development, also for calli production, the response to the different media and to the thermal treatments was genotype-dependent. The cold thermal treatments positively influenced callus production: specifically, F<sup>+</sup> treatment induced the highest average number of calli per Petri dish in C, IE (in combination with P medium), GR (in combination with both P and N6 media); while, the hot thermal treatments, and particularly, the H<sup>+</sup>, seemed to produce a higher callus number in M (in combination with N6) and in TR (in combination with P and N6) (data not shown).

The embryo achieving does not appear to be influenced by the treatments in the same way than calli. Really, with the exception of M, in which the combination of H<sup>+</sup> thermal treatment with N6 medium, was proven to be suitable both for callus and embryo achieving. For the other cultivars, the results were different: the highest average embryo number were recorded in the control, in combination with P medium for C and TR, with N6 medium for GR, and in the H<sup>+</sup> treatment in combination with N6 medium for IE (Fig. 5b-f).

Generally, regarding the effect of the medium composition, P medium induced a little higher percentage of multinucleated microspores and also a higher number of embryos. Not always at the genotype with the very high number of multinucleated microspores corresponded the higher microspore-derived embryo number (for example, IE).

Regarding the stress, many research highlighted its importance in inducing microspore embryogenesis. However, in this study the higher number of embryos was obtained in both P and N6 control (without additional stresses other than 4 °C before isolation), except for IE.

### ***Detection of homozygosity***

Genetic analysis showed that the hazelnut embryos had a single allele at each of the five SSR loci analysed (Figs. 6 and 7). The allele found at each locus matched either one of the 2 alleles of the parent genotype (in the case of the figures 'Minnulara' was the cultivar of origin of the microspores). Being the embryos very small, ploidy level could not be determined; for this reason we can postulate that the embryos are very likely haploids but there is no evidence they may not be double haploids. For sure the embryos analysed originated from a haploid microspore.

The production of homozygous plants in fruit trees of high commercial value like hazelnut is highly desired to help increase the breeding efficiency and reduce the time to obtain new varieties. But, this is not possible in woody plants, like hazelnut, characterized by long juvenile cycle, high degree of heterozygosity, large size and with incompatible systems. The only way to obtain isogenic lines is through haploid technology. Actually, microspore embryogenesis is a promising tool in plant breeding since it permits to obtain full homozygous plants in just one step using heterozygous parents (Germanà 2011 a, b). This technology can be of great interest



to increase the research and plant breeding of *Corylus avellana*, a species not very often subject of improvement (Dunstan and Thorpe 1986).

## Conclusions

In our knowledge, this is the first time that haploid technology was applied to hazelnut and that homozygous early embryos were regenerated through isolated microspore culture. The response to the culture media and treatments, was, however, highly genotype-dependent, as previously reported in experiments regarding tissue culture in hazelnut as well as microspore embryogenesis in other fruit crops. Actually, each genotype presented different behaviour, even if, all of them produced calli and, more important, microspores-derived early embryos. Really, even if it is not possible to individuate the absolute best medium or thermal treatment stress, this result represents a major step forward in the knowledge on gametic embryogenesis in fruit crops. Further studies are, however, necessary to better understand the process of gametic embryogenesis in this species and to optimize the rate of microspore-derived embryos and, in addition, to obtain their germination into plantlets.

## Acknowledgments

Thanks are due to Prof. Basilio Baratta for providing catkins of some halzenut cultivars and to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the scholarships to Marines Marli Gniech Karasawa. Authors wish to thank Prof. Louise Ferguson for English editing of the manuscript.

## References

- Aleza P, Juárez J, Hernández M, Pina JA, Ollitrault P, Navarro L (2009) Recovery and characterization of a *Citrus clementina* Hort. ex Tan. ‘Clemenules’ haploid plant selected to establish the reference whole Citrus genome sequence. BMC Plant Biology 9 (110). doi 10.1189/1471-2229-9-110. <http://www.biomedcentral.com/1471-2229/110>
- Arzate-Fernandez A-M, Nakazaki T, Yamagata H, Tanisaka T (1997) Production of doubled-haploid plants from *Lilium longiflorum* Thunb. anther culture. Plant Science 123:179-187
- Ashok Kumar HG, Murthy HN, Paek KY (2003) Embryogenesis and plant regeneration from anther cultures of *Cucumis sativus* L. Scientia Horticulturae 98:213-222
- Ayed OS, Buyser JD, Picard E, Trifa Y, Amara HS (2010) Effect of pre-treatment on isolated microspores culture hability in durum wheat (*Triticum turgidum* subsp. *durum* Desf.). Journal of Plant Breeding and Crop Science 2 (2):030-038
- Bàrany I, Testillano PS, Mitykó J, Risueño MC (2001) The switch of the microspore developmental program in *Capsicum* involves HSP70 expression and leads to the production of haploid plants. International Journal of Developmental Biology 45 (S1):S39-S40
- Bhowmik P, Dirpaul J, Ferrie AM (2011) A high throughput *Brassica napus* microspore culture system: influence of percoll gradient separation and bud selection on embryogenesis. Plant Cell, Tissue and Organ Culture 106:95-107

286 Binarova P, Hause G, Cenklová V, Cordewener JH, Campagne ML (1997) A short severe heat shock is required  
 287 to induce embryogenesis in late bicellular pollen of *Brassica napus* L. Sexual Plant Reproduction 10:200-208  
 288 Boccacci P., Akkak A., Bassil N.V., Mehlenbacher S.A., Botta R. 2005. Characterization and evaluation of  
 289 microsatellite loci in European hazelnut (*Corylus avellana* L.) and their transferability to other *Corylus* species.  
 290 Molecular Ecology Notes 5: 934-937.  
 291 Boccacci P., Botta R., Rovira M. 2008. Genetic Diversity of Hazelnut (*Corylus avellana* L.) Germplasm in  
 292 Northeastern Spain Hort Science 43(3):667–672.  
 293 Bueno MA, Agundez MD, Gomez A, Carrascosa MJ, Manzanera JA (2000) Haploid origin of cork-oak anther  
 294 embryos detected by enzyme and rapid gene markers. International Journal Plant Sciences 161:363-367  
 295 Bueno MA, Gómez A, Sepúlveda F, Seguí JM, Testillano PS, Manzanera JA, Risueño MC (2003) Microspore-  
 296 derived embryos from *Quercus suber* anthers mimic zygotic embryos and maintain haploidy in long-term anther  
 297 culture. Journal of Plant Physiology 160:953-960  
 298 Bueno MA, Pintos B, Martin A (2006) Induction of embryogenesis via isolated microspore culture in *Olea*  
 299 *europaea* L. Olivebioteq 2006, Novembre, Mazara del Vallo, Marsala, Italy  
 300 Burnett L, Yarrow S, Huang B (1992) Embryogenesis and regeneration from isolated microspores of *Brassica*  
 301 *rapa* L. ssp. *oleifera* Plant Cell Reports 11:215-218  
 302 Cappadocia M, Cheng DSK, Ludlum-Simonette R (1984) Plant regeneration from *in vitro* culture of anthers of  
 303 *Solanum chacoense* Bitt. and inter specific diploid hybrids *S. tuberosum* L and *S. chacoense*. Bitt. Theor Appl  
 304 Genet 69:139–143  
 305 Chen Z (1985) A study on induction of plants from *Citrus* pollen. Fruit Varieties Journal 39:44-50  
 306 Chiancone B, Gniech Karasawa MM, Gianguzzi V, Mohamed A A, Bárány I, Testillano P, Torello Marinoni D,  
 307 Botta R, Germanà MA (2015) Early embryo achievement through isolated microspore culture in *Citrus*  
 308 *clementina* Hort. ex Tan., cvs. ‘Monreal Rosso’ and ‘Nules’. Front. Plant Sci., Vol. 6 11 June 2015. N 00413 |  
 309 <http://dx.doi.org/10.3389/fpls.2015.00413> ISSN=1664-462X  
 310 Chu CG, Xu SS, Friesen TL, Faris JD (2008) Whole genome mapping in wheat doubled haploid population  
 311 using SSRs and TRAPs and the identification of QTL for agronomic traits. Molecular Breeding 22:251-266  
 312 Custers JBM, Cordewener JHG, Nölten Y, Dons HJM, Van Lookeren Campagne Mm (1994) Temperature  
 313 controls both gametophytic and sporophytic development in microspore cultures of *Brassica napus*. Plant Cell  
 314 Reports 13:267–271  
 315 Datta SK (2005) Androgenic haploids: Factors controlling development and its application in crop  
 316 improvement. Current Science 89 (11):1870-1878  
 317 Dias JS (2003) Protocol for broccoli microspore culture. In: Maluszynski M, Kasha KJ, Forster BP, Szarejko I  
 318 (eds) Doubled Haploid Production in Crop Plants: a Manual. Kluwer Academic Publishers, Dordrecht, The  
 319 Netherlands, pp 195–204.  
 320 Dirks R, Van Dun K, De Snoo, Van Den Berg M, Lelivelt CL, Voermans W, Woudenberg L, De Wit JP,  
 321 Reinink K, Schut JW (2009) Reverse breeding: a novel breeding approach based on engineered meiosis. Plant  
 322 Biotechnology Journal 7:837–845  
 323 Doyle J. J., Doyle J. L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue.  
 324 Phytochemical Bulletin 19:11-15

325 Duijs J, Voorrips R, Visser S, Custers J (1992) Microspore culture is successful in most crop types of *Brassica*  
 326 *oleracea* L. Euphytica 60:45-55  
 327 Dunstan DI, Thorpe TA (1986) Regeneration in Forest Trees. Vasil IK (ed) Cell Culture and somatic cell  
 328 genetics of plants v. 3: Plant regeneration and genetic variability. Chapter 11.  
 329 Dunwell JM, Thurling N (1985) Role of sucrose in microspore embryo production in *Brassica napus* spp.  
 330 *oleifera*. Journal of Experimental Botany 36:1478-1491  
 331 FAOSTAT (2015) <http://faostat3.fao.org/>  
 332 Ferrie AMR, Caswell KL (2010) Isolated microspore culture techniques and recent progress for haploid and  
 333 doubled haploid plant production. Plant Cell, Tissue and Organ Culture 104 (3):301-309  
 334 Ferrie AM, Möllers C (2011) Haploids and doubled haploids in *Brassica* spp. for genetic and genomic research.  
 335 Plant Cell Tissue and Organ Culture 104:375–386  
 336 Foster BP, Herberle-Bors E, Kasha KJ, Touraev A (2007) The resurgence of haploids in higher plants. Trends in  
 337 Plant Science 12 (8):368-375  
 338 Froelicher Y, Ollitrault P (2000) Effects of the hormonal balance on *Clausena excavate* androgenesis. Acta  
 339 Horticulturae 535:139-146  
 340 Geraci G, Starrantino A (1990) Attempts to regenerate haploid plants from “in vitro” cultures of Citrus anthers.  
 341 Acta Horticulturae 280:315-320  
 342 Germain E, Dimoulas I (1978) La physiologie de la reproduction chez la noisetier *Corylus avellana* L.:  
 343 Connaissances actuelles, 104ème Congrès National des Sociétés Savantes Bordeaux, Sciences, fasc.II 435-446.  
 344 Germanà MA (2006) Doubled haploid production in fruit tree crops. Plant Cell, Tissue and Organ Culture  
 345 86:131-146.  
 346 Germanà MA (2011a) Gametic embryogenesis and haploid technology as valuable support to plant breeding.  
 347 Plant Cell Reports 30:839-857  
 348 Germanà MA (2011b) Anther culture for haploid and doubled haploid production. Plant Cell, Tissue and Organ  
 349 Culture 104:283-300  
 350 Germanà MA, Chiancone B (2003) Improvement of *Citrus clementina* Hort. Ex Tan. Microspore-derived  
 351 embryoid induction and regeneration. Plant Cell Report 22:181-187  
 352 Germana' MA, Aleza P, Carrera E, Chen C, Chiancone B, Costantino G, Dambier D, Deng X, Federici TC,  
 353 Froelicher Y, Guo W, Ibáñez V, Juárez J, Kwok K, Luro F, Machado Am, Naranjo AM, Navarro L, Ollitrault P,  
 354 Ríos G, Roose Lm, Talon M, Xu Q, Gmitter GF (2013) Cytological and molecular characterization of three  
 355 gametoclones of *Citrus clementina*. BMC Plant Biology, vol. 13, ISSN: 1471-2229. doi: 10.1186/1471-2229-13-  
 356 129. <http://www.biomedcentral.com/1471-2229/13/129>  
 357 Giménez-Abián M, Rozalén A, Carballo J, Botella L, Pincheira J, López-Sáez J, de la Torre C (2004) HSP90  
 358 and checkpoint-dependent lengthening of the G2 phase observed in plant cells under hypoxia and cold.  
 359 Protoplasma 223:191-196  
 360 Hoekstra S, van Zijderveld MH, Louwerse JD, Heidekamp F, van der Mark F (1992) Anther and microspore  
 361 culture of *Hordeum vulgare* L. cv Igri. Plant Science 86:89-96  
 362 Höfer M (2003) *In vitro* androgenesis in apple. In: Malusynski M. et al. (eds) Doubled Haploid Production in  
 363 Crop Plants, Kluwer academic publishers. IAEA, pp 287-292

364 Höfer M (2004) *In vitro* androgenesis in apple-improvement of the induction phase. Plant Cell Reports 22(6):  
 365 365-370  
 366 Höfer M, Touraev A, Heberle-Bors E (1999) Induction of embryogenesis from isolated apple microspores. Plant  
 367 Cell Reports 18:1012-1017  
 368 Hou L, Ullrich SE, Kleinhofs A, Stiff CM (1993) Improvement of anther culture methods for doubled haploid  
 369 production in barley breeding. Plant Cell Reports 12:334-338  
 370 Houssain T, Tausend P, Graham G, Ho J (2007) Registration of IBM2 SYN10 doubled haploid mapping  
 371 population of maize. Journal of Plant Registrations 1:81  
 372 Kafkas S, Dogan Y (2009) Genetic characterization of hazelnut (*Corylus avellana* L.) cultivars from Turkey  
 373 Using Molecular Markers. Hortscience 44(6):1557–1561  
 374 Kamada H, Tachikawa Y, Harada H (1994) Heat stress induction of carrot somatic embryogenesis. Plant tissue  
 375 culture letters 11 (3):229-232  
 376 Kumlehn J (2009) Embryogenic pollen culture: a promising target for genetic transformation. In: Touraev A,  
 377 Forster B, Jain M (eds) Advances in haploid production in higher plants. Springer, Heidelberg, pp 295-305  
 378 Kunzel GL, Korzun L, Meister A (2000) Cytologically integrated physical restriction fragment length  
 379 polymorphism maps for the barley genome based on translocation breakpoints. Genetics 154:397-412  
 380 Labbani Z, Richard N, De Buyser J, Piccard E (2005) Plantes chlorophylliennes de blé dur obtenues par culture  
 381 de microspores isolées: importance des prétraitements. Comptes Rendue Biologies 328: 713-723  
 382 Leeuwen H, Monfort A, Zhang H, Puigdomenech P (2003) Identification and characterization of a melon  
 383 genomic region containing a resistance gene cluster from a constructed BAC library. Microcolinearity between  
 384 *Cucumis melo* and *Arabidopsis thaliana*. Plant Molecular Biology 51(5):703-718  
 385 Ma R, Yang-Dong G, Seppo P (2004) Comparison of anther and microspore culture in the embryogenesis and  
 386 regeneration of rye (*Secale cereale*). Plant Cell, Tissue and Organ Culture 76:147-157  
 387 Martins S, Rovira M, Silva AP, Carnide (2010) Pollen incompatibility in Portuguese hazelnut landraces.  
 388 XXVIII International Horticultural Congress on Science and Horticulture for People (IHC2010). Acta  
 389 Horticulturae, pp 932  
 390 Nitsch C (1977) Culture of isolate microspore. In: Reinert J, Bajaj YPS (eds). Applied and fundamental aspects  
 391 of plant cell, tissue and organ culture. Springer, Berlin, pp 268-278  
 392 Prem D, Solís M-T, Bárány I, Rodríguez-Sanz H, Risueño MC, Testillano PS (2012) A new microspore  
 393 embryogenesis system under low temperature which mimics zygotic embryogenesis initials, expresses auxin and  
 394 efficiently regenerates doubled-haploid plants in *Brassica napus*. BMC Plant Biology 12 (127):  
 395 doi:10.1186/1471-2229-12-127  
 396 Rimberia FK, Sunagawa H, Urasaki N, Ishimine Y, Adaniya S (2005) Embryo induction via anther culture in  
 397 papaya and sex analysis of the derived plantlets. Science Horticulturae 103:199-208.  
 398 Rovira M, Aleta N, Germain E, Arus P (1993) Inheritance and linkage relationships of ten isozyme genes in  
 399 hazelnut. Theoretical and Applied Genetics 86:322–328  
 400 Salas P, Rivas-Sendra A, Proens J, Seguí-Shimarro JM (2012) Influence of the stage for anther excision and  
 401 heterostyly in embryogenesis induction from eggplant anther culture. Euphytica 184:235-250  
 402 Scott RJ, Spielman M, Dickinson HG (2004) Stamen structure and function. The Plant Cell 46:S46-S60

403 Seguí-Simaro JM, Nuez F (2008) Pathways to doubled haploidy: chromosome doubling during androgenesis.  
 404 Cytogenetics and Genome Research 120 (3-4):358-369  
 405 Shariatpanahi ME, Bal U, Heberle-Bors E, Touraev A (2006) Stresses applied for the re-programming of plant  
 406 microspores towards in vitro embryogenesis. *Physiologia Plantarum* 127:519-534  
 407 Silva TD (2012) Microspore embryogenesis. In: Sato, K-I (ed) *Embryogenesis*, In Tec, pp 573–596.  
 408 [www.intechopen.com](http://www.intechopen.com)  
 409 Soriano M, Li H, Boutilier K (2013) Microspore embryogenesis: establishment of the embryo identity and  
 410 pattern in culture. *Plant Reproduction* 26:181-193  
 411 Spory S, Munshi M (1996) Anther culture. In: Mohan, J.M. *et al.* (eds). *In vitro haploid production in higher*  
 412 *plants*, Vol. 1, Kluwer, Dordrecht, pp 145-176  
 413 Szarejko I, Foster BP (2006) Doubled haploidy and induced mutation. *Euphytica* 158:359-370.  
 414 Talón M, Gmitter FG (2008) Citrus genomics. *International Journal of Plant Genomics*.  
 415 <http://dx.doi.org/10.1155/2008/528361>  
 416 Touraev A, Ilham A, Vicente O, Heberle-Bors E (1996a) Stress induced microspore embryogenesis from  
 417 tobacco microspores: an optimized system for molecular studies. *Plant Cell Reports* 15:561–565  
 418 Touraev A, Indrianto A, Wratschko I, Vicente O, Heberle-Bors E (1996b) Efficient microspore embryogenesis  
 419 in wheat (*Triticum aestivum* L.) induced by starvation at high temperatures. *Sexual Plant Reproduction* 9:209–  
 420 215  
 421 Wang M, van Bergen S, Van Duijn B (2000) Insights into a key developmental switch and its importance for  
 422 efficient plant breeding. *Plant Physiol* 124:523–530  
 423 Wang Z, Tarantino G, Yang D, Liu G, Tingey SV, Miao GH, Wang GL (2001) Rice ESTs with disease-  
 424 resistance gene or defense-response gene like sequences mapped to regions containing major resistance genes or  
 425 QTLs. *Molecular Genetics and Genomics* 265:302-310  
 426 Zavattieri MA, Frederico AM, Lima M, Sabino R, Arnhold-Schmitt B (2010) Induction of somatic  
 427 embryogenesis as an example of stress-related plant. *Electronic Journal of Biotechnology* 13(1):1-9  
 428 Zhang KP, Zhao L, Tian JC, Chen GF, Jiang XL, Liu B (2008) A genetic map constructed using a doubled  
 429 haploid population derived from two elite Chinese common wheat varieties. *Journal Integrative Plant Biology*  
 430 50(8):941-950  
 431

## Tables

Table 1 Influence of genotype on isolated microspore development, after twenty months of culture.

Genotype	Uninucleated microspores %	Binucleated microspores (SND) %	Binucleated microspores (ASD) %	Trinucleated microspores %	Multinucleated microspores %
‘Carrello’	27.7 a	28.4 ab	1.9 a	12.9 b	29.1 ab
Tonda G. Romana GR	30.0 a	25.3 bc	2.9 a	11.8 b	30.0 ab
‘Imperatrice Eugenia’	23.0 b	23.5 c	0.1 b	22.0 a	31.4 a
‘Minnulara’	23.4 b	31.1 a	0.6 b	21.6 a	23.3 b
Tonda G. Romana TR	27.6 a	27.1 ab	0.6 b	19.7 a	25.0 b

Per each column, values followed by different letters are statistically different.

One-way ANOVA, Conover-Inman’s test,  $p \leq 0.05$ .

SND: Symmetrical nucleus division

AND: Asymmetrical nucleus division

Table 2 Influence of medium composition and thermal treatment on ‘Carrello’ isolated microspore development, after twenty months of culture.

Factors		Uninucleated microspores %	Binucleated microspores (SND) %	Binucleated microspores (ASD) %	Trinucleated microspores %	Multinucleated microspores %
Medium (M)	P	27.6 a	25.5 b	2.6 a	12.6 a	31.7 a
	N6	27.7 a	31.2 a	1.2 a	13.3 a	26.5 b
M p value		0.962	0.003	0.073	0.593	0.014
Treatment (T)	Co	23.7 b	29.7 a	1.7 a	13.7 a	31.3 a
	H	34.0 a	26.2 a	2.1 a	11.0 a	26.7 a
	H+	29.4 ab	26.4 a	1.4 a	11.4 a	31.4 a
	F	26.7 ab	29.3 a	2.0 a	14.0 a	28.0 a
	F+	24.7 ab	30.3 a	2.3 a	14.7 a	28.0 a
	T p value	0.025	0.398	0.928	0.400	0.412
M X T p value		0.425	0.330	0.658	0.175	0.145

Per each factor and per each column, values followed by different letters are statistically different.

Two-way ANOVA, Tukey’s test,  $p \leq 0.05$ .

SND: Symmetrical nucleus division

AND: Asymmetrical nucleus division

P medium: Germanà et al 1996; N6 medium: Germanà and Chiancone, 2003.

Co: no thermal treatment; H: 35 °C for 30 minutes; H+: 40 °C for 60 minutes; F: -20 °C for 30 minutes; F+: -20 °C for 60 minutes.

Table 3 Influence of medium composition and thermal treatment on ‘Tonda Gentile Romana’ GR isolated microspore development, after twenty months of culture.

Factors		Uninucleated microspores	Binucleated microspores (SND)	Binucleated microspores (ASD)	Trinucleated microspores	Multinucleated microspores
		%	%	%	%	%
Medium (M)	P	32.7 a	22.8 b	3.4 a	11.4 a	29.6 a
	N6	27.2 b	27.7 a	2.4 a	12.2 a	30.5 a
M p value		0.010	0.005	0.320	0.598	0.501
Treatment (T)	Co	29.0 a	26.3 a	2.0 a	14.3 a	28.3 ab
	H	34.8 a	23.8 a	4.6 a	10.9 a	25.9 b
	H+	29.0 a	25.0 a	2.0 a	11.7 a	32.3 ab
	F	28.2 a	25.9 a	3.0 a	11.9 a	30.9 ab
	F+	28.7 a	25.3 a	3.0 a	10.3 a	32.7 a
	T p value	0.050	0.852	0.500	0.523	0.023
M X T p value		0.010	0.272	0.157	0.694	0.000

Per each factor and per each column, values followed by different letters are statistically different.

Two-way ANOVA, Tukey's test,  $p \leq 0.05$ .

SND: Symmetrical nucleus division

AND: Asymmetrical nucleus division

P medium: Germanà at al 1996; N6 medium: Germanà and Chiancone, 2003.

Co: no thermal treatment; H: 35 °C for 30 minutes; H+: 40 °C for 60 minutes; F: -20 °C for 30 minutes; F+: -20 °C for 60 minutes.

Table 4 Influence of medium composition and thermal treatment on ‘Imperatrice Eugenia’ isolated microspore development, after twenty months of culture.

Factors		Uninucleated microspores	Binucleated microspores (SND)	Binucleated microspores (ASD)	Trinucleated microspores	Multinucleated microspores
		%	%	%	%	%
Medium (M)	P	23.2 a	22.9 a	0.0 a	20.3 a	34.5 a
	N6	22.4 a	22.5 a	0.0 a	22.1 a	33.9 a
M p value			0.514	0.173	0.486	0.294
Treatment (T)	Co	19.3 a	23.7 a	0.0 a	20.7 a	36.3 a
	H	23.0 a	21.3 a	0.3 a	20.3 a	37.3 a
	H+	26.7 a	27.3 a	0.0 a	16.3 a	29.7 a
	F	20.0 a	21.7 a	0.0 a	26.7 a	31.7 a
	F+	26.0 a	23.7 a	0.3 a	23.7 a	26.3 a
	T p value		0.337	0.570	0.096	0.209
M X T p value			0.823	0.570	0.544	0.024

Per each factor and per each column, values followed by different letters are statistically different.

Two-way ANOVA, Tukey's test,  $p \leq 0.05$ .

SND: Symmetrical nucleus division

AND: Asymmetrical nucleus division

P medium: Germanà at al 1996; N6 medium: Germanà and Chiancone, 2003.

Co: no thermal treatment; H: 35 °C for 30 minutes; H+: 40 °C for 60 minutes; F: -20 °C for 30 minutes; F+: -20 °C for 60 minutes.

Table 5 Influence of medium composition and thermal treatment on ‘Minnulara’ isolated microspore development, after twenty months of culture.

Factors		Uninucleated microspores	Binucleated microspores (SND)	Binucleated microspores (ASD)	Trinucleated microspores	Multinucleated microspores
		%	%	%	%	%
Medium (M)	P	25.6 a	24.5 b	0.0 a	19.7 a	30.1 a
	N6	29.6 a	28.2 a	1.0 a	19.4 a	21.8 b
M p value		0.067	0.000	0.415	0.245	0.000
Treatment (T)	Co	24.3 a	23.0 a	0.3 a	22.7 a	29.7 a
	H	28.3 a	25.7 a	0.7 a	16.0 b	29.3 a
	H+	31.0 a	28.0 a	0.3 a	18.3 ab	22.3 a
	F	26.3 a	28.1 a	0.4 a	21.2 a	23.9 a
	F+	28.0 a	27.0 a	0.7 a	19.7 ab	24.7 a
T p value		0.059	0.282	0.289	0.010	0.944
M X T p value		0.008	0.234	0.500	0.008	0.714

Per each factor and per each column, values followed by different letters are statistically different.

Two-way ANOVA, Tukey's test,  $p \leq 0.05$ .

SND: Symmetrical nucleus division

AND: Asymmetrical nucleus division

P medium: Germanà et al 1996; N6 medium: Germanà and Chiancone, 2003.

Co: no thermal treatment; H: 35 °C for 30 minutes; H+: 40 °C for 60 minutes; F: -20 °C for 30 minutes; F+: -20 °C for 60 minutes.

Table 6 Influence of medium composition and thermal treatment on ‘Tonda Gentile Romana’ TR isolated microspore development, after twenty months of culture.

Factors		Uninucleated microspores	Binucleated microspores (SND)	Binucleated microspores (ASD)	Trinucleated microspores	Multinucleated microspores
		%	%	%	%	%
Medium (M)	P	25.6 b	24.5 a	0.0 a	19.7 a	30.1 a
	N6	29.6 a	28.2 a	1.0 a	19.4 a	21.8 b
M p value		0.035	0.303	0.204	0.794	0.002
Treatment (T)	Co	24.3 a	23.0 a	0.3 a	22.7 a	29.7 a
	H	28.3 a	25.7 a	0.7 a	16.0 a	29.3 a
	H+	31.0 a	28.0 a	0.3 a	18.3 a	22.3 b
	F	26.3 a	28.1 a	0.4 a	21.2 a	23.9 b
	F+	28.0 a	27.0 a	0.7 a	19.7 a	24.7 b
T p value		0.222	0.221	0.709	0.364	0.007
M X T p value		0.010	0.010	0.954	0.330	0.043

Per each factor and per each column, values followed by different letters are statistically different.

Two-way ANOVA, Tukey's test,  $p \leq 0.05$ .

SND: Symmetrical nucleus division

AND: Asymmetrical nucleus division

P medium: Germanà et al 1996; N6 medium: Germanà and Chiancone, 2003.

Co: no thermal treatment; H: 35 °C for 30 minutes; H+: 40 °C for 60 minutes; F: -20 °C for 30 minutes; F+: -20 °C for 60 minutes.



## Figure legends

Fig. 1: a) Catkins and one anther of *Corylus avellana* L., cv. 'Imperatrice Eugenia', at the developmental stage used for microspore isolation; b) Uninucleated microspore of cv. 'Imperatrice Eugenia' (DAPI staining) (Bar represents 10µm).

Fig. 2: a) Binucleated pollen, symmetrical division (DAPI staining), cv. 'Imperatrice Eugenia'; b) Trinucleated microspore (DAPI staining), cv. 'Imperatrice Eugenia'; c-d) Multinucleated microspore (DAPI staining), cv. 'Carrello' (c) and 'Tonda Gentile Romana' TR (d) (Bars represent 10µm).

Fig. 3: Microspore-derived calli of cv. 'Tonda Gentile Romana' GR (Bars represent 100µm).

Fig. 4 Microspore-derived embryos of cv. 'Tonda Gentile Romana' GR: a-c) different stages of development (Bars represent 20µm).

Fig. 5: a) Influence of the genotype on the regeneration of calli and embryos of five hazelnut genotypes (One way ANOVA, Conover-Inman's test  $p \leq 0.05$ ) Within each parameter (callus or embryo), values followed by different letters are statistically different at  $p \leq 0.05$  according to Conover-Inman's test; b-f): Influence of the medium composition (P and N6) and of thermal treatments (35 °C for 30 minutes (H), 40 °C for 60 minutes (H+), -20 °C for 30 minutes (F), -20 °C for 60 minutes (F+)) on the production of calli and embryos of 'Carrello' (b), 'Tonda Gentile Romana' GR (c); 'Imperatrice Eugenia' (d), 'Minnulara' (e) and 'Tonda Gentile Romana' TR (f).

Fig. 6 Genetic profile at SSR locus CaT-B505 of the cultivar "Minnulara" (above) and of the microspore-derived embryo (below). The embryo shows a single allele (128 bp) shared with the parental genotype.

Fig. 7 Genetic profile at SSR locus CaT-B107 of the cultivar "Minnulara" (above) and of the microspore-derived embryo (below). The embryo shows a single allele (114 bp) shared with the parental genotype.