

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Seasonal changes in the fillet fatty acid profile and nutritional characteristics of wild Trasimeno Lake goldfish (*Carassius auratus* L.)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1620077> since 2017-04-04T12:15:48Z

Published version:

DOI:10.1016/j.foodchem.2011.11.043

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in FOOD CHEMISTRY, 132 (2), 2012, 10.1016/j.foodchem.2011.11.043.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.foodchem.2011.11.043

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0308814611016244>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1620077>

Seasonal changes in the fillet fatty acid profile and nutritional characteristics of wild Trasimeno Lake goldfish (*Carassius auratus* L.)

Alessandro Dal Bosco [†], Cecilia Mugnai, Evangelia Mourvaki, Cesare Castellini
Department of Applied Biology, University of Perugia, XX Giugno 74, 06100 Perugia, Italy

abstract

Goldfish (*Carassius auratus* L.) is very diffuse in Trasimeno Lake, and show great adaptability and have become a dominant fish of the lake ecosystem. The aim of this study was to evaluate the seasonal effect on the fatty acid profile and nutritional characteristics of goldfish caught in Trasimeno Lake. Forty fillets per season were used to evaluate the respective proximate composition, fatty acid profile, nutritional indices and protein and lipid oxidative stability. Season significantly affected the chemical traits of the fillets. A lower level of SFA was observed during spring, while a higher proportion of MUFA was observed during the winter. PUFA exceeded 50% of the total fatty acids content during the autumn, spring and summer. The nutritional indices were very good in comparison to those observed in other livestock animals. Goldfish fillets are characterised by a high nutritional value and good oxidative stability; the best results of this study were obtained in autumn and spring fillets.

Introduction

Trasimeno is one of the largest lakes of the Italian peninsula, with a surface area of 128 km²; the water level significantly fluctuates due to rainfall and seasonal demands from the towns, villages and farms that dot the shore. Trasimeno has always been quite shallow, muddy and very rich in fishes (pike, carp and tench).

Goldfish (*Carassius auratus* L.) is an allochthonous species very diffuse at present in the Trasimeno, showing great adaptability. They have become the dominant fish of the lake ecosystem (Lorenzoni, Corboli, Ghetti, Pedicillo, & Carosi, 2007; Mearelli, Lorenzoni, & Mantilacci, 1990). The high invasive capacity of this species is due to its tolerance to extreme environmental conditions (low temperature, high rate of pollution, cloudy water with a low percentage of oxygen), its high fertility and wide alimentary regimen. The temperature tolerance is the most important factor for success of goldfish; in fact it can survive from 0.3 to 43.6 °C (Ford & Beitingger, 2005). This species lives in shallow ponds, lakes rich in vegetation and slow-moving rivers. During the dry season or winter it burrows in mud (Ford & Beitingger, 2005). It feeds on plants, insect larvae and plankton. For these reasons, goldfish, as well as all the allochthonous species, alter the equilibrium of the fish community of the Trasimeno Lake, to the detriment mainly of carp and tench.

When caught for food uses, the greatest economic disadvantage of this fish is the large number of fine inter-muscular bones the species possesses, which makes it laborious to eat for the unaccustomed. Thus, this meat is used as food only after undergoing

processing that improves the economic value of goldfish. Therefore, it is very unlikely that goldfish production could increase and become important in the international market. On the other hand, there are some advantages of fishing Trasimeno goldfish; in fact, this species is dangerous to the Trasimeno Lake ecosystem, making its capture a solution to reduce its number, especially in springtime before the reproductive phase. Moreover, this solution could increase the income of local fishermen. So, knowledge of this species, also present in other Italian natural freshwater environments, would be valuable.

A previous study (Dal Bosco, Mourvaki, Mugnai, & Castellini, 2010) found that goldfish meat, processed into croquettes, can be considered a good source of n-3 polyunsaturated fatty acid (PUFA), providing approximately 78.6 mg of Eicosapentaenoic fatty acid (C20:5n-3, EPA) and 137 mg of docosahexaenoic fatty acid (C22:6n-3, DHA) per 100 g of consumed product.

An important issue regarding captured goldfish on a semi-commercial scale is the homogeneity of the chemical composition of fish meat. Indeed, the chemical composition of other fish species varies from season to season due to their natural cycling, maturation and feeding. Thus, the aim of this study was to evaluate the seasonal effect on the fatty acid profile and nutritional characteristics of goldfish caught in Trasimeno Lake.

1. Materials and methods

1.1. General

This study was carried out in collaboration with the Fisherman Cooperative "Albatrasimeno" of S. Feliciano (Perugia, Italy), utilizing 20 fished goldfish per season (February, April, July and October, 2010), which, after being caught, were transferred to the laboratory of the cooperative.

The average water temperature and the amount of oxygen dissolved in the above-mentioned periods were, respectively: February 9.5 °C, 13.8 mg/L; April 12.6 °C, 10.5 mg/L; July 23.4 °C, 8.9 mg/L; October 18.9 °C, 9.6 mg/L.

1.2. Fillets and samples preparation

After washing with running water, goldfish with an average live weight of 550 g were gutted; the tails and heads were then removed. The epaxial and ipaxial parts of the fillets were dissected and both utilised for the preparation of 40 individual samples (mean weight of 100 ± 13 g), for analysis for each season.

The samples were immediately transferred to the laboratory of the Department of Applied Biology of the University of Perugia for analysis.

1.3. Chemical composition

Analyses were immediately carried out in duplicate to determine the proximate composition. Moisture, ash and total nitrogen were obtained using AOAC (1995) methods (N. 950.46B, 920.153 and 928.08, respectively). The total protein content was calculated using Kjeldahl nitrogen and a conversion factor of 6.25. The total lipid content was extracted (Folch, Lees, & Sloane-Stanley, 1957) from 5 g of each homogenised sample and determined gravimetrically.

1.4. Fatty acid profile and desaturase activity

Fatty acids contents were determined by gas chromatography after lipid extraction according to the Folch et al. (1957) method. One millilitre of lipid extract was evaporated under a stream of nitrogen, and the residue was derivatised by adding 3 ml of sulphuric acid (3% in methanol). After incubating at 80 °C for 1 h, methyl esters were extracted with petroleum ether, and 1 μ l was injected into the gas chromatograph (Fisons Mega 2 Carlo Erba Gas Chromatograph, model HRGC, Milan, Italy) equipped with a flame ionisation detector.

The separation of fatty acid methyl esters (FAME) was carried out on an Agilent (J&W) capillary column (30 m \times 0.25 mm I.D, CPS Analitica, Milan, Italy) coated with a DB-Wax stationary phase (film thickness of 0.25 μ m). The operating conditions of the column injection were as follows: the temperatures of the injector and detector were 270 and 280 °C, respectively; the detector gas flows were H₂ 50 ml/min and air 100 ml/min. The oven temperature was programmed to give good peak separation; the initial temperature was set at 130 °C and it was then increased at a rate of 4.0 °C/min until reaching a temperature of 180 °C, which was held for 5 min; the temperature was subsequently increased at a rate of 5.0 °C/min until it reached 230 °C, which was held for 5 min. Helium was used as a carrier gas at a constant flow rate of 1.1 ml/min. Individual fatty acid methyl esters were identified with reference to the retention time of tri-decanoic acid (C13:0) methyl ester, added before extraction as an internal standard. The relative proportion of individual fatty acids was expressed as a percentage.

To evaluate the activity of both D⁵-desaturase and D⁶-desaturase, the enzymes catalysing the formation of long-chain n-6 and n-3 PUFA, starting from the precursors C18:2n-6 and C18:3n-3, the following equation was calculated, as suggested by Sirri, Castellini, Roncarati, Franchini, and Meluzzi (2010):

$$\frac{1}{4} \left[\frac{\%C20 : 2n - 6}{C18 : 3n - 3} + \frac{\%C20 : 4n - 6}{C20 : 2n - 6} + \frac{\%C20 : 5n - 3}{C20 : 4n - 6} + \frac{\%C22 : 5n - 3}{C22 : 6n - 3} + \frac{\%C22 : 6n - 3}{C22 : 5n - 3} \right] \times 100$$

1.5. Nutritional characteristics

The mean value of each fatty acid was used to calculate the sum of the saturated

(SFA), monounsaturated (MUFA) and PUFA, and calculate the peroxidability index (PI) according to the equation proposed by Arakawa and Sagai (1986):

$$PI = \frac{1}{4} \times \text{\%monoenoic} + 0.025 \times \text{\%dienoic} + 1 \times \text{\%trienoic} + 2 \times \text{\%tetraenoic} + 4 \times \text{\%pentaenoic} + 6 \times \text{\%hexaenoic} + 8 \times \text{\%heptaenoic}$$

The concentration of each fatty acid (mg/100 g of fish) was obtained from the lipid content of the fish, using the conversion factor 0.91 according to Johansson, Kiessling, Kiesling, and Berglund (2000).

The amount of each fatty acid was used to calculate the indices of atherogenicity (AI) and thrombogenicity (TI), as proposed by Ulbricht and Southgate (1991), and the hypocholesterolaemic/hypercholesterolaemic ratio (HH), as suggested by Santos-Silva, Bessa, and Santos-Silva (2002):

$$\begin{aligned} - AI &= (C12:0 + 4 \times C14:0 + C16:0) / [(P_{MUFA} + P_{(n-6)} + P_{(n-3)})]; \\ - TI &= (C14:0 + C16:0 + C18:0) / [(0.5 \times P_{MUFA} + 0.5 \times (n-6) + 3 \times (n-3) + (n-3)/(n-6)]; \end{aligned}$$

$$HH = \frac{C18:1n-9 + C18:2n-6 + C20:4n-6 + C18:3n-3 + C20:5n-3 + C22:5n-3 + C22:6n-3}{C14:0 + C16:0}$$

1.6. Assessment of oxidative stability

The extent of lipid oxidation was evaluated in fillets as thiobarbituric acid-reactive substances (TBARs), according to the modified method of Ke, Ackman, Linke, and Nash (1977). Oxidation products were quantified as malondialdehyde (MDA) equivalents (mg MDA/kg muscle).

The amount of products arising from oxidative damage to proteins (protein carbonyl groups (PCGs)) were measured spectrophotometrically by reacting with 2,4-dinitrophenylhydrazine (DNPH) according to the method proposed by Lushchak et al. (2005). In detail, samples were homogenised (1:10 w/v) using a Potter-Elvehjem glass homogeniser in 50 mM potassium phosphate buffer, pH 7.0, containing 0.5 mM EDTA. A few crystals of phenylmethylsulphonyl fluoride were added prior to homogenisation to inhibit proteases. A 250 µl aliquot of this homogenate was then mixed with 0.5 ml of 10% (final concentration) trichloroacetic acid and centrifuged for 5 min at 13,000g. The resulting pellets were mixed with 1 ml of 10 mM DNPH in 2 M HCl and incubated for 1 h at room temperature. Control samples were mixed with 1 ml of 2 M HCl. After washing three times with 1 ml of ethanol-butylacetate (1:1 v/v), pellets were dissolved in 1.5 ml of urea (6 M), and PCGs were measured using a spectrophotometer (Hitachi U-2000) set at 370 nm, using a molar extinction coefficient of 22×10^3 M/cm. The values are expressed as nanomoles of PCGs per milligramme of protein in the urea solution. The protein concentration was measured by the Bradford (1976) method with Coomassie Brilliant Blue G-250, using bovine serum albumin as a standard.

The concentration of thiol groups (SH) was spectrophotometrically measured at

412 nm (molar extinction coefficient of 13.6×10^3 M/cm) in supernatants after reaction with 5,5⁰-dithio-bis (2- nitrobenzoic acid), as described by Lushchak and Bagnyukova (2006). Sample homogenates were prepared as described for the PCG assay, and supernatants were obtained after centrifugation at 4 °C for 15 min at 15,000g. The thiol levels are expressed as micromoles of SH-groups per gramme.

1.7. Statistical analysis

The qualitative traits were analysed with a linear model (STATA, 2005) and a fixed effect of season. Least squares means and planned comparisons were used for mean separation when the model was significant ($P < 0.05$).

2. Results and discussion

The chemical characteristics of goldfish captured during different seasons are listed in Table 1. In general, the fillets of Trasimeno Lake goldfish showed protein contents similar to those observed by Hossain and Jauncey (1989) and Takeuchi (1979) in fillets of carp (*Cyprinus carpio*). The season significantly affected the chemical traits of the fillets; in particular, the moisture level was lowest during autumn and did not exhibit significant differences during the other seasons. A higher level of protein was observed during autumn, whereas the lipid contents were lower during winter and spring.

The fatty acid profiles of fillets during the analysed period (Table 2) generally showed some important peculiarities unique to this species. As observed in our previous investigation (Dal Bosco et al., 2010), the major fatty acids in goldfish fillet were palmitic (16:0), oleic (C18:1n-9), linoleic (C18:2n-6), arachidonic (C20:4n-6), EPA and DHA, according to the results of Vornanen, Tiitu, Kkel, and Aho (1999) for crucian carp (*Carassius carassius*). Independent of the season, these results are very interesting when compared with those observed for other species, such as the much more economically relevant sea bass (*Dicentrarchus labrax*) and gilt-head seabream (*Sparus aurata*), whose levels of EPA and DHA reach 10.9% and 10.9% (EPA) and 10.7% and 14.0% (DHA), respectively (Turchetto et al., 1994). Also, a comparison between the two fish species present in the lakes of central Italy, coregon (*Coregonus lavaretus*) and perch (*Perca fluviatilis* L.) (PRAL, 1999–2001), confirm the good nutritional properties of goldfish meat. Poli et al. (1994) analysed the fatty acid profiles of this last species caught from Trasimeno Lake and found values of PUFA equal to 33.3%, with 9.78% of EPA and 19.9% DHA.

A lower level of SFA was observed during spring, while a higher proportion of MUFA was observed during winter. PUFA exceeded 50% of the total fatty acid content in autumn, spring and summer, with very good levels of EPA (6.8%, 6.5% and 8.1%) and DHA (14.6%, 13.3% and 14.5%, respectively). Accordingly, the activities of D⁵-desaturase and D⁶-desaturase were significantly lower during winter, intermediate during spring and higher during autumn and summer.

The seasonal differences in the chemical composition and fatty acid profiles can be ascribed to different factors, such as the availability of aquatic plants, water

temperature and the physiological phases of fish (Haard, 1992). With respect to the first factor, it is useful to remember that the Trasimeno Lake ecosystem is characterised by a winter scarcity of vegetal species, followed by a quick spring resurgence; subsequently, a slow decline takes place, which culminates in the summer. Afterwards, an autumnal recycling occurs but generally does not reach the spring peak. Although Trasimeno Lake has a laminar structure, there are important changes in water temperature and dissolved oxygen that occur throughout the year: the temperature varies from 8 to 9 °C in December and January to 26–27 °C in August (Mearelli et al., 1990). Finally, from a biological point of view, Lorenzoni et al. (2007) reported that, in winter, goldfish reduce their basal metabolism and growth by re-using body resources in order to survive such that their condition worsens until February. At the beginning of spring, their overall body condition starts to improve, due to the weight gain attributed to the development of the gonads but, as the season progresses, there is a new decline that occurs due to spawning and then the emptying of the gonads. The somatic condition, as opposed to the overall body condition, continues to decline during spring, when the fish faces the maximum reproductive effort. The spring weight loss is not only due to the emptying of the gonads but also to the loss of muscle mass and tissue reserves, resulting in significant waste energy attributed to reproduction. During the summer, the increased availability of food and Mendez and Gonzales (1997) also observed a decrease in the lipid content of fillets from southwest Atlantic hake during favourable climatic conditions allow for the rapid growth and accumulation of fat reserves. In this way, goldfish reach an optimal body condition during the autumn season. Then, with respect to physiological status, it is possible to observe two opposite conditions, ranging from a semi-hibernation state during winter to intense swimming activity in spring/summer.

spawning months. During these periods, the protein and lipid contents decreased due to the use of muscle lipids and proteins as energy reserves. In autumn, the protein and lipid contents increased and, during winter, when the hake were not feeding, the lipid content decreased, indicating the use of lipids as fuel during food shortage.

Tocher et al. (2004) and Ninno, De Torrengo, Costuma, and Brenner (1974) pointed out, in rainbow trout (*Oncorhynchus mykiss*) and *Pimelodus maculatus*, respectively, the important role of water temperature in the regulation of the desaturation/elongation processes during β -oxidation in hepatocytes and enterocytes: the activities of D^6 and D^5 desaturase increased many times when the temperature ranged from 30 to 16 °C. The results regarding goldfish seem to contrast with this affirmation, probably because, as has already been mentioned, this is a fish species that has a very slow metabolism during winter, as it exhibits a lower value of D^5 plus D^6 desaturase activity.

In our study, the good fatty acid profile exhibited during spring, summer and autumn is justified by the great availability of vegetation and thus by a higher probable antioxidant intake.

Table 3 shows the seasonal changes of the meat nutritional indices. As expected, due to the highest lipid content, the SFA, sum of n-3, EPA and DHA were highest during the autumn and summer; content of PUFA was highest in the autumn.

The AI and TI showed very low values in comparison to those observed in other livestock animals. In particular, AI and TI were both lower than those reported by Dal Bosco, Castellini, Bianchi, and Mugnai (2004) for rabbit meat (0.70 and 0.99), Dal Bosco, Castellini, and Cardinali (2005) for pigeon meat (0.41 and 0.94) and Castellini, Dal Bosco, Mugnai, and Pedrazzoli (2005) for poultry meat (0.49 and 0.88). The best AI was found during autumn and spring.

The HH ratio observed in autumn and winter was lower than that found by Testi, Bonaldo, Gatta, and Badiani (2006) in the dor- sal (2.93 *vs.* 2.40) and ventral rainbow trout fillets (2.93 *vs.* 2.46). These data confirm the optimal nutritional quality of goldfish fillets (Dal Bosco et al., 2010) as well as the seasonal affect on the studied traits.

Kminkova, Winterova, and Kucera (1993) found similar values of EPA and DHA contents in pond-reared carp muscle only during the summer (149 and 214 mg/100 g tissue, respectively); during the other seasons, the values were very low, probably due to the diets based on wheat, rye, barley and pea. For the wild carp of Ivriz Dam Lake (Turkey), Kalyoncu, Yaman, and Aktumsek (2010) obtained results similar to ours; in particular, these authors found higher values of EPA and DHA and lower of LA in fillets during au- tumn. In fillets of the wild-caught carp of another Turkish lake, Guler, Kiztanir, Aktumsek, Cital, and Ozparlak (2008) found very high values of EPA and DHA during summer.

Vujkovic, Karlovic, Vujkovic, Vörösbaranyic, and Jovanovic (1999) found EPA and DHA contents equal to 150 and 89 mg/ 100 g fillet for silver carp (*Hypophthalmichthys molitrix* Val.) and big- head carp (*Aristichthys nobilis* Val.) harvested in the Despotovo Fishfarm (Vojvodina, Yugoslavia).

From a nutritional point of view, the EPA + DHA content is of great interest because of the role of these fatty acids in the therapy and prevention of cardiovascular diseases (Uauy & Valenzuela, 1992). The mean values for the EPA + DHA contents were 430, 251, 338 and 426 mg per 100 g fillet tissue, respectively, in autumn, winter, spring and summer (data not shown). This means that, for the recommended daily ingestion of 1 g of EPA + DHA (Ackman, 1989), about 250–300 g of goldfish fillets are needed. Moreover, intakes of EPA and DHA of about 250 mg per day are required to maintain normal cardiac function (EFSA Journal, 2010). The best values for the n-6/n-3 ratio were observed during autumn, spring and summer. A decrease in the n-6/n-3 ratio in the human diet is essential to prevent coronary heart disease by reducing plasma lipids and to reduce the risk of cancer (Kinsella, Lokesh, & Stone, 1990). Nutrition advisers recommend an in- creased intake of n-3 in human diets with a reduction of n-6/n-3 ratio to values below 4 (Kark, Kaufmann, Binka, Goldberger, & Berry, 2003). In the present trial, the n-6/n-3 ratio values were al- ways much below 4.

Regarding the oxidative status, TBARs, carbonyl and thiol con- tents were higher during the summer. These results reflect the highest amounts of long-chain n-3 fatty acids. Considering the high degree of unsaturation found during all seasons and the low values of TBARs and PI, the oxidative status was good. This is evident if compared with those found, in our previous study, on rainbow trout (*Oncorhynchus*

mykiss) reared in Valnerina (Italy), where values of TBARs of 0.79 mg of MDA per kg of fresh fillet and a PI of 239, were found. The situation related in the present study could be due to the good capture technique, to the high ingestion of antioxidants during spring and autumn and semi-hibernation during winter.

Some authors (Lushchak & Bagnyukova, 2006; Lushchak et al., 2005), have demonstrated that Goldfish tissues possess high constitutive activities of antioxidant enzymes which might be enough to cope with short-term disturbance of free radical processes.

Finally, the trend found in TBARs, carbonyls and thiols suggests possible relationships among these oxidative stress markers. Hence, it is likely that the production of MDA during lipid oxidation might trigger the protein carbonylation. In nature, a gradual increase in some antioxidant enzyme activities during the transition from cold winter conditions to warmer summer conditions would reflect the needs of an organism to cope with elevated oxidative stress that accompanies enhanced oxygen consumption and metabolic rate at higher environmental temperatures. This hypothesis might be even more probable if based on the fact that goldfish are a wild species with a more accentuated muscular oxidative metabolism (Lushchak & Bagnyukova, 2006). In conclusion, this study reveals that, during different environmental and physiological conditions change, and cause seasons, caught during these periods should increase. This should be advantageous for both healthy characteristics of fillets obtained from fish and for re-equilibration of the fish ecosystem of Trasimeno Lake.

The characterisation and valorisation of this product could represent a good economic opportunity for local fishermen.

Acknowledgements

The authors are grateful to Dr. Mauro Natali (Servizio Programmazione e Gestione Faunistica, Provincia Perugia), Mr. Aurelio Cocchini (Cooperativa Pescatori Alba Trasimeno, S. Feliciano, Perugia) and Mr. Francesco Gonnelli for technical assistance.

References

- AOAC (1995). *Official methods of analysis* (15th ed.). Washington, DC: Association of Official Analytical Chemists.
- Ackman, R. G. (1989). Nutritional composition of fats in seafoods. *Progress in Food and Nutrition Science*, 13, 161–241.
- Arakawa, K., & Sagai, M. (1986). Species differences in lipid peroxide levels in lung tissue and investigation of their determining factors. *Lipids*, 21, 769–775.
- Bradford, M. M. (1976). A rapid and sensitive method for quantization of microgram quantities of protein utilizing the principle of protein-dye-binding. *Analytical Biochemistry*, 72, 248–258.
- Castellini, C., Dal Bosco, A., Mugnai, C., & Pedrazzoli, M. (2005). Oxidative stability and sensory attributes of poultry meat during display according to age and rearing system. *Italian Journal Animal Science*, 5, 355–365.

- Dal Bosco, A., Castellini, C., Bianchi, L., & Mugnai, C. (2004). Effect of dietary α -linoleic acid and vitamin E on fatty acid composition, storage stability and sensory traits of rabbit meat. *Meat Science*, 66, 407–413.
- Dal Bosco, A., Castellini, C., & Cardinali, R. (2005). Effect of dietary administration of rosemary extract on the oxidative stability of pigeon meat. *Italian Journal of Food Science*, 4, 17–26.
- Dal Bosco, A., Mourvaki, E., Mugnai, C., & Castellini, C. (2010). Nutritional evaluation of fillets, Pulp and croquette of wild caught Trasimeno lake goldfish (*Carassius auratus* L.). *Italian Journal of Food Science*, 2, 192–199.
- EFSA Journal (2010). Scientific opinion on the substantiation of health claims related to eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), docosapentaenoic acid (DPA). *EFSA Journal*, 8, 1796.
- Folch, J., Lees, M., & Sloane-Stanley, G. H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biology and Chemistry*, 226, 497–509.
- Ford, T., & Beitinger, T. L. (2005). Temperature tolerance in the goldfish, *Carassius auratus*. *Journal of Thermal Biology*, 30, 147–152.
- Guler, G. O., Kiztanir, B., Aktumsek, A., Citil, O. B., & Ozparlak, H. (2008). Determination of the seasonal changes on total fatty acid composition and n-3/n-6 ratios of carp (*Cyprinus carpio* L.) muscle lipids in Beysehir Lake (Turkey). *Food Chemistry*, 108, 689–694.
- Haard, F. N. (1992). Control of chemical composition and food quality attributes of cultured fish. *Food Research International*, 25, 289–307.
- Hossain, M. A., & Jauncey, K. (1989). Nutritional evaluation of some Bangladeshi oil seed meals as partila substitutes for fish meal in the diets of common carp, *Cyprinus carpio* L. *Aquaculture, Fishery and Management*, 20, 255–268.
- Johansson, L., Kiessling, A., Kiesling, K. H., & Berglund, L. (2000). Effect of altered ration levels on sensory characteristics, lipid content and fatty acid composition of rainbow trout (*Oncorhynchus mikiss*). *Food Quality and Preference*, 11, 247–254.
- Kark, J. D., Kaufmann, N. A., Binka, F., Goldberger, N., & Berry, E. M. (2003). Adipose tissue n-6 fatty acids and acute myocardial infarction in a population consuming a diet high in polyunsaturated fatty acids. *American Journal Clinic and Nutrition*, 77, 796–802.
- Kalyoncu, L., Yaman, Y., & Aktumsek, A. (2010). Seasonal changes on total fatty acid composition of carp (*Cyprinus carpio* L), in Ivriz Dam Lake, Turkey. *African Journal of Biotechnology*, 9, 3896–3900.
- Ke, P. J., Ackman, R. G., Linke, B. H., & Nash, D. M. (1977). Differential lipid oxidation products in various parts of frozen mackerel. *Journal of Food Technology*, 12, 37–47.
- Kinsella, J. E., Lokesh, B., & Stone, R. A. (1990). Dietary n-3 polyunsaturated fatty acids and amelioration of cardiovascular diseases: Possible mechanisms. *American Journal of Clinical Nutrition*, 52, 10–28.
- Kminkova, M., Winterova, R., & Kucera, J. (1993). Fatty acids in lipids of carp (*Cyprinus carpio*) tissues. *Czech Journal of Food Science*, 19, 177–181.
- Lorenzoni, M., Corboli, M., Ghetti, L., Pedicillo, G., & Carosi, A. (2007). Growth and

- reproduction of the goldfish *Carassius auratus*: A case study from Italy. In F. Gherardi (Ed.), *Biological invaders in inland waters: Profiles, distribution, and threats* (pp. 259). Milano, Italy: Springer.
- Lushchak, V. I., Bagnyukova, T. V., Husak, V. V., Luzhna, L. I., Lushchak, O. V., & Storey, K. B. (2005). Hyperoxia results in transient oxidative stress and an adaptive response by antioxidant enzymes in goldfish tissues. *International Journal of Biochemistry and Cellular Biology*, 37, 1670–1680.
- Lushchak, V. I., & Bagnyukova, T. V. (2006). Temperature increase results in oxidative stress in goldfish tissues. 1. Indices of oxidative stress. *Comparative Biochemistry and Physiology Part C*, 143, 30–35.
- Miarelli, M., Lorenzoni, M., & Mantilacci, L. (1990). Il lago Trasimeno. *Rivista di Idrobiologia*, 29, 353–360.
- Mendez, E., & Gonzales, M. E. (1997). Seasonal changes in the chemical and lipid composition of fillets of the Southwest Atlantic hake (*Merluccius hubbsi*). *Food Chemistry*, 59, 213–217.
- Ninno, R. E., De Torrenzo, M. A. P., Costuma, J. C., & Brenner, R. R. (1974). Specificity of 5- and 6- fatty acid desaturases in rat and fish. *Biochemical and Biophysical Acta*, 360, 124–132.
- Poli, B. M., Parisi, G., Dell'Agello, M., Guidotti, P., Bonelli, A., & Geri, G. (1994). Caratteristiche chimiche e qualità dei lipidi di pesci di diversa origine (tipo genetico, allevamento e cattura, acque naturali o riscaldate). In: *Proc. International Congress "Parliamo di acquacoltura", Fossano, CN, Italy 13–14 October, 1994*, pp. 171–180.
- PRAL, Programma Triennale di Ricerca Agricola. Agroambientale ed Agroindustriale della Regione Lazio (1999–2001). Available from: <http://www.regione.lazio.it/binary/agriweb/agriweballegatishedeinformativo/OpuscoloPRALiprodottiitticidelaghidelLaziopesca.1190976625.pdf>.
- Santos-Silva, J., Bessa, R. J. B., & Santos-Silva, F. (2002). Effect of genotype, feeding system and slaughter weight on the quality of light lambs. II. Fatty acid composition of meat. *Livestock Production Science*, 77, 187–194.
- Sirri, F., Castellini, C., Roncarati, A., Franchini, A., & Meluzzi, A. (2010). Effect of feeding and genotype on the lipid profile of organic chicken meat. *European Journal of Lipid Science and Technology*, 112, 994–1002.
- StataCorp. (2005). Stata statistical software: Release 9. Stata Corp. L.P., College Station, TX.
- Takeuchi, T. (1979). Digestibility of hydrogenated fish oils in carp and rainbow trout. *Bulletin Japanese Society of Science and Fishery*, 45, 1521–1525.
- Testi, S., Bonaldo, A., Gatta, P. P., & Badiani, A. (2006). Nutritional traits of dorsal and ventral fillets from three farmed fish species. *Food Chemistry*, 98, 104–111.
- Tocher, D. R., Fonseca-Madrugal, J., Dick, J. R., Wing, N., Bell, J. G., & Campbell, J. (2004). Effect of water temperature and diets containing palm oil on fatty acid desaturation and oxidation in hepatocytes and intestinal enterocytes of rainbow trout (*Onchorhynchus mykiss*). *Comparative Biochemistry and Physiology*, 137, 49–55.

- Turchetto, E., Pregolato, P., Barzanti, V., Maranesi, M., Mordenti, T., & Rogato F. (1994). Caratteristiche qualitative e pregi nutrizionali dei prodotti ittici di allevamento. Realtà e Prospettive dell'acquacoltura. La situazione italiana: Produzione e Qualità. Schede informative 3-4/94, Istituto Scotti Bassani per la ricerca e l'informazione scientifica e nutrizionale, Milano, pp. 31-39.
- Uauy, R., & Valenzuela, A. (1992). Marine oils as a source of omega-3 fatty acids in the diet: How to optimize the health benefits. *Progress in Food Nutrition Science*, 16, 199-243.
- Ulbricht, T. L., & Southgate, D. A. T. (1991). Coronary heart disease: Seven dietary factors. *The Lancet*, 338, 985-992.
- Vornanen, M., Tiitu, V., Kkel, R., & Aho, E. (1999). Effects of thermal acclimation on the relaxation system of crucian carp white myotomal muscle. *Journal of Experimental Zoology*, 284, 241-251.
- Vujkovic, G., Karlovic, D., Vujkovic, I., Vörösbaranyic, I., & Jovanovic, B. (1999). Composition of muscle tissue lipids of silver carp and bighead carp. *JAOCS*, 76, 475-479.

Table 1

Chemical composition of goldfish fillets in different seasons.

		Season				SEM
		Autumn	Winter	Spring	Summer	
Moisture	%	77.54 ^a	78.96 ^b	78.92 ^b	78.17 ^{a,b}	1.25
Protein	%	18.7 ^b	17.5 ^a	17.4 ^a	17.9 ^a	0.69
Lipids	%	2.01 ^b	1.59 ^a	1.70 ^a	1.89 ^b	0.10
Ash	%	1.80	1.95	2.01	2.03	0.48

N = 20 per season.

^{a,b} $P < 0.05$.

Table 2
Fatty acid profile of goldfish fillets in different seasons.

		Season				SEM
		Autumn	Winter	Spring	Summer	
C14:0	%	1.14 ^a	2.58 ^b	2.17 ^b	3.17 ^c	0.25
C16:0	%	22.1 ^b	21.8 ^b	15.2 ^a	17.1 ^a	2.47
C18:0	%	7.19 ^b	4.80 ^a	6.74 ^{a,b}	7.70 ^b	0.98
Others	%	2.81 ^a	3.49 ^{a,b}	4.60 ^c	3.61 ^b	1.01
SFA	%	33.2 ^b	32.6 ^b	29.7 ^a	33.6 ^b	2.08
C14:1n-6	%	0.11	0.32	0.13	0.11	0.22
C16:1n-7	%	2.54 ^a	4.86 ^b	3.56 ^{a,b}	4.47 ^b	1.47
C18:1n-9	%	9.59	11.0	11.3	10.4	2.56
Others	%	4.14	4.76	3.38	3.82	1.45
MUFA	%	16.4 ^a	20.9 ^b	18.3 ^{a,b}	18.8 ^{a,b}	3.01
C18:2n-6 <i>LA</i>	%	13.8 ^{a,b}	17.1 ^c	15.6 ^b	12.5 ^a	1.43
C20:2n-6	%	0.20	0.19	0.18	0.26	0.35
C20:3n-6	%	0.12	0.28	0.20	0.46	0.35
C20:4n-6	%	8.05 ^b	6.78 ^b	5.91 ^a	5.60 ^a	2.41
C18:3n-3 <i>LNA</i>	%	3.13 ^a	2.61 ^a	4.67 ^b	5.31 ^b	0.99
C20:3n-3	%	0.27	0.42	0.21	0.23	0.26
C20:5n-3 <i>EPA</i>	%	6.81 ^{a,b}	5.15 ^a	6.54 ^{a,b}	8.09 ^b	1.11
C21:5n-3	%	0.48 ^a	1.11 ^b	0.10 ^a	0.11 ^a	0.08
C22:5n-3	%	2.41 ^{a,b}	1.84 ^a	3.44 ^b	2.41 ^{a,b}	0.47
C22:6n-3 <i>DHA</i>	%	14.6 ^b	10.6 ^a	13.3 ^b	14.5 ^b	2.08
Others	%	0.61	0.34	0.80	0.14	0.64
PUFA	%	50.4 ^b	46.5 ^a	51.9 ^c	50.0 ^b	4.52
Δ^5 plus Δ^6 *		65.5 ^c	55.5 ^a	58.1 ^b	63.4 ^c	2.15

N = 20 per season.

^{a-c} *P* < 0.05.

* Desaturase index.

Table 3

Quantitative fatty acid contents, nutritional indices and oxidative status of goldfish filets in different seasons.

		Season				SEM
		Autumn	Winter	Spring	Summer	
SFA	mg/100 g	668 ^b	519 ^a	506 ^a	635 ^b	50.4
MUFA	mg/100 g	329	333	312	357	51.1
\sum n-3	mg/100 g	543 ^{b,c}	345 ^a	498 ^b	578 ^c	42.4
\sum n-6	mg/100 g	443	391	371	353	82.4
EPA	mg/100 g	137 ^b	81.9 ^a	111 ^a	153 ^b	14.5
DHA	mg/100 g	293 ^c	169 ^a	226 ^b	273 ^c	29.4
PUFA	mg/100 g	1014 ^c	739 ^a	883 ^b	937 ^b	65.3
n-6/n-3		0.81 ^a	1.13 ^b	0.74 ^a	0.68 ^a	0.22
TBARs	mgMDA/kg	0.20 ^b	0.11 ^a	0.18 ^b	0.27 ^c	0.11
Carbonyls	nmol/mg proteins	3.13 ^b	1.43 ^a	4.22 ^b	6.23 ^c	1.21
Thiols	μ g/g	15.9 ^a	16.7 ^a	15.8 ^a	18.4 ^b	2.45
Peroxidability index		228 ^b	185 ^a	220 ^b	227 ^b	15.9
Atherogenicity index		0.41 ^{a,b}	0.48 ^b	0.35 ^a	0.44 ^b	0.10
Thrombogenicity index		0.30 ^b	0.32 ^b	0.22 ^a	0.24 ^a	0.08
HH*		1.78 ^a	2.19 ^a	3.35 ^b	2.78 ^{a,b}	0.49

N = 20 per season.

^{a-c} $P < 0.05$

* Hypocholesterolaemic/hypercholesterolaemic ratio.