



Faba bean (*Vicia faba* var. *minor*) as a protein source for organic chickens: performance and carcass characteristics

Francesca Perella, Cecilia Mugnai, Alessandro Dal Bosco, Federico Sirri, Elisa Cestola & Cesare Castellini

To cite this article: Francesca Perella, Cecilia Mugnai, Alessandro Dal Bosco, Federico Sirri, Elisa Cestola & Cesare Castellini (2009) Faba bean (*Vicia faba* var. *minor*) as a protein source for organic chickens: performance and carcass characteristics, Italian Journal of Animal Science, 8:4, 575-584, DOI: [10.4081/ijas.2009.575](https://doi.org/10.4081/ijas.2009.575)

To link to this article: <http://dx.doi.org/10.4081/ijas.2009.575>



Copyright 2009 Taylor & Francis Group LLC



Published online: 01 Mar 2016.



Submit your article to this journal [↗](#)



Article views: 30



View related articles [↗](#)



Citing articles: 1 View citing articles [↗](#)



Faba bean (*Vicia faba* var. *minor*) as a protein source for organic chickens: performance and carcass characteristics

Francesca Perella¹, Cecilia Mugnai¹, Alessandro Dal Bosco¹,
Federico Sirri², Elisa Cestola¹, Cesare Castellini¹

¹Dipartimento di Biologia Applicata. Università di Perugia, Italy

²Dipartimento di Scienze degli Alimenti. Università di Bologna, Italy

Corresponding author: Dr. Alessandro Dal Bosco. Dipartimento di Biologia Applicata. Università di Perugia. Via Borgo XX Giugno 74, 06100 Perugia, Italy - Tel. +39 075 5857110 - Fax: +39 075 5857122 - Email: dalbosco@unipg.it

Received March 11, 2009; accepted April 22, 2009

ABSTRACT

The ban of transgenic feed and synthetic amino acids in organic chicken diets has made necessary to find protein-rich feedstuffs which do not alter productive performance. The effect of the faba bean (*Vicia faba* var. *minor*) as a partial substitute for soybean on the performance and carcass traits of slow-growing chickens of both sexes reared under the organic method was assessed. The experiment was carried out in the spring and autumn; one thousand birds per season were split into 8 groups (4 groups of males and 4 groups of females, each duplicated) and fed a double phase diet: a common starter diet for 20 days and two different growing-finisher diets containing either soybean (24%; S) or faba bean (16%; F) till the end of the rearing period (120 d). Average values of final live weight, feed intake and daily weight gain were significantly affected by sex, being higher in males than in females; the diets affected performance only in the 21-60 days of age period when the F group grew less and had poorer feed efficiency compared to the S group. Successively, the compensatory growth eliminated differences in slaughter weight. Regarding carcass traits, the diet affected only the ready-to-cook carcass percentage and the carcass length, both of which were higher in S birds. Faba beans could be a valuable protein source in the diet of organic chickens when used after the critical starter period.

Key words: Faba bean, Organic system, Performance, Carcass.

RIASSUNTO

UTILIZZO DEL FAVINO (*VICIA FABA* VAR. *MINOR*) COME FONTE PROTEICA PER POLLI
BIOLOGICI: PRESTAZIONI PRODUTTIVE E CARATTERISTICHE DELLA CARCASSA

Considerato il divieto di utilizzo di alimenti transgenici e di aminoacidi sintetici nell'avicoltura biologica lo scopo della presente ricerca è stato quello di studiare l'effetto della parziale sostituzione della soia con

favino (Vicia faba var. minor) sulle prestazioni produttive e sulle caratteristiche della carcassa di polli biologici. L'esperimento è stato ripetuto in due stagioni: Primavera e Autunno. Mille polli per stagione sono stati suddivisi in 8 gruppi (4 gruppi di maschi e 4 di femmine, ognuno replicato) ed alimentati con una dieta bi-fase: dieta starter per 20 giorni e due differenti diete di finissaggio contenenti soia (24%; S) o favino (16%; F) fino alla fine del periodo di allevamento (120 giorni). I pesi finali, il consumo di alimento e gli accrescimenti medi giornalieri sono stati influenzati solo dal sesso, mentre la dieta ha prodotto effetti significativi solo nel periodo 21-60 giorni, quando gli animali alimentati con la dieta F sono cresciuti meno ed hanno mostrato peggiori indici di conversione. Alla macellazione i pesi dei gruppi erano gli stessi per effetto di una crescita compensativa. Riguardo al caratteristiche della carcassa, solo la resa in busto e la lunghezza della carena sono risultate superiori negli animali del gruppo S. In conclusione si può affermare che il favino può essere considerato una valida fonte proteica per l'alimentazione di polli biologici a patto che venga utilizzato dopo il primo periodo critico di accrescimento.

Parole chiave: Favino, Polli biologici, Prestazioni produttive, Carcassa.

Introduction

The organic livestock system is governed by EU-Regulation 1804/1999 which provides specifications for housing conditions, rearing and animal care, disease prevention, veterinary treatment and nutrition. Since the synthetic forms of amino acids and GMO ingredients are banned from organic diets, the possible alternatives for poultry producers include the use of slow-growing genotypes which have lower protein and amino acids requirements (especially during the starter phase) (Leclercq *et al.*, 1993; Pesti *et al.*, 1994; Rosa *et al.*, 2001) and feed formulations utilizing lesser amount of soybean which has a high risk of GMO contamination (Hanson *et al.*, 2004). Slow-growing birds are recommended for better adaptation to the outdoor environment and to a longer rearing period (Castellini *et al.*, 2002; Fanatico *et al.*, 2005); moreover they are healthier birds with fewer heart, leg and metabolic problems (Julian, 1998).

One of the most studied legumes as an alternative to soybean is the faba bean (*Vicia faba var. minor*), which has received considerable interest as an indigenous source of protein for Mediterranean countries since it can grow easily in dry areas. Unfortunately, there have been reports (Sterling *et al.*, 2002; Attia *et al.*, 2005) of low growth rates

and food intake of broilers fed diets containing high levels of faba bean. The presence of toxic constituents and the low sulphur amino acid content of faba bean proteins have been claimed to be the main reasons for its low nutritional value (Rubio *et al.*, 1990). Van Leeuwen *et al.* (2000) showed that the faba bean has the lowest digestibility of the essential amino acids (mainly methionine) among several plant sources (maize, wheat, lupin, soybean), rendering the total substitution of soybean with faba bean not feasible.

However, heat treatment can improve the nutritional quality of faba beans by removing or destroying other thermo labile anti-nutritive factors responsible for reduced chick weight, feed efficiency and retention of dry matter, protein and crude fibre (Marquardt *et al.*, 1976; Ward *et al.*, 1977; Diaz *et al.*, 2006).

The aim of this study was to verify the effects of the faba bean (*Vicia faba var. minor*) as a partial substitute for soybean in growing-finisher diets on the performance of slow-growing chickens reared under the organic system.

Material and methods

Animals and experimental design

The experiment was carried out at the Agricultural farm of the University of Peru-

gia. The effect of the partial substitution of soybean with faba bean was tested in poultry diets when administered after a common starter diet. The trials were repeated in two different seasons (autumn and spring).

All the animals were reared according to EU Regulation 1804/99 and Italian directives (Italian Republic, 1992) on animal welfare for experimental and other scientific purposes.

For each season, one thousand day-old slow-growing chicks (Gaina, Avicola Berlanda, Padua, Italy) were divided into 8 experimental groups of 125 birds each (4 groups of males and 4 groups of females, each duplicated). They were fed the same starter soybean-based diet from 1 to 20 days of life. From day 21 until slaughter (120 d), they were fed 2 different growing-finisher diets: soybean based growing diet (S) or partially replaced with extruded faba bean (F). Diets were formulated according to NRC (1994) recommendations for slow-growing chickens (Table 1). Feed and water were provided *ad libitum*.

From hatching to 20 days of age all birds were kept in an environmentally controlled poultry house with temperature and relative humidity ranging from 30 to 32°C and from 65 to 70%, respectively. At 21 days of age they were transferred to a straw bedded poultry house (0.10 m²/bird), equipped with feeders and drinkers and with free access to open air runs (4 m²/bird).

The chicks were vaccinated against Marek and New Castle disease and coccidiosis (Paracox®).

Individual body weight (20% of the animals in each group were sampled) was recorded every 10 days and daily weight gain (DWG) and feed efficiency (FE) were calculated accordingly. Feed intake (FI) and bird mortality was recorded daily. At 120 days, the birds in all the experimental groups were slaughtered.

Analytical determinations of diets

The analyses of the chemical composition of the diets were carried out according to AOAC methods (1995) and the metabolizable energy was estimated according to the method described by Carrè and Rozo (1990).

The extraction and purification of vicine and convicine of diets were carried out by the reference method HPLC adapted according to Quemener (1988). Faba bean seeds were ground in an IKA-Universal M20 mill for 3 min to a particle size of less than 0.5 mm. A 500 mg quantity of sample was weighed in a centrifuge tube and 15 mL of aqueous uridine solution (0.4 mg mL⁻¹) was added. The tube contents were homogenized for 2 min with a Polytron homogenizer equipped with 10S Model axis and centrifuged at 25000 g for 5 min. Portions of 10 mL of supernatant were transferred to another centrifuge tube and 60 µL of 1 N HCl was added in order to precipitate most of the proteins at their isoelectric pH (pHi~4.2). The mixture was shaken and centrifuged at 2,500 g for 5 min. An aliquot of 100 µL of supernatant was diluted six times with ultra-pure water and maintained at 4°C until glycoside HPLC analysis.

The HPLC system consisted of an HP series 1100, Rheodyne LooT injector, Supelcosil LC PAH (15 cm x 4.6 mm) Column. Ultra-pure water was eluent at flow rate of 1 mL min⁻¹; the volume of injection was 20 µL. Concentration of vicine and convicine was quantified by relative UV response factors to uridine 0.803 and 0.601, respectively, measured at wavelength of 273 nm.

Carcass dissection

At 120 days, a representative sample of 10 birds per sex and diet for both seasons (weight ranging ±10% of the mean weight of all the birds) were slaughtered 12 hours after feed withdrawal. Chickens were electrically stunned (110 V; 350 Hz) before killing.

Table 1. Formulation (%), chemical composition (% DM except dry matter), energetic value (MJ kg⁻¹), amino acids, vicine and convicine (%) of the starter and finisher diets.

	Starter	Growing/Finisher	
		S	F
Ingredients:			
Maize	31.00	46.00	42.00
Wheat	13.50	13.50	13.50
Extruded soybean flakes	36.00	24.00	13.50
Pea	7.00	7.00	7.00
Extruded Faba bean	-	-	16.00
Gluten feed	9.00	-	-
Wheat bran	-	6.00	4.00
Soybean oil	-	-	0.50
Salt	0.20	0.20	0.20
Calcium phosphate	1.10	1.10	1.10
Calcium carbonate	1.60	1.60	1.60
Minerals and vitamins*	0.60	0.60	0.60
Chemical composition:			
Dry matter	%	88.49	88.00
Crude protein		20.47	16.25
Ether extract		8.22	6.65
Crude fibre		4.54	4.20
Ash		6.18	5.65
Metabolizable Energy**		13.14	12.66
Lysine ***		1.15	0.87
Methionine + cystine ***		0.72	0.58
Threonine		0.80	0.62
Tryptophan		0.24	0.19
Antinutritional factors:			
Vicine	nd	nd	0.18
Convicine	nd	nd	0.09

S: soybean; F: faba bean.

*Amounts per kg: Vit. A 11,000 U; Vit. D₃ 2000 U; Vit. B₁ 2.5 mg; Vit. B₂ 4 mg; Vit. B₆ 1.25 mg; Vit. B₁₂ 0.01 mg; α -tocopheryl acetate 30 mg; Biotin 0.06 mg; Vit. K 2.5 mg; Niacin 15 mg; Folic acid 0.30 mg; Panthotenic acid 10 mg; Choline chloride 600 mg; Mn 60 mg; Fe 50 mg; Zn 15 mg; I 0.5 mg; Co 0.5 mg.

**Estimated by Carrè and Rozo (1990).

***Calculated values.

nd=not detected.

The carcasses were plucked, eviscerated (non-edible viscera: intestines, paraventriculus, gall bladder, spleen, oesophagus and full crop) and individually weighed before and after refrigeration for 24 hours at +4°C. The carcass weight losses occurring during refrigeration were calculated as the difference in weight before and after refrigeration.

Full and empty gastrointestinal tracts were also individually weighed and the weight of the *ingesta* was calculated by difference.

The head, neck, legs, edible viscera (heart, liver, gizzard) and fat (perivisceral, perineal and abdominal) were removed and weighed; the ready-to-cook carcasses (RCC) were then weighed. Biometric measurements were carried out following the procedure described by ASPA (1996). The breasts and drumsticks were also removed from the carcasses and weighed, and their yields were calculated as a percentage of the RCC.

Statistical analyses

Data were analysed with a 3-way linear model including the effects of dietary regimen, sex and season. The effect of season and the interactions were not significant and were thus omitted.

The significance of differences was evaluated by the t-test (SAS, 1999). Mortality was analysed with CHISQ proc. (Pearson's chi-square).

The effect of diet on feed intake, feed efficiency and daily weight gain were also tested in the single sub periods.

The growth curve of the birds was also evaluated with the Gompertz equation (Roush *et al.*, 2006).

Results and discussion

Formulation (%), chemical composition (% d.m.), energetic value (MJ kg⁻¹), amino acids, vicine and convicine of the diets (%

d.m.) are reported in Table 1.

The amounts of vicine and convicine in the F diet were 0.18 and 0.09% d.m., respectively. These values agree with those observed by Gutierrez *et al.* (2006) who determined their contents in different faba bean cultivars (ranging from 0.02 and 0.01% d.m. to 0.33 and 0.21% d.m. for vicine and convicine, respectively). According to Sixdenier *et al.* (1996), the total amount of vicine and convicine should be considered high when the values are greater than 0.15% d.m., while values lower than 0.05% d.m. are classified as low; in our study the amount was 0.28% and should thus be considered high. These high levels in the diet showed that extrusion was not able to reduce the glucoside content. Although heat treatment inactivates part of the antinutritional factors (Ward *et al.*, 1977; Diaz *et al.*, 2006) contained in the seeds, vicine and convicine are thermostable (El-Sayed *et al.*, 1996; Elkowicz and Sosulski, 1982).

The effects of diet and sex on productive performance are reported in Table 2. The average values of final live weight, feed intake and daily weight gain were significantly affected only by sex, being higher in males than in females; diet did not influence performance traits.

A similar trend was observed in the productive performance in the different sub periods of the rearing cycle (Table 3), except from 21 to 60 days when the F group grew less and had a poorer feed efficiency compared to the S group. Afterwards, these differences between groups disappeared. Trends of DWG, FE and FI are ascribable to a compensatory growth phenomenon. It has been hypothesized that birds have a specific set point for body weight at a certain age and when they are behind their scheduled growth curve, they try to reach this weight in a shorter time (Wilson and Osbourn, 1960; Mosier, 1986). When the early nutri-

Table 2. Productive performance of organic broilers fed with different growing-finisher diets.

		Diet		Sex		Significance		Pooled SE / χ^2
		S	F	M	F	Diet	Sex	
Final live weight (LW)	g	2,583	2,458	2,767	2,274	ns	**	264
Feed intake	g/d	80.0	75.6	85.6	70.0	ns	**	11.0
Daily weight gain	g/d	21.3	20.1	22.8	18.6	ns	*	2.4
Feed efficiency		3.76	3.76	3.75	3.76	ns	ns	2.2
Mortality	%	9.6	10.4	11.0	9.0	ns	ns	3.1

S: soybean; F: faba bean.

n=400; *= $P < 0.05$; **= $P < 0.001$; ns: not significant.


Table 3. Influence of protein sources on the feed intake, daily weight gain and feed efficiency of the chickens.

		Diet		Pooled SE
		S	F	
Feed intake:				
1-20 d	g/d	12.5	12.2	2.0
21-40 d	"	48.9	45	3.2
41-60 d	"	85.4	79.8	7.3
61-80 d	"	108.9	104	14.1
81-100 d	"	115.4	110.5	8.7
101-120 d	"	108.9	102.3	10.2
Daily weight gain:				
1-20 d	g	8.3	8.7	1.8
21-40 d	"	22.1 ^b	17.9 ^a	2.5
41-60 d	"	29.1 ^b	24.5 ^a	1.8
61-80 d	"	26.2	25.9	4.0
81-100 d	"	23.4	24.0	3.2
101-120 d	"	18.9	19.8	2.3
Feed efficiency:				
1-20 d		1.5	1.4	0.2
21-40 d		2.2 ^a	2.6 ^b	0.6
41-60 d		2.9 ^a	3.3 ^b	0.6
61-80 d		4.2	4.0	1.1
81-100 d		4.9	4.6	1.2
101-120 d		5.7	5.4	1.4

S: soybean; F: faba bean.

n=400; a,b= $P < 0.05$.

Figure 1. Growth curve of control and faba bean fed chickens.



S: soybean diet; F: faba bean diet.

Table 4. Carcass characteristics of organic broilers fed with different growing-finisher diets.

		Diet		Sex		Significance		Pooled SE
		S	F	M	F	Diet	Sex	
Eviscerated carcass (EC)	g	2097	2018	2225	1890	ns	**	161
Dressing out	EC/LW	81.2	82.1	80.4	83.1	ns	**	2.8
Refrigeration losses	%	0.13	0.12	0.13	0.12	ns	ns	0.02
Gastro-intestinal tract	"	11.52	11.58	11.60	11.50	ns	ns	1.81
Gastro-intestinal content	"	5.84	5.85	5.90	5.79	ns	ns	0.15
Abdominal fat	"	1.00	1.10	1.00	1.12	ns	ns	0.10
Ready-to-cook carcass (RCC)	g	1589	1531	1715	1405	*	**	150
Dressing out	RCC/LW	0.62	0.62	0.62	0.62	ns	ns	1.81
Breast yield	% RCC	16.00	16.20	15.40	16.80	ns	*	2.15
Thigh yield	"	17.20	17.20	17.70	16.70	ns	*	0.31
Biometric measurements:								
Tibia length	cm	14.10	14.30	14.50	13.90	ns	*	0.41
Sternum	"	13.80	13.80	14.20	13.40	ns	*	1.11
Carena	"	11.18	10.60	11.20	10.56	*	*	0.36
Rostrum	"	2.77	2.75	2.92	2.60	ns	*	0.22
Breast width	"	5.90	5.90	6.00	5.80	ns	ns	0.84
Breast layer thickness	"	2.20	2.20	2.30	2.10	ns	ns	0.34
Meat-to-bone ratio	"	2.94	2.96	2.85	3.05	ns	ns	0.49

S: soybean; F: faba bean.

n=160; * = P < 0.05; ** = P < 0.001; ns: not significant.

tional requirements of the birds were met by a soybean-based starter diet, the successive compensatory growth, occurring after 81 days of age, was able to reach similar slaughter weights (Figure 1).

Moschini *et al.* (2005) showed that 479 g/kg of raw faba bean did not have any negative effects on DWG and FE during the growing period (1-21) of conventionally reared Ross chickens. It is important to emphasize that in their trial supplementation of synthetic amino acids was carried out, while in our experiment synthetic amino acids were absent since they are not allowed by organic rules.

Regarding carcass traits (Table 4), the diet affected only the RCC percentage and the carena length, both of which were higher in S birds. Sex showed a strong effect on most of the carcass traits, including the eviscerated carcass, dressing out, ready-to-cook carcass, breast and thigh yields as well as the biometric measurements, due to the pronounced sexual dimorphism of slow-growing birds (Fanatico *et al.*, 2005). In chickens fed 500 g/kg of raw or extruded faba bean (Diaz

et al., 2006) observed a higher breast meat percent yield with respect to the standard diet independent of processing.

Conclusions

Data obtained in this experiment prove the effectiveness of the faba bean (*Vicia faba* var. *minor*) in poultry diets as a protein source as partial substitute for soybean. Chickens fed a growing-finisher diet containing faba beans as a partial substitute for soybean, after a transition period of lower DWG, showed compensatory growth in the later period, reaching a final (120 d) live weight similar to the S group and reducing the diet-dependent differences in the carcass traits. Thus the F diet without synthetic amino acids, if administered after a starter period, is able to meet the requirement of slow-growing birds.

This study was supported by the E.Q.U.I.ZOO. BIO project (Efficiency, Quality and Innovation in Organic Livestock).

REFERENCES

- AOAC, 1995. Official Methods of Analysis. 15th ed. Association of Official Analytical Chemists, Washington, DC, USA.
- Attia, Y.A., Hassan, R.A., Shehatta, M.H., Abd El-Hady, S.B., 2005. Growth, carcass quality and serum constituents of slow growing chicks as affected by betaine addition to diets containing 2 different levels of methionine. *Int. J. Poultry Sci.* 4:856-865.
- Carré, B., Rozo, E., 1990. La prédiction de la valeur énergétique des matières premières destinées à l'aviiculture. *Prod. Anim.* 3:163-169.
- Castellini, C., Dal Bosco, A., Mugnai, C., Bernardini, M., 2002. Performance and behaviour of chickens with different growing rate reared according to organic system. *Ital. J. Anim. Sci.* 1:291-300.
- Diaz, D., Morlacchini, M., Masoero, F., Moschini M., Fusconi, G., Piva, G., 2006. Pea seeds (*Pisum sativum*), Faba bean (*Vicia Faba* var. *minor*) and lupin seeds (*Lupinus albus* var. *multitalia*) as protein sources in broiler diet: effect of extrusion on growth performance. *Ital. J. Anim. Sci.* 5:43-53.
- Elkowicz, K., Sosulski, F.W., 1982. Antinutritive factors in eleven legumes and their air-classified protein and starch fractions. *J. Food Sci.* 47:1301-1304.
- El-Sayed, M.A., Sosulski, F.W., Shehata, A.A.Y., Youssef, M.M., 1996. Nutritional, functional and sensory properties of wheat, rice and faba bean

- blends texturized by drum drying. *Int. J. Food Sci. Tech.* 31:257-266.
- European Commission, 1999. Council Regulation n.1804/99 of July 1999 supplementing Regulation (ECC) n. 2092/91 on organic production of agricultural products. In: Official Journal, L 222, 24/08/1999, pp 1-28.
- Fanatico, A.C., Pillai, P.B., Cavitt, L.C., Owens, C.M., Emmert, J.L., 2005. Evaluation of slower-growing broiler genotypes grown with and without outdoor access: growth performance and carcass yields. *Poultry Sci.* 84:1321-1327.
- Gutierrez, N., Avila, C.M., Duc, G., Marget, P., Suso, M.J., Moreno, M.T., Torres, A.M., 2006. CAPs markers to assist selection for low vicine and convicine contents in faba bean (*Vicia faba* L.). *Theor. Appl. Genet.* 114:59-66.
- Hanson, J., Dismukes, R., Chambers, W., Greene, C., Kremen, A., 2004. Risk and risk management in organic agriculture: views of organic farmers. *Renew. Agr. Food Syst.* 19:218-227.
- Italian Republic, 1992. Legislative Decree no. 116 of 27 January 1992 implementing the European Directive 86/609/EEC on the protection of animals used for experimental and other scientific purposes. In: Official Gazette n. 40, 18/02/1992, Ordinary Suppl., pp 1-12.
- Julian, R.J., 1998. Rapid growth problems: ascites and skeletal deformities in broilers. *Poultry Sci.* 77:1773-1780.
- Leclercq, B., Chagneau, A.M., Cochard, T., Hamzaoui, S., Larbier, M., 1993. Comparative utilisation of sulphur-containing amino acids by genetically lean or fat chickens. *Brit. Poultry Sci.* 34:383-391.
- Marquardt, R.R., Campbell, L.D., Ward, T., 1976. Studies with chicks on the growth depressing factor(s) in faba bean (*Vicia faba* var. *minor*). *J. Nutr.* 106:275-284.
- Moschini, M., Masoero, F., Prandini A., Fusconi, G., Morlacchini, M., Piva, G., 2005. Raw Pea (*Pisum sativum*), raw Faba bean (*Vicia faba* var. *minor*) and raw Lupin (*Lupinus albus* var. *multitalia*) as alternative protein sources in broiler diets. *Ital. J. Anim. Sci.* 4:59-69.
- Mosier, H.D., 1986. The control of catch-up growth. *Acta Endocrinol-Cop.* 113:1-7.
- National Research Council, 1994. Nutrient requirements of poultry. 9th rev. ed. National Academy Press, Washington, DC, USA.
- Pesti, G.M., Leclercq, B., Chagneau, A.M., Cochard, T., 1994. Comparative response of genetically lean or fat chicken to lysine, arginine and non essential amino acids supply. II. Plasma amino acids responses. *Brit. Poultry Sci.* 35:697-707.
- Quemener, B., 1988. Improvements in the high-pressure liquid chromatographic determination of amino acids sugar and α -galactosides in faba bean, lupin and pea. *J. Agr. Food Chem.* 36:754-759.
- Romboli, I., Cavalchini, L., Gualtieri, M., Franchini, A., Nizza, A., Quarantelli, A., 1996. Metodologie relative alla macellazione del pollame, alla valutazione e dissezione delle carcasse e delle carni avicole. *Zoot. Nutr. Anim.* 22:177-180.
- Rosa, A.P., Pesti, G.M., Edwards, H.M., Jr., Bakalli, R.I., 2001. Threonine requirements of different broiler genotypes. *Poultry Sci.* 80:1710-1717.
- Roush, W.B., Dozier III, W.A., Branton, S.L., 2006. Comparison of Gompertz and neural network models of broiler chickens. *Poultry Sci.* 85:794-797.
- Rubio, L.A., Brenes, A., Castaño, M., 1990. The utilization of raw and autoclaved faba bean (*Vicia faba* L., var. *minor*) and faba bean fractions in diets for growing broiler chickens. *Brit. J. Nutr.* 63:419-430.
- SAS, 1999. User's Guide Statistics. Version 8.0. SAS Institute, Inc., Cary, NC, USA.
- Sixdenier, G., Cassecuelle, F., Guillaumin, L., Duc, G., 1996. Rapid spectrophotometric method for reduction of vicine and convicine in faba bean seed. *FABIS Newslett.* 38/39:42-44.
- Sterling, K.G., Costa, E.F., Henry, M.H., Pesti, G.M., Bakalli, R.I., 2002. Responses of broiler chickens to cottonseed- and soybean meal-based diets at several protein levels. *Poultry Sci.* 81:217-226.
- Van Leeuwen, P., Babinszky, L., Verstegen, M.W.A., Tossenberger, J., 2000. A procedure for ileostomisation of adult roosters to determine apparent

- ileal digestibility of protein and amino acids of diets: Comparison of six diets in roosters and growing pigs. *Livest. Prod. Sci.* 67:101-111.
- Ward, A.T., Marquardt, R.R., Campbell, L.D., 1977. Further studies on the isolation of the thermolabile growth inhibitor from the faba bean (*Vicia faba L. var. minor*). *J. Nutr.* 107:1325-1334.
- Wilson, P.K., Osbourn, D.F., 1960. Compensatory growth after undernutrition in mammals and birds. *Biol. Rev.* 35:325-363.