

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Resource driven community dynamics of NH₄⁺ assimilating and N₂O reducing archaea in a temperate paddy soil

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1633796> since 2017-08-09T18:05:56Z

Published version:

DOI:10.1016/j.pedobi.2017.02.001

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in PEDOBIOLOGIA, 62, 2017, 10.1016/j.pedobi.2017.02.001.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.pedobi.2017.02.001

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0031405616300762>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1633796>

1 **Type of article:** original article

2

3 **Resource driven community dynamics of NH₄⁺ assimilating and N₂O reducing archaea in a**
4 **temperate paddy soil**

5

6 **Maria Alexandra Cucu^{1,2,3*}**, Sven Marhan², Daniel Said-Pullicino¹, Luisella Celi¹, Ellen Kandeler²
7 and Frank Rasche³

8

9 ¹Rice Agro-ecosystem and Environmental Research Group, Department of Agricultural, Forest and
10 Food Sciences, University of Turin, Italy

11 ²Institute of Soil Science and Land Evaluation, Soil Biology Section, University of Hohenheim,
12 Stuttgart, Germany

13 ³Institute of Plant Production and Agroecology in the Tropics and Subtropics, University of
14 Hohenheim, Stuttgart, Germany

15

16 **Running title:** Archaeal N cycling in temperate paddy soils

17

18 ***Corresponding author:**

19 Dr. Maria Alexandra Cucu

20 Phone: +39 011 670 8544

21 Fax: +39 011 670 8692

22 Email: mariaalexandra.cucu@unito.it

23

24

25

26

27 **Abstract.**

28 In fertilized paddy soils, the role of resource availability on ammonium (NH_4^+) assimilation and
29 immobilization by archaea requires advanced understanding as this may have considerable
30 implications on subsequent catalytic steps in the soil N cycle including archaeal nitrous oxide (N_2O)
31 reduction. To gain a deeper understanding about these process links, we incubated a temperate
32 paddy soil under submerged conditions with or without straw and fertilized with either ^{15}N -enriched
33 (99 atom% ^{15}N) or non-enriched $(\text{NH}_4)_2\text{SO}_4$. Notably, a variation in community structure and a
34 higher abundance of archaeal N_2O reductase (*arc-nosZ*) genes in the no straw treatment than in the
35 straw one was observed. This was attributed to NH_4^+ assimilation by N_2O reducing archaea as was
36 further corroborated by a considerable ^{15}N -enrichment of archaeal glutamate dehydrogenase (*gdhA*)
37 genes. Moreover, indications were found that denitrifying archaea controlled their chemotrophic
38 and heterotrophic metabolisms in response to different availabilities of inorganic and organic N and
39 C resources. However, in the presence of straw, bacterial *nosZ* genes may have also contributed to
40 the completion of denitrification. Our results suggested that N assimilation contributed to the last
41 step of archaeal denitrification. Therefore, archaea may play a key role in regulating major N fluxes
42 in fertilized paddy soils, especially in the absence of rice straw.

43

44 **Keywords:** *arc-nosZ* gene; total prokaryotic *nosZ* clade I gene; *gdhA* gene; ^{15}N -DNA-based stable
45 isotope probing.

46 **1. Introduction**

47 Fertilized rice paddy soils generally exhibit lower nitrogen (N) recovery rates (30-40%) with
48 respect to upland soils (Cassman et al., 2002). Incorporation of crop residues such as rice straw
49 generally leads to N immobilization (assimilation to sustain their own metabolism) which reduces
50 fertilizer use efficiency to a large extent (Reddy 1982; Eagle et al., 2000; Bird et al., 2001). In
51 previous studies we observed that addition of rice straw induced, in comparison to no straw
52 addition, an 80% increase of immobilization of applied ammonium (NH_4^+) along with a 65%
53 decrease in gaseous N losses (Cucu et al., 2014; Said-Pullicino et al., 2014). On the other hand,
54 incorporation of rice straw in paddies was shown to stimulate N gaseous losses (i.e., N oxide gases
55 and N_2) from soil-plant systems since up to 70% of applied N may be lost through coupled
56 nitrification-denitrification (Cassman et al., 1998; Ghosh and Bhat, 1998; Majumdar, 2013).
57 Significant amounts of nitrous oxide (N_2O), an important greenhouse gas, have been shown to be
58 emitted under aerobic conditions (Zou et al., 2007; Ishii et al., 2011a). To mitigate this climate
59 variation promoting process, it was acknowledged that in continuously submerged paddies complete
60 denitrification via microbial nitrous oxide reduction can act as a considerable sink for N_2O which
61 favors N_2 losses (Davidson et al., 1986; Conrad 1995, 1996; Ferrè et al., 2012).

62 Both N assimilation and denitrification are mediated by specific microbial communities which are
63 well adapted to the anaerobic conditions of paddy environments (Liesack et al., 2000; Kögel-
64 Knabner et al., 2010). These ecosystems are characterized by a dynamic microbial community in
65 which archaea may predominate over bacteria (Cabello et al., 2004, Rush, 2016), and may also
66 exhibit different metabolic pathways and resource utilization. Denitrification *sensu lato*, as the
67 pathway of nitrate (NO_3^-) respiration, can be associated with dissimilatory and assimilatory
68 branches of microbial metabolism (Zumft, 1997). Under anaerobic conditions, bacterial denitrifiers
69 typically adopt a dissimilatory metabolism for their energy gain (Zumft, 1992), while their potential
70 assimilatory NO_3^- reduction pathway is suppressed when NH_4^+ is the predominant inorganic N form
71 (Rice and Tiedje, 1989; Cabello et al., 2004).

72 Archaeal denitrifiers can simultaneously perform dissimilatory and assimilatory reactions (Rusch,
73 2013) by using N not only for energy but also electron sink and detoxification (Zumft, 1997). The
74 gained N is also utilized to build up nitrogenous compounds (e.g., DNA, amino sugars, proteins)
75 (Devêvre and Horwáth, 2001; Nannipieri and Paul, 2009), which ultimately contribute to the soil
76 organic N pool (Kögel-Knabner et al., 2010). However, assimilatory NO_3^- reduction is less frequent
77 than respiratory NO_3^- reduction in the archaeal domain (Martínez-Espinosa et al., 2001). Instead,
78 Cabello et al. (2004) highlighted the importance of the archaeal NH_4^+ assimilation pathway
79 catalyzed by glutamate dehydrogenase (GDH). Encoded by the *gdhA* gene, GDH generally shows a
80 high activity under non-limiting NH_4^+ soil conditions (Bonete et al., 2008). This may be the case for
81 fertilized rice agro-ecosystems characterized by high NH_4^+ availability as a result of mineral N
82 addition and mineralization of organic matter (OM) including freshly applied rice straw (Sahrawat,
83 2004; Cucu et al., 2014). It needs to be pointed out that NH_4^+ assimilation by archaea in paddy soils
84 requires explicit understanding as this may have considerable implications on subsequent catalytic
85 steps in the soil N cycle including the last step of denitrification (i.e., N_2O reduction). Although N
86 fertilization is recognized as the primary factor influencing denitrification under reduced conditions
87 (Garcia and Tiedje 1982; Nogales et al., 2002; Prieme et al., 2002), the role of rice straw and the
88 interactive effects of available N and C resources on archaeal NH_4^+ assimilation and N_2O reduction
89 remains uncertain.

90 Compared to bacterial denitrification (Philippot et al., 2002; Henry et al., 2006; Ishii et al., 2011b),
91 only limited knowledge exists about this process in archaea. Likewise, only a few archaeal
92 denitrification genes and related enzymes have been investigated so far (de Vries and Schroder,
93 2002; Cabello et al., 2004). Moreover, information on the reduction of N_2O to N_2 is only available
94 for very few hypertermophilic and halophilic archaeal species including *Haloferax denitrificans* and
95 *Pyrobaculum aerophilum* (Tomlinson et al., 1986; de Vries and Schroder, 2002). Jones et al. (2013)
96 assigned several archaeal species to *nosZ* groups I and II covering the bacterial and archaeal
97 domains. Recently, Rusch (2013) has introduced a complementary marker for the *arc-nosZ* gene

98 that targets the archaeal N₂O reducing community. The *arc-nosZ* gene is part of the total
99 prokaryotic *nosZ* gene clades I and II. Although a phylogenetic bias based on hypertermophilic and
100 halophilic archaeal sequences has to be taken into account due to limited database information,
101 Rusch (2013) considered the *arc-nosZ* gene marker suitable for the full range of applications
102 targeting archaeal denitrifiers. While the crucial role of archaeal species in methanogenesis and
103 methanotrophy in rice paddies (Fazli et al 2013) is well known, their involvement in denitrification
104 remain less understood.

105 The primary objective of this study was to provide advanced understanding about the role of
106 archaea involved in N assimilation coupled with the final step of denitrification (i.e., N₂O
107 reduction) as a function of C and N resource availability in a temperate paddy soil with no NH₄⁺
108 limitation. To achieve this goal, we tested the following hypotheses: i) archaeal N assimilation is
109 controlled by the availability of labile organic C substrates (i.e., rice straw addition); ii) a fraction of
110 the assimilated N is used for proliferation of archaeal community members harboring the *arc-nosZ*
111 gene; iii) in absence of rice straw, the relative contribution of archaea to denitrification is more
112 important than that of their bacterial counterparts.

113

114 **2. Materials and methods**

115 2.1. Incubation experiment

116 Soil was collected from the Ap horizon (0-15 cm) of a long-term field experiment in Vercelli,
117 Northwest Italy (45°17'47"N, 8°25'51"E) in February 2012 before rice straw incorporation. The
118 paddy soil, classified as a Haplic Gleysol (WRB, 2007), has been under continuous single-cropped
119 rice cultivation for the last 30 years. The soil had a pH around 6, a sandy loam texture (7% clay,
120 41% silt, 52% sand) and relatively low contents of organic C and total N (11.6 and 1.1 g kg⁻¹,
121 respectively). The amount of clay in conjunction with soil organic matter content accounted for the
122 low cation exchange capacity (CEC = 6.7 cmol (+) kg⁻¹). Further details on soil properties have been
123 previously reported in Cucu et al. (2014) and Said-Pullicino et al. (2014). After manual removal of

124 remaining vegetal residues, field moist soil was homogenized after passing through a 2 mm sieve.
125 Rice straw (*Oryza sativa* L. cv. Sirio CL) was sampled from the same experimental site in October
126 2011 after grain harvest, dried at 40°C and cut into 1 cm segments. Total C and N contents of the
127 rice straw were 400 and 6.6 g kg⁻¹ respectively, resulting in a C/N ratio of 61.

128 A laboratory incubation experiment was carried out adopting a setup similar to that described by
129 Cucu et al. (2014). The experimental design consisted of a completely randomized arrangement of
130 soil microcosms incubated under submerged conditions with or without the addition of rice straw
131 (application dose of 4.3 g straw kg⁻¹ soil, equivalent to 10 Mg dry weight ha⁻¹). After a pre-
132 incubation period with or without straw addition for 14 days at 50% soil water holding capacity to
133 reestablish the microbial equilibrium, inorganic N (application dose of 56 mg N kg⁻¹ soil, equivalent
134 to 130 kg N ha⁻¹, generally applied in the field) was added to the soil samples and immediately
135 submerged under 5 cm of deionized water. Inorganic N was added as ¹⁵N-labelled (NH₄)₂SO₄ (99
136 atom% ¹⁵N) to assess active N assimilatory prokaryotes by stable isotope probing (SIP, see below).
137 A second set of soil samples was treated similarly with non-labeled (NH₄)₂SO₄ as control for SIP
138 studies (Buckley et al., 2007b; Neufeld et al., 2007). The experiment was run with six replicates per
139 treatment and time point. Incubation under submerged conditions was carried out for 60 days at
140 25°C in the dark. After 1, 5, 10, 20, 30 and 60 days from fertilizer application, soil microcosms
141 were destructively sampled. The fresh soil sample was immediately used for DNA extraction as
142 well as for inorganic N determination (see below). Submergence water was decanted, total volume
143 recorded, filtered at 0.45 µm and used for inorganic N determination. Afterwards, both soil and
144 water samples have been frozen at -20 °C for further analysis.

145

146 2.2. Chemical analyses

147 N₂O emissions were monitored daily during the first 4 weeks of incubation. On each measurement
148 day, soil microcosm headspace was sampled twice: immediately and 30 min after closure of the
149 lids. Afterwards the gas sample was transferred into pre-evacuated exetainers (5.9 ml, Labco Ltd.,

150 Lampeter, UK), and analyzed for N₂O using a gas chromatograph equipped with an electron capture
151 detector (Agilent 7890, Santa Clara, CA, USA). Three standard gases with known concentrations
152 were used for gas flux calibration and calculation. N₂O fluxes were calculated by linear regression
153 as described by Hutchinson and Mossier (1981).

154 Soil pH was measured potentiometrically in H₂O. Submergence water samples as well as 0.5 M
155 K₂SO₄ soil extracts (1:4 w/v soil-to-extractant ratio) were analyzed for inorganic N (NH₄⁺, NO₃⁻)
156 (Auto-analyzer 3, Bran & Luebbe, Norderstedt, Germany), total dissolved nitrogen (TDN) and
157 dissolved organic carbon (DOC) after sample acidification (TOC/TN analyzer Multi NC 2100S,
158 Analytic Jena GmbH, Jena, Germany).

159

160 2.3. Molecular analyses

161 2.3.1. DNA extraction

162 At each time point, total DNA was extracted from each fresh individual replicate soil sample using
163 the FastDNA[®] Spin kit for soil (MP Biomedicals, Solon, Ohio, USA) according to the
164 manufacturer's instructions. DNA quantification was performed with a NanoDrop ND-2000
165 (NanoDrop Technologies, Wilmington, DE, USA).

166

167 2.3.2. Stable isotope probing

168 *Measurement of ¹⁵N-enrichment in DNA extracts*

169 To measure the ¹⁵N-enrichment of soil DNA extracts according to España et al. (2011), a tin capsule
170 (Hekatech, Wegberg, Germany) was filled with 10 mg sorbsil (May & Baker, Dagenham, United
171 Kingdom) plus 2 µg of the DNA extracts. Twenty µg N of dissolved unlabeled ammonium sulphate
172 (Roth, Karlsruhe, Germany) were then added to the capsule and dried at 50°C overnight. All DNA
173 samples were analyzed with an elemental analyzer (Euro EA 3000; Hekatech) coupled with an
174 isotope ratio mass spectrometer (DeltaPlus XP, Thermo Scientific, Waltham, USA). ¹⁵N-enrichment

175 of DNA extracts was subsequently calculated by taking into account the isotopic signature of the
176 ammonium sulphate spike.

177 ¹⁵N-DNA-based stable isotope probing (SIP) was carried out on samples incubated for 30 days,
178 corresponding to distinct alteration of microbial activity (i.e., ¹⁵N enrichment of soil DNA extracts)
179 between “straw” and “no straw” treatments. To reach a high concentration and volume for the
180 further centrifugation, two replicates per DNA samples were pooled together. Finally, four DNA
181 samples were selected for SIP: two with highest ¹⁵N label (straw: 6.83 atom% ¹⁵N, no straw: 5.02
182 atom% ¹⁵N) and two respective ¹⁴N controls. Centrifugation mixtures were prepared by adding 5
183 mg of DNA diluted in 1 mL gradient buffer (0.1 M Tris, 0.1 M KCl, 1 mM EDTA) to 4.9 ml of a
184 7.163 M CsCl (Calbiochem/Merck, Darmstadt, Germany) solution. Isopycnic fractionation was
185 performed in 5.1 mL polyallomer centrifuge tube (13×51mm) placed in a Vti 65.2 vertical rotor
186 (both Beckman Coulter, Krefeld, Germany). Tubes were then centrifuged in an Optima™ L-90K
187 ultracentrifuge (Beckman Coulter) was performed with 140,000 × *g* at 20°C for 69 h (España et al.,
188 2011). After centrifugation, each gradient was fractionated into 16 individual fractions (312 µl
189 each) using a syringe pump (NE-1000, New Era Pump Systems, New York, NY, USA). Buoyant
190 density was adjusted prior to centrifugation (AR200 digital refractometer, Reichert, New York,
191 USA) and calculated for each fraction according to Buckley et al. (2007b). SIP fractions were
192 precipitated with 20 µg of glycogen (Roche Diagnostics GmbH, Penzberg, Germany) and 1 ml of a
193 30% polyethylene glycol 6000 solution (Carl Roth GmbH, Karlsruhe, Germany), washed (70%
194 ethanol) and suspended in 30 µl of Tris-EDTA buffer (pH 8.0) (Neufeld et al., 2007).

195

196 2.3.3. Microbial abundance

197 Abundance of total bacterial and archaeal communities were done by quantifying the respective 16S
198 rRNA genes (Rasche et al., 2011). Description of primer sets and amplification details used for
199 quantitative PCR are specified in Table 1. Quantification of *nosZ* clade I was performed as
200 described by Henry et al. (2006). The quantification of *arc-nosZ* genes was performed using

201 primers published by Rusch, (2013). Primer accuracy was confirmed (Rusch, personal
202 communication) and preliminary tests (data not shown) were carried out to assess the specificity of
203 the primer set for the studied soil. To confirm the targeted archaeal NH_4^+ assimilation process and
204 the implied labeling success, primers targeting N assimilation in *Methanobrevibacter smithii* by
205 means of the GDH system encoded by the *gdhA* gene were used. *Methanobrevibacter* sp. have been
206 earlier found in paddy soils (Fetzer et al., 1993) and bioreactors (i.e., *Methanobrevibacter* related
207 species), here with potential denitrifying abilities (Chuang et al., 2014)

208 Abundance of bacterial and archaeal 16S rRNA genes and of the three functional genes (i.e., total
209 prokaryotic *nosZ* clade I, *arc-nosZ*, archaeal *gdhA*) was determined by quantitative PCR (qPCR)
210 using a StepOnePlus™ Real-Time PCR System (Applied Biosystems, Foster City, CA, USA) for
211 bulk soil DNA as well as for *gdhA* genes in SIP fractions (Table 1). For standard preparation,
212 amplicons from each target gene were generated, purified (Invisorb Fragment CleanUp, Stratec
213 Molecular GmbH, Berlin, Germany), ligated into the Strata-Clone PCR cloning vector pSC-A
214 (Strataclone PCR Cloning Kit, Agilent Technologies Inc.) and ligation products were transformed
215 into StrataClone SoloPack competent cells (Stratagene). Specificity of clones used as quantitative
216 PCR (qPCR) standards were checked via sequencing at LGC Genomics GmbH (Berlin, Germany)
217 and BLAST analysis. Plasmid DNA was isolated (GenElute™ Plasmid Miniprep Kit, Sigma-
218 Aldrich, St. Louis, MO, USA) from standard clones and quantified as described above.

219 As assessed in preliminary tests and qPCR reaction optimization, each 25 μl qPCR cocktail
220 contained 5 ng DNA (16S rRNA genes) or 10 ng DNA (total prokaryotic *nosZ* clade I and archaeal
221 *arc-nosZ* gene), 1x Power SYBR green master mix (Applied Biosystems), 0.15 μM of each primer
222 (Table 1), 0.25 μl of T4 gene 32 protein (500 $\mu\text{g ml}^{-1}$, MP Biomedicals). For *gdhA* gene, the 25 μl
223 qPCR cocktail contained 1x FastStart Universal SYBR Green Master (ROX) (Roche), 0.2 μM of
224 each primer (Table 1), 0.625 U Uracil-DNA Glycosylase (Roche) and the DNA template (10 ng
225 from extracted soil DNA, 10 μl DNA from SIP fractions). Each sample was quantified in triplicate
226 across plates, while standards in 10-fold serial dilutions were run in duplicate. The optimal dilution

227 of DNA extracts was tested to compensate any reaction inhibition by organic compounds (e.g.,
228 humic acids) co-extracted during DNA isolations (data not shown). Melting curves of amplicons
229 were generated to ensure that fluorescence signals originated from specific amplicons and not from
230 primer dimers or other artifacts. This was confirmed by checking the amplification products on 1%
231 agarose gel. For quality control, melting curve analyses were performed. Amplification efficiency
232 ranged from 80% (archaeal 16S rRNA, total prokaryotic *nosZ* clade I and *arc-nosZ* genes) to 103%
233 (bacterial 16S rRNA and *gdhA* genes). The slopes were between -3.890 and -3.235 and $R^2 \geq 0.98$.
234 Gene copy numbers were calculated with StepOne™ software version 2.2 (Applied Biosystems).
235 The data were normalized and presented in figures as copies g^{-1} dry soil.
236 To determine the relative abundance of *arc-nosZ* gene with respect to the total prokaryotic *nosZ*
237 gene clade I, the *arc-nosZ* to *nosZ* gene abundance ratio was calculated. Values > 0.5 indicate a
238 higher relative abundance of archaea with respect to their bacterial counterparts within total
239 prokaryotic *nosZ* clade I.

240

241 2.3.4. Microbial community structure

242 The community structure of archaeal 16S rRNA, total prokaryotic *nosZ* clade I, and archaeal *arc-*
243 *nosZ* genes in bulk DNA soil, as well as in SIP fractions (total prokaryotic *nosZ* clade I and *arc-*
244 *nosZ* genes) was studied by terminal restriction fragment length polymorphism (T-RFLP) analysis.
245 Description of primer sets, PCR ingredients and amplification details used for T-RFLP analysis are
246 specified in Table 2. Prior to digestion, all genes were amplified using PCR cocktails and
247 amplification according to the details described in Table 2. For T-RFLP analysis, each forward
248 primer was labeled with the fluorescent dye 6-FAM. Replicate amplicons were pooled, purified
249 (Sephadex G-50, GE Healthcare Biosciences, Waukesha, WI, USA) according to Rasche et al.
250 (2006), and 200 ng of each amplicon were digested with 5 U *AluI* (archaeal 16S rRNA; New
251 England Biolabs (NEB) Inc., Ipswich, MA, USA), overnight at 37°C (Rasche et al., 2011). For the
252 digestion of *nosZ* and *arc-nosZ* gene amplicons, several restriction enzymes and their combinations

253 were tested (data not shown). Finally, 10 U HpyCH4IV (total prokaryotic *nosZ* clade I; NEB) and a
254 10 U cocktail of *AluI*, *RsaI*, and HpyCH4IV (*arc-nosZ*; NEB) revealed representative T-RFLP
255 fingerprints of both genes. Prior to T-RFLP analysis, digests were purified with Sephadex™ G-50
256 (Rasche et al., 2006) and a 2 µl aliquot was mixed with 17.75 µl HiDi formamide (Applied
257 Biosystems) and 0.25 µl internal size standard (500 ROX Size Standard, Applied Biosystems).
258 Labeled terminal-restriction fragments (T-RFs) were denatured at 95°C for 3 min, chilled on ice and
259 detected on an ABI 3130 automatic DNA sequencer (Applied Biosystems). Electropherograms were
260 compiled into numeric data using the Peak Scanner software (version 1.0, Applied Biosystems) and
261 fragment length with peak height >50 fluorescence units were used for profile comparison. Raw
262 data were normalized prior to statistical analysis (Dunbar et al., 2000).

263

264 2.4. Statistical data analysis

265 Data on abundance of the studied genes and soil chemical properties were subjected to ANOVA
266 using Statistical Analysis Software program (SAS V 9.2, SAS Institute Inc., North Carolina, USA).
267 For statistical analysis of the gene abundance data, a generalized linear mixed model with a
268 negative binomial distribution error and a log link function was used. Effects of straw addition
269 (factor “straw”) and incubation time (factor “time”) as well as the interaction between both factors
270 on abundance of all studied genes and on chemical properties were tested by two-way analysis of
271 variance (ANOVA). The data were checked for normality and homoscedasticity on model residuals
272 using quantile–quantile (Q–Q) plots, histograms and studentized residual plots.

273 Pearson linear correlation analyses were conducted and visualized for linearity in the SAS COR
274 procedure (data not shown) to relate the abundance of the target genes (dependent variables) to the
275 soil physico-chemical properties (independent variables).

276 Co-occurrence/co-exclusion analysis between the abundances of *arc-nosZ* and *gdh* genes and
277 chemical properties (NH₄⁺, DON) was carried out by using the “corrplot” package of R
278 (<http://www.r-project.org>). The analysis was performed controlling the “False Discovery Rate”

279 (FDR), as described by Benjamini and Hochberg (1995). The adjustment methods include the
280 Bonferroni correction ("bonferroni") in which the p-values are multiplied by the number of
281 comparisons.

282 TRFLP data sets were analyzed using Bray-Curtis similarity coefficients (Bray and Curtis, 1957). A
283 similarity matrix was generated for 16S archaea, total prokaryotic *nosZ* (*nosZ*) and archaeal *nosZ*
284 (*arc-nosZ*) target genes. This similarity matrix was used for one-way analysis of similarity
285 (ANOSIM) statistics (Clarke, 1993) to test if the composition of target communities was altered by
286 factors "straw", "time" and by "straw x time" interaction. ANOSIM is based on rank similarities
287 between the sample matrix and produces a test statistic 'R' (Rees et al., 2005). A 'global' R was
288 first calculated in ANOSIM, which evaluated the overall effect of a factor in the data set. This step
289 was followed by a pair wise comparison, whereby the magnitude of R indicated the degree of
290 separation between two tested communities. An R score of 1 indicated a complete separation, while
291 0 indicated no separation (Rees et al., 2005).

292 Statistical analysis of T-RFLP profiles was performed with Primer V6.1.13 software (Primer E,
293 Plymouth, UK) according to Rasche et al. (2011).

294 Additionally, to test the influence of NH₄ on the structure of *nosZ* and *arc-nosZ* genes among straw
295 and no straw treatments over the incubation time, principal component analysis (PCA) was
296 performed using the "dudi.pca" function of in R-package ade4 (Thioulouse et al. 1997; Dray et al.
297 2007). Thus, to indicate a redundancy in the data, a correlation matrix was constructed, where the
298 presence or absence as well as relative height of T-RFs were used as distinct data, whereas NH₄⁺
299 data were included in the analysis as 'environmental' variable. The data were grouped by straw and
300 no straw treatments over the incubation time. The graph of the PCA represents the differences
301 induced by NH₄⁺ in the community structures of *nosZ* and *arc-nosZ* genes across the treatments at
302 each sampling time: 1, 5, 10, 20, 30, and 60 days, according to the axes 'x' and 'y', which represent
303 eigen vectors with the greater variance.

304

305 3. Results

306 3.1. Soil chemical properties

307 After submergence, soil pH showed a progressive increase from 5.8 to 7.0 ($p < 0.01$) after 60 days of
308 incubation for both treatments with or without straw. This was in line with our previous results
309 confirming the onset of anaerobic conditions with a decrease in redox potential (Eh) from +293 mV
310 to -180 and +110 mV over the first 30 days of incubation, in the presence or absence of straw,
311 respectively (Cucu et al., 2014).

312 N_2O fluxes from soils receiving straw were negligible throughout the incubation period, while soils
313 not receiving straw showed a peak in N_2O emissions on the first three days immediately after
314 mineral N addition, followed by negligible emissions for the remaining incubation period (Fig.1).

315 Under submerged conditions, NH_4^+ was the main inorganic N form (Figure 2a). In the presence of
316 straw, NH_4^+ decreased from 57 mg N kg^{-1} to 41 mg N kg^{-1} in the first 20 days and subsequently
317 increased to values similar to the initial concentration by the end of the incubation (time \times straw
318 interaction, $p < 0.001$). In contrast, in the absence of straw, NH_4^+ concentrations decreased steadily to
319 23 mg N kg^{-1} by day 60 ($p < 0.001$).

320 Soils receiving straw showed NO_3^- concentrations below detection limit throughout the incubation
321 period (Fig. 2b). In the absence of straw, NO_3^- present at the beginning of the incubation (5 mg N
322 kg^{-1}) rapidly disappeared within the first 5 days, followed by an increase with time reaching 4 mg N
323 kg^{-1} at day 30 and 3 mg N kg^{-1} at day 60 ($p < 0.001$).

324 TDN concentrations in soils receiving straw increased from 66 to 76 mg N kg^{-1} with time ($p < 0.001$;
325 Fig. 2c), and DON, calculated as the difference between TDN and mineral N (i.e., $NH_4^+-N + NO_3^-$ -
326 N), increased from 10 to 20 mg N kg^{-1} at day 60. On the other hand, in soils without straw addition,
327 TDN decreased from 66 to 32 mg N kg^{-1} by 60 days (time \times straw interaction, $p < 0.001$) mainly due
328 to a loss of inorganic N-forms, since DON only varied from 10 to 6 mg kg^{-1} over the same time
329 period.

330 During the entire incubation, soils with straw showed higher DOC concentrations with respect to
331 soils without straw ($p < 0.01$), with initial concentrations of approximately 155 mg and 79 mg C kg⁻¹,
332 respectively (Fig. 2d). In both treatments, DOC contents decreased rapidly within the first 5 days of
333 the incubation and increased thereafter reaching highest values by day 60 (192 and 110 mg C kg⁻¹ in
334 straw and no straw treatments, respectively). Increase of DOC was more pronounced for soils
335 receiving straw (time × straw interaction, $p < 0.01$).

336

337 3.2. Microbial abundance

338 Abundance of bacterial 16S rRNA genes generally decreased with time, particularly for soils
339 receiving straw (time × straw interaction; $p < 0.001$), with values ranging between 1.0×10^{10} and
340 2.7×10^{10} copy numbers g⁻¹ soil (Fig. S1a). Variations in archaeal 16S rRNA gene abundance with
341 time generally followed a trend similar to that observed for bacterial 16S rRNA gene abundance
342 (time × straw interaction; $p < 0.001$), also exhibiting a higher gene copy number in soils without
343 straw (8.5×10^9) with respect to those with straw (6.5×10^9). However, soil samples receiving straw
344 showed a more pronounced decrease in archaeal 16S rRNA gene copy numbers from day 5 onwards
345 with respect to bacterial 16S rRNA genes. After 60 days of incubation, archaeal 16S rRNA gene
346 abundance declined to 1.1×10^9 and 5.5×10^8 copy numbers g⁻¹ soil for the no straw and straw
347 treatments, respectively (Fig. S1b).

348 Abundance of *gdhA* genes ranged from 1.3×10^9 to 2.0×10^9 copy numbers g⁻¹ soil in the straw
349 treatment, and from 1.1×10^9 to 1.8×10^9 copy numbers g⁻¹ soil in the no straw treatment (Fig. S2). In
350 both treatments, *gdhA* gene abundance decreased during the first 5 days of incubation, but
351 subsequently increased with time, generally showing higher values for soils receiving straw than
352 those not receiving straw by the end of the incubation time (time × straw interaction; $p < 0.001$).

353 Abundance of total prokaryotic *nosZ* clade I genes ranged from 2.3×10^7 to 8.6×10^7 copy numbers
354 g⁻¹ soil in the straw treatment and from 2.5×10^7 to 6.2×10^7 copy numbers g⁻¹ soil in the no straw
355 treatment (Fig. 3a), while the abundance of *arc-nosZ* genes ranged from 7×10^6 to 6.3×10^7 copy

356 numbers g^{-1} soil and from 1.7×10^7 to 6.4×10^7 copy numbers g^{-1} soil for the straw and no straw
357 treatments, respectively (Fig. 3b). For both genes, the abundance tended to decrease during the first
358 10 days of incubation, particularly for the *arc-nosZ* genes in the straw treatment (time \times straw
359 interaction; $p < 0.001$) and increased again during the later phase of incubation (between 20-60
360 days).

361 The relative abundance of *arc-nosZ* genes with respect to total prokaryotic *nosZ* clade I genes,
362 expressed as *arc-nosZ/nosZ* ratio, varied during the incubation time from 0.6 to 1 and from 1 to 0.2
363 in the no straw and straw treatments, respectively (data not shown).

364 Soils not receiving straw showed stronger positive correlations between gene abundance and
365 chemical properties with correlation coefficients ranging from $r = 0.216$ ($p < 0.01$) to 0.879
366 ($p < 0.001$) (Table 3). On the other hand, soils receiving straw showed correlation coefficients
367 ranging between $r = 0.153$ ($p < 0.01$) and 0.574 ($p < 0.001$).

368 The co-occurrence and co-exclusion between *arc-nosZ* and *gdhA* gene abundance and NH_4^+ and
369 DON were investigated by considering the significant correlations at False Discovery Rate (FDR) $<$
370 0.05 (Fig. S3). Abundance of *arc-nosZ* genes showed positive correlations with the abundance of
371 the *gdhA* gene and NH_4^+ content, while a negative correlation was found for DON contents (FDR $<$
372 0.05).

373

374 3.3. Microbial community structure

375 ANOSIM was used to compare effects of “straw” and “time” on the community structure of
376 archaeal 16S rRNA gene, *nosZ* clade I genes and *arc-nosZ* genes (Table 4). Factor “time” had a
377 greater influence than factor “straw” on all studied genes ($p < 0.001$). ANOSIM detected significant
378 interactions between both factors (archaeal 16S rRNA ($R = 0.724$), total prokaryotic *nosZ* clade I
379 ($R = 0.691$), *arc-nosZ* ($R = 0.735$) genes; $p < 0.001$)).

380 NH_4^+ had a distinct effect on the determined shifts in the community structures of *nosZ* and *arc-*
381 *nosZ* genes across the treatments during the incubation time (Fig. 4 and 5). The PCA clearly showed

382 a clustering of *nosZ* straw treated samples (Fig. 4a) from 10, 20 and 30 days which were well
383 separated from samples belonging to 1 and 5 and 60 days. A different pattern was revealed by the
384 PCA of *nosZ* no straw treated samples (Fig. 4b) which clustered in two different groups: samples
385 from 1 and 5 days and samples belonging to 10, 20, 30 and 60 days.

386 The PCA of *arc-nosZ* straw samples (Fig. 5a) showed a separation of samples from 10, 20, 30 and
387 60 days from samples belonging to 1 and 5 days, while the PCA of *arc-nosZ* no straw samples (Fig.
388 5b) revealed community differences between samples from 20, 30 and 60 days, separated from
389 samples belonging to 1, 5 and 10 days.

390

391 3.4. DNA ¹⁵N-enrichment

392 Addition of labeled (99 atom% ¹⁵N) (NH₄)₂SO₄ resulted in a progressive ¹⁵N-enrichment of DNA
393 with respect to natural ¹⁵N abundance (0.3663 atom% ¹⁵N) (Fig. S4). The highest enrichment was
394 obtained after 30 days of incubation, corresponding to 6.8 and 5.4 atom% ¹⁵N excess in the straw
395 and no straw treatments, respectively (time × straw interaction; p<0.001).

396

397 3.5. ¹⁵N-SIP analysis

398 T-RFLP fingerprints of *arc-nosZ* genes were generated from heaviest SIP gradient fractions (1.70 to
399 1.76 g ml⁻¹) to identify those community members which most efficiently incorporated the ¹⁵N
400 tracer (Fig. 6). As expected, community structure in fractions showed a lower diversity as compared
401 to the bulk samples, since only a part of the community was active in ¹⁵N assimilation. In all
402 fractions, with or without straw, the *arc-nosZ* gene community was dominated by two T-RFs (i.e.,
403 191 and 273 bp), but also by other T-RFs (e.g., 107, 117, 164, 185 bp) which were detected in
404 several fractions. T-RFLP fingerprinting revealed a clear distinction in T-RF allocation between ¹⁴N
405 (Fig. 6a and 6c) and ¹⁵N SIP gradients (Fig. 6b and 6d) suggesting a successful ¹⁵N enrichment in
406 the heaviest fractions.

407 Conversely, total *nosZ* gene clade I T-RFLP fingerprinting analysis showed no ¹⁵N-enrichment in
408 all fractions of both the straw and no straw treatments (data not shown).
409 Incorporation of the ¹⁵N tracer in *arc-nosZ* genes was further corroborated by quantification of
410 archaeal *gdhA* genes performed on heavy CsCl gradient fractions in both treatments, with or
411 without straw. In the straw treatment (Fig. 7a), label incorporation was evident in the ¹⁵N labeled
412 gradient, where the gene abundance peak shifted to a buoyant density (BD) of 1.7594 g ml⁻¹
413 compared to that of the ¹⁴N control (1.7570 g ml⁻¹). This was similar in the no straw treatment,
414 where the gene abundance peak in the ¹⁵N gradient shifted to a BD of 1.7157 g ml⁻¹ compared to the
415 ¹⁴N control with 1.6987 g ml⁻¹ (Fig. 7b). The shift in BD of approximately 0.02 g ml⁻¹ corresponded
416 to 50% of the density shift of 0.04 g ml⁻¹ expected for 100% label incorporation (Lueders et al.,
417 2004). In addition, the presence of ¹⁵N labeled DNA into even heavier fractions (BD of 1.7365 to
418 1.7560 g ml⁻¹) indicated an overall higher label incorporation in the no straw with respect to straw
419 ¹⁵N-gradient.

420

421 **4. Discussion**

422 *4.1. Archaeal N assimilation*

423 Rice straw addition to submerged rice paddy soils represented the key factor for observed diverse
424 responses of total bacterial and archaeal communities. This suggested that soil N immobilization
425 and gaseous losses – as quantified in the previous experiment as a function of straw application
426 (Cucu et al., 2014) – were controlled by both bacteria and archaea. However, under our
427 experimental conditions, the archaeal community showed a clear domination over bacteria,
428 especially in the absence of straw. In contrast to denitrifying bacteria, which usually use a
429 dissimilatory metabolism under non-limiting NH₄⁺ conditions, archaeal denitrifiers rely on an
430 assimilatory pathway to acquire N for formation of nucleic and amino acids as well as proteins
431 (Cabello et al., 2004; Nannipieri and Paul, 2009; Rusch, 2013). This was confirmed by the high
432 abundance of *gdhA* gene and progressive ¹⁵N-enrichment of DNA that displayed the effective biotic

433 immobilization of N during incubation. These observations suggested that independently of straw
434 addition, archaeal N assimilation was an important process in the studied soil. Moreover, the similar
435 increase in *gdhA* gene abundance in both straw-treated and non-treated soils suggested that archaeal
436 assimilation under non-NH₄⁺ limiting environments was hardly influenced by the presence of labile
437 organic C. However, the negative correlation between *gdhA* gene abundance and DOC indicated a
438 stronger influence of soil derived C sources with respect to that of added straw. Moreover, although
439 *gdhA* gene abundance was positively correlated with NH₄⁺ content in both treatments, this
440 relationship was stronger for soils not receiving straw with respect to straw treated soils. On the
441 other hand, the stronger correlation between *gdhA* gene abundance and TDN (i.e., the sum of
442 inorganic and organic N) with respect to NH₄⁺ for soils receiving straw suggested that, although
443 archaeal N assimilation was driven by NH₄⁺ availability, archaea could have modified their
444 metabolism with time to obtain N from other easily accessible sources (e.g., dissolved organic N
445 (DON) from labile OM) (Offre et al., 2013). Support for this was given by the negative correlation
446 between *gdhA* gene abundance and DON which was in line with a rapid consumption and turnover
447 of this labile N pool and the consequent increase in inorganic available N forms (Cucu et al., 2014).
448 In addition, the slightly higher *gdhA* gene abundance in the presence of labile organic matter (OM)
449 could be most likely attributed to an enhanced supply of NH₄⁺ resulting from the mineralization of
450 added straw and dissolved OM released under anoxic conditions (Cucu et al., 2014).

451

452 4.2. N assimilation as support for archaeal denitrification

453 ¹⁵N-DNA-based SIP revealed specific archaeal community members involved in both N
454 assimilation and denitrification processes. The ¹⁵N-labeled NH₄⁺ was actively assimilated via *gdhA*
455 gene for subsequent utilization (e.g., N₂O reductase synthesis), N immobilization and community
456 growth. Although labeling success of DNA was apparently low in comparison to other studies (e.g.,
457 Buckley et al., 2007a; España et al., 2011), a clear labeling effect was observed in *arc-nosZ* gene
458 community fingerprints of ‘heavy’ fractions in ¹⁵N-enriched samples compared to control

459 treatments. Moreover, NH_4^+ assimilation was confirmed by a higher abundance of *gdhA* gene in
460 heavy ^{15}N labeled fractions, especially in the absence of straw.

461 Total prokaryotic *nosZ* clade I gene fingerprinting in the SIP fractions showed no evident ^{15}N
462 assimilation (data not shown). This result indicated especially in the straw treatment the shift in the
463 total prokaryotic *nosZ* community towards bacterial members and suggested that bacterial
464 denitrifiers were most probably using only N as an electron sink and for energy conservation
465 (Zumft 1997). Archaeal N assimilation resulted in an increase in the abundance of assayed N_2O
466 reduction genes (*arc-nosZ*), indicating the predominance of denitrifying archaea compared to their
467 bacterial counterparts, especially in soils not receiving straw. This was confirmed by the high
468 relative abundance of *arc-nosZ* with respect to total prokaryotic *nosZ* clade I genes, as evidenced by
469 *arc-nosZ/nosZ* ratios. Since the abundance of total prokaryotic *nosZ* clade II genes is in the same
470 range of clade I (Jones et al., 2013), quantification of clade II may result in an underestimation of
471 the archaeal proportion of the total N_2O reducing community. However, archaeal members may still
472 have a significant role in completing the denitrification process. The abundance of total prokaryotic
473 *nosZ* clade I and specifically *arc-nosZ* genes was reflected in negligible N_2O emission fluxes over
474 the whole incubation period in both treatments. This confirmed that the gaseous losses of applied N
475 (up to 60%) in the absence of straw (c.f. only 20% lost in the presence of straw), reported by Cucu
476 et al (2014) could be mainly attributed to N_2 emissions. These findings were in line with previous
477 studies reporting high N_2O reduction under continuous rice soil flooding (DeDatta 1995; Ussiri and
478 Lal, 2013; Peyron et al., 2016).

479

480 4.3. Resource driven archaeal NH_4^+ assimilation and denitrification

481 It is generally assumed that net ammonification and consequently N availability are greater under
482 anaerobic soil conditions due to low metabolic N requirements of anaerobic microorganisms
483 (Reddy and DeLaune, 2008). However, our findings indicated that archaeal denitrifiers were able to

484 assimilate N under non-limiting NH_4^+ conditions and that this metabolic modification was most
485 likely a function of C and N availability.

486 Observed alterations of *arc-nosZ* gene community dynamics in soils without straw suggested a
487 greater efficiency in NH_4^+ assimilation and N immobilization than in the presence of straw. This
488 was probably due to a more competitive and versatile metabolism of chemoautotrophic
489 microorganisms (e.g., archaea) being well-adapted to resource limitation. Although it is generally
490 accepted that increased C substrate availability increases microbial population size (Anderson and
491 Domsch 1978; Fontaine et al, 2003) and thus activity, our results clearly indicated that under C
492 limiting conditions archaeal denitrifiers were most likely equipped with a unique metabolic
493 flexibility to scavenge alternative nutrient sources (Müller et al., 2014). Accordingly, in the absence
494 of alternative labile organic C (e.g., rice straw), the microbial communities might utilized carbon
495 dioxide as their main C source (Bock et al., 1991), reduced compounds (e.g., ammonium, iron(II),
496 and sulfide) as electron donors (Liesack et al., 2000; Megonigal et al., 2004), or N_2O as sole
497 electron acceptor (Stres et al., 2004; Strohm et al., 2007; Braker and Conrad, 2011). This suggested
498 a contrast with the general assumption that prokaryotic denitrifiers are predominantly heterotrophic
499 (Parkin 1987).

500 Independent of the adopted agricultural management, our results might display that rice paddy soils
501 act as temporary sink for N_2O . Furthermore, abundance of both denitrifying genes was positively
502 correlated to NH_4^+ and TDN concentrations highlighting the co-occurrence of both complete
503 denitrification and ammonification in the same genome (Tiedje et al., 1988; Sanford et al., 2012).
504 Support for this was also given through the negative correlation of *arc-nosZ* gene abundance with
505 DON contents. Likewise, mineralization promotes the recycling of immobilized N which may then
506 enter further processes including denitrification (Nannipieri et al., 2003).

507 As expected, the higher availability of labile C with straw addition under non-limiting N conditions
508 induced N assimilation but also a lower abundance of total *nosZ* clade I and *arc-nosZ* genes, as well
509 as distinct variation in the community structure of these genes. Lower abundance of *nosZ* genes in

510 the presence of straw was in contrast with other studies (Chen et al., 2012a,b). However, at the later
511 stages of incubation, when most of added straw was probably decomposed, progressively higher
512 abundance of *nosZ* and *arc-nosZ* genes was positively related to DOC content which was in line
513 with the findings of Kandeler et al., (2006) and Philippot et al., (2009). Labile organic C served as
514 resource for denitrifying bacteria that outcompeted archaeal counterparts, as evidenced by a
515 decreasing *arc-nosZ/nosZ* ratio with incubation time. This was in agreement with Ishii et al.
516 (2011b) who, by using functional single-cell (FSC) and DNA-based SIP with ¹³C-labeled succinate
517 as electron donor and N₂O as electron acceptor showed that under their experimental conditions,
518 most N₂O reducers are bacterial denitrifiers. Examining the N₂O reduction rates of the isolated
519 strains, the authors confirmed the growth of putative bacterial denitrifiers reciprocally to N₂O
520 reduction in rice field soils, although many bacteria have only partial pathways of denitrification
521 (Shapleigh, 2013).

522

523 **5. Conclusions and outlook**

524 Nitrogen immobilization based on archaeal NH₄⁺ assimilation was shown to represent an important
525 step for proliferation and dynamics of those microbial community members harboring the *arc-nosZ*
526 gene encoding N₂O reductase enzyme, irrespective of the presence or absence of labile OM. Under
527 our experimental conditions, the relative abundance of *arc-nosZ* genes with respect to total
528 prokaryotic *nosZ* from clade I genes was based on an adaptable metabolic portfolio. However, in
529 the presence of straw, bacterial *nosZ* genes may also contribute to the completion of denitrification.
530 Based on these considerations, we developed a conceptual model to represent the archaeal N
531 assimilation and denitrification pathways driven by different resources under anaerobic conditions
532 (Fig. 8). The response of *arc-nosZ* genes to different C availability under non-limiting N conditions
533 may have important ecological implications in controlling the immobilization and loss of N from
534 fertilized paddy soils. Furthermore, the high adaptability of archaea to drive denitrification to
535 completion may be a key feature in mitigating N₂O emissions.

536 The present study suggested the potential contribution of archaea to the last step of denitrification
537 when favorable environmental conditions are given as complement to the bacterial counterparts
538 from *nosZ* clade I. However, although we have provided important insights, the full understanding
539 of archaeal involvement in the denitrification process remains still incomplete. This accounts
540 particularly for the currently limited phylogenetic information on *arc-nosZ* gene synthesizing
541 archaea in soils (Rusch 2013). Hence, we strongly suggest the generation of *arc-nosZ* gene libraries
542 to enhance phylogenetic knowledge on this particular gene. In addition, prospective research should
543 also consider the unconsidered *nosZ* gene (e.g., clade II, atypical *nosZ*) to better understand the
544 relative role of archaea in acting as N₂O sink.

545 As N assimilation processes are regulated by resource availability, we recommend evaluating the
546 effects of different N fertilizer and organic residue types with contrasting C/N ratio. In this respect,
547 ¹⁵N and ¹³C-label SIP-based studies may be appropriate to evaluate the metabolic adaptability of
548 archaea for mineral and organic substrate consumption and their implication in mitigating emissions
549 of climate relevant gases from paddy soils.

550

551 **Funding**

552 This work was partly supported by the Italian Ministry of Agriculture, Food and Forestry
553 (MiPAAF) under the project “Sustaining the National Rice Industry through Research, Technology,
554 Innovation and Formation (POLORISO)” and International Humic Substance Society (IHSS).

555

556 **Acknowledgements**

557 We gratefully thank R. Gorra (Department of Agricultural, Forest and Food Sciences, University of
558 Turin, Italy) for fruitful discussions; J. Laso (Department of Bioinformatics, University of
559 Hohenheim, Germany) and I. Ferrocino (Department of Agricultural, Forest and Food Sciences,
560 University of Turin, Italy) for their support in statistical analysis. The authors are further grateful to
561 L. Hölzle (Institute of Animal Science, University of Hohenheim) for access to his ABI 3130

562 Genetic Analyzer and C. Röhl (Institute of Plant Production and Agroecology in the Tropics and
563 Subtropics, University of Hohenheim) for her technical support.

564 **References**

- 565 Anderson JPE, Domsch KH (1978) A physiological method for the quantitative measurement of
566 microbial biomass in soils. *Soil Biol Biochem*: 10: 215–221
- 567 Bird JA, Horwáth WR, Eagle AJ, van Kessel (2001) Immobilization of fertilizer nitrogen in rice.
568 *Soil Sci Soc Am J* 65:1143-1152
- 569 Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful
570 approach to multiple testing. *J R Stat Soc Series B* 57: 289–300
- 571 Bray JR, Curtis JT (1957) An ordination of the upland forest communities of southern Wisconsin.
572 *Ecol Monogr* 27:325-349
- 573 Bock E, Koops HP, Harms H, Ahlers B (1991) The biochemistry of nitrifying organisms. In:
574 Shively JM, Barton LL (eds) *Variations in autotrophic life*, London, Academic Press, pp
575 171-200
- 576 Bonete MJ, Martínez-Espinosa RM, Pire C, Zafrilla B, Richardson DJ (2008) Nitrogen metabolism
577 in haloarchaea. *Saline Systems* 4:9
- 578 Braker G and Conrad R. Diversity, structure, and size of N₂O-producing microbial communities in
579 soils—what matters for their functioning? (2011) In: Laskin I, Sariaslani S, Gadd GM (eds)
580 *Advances in Applied Microbiology*, vol.75, SanDiego, Elsevier Academic Press Inc., pp 33–
581 70
- 582 Buckley DH, Huangyutitham V, Hsu SF, Nelson TA (2007a) Stable isotope probing with ¹⁵N₂
583 reveals novel non-cultivated diazotrophs in soil. *Appl Environ Microbiol* 73:3196-3204
- 584 Buckley DH, Huangyutitham V, Hsu SF, Nelson TA (2007b) Stable isotope probing with ¹⁵N
585 achieved by disentangling the effects of genome G+C content and isotope enrichment on
586 DNA density. *Appl Environ Microbiol* 73:3189-3195
- 587 Cabello P, Roldán MD, Moreno-Vivían C (2004) Nitrate reduction and the nitrogen cycle in
588 archaea. *Microbiol* 150:3527-3546

- 589 Cassman KG, Peng S, Olk DC, Ladha JK, Reichardt W, Dobermann A, Singh U (1998)
590 Opportunities for increased nitrogen-use efficiency from improved resource management in
591 irrigated rice systems. *Field Crop Res* 56:7-39
- 592 Cassman KG, Dobermann A, Walters DT (2002) Agro-ecosystems, nitrogen-use efficiency, and
593 nitrogen management. *Ambio* 31:132-140
- 594 Chen Z, Liu J, Wu M, Xie X, Wu J, Wei W (2012a) Differentiated response of denitrifying
595 communities to fertilization regime in paddy soil. *Microb Ecol* 63:446-459
- 596 Chen Z, Hou H, Zheng Y, Qin H, Zhu Y, Wu J, Wei W (2012b) Influence of fertilization regimes
597 on a *nosZ*-containing denitrifying community in a rice paddy soil. *J Sci Food Agr* 92:1064-
598 1072
- 599 Chuang H-P, Wu J-H, Ohashi A, Abe K, Hatamoto M (2014) Potential of nitrous oxide conversion
600 in batch and down-flow hanging sponge bioreactor systems. *Sustain Environ Res* 24:117-
601 128
- 602 Clarke KR (1993) Non-parametric multivariate analyses of changes in community structure. *Aust J*
603 *Ecol* 18: 117-143
- 604 Conrad R (1995) Soil microbial processes involved in production and consumption of atmospheric
605 trace gasses. In: Jones JG (ed) *Advances in Applied Microbiology*, New York. Plenum
606 Press, pp 207-250
- 607 Conrad R (1996) Soil microorganisms as controllers of atmospheric trace gases (H_2 , CO , CH_4 ,
608 OCS , N_2O , and NO). *Microbiol Rev* 60:609-640
- 609 Cucu MA, Said-Pullicino D, Maurino V, Bonifacio E, Romani M, Celi L (2014) Influence of redox
610 conditions and rice straw incorporation on nitrogen availability in fertilized paddy soils.
611 *Biol Fertil Soil* 50:755-764
- 612 Davidson EA, Swank WT, Perry TO (1986) Distinguishing between nitrification and denitrification
613 as sources of gaseous nitrogen production in soil. *Appl Environ Microb* 52:1280-1286
- 614 DeDatta SK (1995) Nitrogen transformations in wetland rice ecosystem. *Fert Res* 42:193-203

615 de Vries S, Schröder I (2002) Comparison between the nitric oxide reductase family and its aerobic
616 relatives, the cytochrome oxidases. *Biochem SocTrans* 30:662-667

617 Devêvre OC, Horwáth WR (2001) Stabilization of fertilizer nitrogen-15 into humic substances in
618 aerobic vs. waterlogged soil following straw incorporation. *Soil Sci Soc Am J* 65:499-510

619 Dray S, AB Dufour (2007) The ade4 package implementing the duality diagram for ecologist. *J Stat*
620 *Softw* 22:1-20

621 Dunbar J, Ticknor LO, Kuske CR (2000) Assessment of microbial diversity in four south western
622 United States soils by 16S rRNA gene terminal restriction fragment analysis. *Appl Environ*
623 *Microb* 66:2943-2950

624 Eagle AJ, Bird JA, Horwáth WR, Linquist BA, Brouder SM et al (2000) Rice yield and nitrogen
625 utilization efficiency under alternative straw management practices. *Agronom J* 92:1096-
626 1103

627 España M, Rasche F, Kandeler E, Brune T, Rodriguez B et al (2011) Identification of active
628 bacteria involved in decomposition of complex maize and soybean residues in a tropical
629 Vertisol using ¹⁵N-DNA stable isotope probing. *Pedobiologia* 54:187-193

630 Fazli P, Hasfalina CM, UmiKalsom MS, Azni I (2013) Review Article: Characteristics of
631 methanogens and methanotrophs in rice fields. *AsPac J Mol Biol Biotechnol* 21: 3-17

632 Ferrè C, Zechmeister-Boltenstern S, Comolli R, Andersson M, Seufert G (2012) Soil microbial
633 community structure in a rice paddy field and its relationships to CH₄ and N₂O fluxes. *Nutr*
634 *Cycl Agrosyst* 93:35-50

635 Fetzer S, Bak F, Conrad R (1993) Sensitivity of methanogenic bacteria from paddy soil to oxygen
636 and desiccation. *FEMS Microbiol Ecol* 12:107–115

637 Fontaine S, Mariotti A, Abbadie L (2003) The priming effect of organic matter: a question of
638 microbial competition? *Soil Biol Biochem* 35:837-843

639 Garcia JL, Tiedje JM (1982) Denitrification in rice soils. In: Dommergues YR, Diem HG (eds)
640 Microbiology of tropical soils and plant productivity. Springer Netherlands, Dordrecht, pp
641 187-208.

642 Ghosh BC, Bhat R (1998) Environmental hazards of nitrogen loading in wetland rice fields.
643 Environ Pollut 102:123-126

644 Henry S, Bru D, Stres B, Hallet S, Philippot L (2006) Quantitative detection of the *nosZ* gene,
645 encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, *narG*,
646 *nirK*, and *nosZ* genes in soils. Appl Environ Microbiol 72:5181-5189

647 Hutchinson GL, Mosier AR (1981) Improved soil cover method for field measurement of nitrous
648 oxide fluxes. Soil Sci Soc Am J, 45: 311–316

649 Ishii S, Ikeda S, Minamisawa K, Senoo K (2011a) Nitrogen cycling in rice paddy environments:
650 past achievements and future challenges. Microbes Environ J 26:282-292

651 Ishii S, Ohno H, Tsuboi M, Otsuka S, Senno K (2011b) Identification and isolation of active N₂O
652 reducers in rice paddy soil. ISME J 5:1936-1945

653 Jones CM, Graf DRH, Bru D, Philippot L, Hallin S (2013) The unaccounted yet abundant nitrous
654 oxide-reducing microbial community: a potential nitrous oxide sink. ISME J 7:417-426

655 Kandeler E, Deiglmayr K, Tscherko D, Bru D, Philippot L (2006) Abundance of *narG*, *nirS*, *nirK*,
656 and *nosZ* genes of denitrifying bacteria during primary successions of a glacier foreland.
657 Appl Environ Microbiol 72:5957-5962

658 Kögel-Knabner I, Amelung W, Cao Z, Fiedler S, Frenzel P, Jahn R et al (2010) Biogeochemistry of
659 paddy soils. Geoderma 157:1-14

660 Lane D (1991) 16S/23S rRNA sequencing. In: Stackebrandt A and Goodfellow M (eds) Nucleic
661 Acid Techniques Systematics, UK, John Wiley, West Sussex, pp 115–175

662 Liesack W, Schnell S, Revsbech NP (2000) Microbiology of flooded rice paddies. FEMS Microbiol
663 Rev 24:625-645

- 664 Lueders T, Friedrich M (2000) Archaeal population dynamics during sequential reduction processes
665 in rice field soil. *Appl Environ Microbiol* 66:2732-2742
- 666 Lueders T, Manefield M, Friedrich MW (2004) Enhanced sensitivity of DNA and rRNA based
667 stable isotope probing by fractionation and quantitative analysis of isopycnic centrifugation
668 gradients. *Environ Microbiol* 6:73-78
- 669 Majumdar D (2013) Biogeochemistry of N₂O uptake and consumption in submerged soils and rice
670 fields and implications in climate change. *Crit Rev Env Sci Technol* 43:2653-2684
- 671 Martínez-Espinosa RM, Marhuenda-Egea FC, Bonete MJ (2001) Assimilatory nitrate reductase
672 from the haloarchaeon *Haloferax mediterranei*: purification and characterization. *FEMS*
673 *Microbiol Lett* 204:381–385
- 674 Megonigal JP, Hines ME, Visscher PT (2004) Anaerobic metabolism: linkages to trace gases and
675 aerobic processes. In: Schlesinger WH (ed) *Biogeochemistry*, Oxford, UK, Elsevier-
676 Pergamon, pp 317-424
- 677 Mulec MI, Ausec L, Danevčič T, Levičnik-Höfferle Š, Jerman V, Kraigher B (2014) Microbial
678 community structure and function in peat soil. *Food Technol Biotechnol* 52:180-187
- 679 Müller J, Hense BA, Marozava S, Kuttler Ch, Meckenstock RU (2014) Model selection for
680 microbial nutrient uptake using a cost-benefit approach. *Math Biosci* 255:52-70
- 681 Muyzer G, Dewaal EC, Uitterlinden AG (1993) Profiling of complex microbial populations by
682 denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified
683 genes coding for 16S rRNA. *Appl Environ Microbiol* 59:695-700
- 684 Nannipieri P, Ascher J, Ceccherini MT, Landi L, Pietramellara G, Renella G (2003) Microbial
685 diversity and soil functions. *Europ J Soil Sc* 54:655-670
- 686 Nannipieri P, Paul EA (2009) The chemical and functional characterization of soil N and its biotic
687 components. *Soil Biol Biochem* 41:2357-2369
- 688 Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M et al (2007) DNA stable-isotope
689 probing. *Nat Protoc* 2:860-866

690 Nogales B, Timmis KN, Nedwell DB, Osborn AM (2002) Detection and diversity of expressed
691 denitrification genes in estuarine sediments after reverse transcription-PCR amplification
692 from mRNA. *Appl Environ Microbiol* 68:5017-5025

693 Offre P, Spang A, Schleper C (2013) Archaea in biogeochemical cycles. *Ann Rev Microbiol* 67:
694 437-457

695 Parkin TB (1987) Soil microsites as a source of denitrification variability. *Soil Sci Soc Am J* 51:
696 1194-1199

697 Peyron M, Bertora C, Pelissetti S, Said-Pullicino D, Celi L, Miniotti E, Romani M, Sacco D (2016)
698 Greenhouse gas emissions as affected by different water management practices in temperate
699 rice paddies. *Agric Ecosyst Environ* 232:17-28

700 Philippot L (2002) Denitrifying genes in bacterial and archaeal genomes. *Biochim Biophys Acta*
701 1577:355-376

702 Philippot L, Čuhel J, Saby NPA, Chèneby D, Chroňáková A, Bru D et al (2009) Mapping field-
703 scale spatial distribution patterns of size and activity of the denitrifiers community. *Environ*
704 *Microbiol* 11:1518-1526

705 Prieme A, Braker G, Tiedje JM (2002) Diversity of nitrite reductase (*nirK* and *nirS*) gene fragments
706 in forested upland and wetland soils. *Appl Environ Microbiol* 68:1893-1900

707 Rasche F, Hödl V, Poll C, Kandeler E, Gerzabek MH, Van Elsas JD et al (2006) Rhizosphere
708 bacteria affected by transgenic potatoes with antibacterial activities compared with the
709 effects of soil, wild type potatoes, vegetation stage and pathogen exposure. *FEMS Microbiol*
710 *Ecol* 56: 219-235

711 Rasche F, Knapp D, Kaiser C, Koranda M, Kitzler B, Zechmeister-Boltenstern S et al (2011)
712 Seasonality and resource availability control bacterial and archaeal communities in soils of a
713 temperate beech forest. *ISME J* 5:389-402

714 Rees GN, Baldwin DS, Watson GO, Perryman S, Nielsen DL (2005) Ordination and significance
715 testing of microbial community composition derived from terminal restriction fragment

716 length polymorphisms: application of multivariate statistics. *Ant Van Leeuwenhoek* 86:
717 339-347. <http://dx.doi.org/10.1007/s10482-005-0498-5>

718 Reddy KR (1982) Nitrogen cycling in a flooded-soil ecosystem planted to rice (*Oryza sativa* L.).
719 *Plant Soil J* 67:209-220

720 Reddy KR, deLaune RD (2008) *Biogeochemistry of wetlands: science and applications*, USA, CRC
721 Press, Boca Raton, FL

722 Rice CW, Tiedje JM (1989) Regulation of nitrate assimilation by ammonium in soils and in isolated
723 soil microorganisms. *Soil Biol Biochem* 21:597-602

724 Rusch A (2013) Molecular tools for the detection of nitrogen cycling. Article ID 676450, *Archaea*
725 doi:10.1155/2013/676450

726 Rusch A (2016) *Archaea*. In Kennish MJ (ed) *Encyclopedia of Estuaries*, Chapter: 287, Springer
727 Science, Business Media Dordrecht, doi: 10.1007/978-94-017-8801-4_287, pp 35-37

728 Sahrawat KL (2004) Ammonium production in submerged soils and sediments: the role of
729 reducible iron. *Commun Soil Sci Plant Anall* 35:399-411

730 Said-Pullicino D, Cucu MA, Sodano M, Birk JJ, Glaser B, Celi L (2014) Nitrogen immobilization
731 in paddy soils as affected by redox conditions and rice straw incorporation. *Geoderma* 228-
732 229:44-53

733 Samuel BS, Hansen EE, Manchester JK, Coutinho PM, Henrissat B, Fulton R et al (2007) Genomic
734 and metabolic adaptations of *Methanobrevibacter smithii* to the human gut. *PNAS J* 104:
735 10643-10648

736 Sanford RA, Wagner DD, Wu Q, Chee-Sanford JC, Thomas SH, Cruz-Garcia C et al (2012)
737 Unexpected nondenitrifier nitrous oxide reductase gene diversity and abundance in soils.
738 *PNAS J* 48:19709-19714

739 Shapleigh JP. Denitrifying prokaryotes (2013). In: Rosenberg E et al (eds) *Prokaryotes. Prokaryotic*
740 *Physiology and Biochemistry*, Berlin Heidelberg, Springer-Verlag, pp 405-425

741 Stres B, Mahne I, Avgustin G, Tiedje JM (2004) Nitrous oxide reductase (*nosZ*) gene fragments
742 differ between native and cultivated Michigan soils. *Appl Environ Microbiol* 70:301–309

743 Strohm TO, Griffin B, Zumft WG, Schink B (2007) Growth yields in bacterial denitrification and
744 nitrate ammonification. *Appl Environ Microbiol* 73:1420–1424

745 Thioulouse J, Chessel D, Doledec S, Olivier JM (1997) ADE-4: a multivariate analysis and
746 graphical display software. *Stat Comput* 7:75–83

747 Tiedje JM (1988) Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In:
748 Zehnder AJB (ed) *Biology of Anaerobic Microorganisms*, New York, John Wiley & Sons,
749 pp 179-244

750 Tomlinson GA, Jahnke LL, Hochstein LI (1986) *Halobacterium denitrificans* sp. nov., an extremely
751 halophilic denitrifying bacterium. *Int J Syst Bacteriol* 36:66-70

752 Ussiri D, Lal R (2013) Nitrous oxide emissions from rice fields. In: *Soil emission of nitrous oxide*
753 *and its mitigation*, Netherlands, Springer, pp 213-242

754 WRB, IUSS Working Group WRB, 2007. *World Reference Base for Soil Resources 2006*, first
755 update 2007. *World Soil Resources Reports No. 103*. FAO, Rome, Italy

756 Zou J, Huang Y, Zheng X, Wang Y (2007) Quantifying direct N₂O emissions in paddy fields during
757 rice growing season in mainland China: dependence on water regime. *Atm Environ* 41:
758 8030-8042

759 Zumft WG (1992) The denitrifying prokaryotes. In: Balows A, Trüper HG, Dworkin M, Harder W,
760 Schleifer K-H (eds) *The prokaryotes. A handbook on the biology of bacteria:*
761 *ecophysiology, isolation, identification, application*, New York, Berlin, Heidelberg,
762 Springer-Verlag, 2nd edn, vol 1, pp 554-582

763 Zumft WG (1997) Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 61:
764 533-616

766 **Figure Captions**

767

768 **Fig. 1** Daily nitrous oxide (N₂O) flux over time in soils incubated with or without straw addition.

769 The error bars (n=3) represent the least significant difference at p=0.05

770

771 **Fig. 2** Variations in concentrations of ammonium (sum of submergence water and soil extractable

772 NH₄⁺) concentrations (a), nitrate (sum of submergence water and soil extractable NO₃⁻) (b), total

773 dissolved N (sum of submergence water and soil water-extractable N) (c), and dissolved organic C

774 (sum of submergence water and soil water-extractable C) (d) with time for soils incubated with or

775 without straw addition. The error bars (n=6) represent the least significant difference at p=0.05

776

777 **Fig. 3** Abundance of total prokaryotic *nosZ* clade I (a) and *arc-nosZ* (b) genes over time in soils

778 incubated with or without straw addition (n=6, means with standard errors). Different letters above

779 bars indicate significant differences at p<0.05 between straw and no straw treatments at every

780 sampling time (lowercase letters) as well as significant differences among different sampling times

781 (uppercase letters).

782

783 **Fig. 4** Principal component analysis (PCA) of TRFLP profiles using *nosZ* gene T-RF relative

784 abundance data obtained from straw (a) and no straw (b) treatments in function of NH₄⁺

785 concentration. Sampling time are indicated with colors (T 1 day – blue; T 5 days – red; T 10 days

786 – purple; T 20 days – pink; T 30 days – dark grey; T 60 days – green)

787

788 **Fig. 5** Principal component analysis (PCA) of TRFLP profiles using *arc-nosZ* gene T-RF relative

789 abundance data obtained from straw (a) and no straw (b) treatments in function of NH₄⁺

790 concentration. Sampling time are indicated with colors (T 1 day – blue; T 5 days – red; T 10 days

791 – purple; T 20 days – pink; T 30 days – dark grey; T 60 days – green)

792

793 **Fig. 6** T-RFLP fingerprints of *arc-nosZ* genes generated from density resolved SIP fractions of the
794 straw ^{14}N - $(\text{NH}_4)_2\text{SO}_4$ (a), ^{15}N - $(\text{NH}_4)_2\text{SO}_4$ (b) and the no straw ^{14}N - $(\text{NH}_4)_2\text{SO}_4$ (c), and ^{15}N -
795 $(\text{NH}_4)_2\text{SO}_4$ (d) treatment. Buoyant densities (g ml^{-1}) of analyzed fractions are given in parentheses

796

797 **Fig. 7** Quantification of archaeal *gdhA* genes in comparative CsCl density fractions of DNA
798 extracted from straw (a) and non-straw (b) treatments fertilized with either ^{14}N - or ^{15}N - ^{15}N -
799 $(\text{NH}_4)_2\text{SO}_4$

800

801 **Fig. 8** Schematic relationship between C and N resource availability and nitrogen assimilatory
802 archaeal denitrification in fertilized paddy soils suggesting the ecological importance of this
803 pathway

804

Graphical abstract




Figure 1




Figure 2




Figure 3




Figure 4





Figure 5





Figure 6




Figure 7





Figure 8



1 **Table 1** Description of primer sets and amplification details used for quantitative PCR.

Target group	Primer (reference)	Amplification details
All bacteria	Eub338 (Lane 1991)	40 cycles
(16S rRNA gene)	Eub518 (Muyzer et al. 1993)	95°C 30s, 55°C 35s, 72°C 45s
All archaea	Ar109f (Lueders and Friedrich 2000)	40 cycles
(16S rRNA gene)	Ar912r (Lueders and Friedrich 2000)	95°C 30s, 52°C 35s, 72°C 45s, 78°C 20s
<i>nosZ</i> gene clade I	<i>nosZ</i> -2f (Henry et al. 2006)	6 touch down cycles
	<i>nosZ</i> -2r (Henry et al. 2006)	95°C 15s, 65°C 30s (-1), 72°C 30s
		40 cycles
		95°C 15s, 60°C 30s, 72°C 30s, 81°C 30s
<i>arc-nosZ</i> gene	<i>arc-Nos-f</i> (Rusch 2013)	6 touch down cycles
	<i>arc-Nos-r</i> (Rusch 2013)	95°C 15s, 65°C 30s (-1), 72°C 30s
		40 cycles
		95°C 15s, 60°C 45s, 72°C 60s, 81°C 35s
<i>gdhA</i> gene	MSM0888f (Samuel et al. 2007)	40 cycles
	MSM0888r (Samuel et al. 2007)	95°C 30s, 58°C 35s, 74°C 30s

2

3 **Table 2** Description of primer sets, PCR ingredients and amplification details used for T-RFLP analysis.

Target group	Primer (reference)	DNA (ng)	Taq (U)	MgCl ₂ (mM)	Primer (mM)	dNTPs (mM)	Amplification details
All archaea	Ar109f (Lueders and Friedrich 2000) Ar912r (Lueders and Friedrich 2000)	5	2	1.5	0.15	0.2	35 cycles 95°C 5 m, 95°C 60s, 52°C 30s, 72°C 60s, 72°C 10 m
<i>nosZ</i> gene clade I	nosZ-2f (Henry et al. 2006) nosZ-2r (Henry et al. 2006)	10	2	1.5	0.2	0.2	6 touch down cycles 94°C 30s, 65°C 30s (-1), 72°C 30s 40cycles 94°C 30s, 60°C 30s, 72°C 30s
<i>arc-nosZ</i> gene	arc-Nos-f (Rusch 2013) arc-Nos-r (Rusch 2013)	10	0.5	-	0.4	0.2	40 cycles 95°C 60s, 95°C 20s, 72°C 20s, 72°C 90s, 72°C 15 m

4

6 **Table 3** Linear correlation coefficients (Pearson correlation coefficients, n=36) between microbial abundance and chemical data

Property	All bacteria		All archaea		total <i>nosZ</i> clade I		<i>arc-nosZ</i>		<i>gdhA</i>	
	straw	no straw	straw	no straw	straw	no straw	straw	no straw	straw	no straw
NH ₄ ⁺	0.351*	0.709***	0.469**	0.879***	0.414*	0.854***	0.457**	0.796***	0.313*	0.706***
TDN	ns	0.593***	ns	0.662***	ns	0.439**	ns	0.550***	0.574***	0.624***
NO ₃ ⁻	ns	ns	ns	ns	ns	0.319*	ns	0.272*	ns	ns
DOC	-0.464**	-0.450**	-0.470**	-0.742***	0.153*	0.216*	0.237*	0.246*	ns	-0.427**

7

8 Significance levels: not significant-ns: p>0.05; *p<0.05; **p<0.01; ***p<0.001.

Table 4 Analysis of similarity (ANOSIM) revealing the treatments effect and time on soil archaeal, total prokaryotic *nosZ* (*nosZ*) clade I and archaeal *nosZ* (*arc-nosZ*) denitrifying community structure

Factor	Global R		
	All archaea	<i>nosZ</i> clade I	<i>arc-nosZ</i>
straw	ns	0.173***	0.472***
time	0.479***	0.358***	0.567***
straw × time	0.724***	0.691***	0.735***

Levels of significance between two tested populations: not significant-ns: $p > 0.05$; *** $p < 0.001$.

Highlights

- Archaeal N assimilation was not affected by C availability
- N assimilation and immobilization preceded the archaeal denitrification
- In the presence of labile C denitrifying bacteria outcompeted archaeal counterparts
- Metabolic resilience induced a significant role of archaea in N₂O reduction
- A conceptual model of archaeal assimilatory denitrification was proposed

Supplementary Figure 1




FIG. S1 Abundance of bacterial (a) and archaeal (b) 16S rRNA genes over time in soils incubated with or without straw addition (n=6, means with standard errors).

Supplementary Figure 2




FIG. S2 Abundance of archaeal *gdhA* genes over time in soils incubated with or without straw addition (n=6, means with standard errors).

Supplementary Figure 3




FIG. S3 Significant co-occurrence and co-exclusion relationships between *arc-nosZ* and *gdhA* genes abundances and chemical properties (NH_4^+ , DON). Spearman's rank correlation matrix of genes abundance and chemical properties values. Strong correlations are indicated by large circles, whereas weak correlations are indicated by small circles. The colours of the scale bar denote the nature of the correlation, with 1 indicating a perfectly positive correlation (dark blue) and -1 indicating a perfectly negative correlation (dark red). Only significant correlations ($\text{FDR} < 0.05$) are shown.

Supplementary Figure 4




FIG. S4 Development of ^{15}N enrichment of DNA (at% ^{15}N) during incubation time and natural abundance of 0.336 at % ^{15}N of ammonium sulphate (dashed line) ($n=3$, means with standard errors).