

PRELIMINARY STUDIES ON THE EFFECT OF Fe-nanosponge COMPLEX IN HORTICULTURE

Monica Vercelli¹, Walter Gaino¹, Valeria Contartese²,
Lorenzo Gallo², Stefano Di Carlo³, Vander Tumiatti³,
Federica Larcher¹, Valentina Scariot¹

¹University of Torino, Italy

²Green Has Italia s.p.a.

³Sea Marconi Technologies s.a.s.

Abstract. Iron deficiency is a major nutritional disorder in crops. Inorganic Fe-compounds and synthetic Fe-chelates are commonly used to control chlorosis but their use arise environmental concerns. We recently developed a new iron fertilizer using a β -cyclodextrin-based nanosponge complex (Fe-NS). In this study, a pilot trial was performed on hydroponically cultivated horticultural plants in order to evaluate the effect of Fe-NS. Sweet corn and tomato were used as model plants analyzing chlorophyll, dry matter and Fe content. Fe-NS effect was compared to FeSO₄ and Fe-DTPA. Fe-NS had a positive effect on re-greening and growth in sweet corn and tomato plants.

Key words: iron chlorosis, Fe-fertilizers, nanosponges, sweet corn, tomato

INTRODUCTION

Mineral nutrients are essential for plant growth and development in soil and inert media [Marschner 1995, Chohura and Komosa 2003]. Depending on how great the growth requirement for a given nutrient, the nutrient is referred to as either a macronutrient or a micronutrient. Among the micronutrients, iron is an essential nutrient for plant growth and development, since it participates in life-sustaining processes, such as respiration and photosynthesis [Martínez-Cuenca et al. 2013]. Plants have developed two different strategies in response to Fe shortage: Strategy I, which occurs in dicotyledonous and non-grass monocotyledonous species and Strategy II, which occurs in

Corresponding author: Monica Vercelli, Department of Agricultural, Forest and Food Sciences, University of Torino, Largo Paolo Braccini 2, 10095 Grugliasco, TO, Italy, e-mail: monica.vercelli@unito.it

© Copyright by Wydawnictwo Uniwersytetu Przyrodniczego w Lublinie, Lublin 2015

Poaceae species. Strategy I plants can mobilize and take up Fe^{3+} ions from soil particles by acidification of the rhizosphere. Strategy II consists of the roots releasing high affinity chelating compounds (phytosiderophores), which are able to form stable complexes with the micronutrient [Marschner et al. 1986]. In both strategies, Fe deficiency induces several mechanisms aimed to increase Fe uptake from the soil [López-Millán et al. 2009].

Increasing the amount of crop-available Fe has long been carried out by means of iron-fertilizer application to soil and irrigation water. There is a very large number of Fe-contain fertilizers, with different degrees of effectiveness due to many different factors [Abadía et al. 2004]. Iron fertilizers are grouped into three main classes: inorganic Fe-compounds (Fe salts), synthetic Fe-chelates and natural Fe-complexes. Most used fertilizers are synthetic Fe-chelates [Lucena 2007]; this fertilizers are very effective but are expensive and their persistence in soil and groundwater arises environmental concerns [Sánchez-Alcalá et al. 2012]. Due to environmental risks associated with their use, alternative strategies have been suggested [Rombolà et al. 2003, Rombolà and Tagliavini 2007]. In the last years, research on Fe fertilizers has focused on development of new fertilizers, including slow-release and environmentally-friendly [Bhattacharya et al. 2007, Chandra et al. 2009]. Good practices to assess the efficiency of Fe-fertilizers are provided by the current knowledge on plant physiology and biochemistry, which not always taken into account in Fe-fertilizer efficiency agronomical studies [El-Jendoubi 2011].

In this context we developed a new iron fertilizer using β -cyclodextrin-based nanosponge complex (Fe-NS). Cyclodextrin-based nanosponges (NS) are biocompatible nanoporous nanoparticles synthesized by the carbonyl or dicarboxylate hyper-cross-linking of cyclodextrins, as reported in the patent [Trotta and Tumiatti 2003]. The presence of cross-linking and cyclodextrin cavities in their structure favors interaction with active molecules. Different type of molecules, either lipophilic and hydrophilic, may be encapsulated, forming either inclusion or non-inclusion complexes, and carried within the NS structure [Ma and Li 1999]. Recently, nanosponge complexes have been developed to deliver new anti-ethylene compounds to improve the cut flower vase life; and phytohormones to improve the regeneration of ornamental plants *in vitro* [Seglie et al. 2008, Seglie et al. 2011, Seglie et al. 2013]. In horticulture NS are relatively unexplored.

The aim of this study was to investigate the effect of Fe-NS in the mineral nutrition of plants. Agronomic trials were performed on hydroponically cultivated horticultural plants. Sweet corn (Strategy II) and tomato (Strategy I) has been selected as model plants [Sharma and Sanwal 1992, Hell and Stephan 2003, Zuchi et al. 2009].

MATERIALS AND METHODS

FE-NS complex synthesis. Fe-NS were developed as iron delivery system. For this purpose, β -cyclodextrins (WeckerChemie) were cross linked with pyromellitic anhydride (Sigma Aldrich) (1 : 5 molar ratio) at 90°C for 2 hours. The β -cyclodextrin was dried at 120°C for constant weight and the pyromellitic anhydride was used without further purification. Dimethyl Sulfoxide (DMSO) was used as solvent (two times the

total mass of the reagents) and triethyl amine as catalyst. The compact and dense final product (β -cyclodextrin-based nanosponges, NS), that has the tendency to become a gel in polar solvents, was smashed, purified with acetone (reflux in soxhlet for 12 hours) and milled to a fine powder.

Fe-NS was prepared stirring 18.8 g NS in 100 ml 9.4% $\text{FeSO}_4 \times 7\text{H}_2\text{O}$ water solution for 24 h. The solution was preventively acidified with HNO_3 (pH3), to prevent precipitation of Fe hydroxides.

Infrared transmission E.S.P. spectra was performed with a Nicolet Avatar 330 FT-IR through a KBr disc (4% NS w/w). pH stability was assessed by adding 0.1 M HCl and 0.1 M NaOH to a 1.85% (w/w) suspension in distilled water, measuring pH until NS complete dissolution. Thermal stability in air was investigated through a TGA Q50 V20.10 Build 36. instrument.

Considering the stability in the nutrient solution, the Fe-NS synthesis do not generate by products [Roggero et al. 2013].

Agronomic trials. Three separate experiments in hydroponical cultivation were carried out on sweet corn (*Zea mays* L., Challenger F1) and tomato (*Lycopersicon esculentum* Mill., Elegance F1). Seeds were germinated and grown in perlite. Seedlings were then transferred in glass pots contained 0.45 L of Hoagland nutrient solution (one seedling per pot). Plants were hydroponically grown in a growth chamber with a photosynthetic flux density at leaf height of $300 \mu\text{mol m}^{-2}\text{s}^{-1}$ photosynthetic active radiation and a 16 h–24°C/8 h–24°C h day/night regime. Preliminary test was conducted on plants using NS + FeSO_4 in the nutrient solution (experiment 1). Then, plant fertilization was performed by adding to nutrient solution $75.8 \mu\text{M}$ Fe as: + $\text{FeSO}_4 \times 7\text{H}_2\text{O}$ (20% Fe), +Fe-DTPA (11% Fe) and +Fe-NS (7% Fe). A randomized experimental design was organized. Each experiment followed a specific protocol according to the different aims and species and was conducted in duplicate, at once. The effect of Fe-NS was evaluated on chlorophyll, dry matter and Fe content. In all experiments, nanosponges (both NS and Fe-NS) were provided at the same concentration.

Experiment 1. NS test on sweet corn. Preliminary NS test on sweet corn was carried out to exclude problems for NS use. A total of 20 sweet corn seedlings, germinated for 12 days, were transplanted to nutrient solution added with $75.8 \mu\text{M}$ Fe as FeSO_4 . Then Half plants were cultivated in the presence of (+NS), the others in absence (-NS). Both shoots and roots were postharvest 26 days after sowing.

Experiment 2. Fertilization test on sweet corn. A total of 24 sweet corn seedlings were germinated for 12 days. Then, plant fertilization was performed by adding to nutrient solution $75.8 \mu\text{M}$ Fe as: + $\text{FeSO}_4 \times 7\text{H}_2\text{O}$ (20% Fe), +Fe-DTPA (11% Fe) and +Fe-NS (7% Fe). Root and shoot dry matter was evaluated (26 days after sowing).

Experiment 3. Fertilization test on tomato. A total of 24 tomato seedlings were germinated for 14 days. Seedlings were grown for more 21 days in a nutrient solution with $37.9 \mu\text{M}$ FeSO_4 to induce Fe-deficiency and then plant fertilization was performed by adding to nutrient solution $75.8 \mu\text{M}$ Fe as: + $\text{FeSO}_4 \times 7\text{H}_2\text{O}$ (20% Fe), +Fe-DTPA (11% Fe) and +Fe-NS (7% Fe). At the end of the growing period (49 days) dry matter (root and shoot) was measured.

Data collection. During the cultivation period biometric parameters were measured on all plants twice a week. The chlorophyll content and the re-greening effect of the

different treatment was evaluated weekly after treatments by SPAD index values of fully expanded young leaves using a portable SPAD-502 meter (Minolta, Osaka, Japan). SPAD provided a sensitive and accurate index of plant response to Fe treatment [Zuchi et al. 2009]. pH level was regularly monitored to prevent increase in alkalinity level. Shoot and root dry weight was assessed at the end of the trial (30 days after sowing) for biomass evaluation. Fe concentration in leaf tissues (dry weight) of sweet corn and tomato plants were determined by ICP-AES after digestion with concentrated HNO₃.

Statistical analysis. In exp. 1 SPAD index values and dry weights were subjected to Tukey's mean comparison test. In exp. 2 and exp. 3 data were subjected to one-way analysis of variance (ANOVA) and the Ryan-Einot-Gabriel-Welsch (F) post hoc test for multiple comparisons ($P < 0.05$). All the data were computed by means of the SPSS statistical package (version 21.0; SPSS Inc., Chicago, USA).

RESULTS AND DISCUSSION

FE-NS complex synthesis. A new iron fertilizer were obtained using β -cyclodextrin-based nanosponge complex (Fe-NS). Fe-NS resulted to be chemically and physically stable between 3–10 pH range and to air heat until 300°C. FT-IR transmission spectra showed the stretching ester-carbonyl band at 1740 cm⁻¹ and the stretching single bond Carbon-Oxygen band at 1200 cm⁻¹ [Roggero et al. 2013].

Agronomic trials. Before treatments Fe-deficiency symptoms were observed in plants. Young leaves showed chlorosis and plant growth was compromised. The application of Fe-fertilizers was provided to correct Fe deficiency symptoms and a better plant development, validating the importance of Fe role in plant nutrition [Abadía 2011].

Experiment 1. NS test on sweet corn. Plants cultivated in nutrient solution with FeSO₄, added or not with NS, grew similarly and did not show any visual damages. The SPAD index value of (+NS) plants was higher than (-NS) plants (39.44 and 33.49, respectively). No significant differences were observed. Also in terms of dry weight no statistical differences were remarked among treatments (shoot: (+NS) 0.91 g, (-NS) 0.92 g; root: (+NS) 0.91 g, (-NS) 0.92 g).

Experiment 2. Fertilization test on sweet corn. Initially, plants were kept in Fe-deficiency conditions and showed leaf chlorosis (mean SPAD index value 12.03). When Fe-fertilizers were applied, SPAD index value began to increase up to 21.9 (day 14). This type of time-course of leaf SPAD well represents the re-greening and it is comparable to the one observed in other crops [El-Jendoubi 2011]. No significant differences were observed among treatments according to the different Fe compounds (tab. 1). However, different trends were highlighted. Seven days after treatments, the highest increase in SPAD index values was observed in Fe-NS (20.69) while the lowest in FeSO₄ (15.32). After 14 days, a further increase for Fe-NS and FeSO₄ was registered, while Fe-DTPA decreased. Regarding dry matter values, significant differences were observed among treatments (tab. 1). The dry weight of Fe-NS plants was higher than the one of plants in FeSO₄ (shoot and root) and Fe-DTPA (shoot). Statistical differences were remarked also in Fe content. Fe-NS and FeSO₄ leaves showed higher Fe content (152 and 149 ppm, respectively) than Fe-DTPA (77 ppm) (tab. 1).

Table 1. SPAD index value (7 and 14 days after treatment), dry matter and Fe content (after 26 days) on sweet corn plants grown with +FeSO₄, +Fe-DTPA, +Fe-NS in the nutrient solution

Treatment	SPAD index value		Dry matter (g)		Fe content (ppm)
	7 days	14 days	root	shoot	leaf
+FeSO ₄	15.32	16.31	0.26 b	0.56 b	149 a
+Fe-DTPA	17.19	14.39	0.47 a	0.52 b	77 b
+Fe-NS	20.69	21.90	0.48 a	1.01 a	152 a
<i>P</i> [†]	ns	ns	*	*	**

† ns, * and **, non significant or significant at $P \leq 0.05$ or $P \leq 0.01$, respectively

Experiment 3. Fertilization test on tomato. The imposition of Fe-deficiency caused visible symptoms of Fe-deficiency (at leaf level: yellowing of full expanded apical leaves (mean SPAD index value 29.98); at root level: proliferation of lateral roots, increase in the diameter of the sub-apical zone and amplified root hair formation) [Tomasi et al. 2009]. When Fe-fertilizers were applied, SPAD index values began to increase. No significant differences were observed among treatments both at 7 and 14 days (tab. 2). On the contrary, significant differences were highlighted in shoot dry matter and Fe content. Fe-NS shoot dry weight was statistically different from Fe-DTPA treatments (1.88 g). No statistical differences were observed in root dry weight. About Fe content Fe-SO₄ and Fe-NS plants and leaves showed higher Fe content than Fe-DTPA (tab. 2).

Table 2. SPAD index value (7 and 14 days after treatment), dry matter and Fe content on tomato plants (after 49 days) grown with +FeSO₄, +Fe-DTPA, +Fe-NS in the nutrient solution

Treatment	SPAD index value		Dry matter (g)		Fe content (ppm)
	7 days	14 days	root	shoot	leaf
+FeSO ₄	40.28	33.48	0.53	1.63 ab	182 a
+Fe-DTPA	40.97	35.29	0.43	1.53 b	120 c
+Fe-NS	41.63	37.70	0.44	1.88 a	170 b
<i>P</i> [†]	ns	ns	ns	*	**

† ns, * and **, non significant or significant at $P \leq 0.05$ or $P \leq 0.01$, respectively

Overall, in the fertilization tests, FeSO₄ treatment was effective. Other Researchers demonstrated that the addition of FeSO₄ increases grain yield of corn grown on Fe-deficient soil too [Mathers 1970, Hergert et al. 1996]. Under the condition of the experiments FeSO₄ probably remained available to the plants and induced a better plant growth and a higher Fe content in leaves. The effectiveness of Fe sulfate may be im-

proved by the addition of organic substrates able to complex the Fe [Rombolà and Tagliavini 2007]. As we observed in Exp.1 +NS could act complexing FeSO₄.

Fe-DTPA was less effective than inorganic forms in plant growth and Fe-content and more expensive especially for low-value crops such as corn because treatments often have to be repeated several times during a growing season [Godsey et al. 2003]. Fe-DTPA is a stable complex. In this study Fe was not available for plants even if Fe-fertilizer was much more present in the nutrient solution. Furthermore their use has significant environmental limitations because they have a little residual effect due to their high solubility and eventual leaching [Nowack 2002, Sánchez-Alcalá et al. 2012]. Although the tested species have two different Fe uptake strategies, Fe-NS supplied in the nutrient solution had a positive effect on re-greening, dry matter and Fe-content both in sweet corn and in tomato plants. Moreover Fe-NS ensure a good plant growth probably due to gradually release nutrients and microelements according with plant needs.

CONCLUSIONS

In conclusion, Fe-NS can be considered as an alternative Fe-fertilizer in horticulture. This pilot investigation opens the possibilities of using Fe-NS as new alternative fertilizer in horticulture, although the comprehension of the mechanism of action needs further elucidations. Future commercial use of Fe-NS as fertilizer will also require studies on different NS types to evaluate optimization. The use of NS complexes could also be broadened for transporting other mineral nutrients.

ACKNOWLEDGMENTS

A PhD scholarship for the first author was provided by Fondazione CRT and Sea Marconi Technologies s.a.s. We thank Prof. Marco Devecchi, University of Torino (Italy) for precious advices.

REFERENCES

- Abadía, J., Vázquez, S., Rellán-Álvarez, R., El-Jendoubi, H., Abadía, A., Álvarez-Fernández, A., López-Millán, A.F. (2011). Towards a knowledge-based correction of iron chlorosis. *Plant Physiol. Biochem.*, 49(5), 471–482.
- Abadía, J., Álvarez-Fernández, A., Rombolà, A.D., Sanz, M., Tagliavini, M., Abadía, A. (2004). Technologies for the diagnosis and remediation of Fe deficiency. *Soil Sci. Plant Nutr.* 50, 965–971.
- Bhattacharya, I., Bandyopadhyay, S., Varadachari, C., Ghosh, K. (2007). Development of a novel slow-releasing iron-manganese fertilizer compound. *Ind. Eng. Chem.*, 46, 2870–2876.
- Chandra, P.K., Ghosh, K., Varadachari, C. (2009). A new slow-releasing iron fertilizer. *Chem. Eng. J.*, 155, 451–456.

- Chohura, P., Komosa, A. (2003). Nutrition status of greenhouse tomato in nutrition status of greenhouse tomato grown in inert media. Part II. Microelements. *Acta Sci. Pol. Hortorum Cultus*, 2(2), 3–13.
- El-Jendoubi, H., Melgar, J.C., Álvarez-Fernández, A., Sanz, M., Abadía, A., Abadía, J. (2011). Setting good practice to assess the efficiency of iron fertilizers. *Plant Physiol. Biochem.*, 49 (5), 483–488.
- Godsey, C.B., Schmidt, J.P., Schlegel, A.J., Taylor, R.K., Thompson, C.R., Gehl, R.J. (2003). Correcting Fe deficiency in corn seed row-applied Fe sulfate. *Agron. J.* 95, 160–166.
- Hell, R., Stephan, U.W. (2003). Iron uptake, trafficking and homeostasis in plants. *Planta*, 216, 541–555.
- Hergert, G.W., Nordquist, P.T., Petersen, J.L., Skates, B.A. (1996). Fertilizer and crop management practices for improving maize yields on high pH soil. *J. Plant Nutr.*, 19, 1223–1233.
- López-Millán, A.F., Morales, F., Gogorcena, Y., Abadía, A., Abadía, J. (2009). Metabolic responses in iron deficient tomato plants. *J. Plant Physiol.*, 166, 375–384.
- Lucena, J.J. (2007). Synthetic Iron Chelates to correct iron deficiency in plants. In: Barton L.L., Abadía J. (eds). *Iron nutrition in plants and rhizospheric microorganisms*. Springer, Dordrecht, The Netherlands, pp. 103–128.
- Ma, M., Li, D. (1999). New organic nanoporous polymers and their inclusion complexes. *Chem. Mater.*, 11, 872–876.
- Marschner, H., Römheld, V., Kissel, M. (1986). Different strategies in higher-plants in mobilization and uptake of iron. *J. Plant Nutr.*, 9, 695–713.
- Marschner, H. (1995). *Mineral nutrition of higher plants*. Academic Press, Cambridge, U.K., 313–324.
- Martínez-Cuenca, M.R., Forner-Giner, M.A., Domingo, J.I., Primo-Millo, E., Legaz, F. (2013). Strategy I responses to Fe-deficiency of two *Citrus* rootstocks differing in their tolerance to iron chlorosis. *Sci. Hortic.*, 153, 56–63.
- Mathers, A.C. (1970). Effect of Ferrous sulfate and sulfuric acid on grain sorghum yields. *Agron. J.*, 62, 555–556.
- Nowack, B. (2002). Environmental chemistry of aminopolycarboxylate chelating agents. *Environ. Sci. Technol.*, 36, 4009–4016.
- Roggero, C.M., Dicarolo, S., Tumiatti, V., Tumiatti, M., Devecchi, M., Scariot, V., Kapila, S. (2013). WO 2013046165 A1. Use of functionalised nanosponges for the growth, conservation, protection and disinfection of vegetable organisms.
- Rombolà, A.D., Toselli, M., Carpintero, J., Ammari, T., Quartieri, M., Torrent, J., Marangoni, B. (2003). Prevention of iron-deficiency induced chlorosis in kiwifruit (*Actinidia deliciosa*) through soil application of synthetic vivianite in a calcareous soil. *J. Plant Nutr.*, 26, 2031–2041.
- Rombolà, A. D., Tagliavini, M. (2007). Iron nutrition of fruit tree crops. In: Barton, L.L., Abadía, J. (eds). *Iron nutrition in plants and rhizospheric microorganisms*. Springer, Dordrecht, The Netherlands, 61–83.
- Sánchez-Alcalá, I., Bellón, F., del Campillo, M.C., Barrón, V., Torrent, J. (2012). Application of synthetic siderite (FeCO₃) to the soil is capable of alleviating iron chlorosis in olive trees. *Shi. Hortic.*, 138, 17–23.
- Sharma, S., Sanwal, G.G. (1992). Effect of Fe deficiency on the photosynthetic system of maize. *J. Plant Physiol.*, 140(5), 527–530.
- Seglie, L., Devecchi, M., Trotta, F., Chiavazza, P.M., Dolci, M. (2008). Effects of nanosponges including new anti-ethylene compounds on post harvest longevity of cut flowers and of nanosponges including phytohormones on *in vitro* regeneration of ornamental species. *Proceeding Book of “The 14th International Cyclodextrins Symposium”*, 8–11 May, Kyoto, Japan.

- Seglie, L., Martina, K., Devecchi, M., Roggero, C., Trotta, F., Scariot, V. (2011). β -Cyclodextrin-based nanosponges as carriers for 1-MCP in extending the postharvest longevity of carnation cut flowers: an evaluation of different degrees of cross-linking. *Plant Growth Regul.*, 65(3), 501–511.
- Seglie, L., Devecchi, M., Trotta, F., Scariot, V. (2013). β -cyclodextrin-based nanosponges improve 1-MCP efficacy in extending the postharvest quality of cut flowers. *Shi. Hortic.*, 159, 162–165.
- Tomasi, N., Rizzardo, C., Monte, R., Gottardi, S., Jelali, N., Terzano, R., Vekemans, B., De Nobili, M., Varanini, Z., Pinton, R., Cesco, S. (2009). Micro-analytical, physiological and molecular aspects of Fe acquisition in leaves of Fe-deficient tomato plants re-supplied with natural Fe complexes in nutrient solution. *Plant Soil*, 325, 25–38.
- Trotta, F., Tumiatti, V. (2003). Cross-linked polymers based on cyclodextrin for removing polluting agents. WO 03/085002.
- Zuchi, S., Cesco, S., Varanini, Z., Pinton, R., Astolfi, S. (2009). Sulphur deprivation limits Fe-deficiency responses in tomato plants. *Planta*, 230(1), 85–94.

WSTĘPNE BADANIA DOTYCZĄCE WPŁYWU KOMPLEKSU NANOGĄBKII Fe W OGRODNICTWIE

Streszczenie. Niedobór żelaza jest głównym zaburzeniem odżywczym roślin uprawnych. Nieorganiczne związki Fe oraz syntetyczne chelaty Fe są powszechnie używane do walki z chlorozą, co niepokoi jednak ekologów. Opracowano więc nowy nawóz żelazowy przy użyciu kompleksu nanogąbki w oparciu o β -cyklodekstrynę (Fe-NS). W niniejszym badaniu przeprowadzono test pilotażowy na roślinach ogrodniczych w celu oceny wpływu Fe-NS. Modelowymi roślinami były słodka kukurydza i pomidor, w których przeanalizowano zawartość chlorofilu, suchej masy oraz Fe. Wpływ Fe-SN porównano z FeSO_4 i Fe-DTPA. Fe-NS miał pozytywny wpływ na nowe zazielenienie oraz wzrost roślin słodkiej kukurydzy i pomidora.

Słowa kluczowe: chloroza żelazowa, nawozy Fe, nanogąbki, słodka kukurydza, pomidor

Accepted for print: 2.02.2015

For citation: Vercelli, M., Gaino, W., Contartese, V., Gallo, L., Di Carlo, S., Tumiatti, V., Larcher, F., Scariot, V. (2015). Preliminary studies on the effect of Fe-nanosponge complex in horticulture. *Acta Sci. Pol. Hortorum Cultus*, 14(2), 51–58.