

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Pointwise decay and smoothness for semilinear elliptic equations and travelling waves

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1622543> since 2019-08-29T17:34:10Z

Published version:

DOI:10.1016/j.jmaa.2016.05.009

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

POINTWISE DECAY AND SMOOTHNESS FOR SEMILINEAR ELLIPTIC EQUATIONS AND TRAVELLING WAVES

MARCO CAPPIELLO AND FABIO NICOLA

ABSTRACT. We derive sharp decay estimates and prove holomorphic extensions for the solutions of a class of semilinear nonlocal elliptic equations with linear part given by a sum of Fourier multipliers with finitely smooth symbols at the origin. Applications concern the decay and smoothness of travelling waves for nonlinear evolution equations in fluid dynamics and plasma physics.

1. INTRODUCTION AND STATEMENT OF THE RESULTS

The aim of this note is to derive pointwise decay estimates for a class of semilinear elliptic equations in \mathbb{R}^d of the form

$$(1.1) \quad p(D)u = F(u),$$

where $F : \mathbb{C} \rightarrow \mathbb{C}$ is any measurable function satisfying

$$(1.2) \quad |F(u)| \leq C_K |u|^p$$

for some $p > 1$, uniformly for u in compact subsets $K \subset \mathbb{C}$, and $p(D)$ is a Fourier multiplier with symbol of the form

$$(1.3) \quad p(\xi) = \sum_{j=0}^h p_{m_j}(\xi)$$

where $p_{m_j} \in C^\infty(\mathbb{R}^d \setminus \{0\})$ are positively homogeneous functions of degree m_j , with $0 = m_0 < m_1 < m_2 < \dots < m_h = M$ and $h \geq 1$.

Equations of this form frequently appear in Mathematical Physics, in particular in the theory of travelling wave solutions of nonlinear evolution equations in fluid dynamics (KdV-type, long-wave-type, Benjamin-Ono and many others) or in the frame of stationary Schrödinger equations. For $d = 1$, in [4, 5], Bona and Li proved that when the symbol $p(\xi)$ is smooth and satisfies $|p(\xi)| \geq C \langle \xi \rangle^M$ for all $\xi \in \mathbb{R}$ and some $C > 0$, then every solution u of (1.1) which tends to 0 at infinity indeed exhibits an exponential decay of the form $e^{-c|x|}$ for some positive constant c . In addition, if $p(\xi)$ satisfies uniform analytic estimates on \mathbb{R} , then also u is analytic and it extends to a holomorphic function in a strip of the form

Key words and phrases. Algebraic decay, holomorphic extensions, nonlocal semilinear equations, travelling waves.

$\{x + iy \in \mathbb{C} : |y| < T\}$ for some $T > 0$. Similar results have been proved later for more general pseudodifferential equations in arbitrary dimension, see [3, 6, 7, 9, 10]. More recently in [8] the first author et al. considered the case when the functions $p_{m_j}(\xi)$ in (1.3) may present some *finite* smoothness at $\xi = 0$. As a difference with respect to the case of C^∞ symbols, a finite smoothness of the symbol may determine a loss of the exponential decay observed in [3, 4, 5, 6, 7, 9, 10] and the appearance of an algebraic decay. This phenomenon has been observed in [1, 2, 14, 15] on the travelling waves of the Benjamin-Ono equation and its generalizations, which solve equations of the form (1.1). Recently, in [8], under the assumption that $F(u)$ is a polynomial in u , the first author et al. proved L^2 -estimates for the solutions u of (1.1) such that $\langle x \rangle^{\varepsilon_0} u \in H^s(\mathbb{R}^d)$ for some $s > d/2$ and for some $\varepsilon_0 > 0$. In [11] we improved the latter result by proving sharp *pointwise* decay estimates under less restrictive assumptions. Namely, we proved that if

$$(1.4) \quad |p(\xi)| \geq C \langle \xi \rangle^M, \quad \xi \in \mathbb{R}^d, \quad \xi \neq 0,$$

for some $C > 0$ (as usual $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$), then every solution u of (1.1) such that $\langle x \rangle^{\varepsilon_0} u \in L^\infty(\mathbb{R}^d)$ for some $\varepsilon_0 > 0$ actually satisfies

$$(1.5) \quad \langle x \rangle^{d+m} u \in L^\infty(\mathbb{R}^d),$$

where m is the smallest m_j for which p_{m_j} is not smooth. This result however has been proved under the additional condition $m > 0$, namely when the zero order term $p_0(\xi)$ is smooth in \mathbb{R}^d and hence constant: $p_0(\xi) = p_0 \in \mathbb{C}$ (and $p_0 \neq 0$ by (1.4)). The decay estimate (1.5) is sharp as can be checked on some basic examples, cf. [8, 11].

It is natural to look for similar results in the presence of a non constant zero order term $p_0(\xi)$, which corresponds to $m = m_0 = 0$ in the above discussion (by letting $\xi \rightarrow 0$ in (1.4) we see that $p_0(\xi) \neq 0$ for $\xi \neq 0$). This situation occurs for instance in the study of travelling wave type solutions to nonlinear equations from fluid dynamics and plasma physics. To be definite let us consider the following equation

$$(1.6) \quad v_t + \mu H v_x + \beta H v_{xx} - \nu v_{xx} + 2v v_x = 0, \quad t \geq 0, x \in \mathbb{R},$$

where β, μ, ν are real parameters and $H = H(D)$ stands for the Hilbert transform, namely the Fourier multiplier with symbol $h(\xi) = -i \operatorname{sign} \xi$. The equation (1.6) has been treated in [13] in view of its connections with the modelization of plasma turbulence. Observe moreover that when $\beta = 1$ and $\mu = \nu = 0$, the equation (1.6) reduces to the Benjamin-Ono equation (cf. [2, 15]), whereas for $\beta = 0$ and $\mu = -1$, we obtain the Sivashinsky equation

$$(1.7) \quad v_t - H v_x - \nu v_{xx} + 2v v_x = 0, \quad t \geq 0, x \in \mathbb{R},$$

which governs the dynamics of wrinkled flame fronts, cf. [16, 18]. When looking for solutions of (1.6) of the form $v(t, x) = u(x - ct)$, $c \in \mathbb{R}$, we are reduced to consider

the equation

$$(1.8) \quad -cu' + \mu Hu' + \beta Hu'' - \nu u'' + 2uu' = 0,$$

which in turn can be written in the conservative form

$$(1.9) \quad -cu + \mu Hu + \beta Hu' - \nu u' = -u^2.$$

Notice that (1.9) is of the form (1.1), (1.3) with $F(u) = -u^2$ and

$$(1.10) \quad p(\xi) = -c - i\mu \operatorname{sign} \xi + \beta|\xi| - i\nu\xi.$$

By direct computation one can verify that if $\mu < 0, \nu > 0$ and $c = \beta\mu/\nu$, the equation (1.9) admits the solution

$$(1.11) \quad u(x) = -\frac{2\nu x + 2b\beta}{x^2 + b^2},$$

with $b = -\nu/\mu$. This solution decays like $O(|x|^{-1})$ for $|x| \rightarrow \infty$ and admits a holomorphic extension in the strip $\{x + iy \in \mathbb{C} : |y| < b\}$. Notice that the symbol $p(\xi)$ in (1.10) is given by

$$p(\xi) = \beta \left(-\frac{\mu}{\nu} + |\xi| \right) - i(\nu\xi + \mu \operatorname{sign} \xi),$$

which satisfies the condition (1.4) if, in addition to $\mu < 0, \nu > 0$, we have $\beta \neq 0$.

Motivated by the example above, in this note we prove pointwise decay estimates and holomorphic extensions for the solutions of the general equation (1.1), (1.2), (1.3), (1.4), extending the results proved in [11] to the case $m = m_0 = 0$.

Namely, we have the following results.

Theorem 1.1. *With the above notation, assume (1.2), (1.3), (1.4). Let u be a distribution solution of $p(D)u = F(u)$, satisfying $\langle x \rangle^{\varepsilon_0} u \in L^\infty(\mathbb{R}^d)$ for some $\varepsilon_0 > 0$. Then $\langle x \rangle^d u \in L^\infty(\mathbb{R}^d)$.*

We have also the following estimates for the derivatives, when F is smooth.

Theorem 1.2. *Assume (1.3), (1.4) and let $F \in C^\infty(\mathbb{C})$, with $F(0) = 0, F'(0) = 0$. Let u be a distribution solution of $p(D)u = F(u)$, satisfying $\langle x \rangle^{\varepsilon_0} u \in L^\infty(\mathbb{R}^d)$ for some $\varepsilon_0 > 0$.*

Then u is smooth and satisfies the estimates

$$(1.12) \quad \|v_{d+|\alpha|-\varepsilon} \partial^\alpha u\|_{L^\infty} < \infty, \quad \alpha \in \mathbb{N}^d,$$

for every $\varepsilon > 0$, where $v_r(x) = \langle x \rangle^r$ ($r \geq 0$).

In dimension $d = 1$, we have the slightly stronger estimate

$$(1.13) \quad \|v_{1+\alpha} \partial^\alpha u\|_{L^\infty} < \infty, \quad \alpha \in \mathbb{N}.$$

Here it is meant that F is smooth with respect to the structure of \mathbb{C} as a real vector space, and with F' we denote its differential. Observe, in particular, that the above assumptions on F imply (1.2).

In [11] we obtained the sharp decay for the derivatives of the solutions in every dimension, namely $\langle x \rangle^{d+m+|\alpha|} \partial^\alpha u \in L^\infty$, provided $m > 0$. In the present case ($m = 0$) the situation is more complicated since this case represents a critical threshold for the mapping properties of the convolution operators involved in the sequel, cf. the next Proposition 2.5. Nevertheless, the above result gives the optimal decay for the derivatives at least in dimension $d = 1$, which includes the equation (1.8), as it is evident from (1.11).

In the case when the nonlinear term $F(u)$ is a polynomial, we can also prove the analyticity of the solution and its holomorphic extension in a strip in the complex domain. We do not include the proof of this result since it can be obtained as in [11] without any modification.

Theorem 1.3. *Assume (1.3), (1.4) and let F be a polynomial in u, \bar{u} , with $F(0) = 0$, $F'(0) = 0$. Let u be a distribution solution of $p(D)u = F(u)$, satisfying $\langle x \rangle^{\varepsilon_0} u \in L^\infty(\mathbb{R}^d)$ for some $\varepsilon_0 > 0$.*

Then there exists $\varepsilon > 0$ such that u extends to a bounded holomorphic function $u(x + iy)$ in the strip $\{z = x + iy \in \mathbb{C}^d : |y| < \varepsilon\}$.

2. NOTATION AND PRELIMINARY RESULTS

In the sequel the Fourier transform of a function or temperate distribution f is normalized as

$$\widehat{f}(\xi) = \mathcal{F}f(\xi) = \int_{\mathbb{R}^d} e^{-ix\xi} f(x) dx.$$

We have already used in the Introduction the notation $p(D)$ for the Fourier multiplier

$$p(D)f = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ix\xi} p(\xi) \widehat{f}(\xi) d\xi$$

with symbol $p(\xi)$. Such an operator is a convolution operator

$$p(D)f = K * f$$

with the convolution kernel $K = \mathcal{F}^{-1}(p)$.

Let $w(x) > 0$ be a measurable function. We define the weighted Lebesgue spaces

$$L_w^\infty = \{u \in L^\infty(\mathbb{R}^d) : \|u\|_{L_w^\infty} := \|wu\|_{L^\infty} < \infty\}.$$

In particular, as already anticipated in the statement of Theorem 1.2, we will deal with the weight functions

$$v_r(x) = \langle x \rangle^r, \quad x \in \mathbb{R}^d, \quad r \geq 0.$$

We now recall the following boundedness results on weighted L^∞ spaces (see [11, Proposition 5] and its proof).

Proposition 2.1. *Let $p(D)$ be a Fourier multiplier as in (1.3), (1.4). Suppose, in addition that $p_0(\xi) = p_0$ is constant.*

Suppose that there exists $j > 0$ such that p_{m_j} is not smooth and let m be the minimum value of such m_j (in particular $m > 0$). Then

$$p(D)^{-1} : L_{v_r}^\infty \rightarrow L_{v_r}^\infty$$

continuously for every $0 \leq r \leq m + d$.

If p_{m_j} is smooth for every $j = 1, \dots, h$, the above conclusion holds for every $r \geq 0$.

Proposition 2.2. *Let $q(\xi)$ be a symbol in the class $S^{-M}(\mathbb{R}^d)$, $M > 0$, i.e. satisfying the estimates¹*

$$|\partial^\alpha q(\xi)| \leq C_\alpha \langle \xi \rangle^{-M-|\alpha|}, \quad \alpha \in \mathbb{N}^d, \xi \in \mathbb{R}^d$$

for some constants $C_\alpha > 0$. Then

$$q(D) : L_{v_r}^\infty \rightarrow L_{v_r}^\infty$$

if $r \geq 0$.

We also need the following classical result about the Fourier transform of homogeneous distributions (for a proof see e.g. [11, Proposition 4]).

Proposition 2.3. *Let $f \in C^\infty(\mathbb{R}^d \setminus \{0\})$ be positively homogeneous of degree $r \geq 0$ and $\chi \in C_0^\infty(\mathbb{R}^d)$. There exists a constant $C > 0$ such that*

$$|\widehat{\chi f}(\xi)| \leq C(1 + |\xi|)^{-d-r}, \quad \xi \in \mathbb{R}^d.$$

Actually the result in [11, Proposition 4]) was stated for positively homogeneous functions of degree $r > 0$ but exactly the same proof holds for $r = 0$.

The following result is proved in [12, Proposition 11.1.3(b)] in the case $s = r > d$, see also [11, Proposition 3].

Proposition 2.4. *Let $Af = K * f$ be a convolution operator with integral kernel $K \in L_{v_s}^\infty$, with $s > d$. Then A is bounded on $L_{v_r}^\infty$ for every $0 \leq r \leq s$.*

In the limiting case $s = d$ we have the following result, which is also a key estimate in the sequel.

Proposition 2.5. *Let $r > 0$. Let $Af = K * f$ be a convolution operator with integral kernel $K \in L_{v_d}^\infty$. Then*

$$A : L_{v_r}^\infty \rightarrow L_{w_r}^\infty$$

¹We set $\mathbb{N} = \{0, 1, 2, \dots\}$.

continuously, with

$$(2.1) \quad w_r(x) = \min\{\langle x \rangle^r / \log(1 + \langle x \rangle), \langle x \rangle^d\} \asymp \begin{cases} \langle x \rangle^r / \log(1 + \langle x \rangle) & \text{if } 0 < r \leq d, \\ \langle x \rangle^d & \text{if } r > d. \end{cases}$$

Proof. It is sufficient to prove that, for $r > 0$ there exists a constant $C > 0$ such that

$$\int_{\mathbb{R}^d} \frac{1}{\langle x - y \rangle^d} \frac{1}{\langle y \rangle^r} dy \leq C \max\left\{ \frac{\log(1 + \langle x \rangle)}{\langle x \rangle^r}, \frac{1}{\langle x \rangle^d} \right\}, \quad x \in \mathbb{R}^d.$$

We split the integral in the left-hand side in the three regions $|y| \leq |x|/2$, $|x|/2 \leq |y| \leq 2|x|$ and $|y| \geq 2|x|$.

When $|y| \leq |x|/2$ we have $|x - y| \asymp |x|$, and therefore

$$\int_{|y| \leq |x|/2} \frac{1}{\langle x - y \rangle^d} \frac{1}{\langle y \rangle^r} dy \lesssim \frac{1}{\langle x \rangle^d} \int_{|y| \leq |x|/2} \frac{1}{\langle y \rangle^r} dy.$$

Now,

$$\int_{|y| \leq |x|/2} \frac{1}{\langle y \rangle^r} dy \lesssim \begin{cases} \max\left\{1, \frac{1}{\langle x \rangle^{r-d}}\right\} & \text{if } r \neq d \\ \log(1 + \langle x \rangle) & \text{if } r = d. \end{cases}$$

Hence we obtain

$$\int_{|y| \leq |x|/2} \frac{1}{\langle x - y \rangle^d} \frac{1}{\langle y \rangle^r} dy \lesssim \begin{cases} \max\left\{\frac{1}{\langle x \rangle^d}, \frac{1}{\langle x \rangle^r}\right\} & \text{if } r \neq d \\ \log(1 + \langle x \rangle) / \langle x \rangle^d & \text{if } r = d. \end{cases}$$

When $|x|/2 \leq |y| \leq 2|x|$ we have $|y| \asymp |x|$, and therefore

$$\begin{aligned} \int_{|x|/2 \leq |y| \leq 2|x|} \frac{1}{\langle x - y \rangle^d} \frac{1}{\langle y \rangle^r} dy &\lesssim \frac{1}{\langle x \rangle^r} \int_{|x|/2 \leq |y| \leq 2|x|} \frac{1}{\langle x - y \rangle^d} dy \\ &\leq \frac{1}{\langle x \rangle^r} \int_{|y-x| \leq 3|x|} \frac{1}{\langle x - y \rangle^d} dy \\ &= \frac{1}{\langle x \rangle^r} \int_{|y| \leq 3|x|} \frac{1}{\langle y \rangle^d} dy \lesssim \frac{\log(1 + \langle x \rangle)}{\langle x \rangle^r}. \end{aligned}$$

Finally for $|y| \geq 2|x|$ we have $|x - y| \asymp |y|$ and using the assumption $r > 0$ we obtain

$$\int_{|y| \geq 2|x|} \frac{1}{\langle x - y \rangle^d} \frac{1}{\langle y \rangle^r} dy \lesssim \int_{|y| \geq 2|x|} \frac{1}{\langle y \rangle^{r+d}} dy \lesssim \frac{1}{\langle x \rangle^r}.$$

□

The following result shows that similar boundedness estimates hold for $p(D)^{-1}$, if $p(D)$ satisfies (1.3) and (1.4).

Proposition 2.6. *Let $p(D)$ be a Fourier multiplier satisfying (1.3), (1.4), and let $r > 0$. Then*

$$p(D)^{-1} = p^{-1}(D) : L_{v_r}^\infty \rightarrow L_{w_r}^\infty$$

continuously, with w_r as in (2.1).

Proof. We consider separately the low and high frequency components of $p(\xi)$. Namely, let $\chi \in C_0^\infty(\mathbb{R}^d)$, $\chi = 1$ in a neighborhood of 0. We write

$$p(\xi)^{-1} = \underbrace{(1 - \chi(\xi))p(\xi)^{-1}}_{q_1(\xi)} + \underbrace{\chi(\xi)p(\xi)^{-1}}_{q_2(\xi)}.$$

By (1.4) we have $q_1 \in S^{-M}$ and therefore by Proposition 2.2 we have

$$q_1(D) : L_{v_r}^\infty \rightarrow L_{v_r}^\infty \hookrightarrow L_{w_r}^\infty.$$

Concerning $q_2(D)$ we write

$$q_2(\xi) = \frac{\chi(\xi)}{p(\xi)} = \frac{\chi(\xi)}{p_0(\xi)} \cdot \frac{1}{1 + \sum_{j=1}^h p_{m_j}(\xi)/p_0(\xi)}$$

(as observed in Introduction, (1.4) implies $p_0(\xi) \neq 0$ for $\xi \neq 0$).

By Proposition 2.1 there exists $m > 0$ such that the second factor in the right-hand side gives rise to a bounded operator $L_{v_r}^\infty \rightarrow L_{v_r}^\infty$ if $0 < r \leq d+m$. On the other hand by Proposition 2.3 the first factor $\chi(\xi)/p_0(\xi)$ has (inverse) Fourier transform $K \in L_{v_d}^\infty$ and therefore the corresponding convolution operator is bounded $L_{v_r}^\infty \rightarrow L_{w_r}^\infty$ by Proposition 2.5. This gives

$$q_2(D) : L_{v_r}^\infty \rightarrow L_{w_r}^\infty \quad \text{for } 0 < r \leq d + m.$$

The case $r > m + d$ follows from the previous case, using the inclusions of the weighted L^∞ spaces:

$$q_2(D) : L_{v_r}^\infty \hookrightarrow L_{v_{d+m}}^\infty \rightarrow L_{w_{d+m}}^\infty = L_{w_r}^\infty \quad \text{for } r > d + m.$$

□

Remark 2.7. *Observe that Proposition 2.6 does not extend to symbols $p(D)$ of order 0, namely for $p(D) = p_0(D)$. Consider for example, in dimension 1, the symbol $p(\xi) = -i \operatorname{sign} \xi$. The corresponding operator $p(D)$ is the Hilbert transform, that is the convolution with the principal value distribution P.V. $1/x$ (up to a multiplicative constant). Now, if this operator were bounded $L_{v_r}^\infty \rightarrow L^\infty$ for some $r > 0$, then for every compact $K \subset \mathbb{R}^d$ we would have the estimate*

$$\|\operatorname{P.V.} \frac{1}{x} * f\|_{L^\infty} \leq C_K \|f\|_{L^\infty}$$

for all test functions $f \in C_0^\infty(K)$. This implies that the distribution P.V. $1/x$ has order 0, and this is not the case.

Hence, it is essential for $p(D)$ in (1.3) to have order $M > 0$.

We end this section with the the following well-known interpolation inequality, which is proved e.g. in [11, Proposition 1].

Proposition 2.8. *Given $0 \leq l \leq n$, there exists a constant $C > 0$ such that the following inequality holds. Let $I = I_1 \times \dots \times I_d \subset \mathbb{R}^d$, where each I_j , $j = 1, \dots, d$, is an interval of the form $[a, +\infty)$ or $(-\infty, a]$. Then*

$$(2.2) \quad \|D^l u\|_{L^\infty(I)} \leq C \|u\|_{L^\infty(I)}^{1-l/n} \|D^n u\|_{L^\infty(I)}^{l/n},$$

where we set $\|D^k u\|_{L^\infty(I)} := \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^\infty(I)}$, $k \in \mathbb{N}$.

In particular, if $u \in L_{v_s}^\infty$ and $D^n u \in L_{v_r}^\infty$ then $D^l u \in L_{v_\nu}^\infty$ with $\nu = (1 - l/n)s + (l/n)r$.

3. PROOFS OF THE RESULTS

Proof of Theorem 1.1. We write the equation in integral form as

$$u = p(D)^{-1}(F(u)).$$

Since the solution u is by assumption in $L_{v_{\varepsilon_0}}^\infty$, $\varepsilon_0 > 0$, we have $F(u) \in L_{v_{p\varepsilon_0}}^\infty$. Now, we can apply Proposition 2.6 with $r = p\varepsilon_0$ and we obtain $u \in L_{w_r}^\infty$. If $r > d$ then $w_r(x) \asymp \langle x \rangle^d$ and the desired result is proved. If instead $r \leq d$ we have $u \in L_{w_r}^\infty \subset L_{v_s}^\infty$, with $s = \varepsilon_0 + (p-1)\varepsilon_0/2$ (we used $\log(1 + \langle x \rangle) \lesssim \langle x \rangle^{(p-1)\varepsilon_0/2}$).

By applying this argument repeatedly, after a finite number of steps we arrive to $u \in L_{w_r}^\infty$, for some $r > d$, which concludes the proof. \square

Proof of Theorem 1.2. The proof of (1.12) is similar to that of [11, Theorem 1.2], but we provide the details for the benefit of the reader.

First of all we show that $u \in H^s$ for any $s \in \mathbb{R}$. Observe that we can write

$$u = p(D)^{-1}(F(u)).$$

By Theorem 1.1, we have $u \in L^\infty \cap L^2$, which gives $F(u) \in L^2$. Moreover the operator $p(D)^{-1}$ has symbol $p(\xi)^{-1}$, hence by (1.4) it maps continuously H^s into H^{s+M} for every $s \in \mathbb{R}$. This gives $u \in H^M \cap L^\infty$. Iterating this argument we obtain $u \in H^s$ for every $s \in \mathbb{R}$.

To prove (1.12) we argue by induction on $|\alpha|$. For $|\alpha| = 0$ we have in fact the stronger information $u \in L_{v_d}^\infty$ by Theorem 1.1. Assume now (1.12) to be true for $|\alpha| \leq N-1$ and let us prove it for $|\alpha| = N$. First of all, since $u \in \bigcap_{s \geq 0} H^s \subset L^\infty$, by Proposition 2.8 with $l = 1$, n large enough and u replaced by $D^{N-1}u$ we have that

$$(3.1) \quad D^N u \in L_{v_{d+N-1-\varepsilon}}^\infty, \quad |\alpha| = N$$

for every $\varepsilon > 0$. Let $\chi \in C_0^\infty(\mathbb{R}^d)$, $\chi = 1$ around the origin. By introducing commutators in the equation (1.1) we have, for $|\beta| \leq |\alpha| = N$:

$$(3.2) \quad x^\beta \partial^\alpha u = \underbrace{p(D)^{-1}[(\chi p)(D), x^\beta] \partial^\alpha u}_{=:q_1(D)u} + \underbrace{p(D)^{-1}[(1-\chi)p(D), x^\beta] \partial^\alpha u}_{=:q_2(D)u} + p(D)^{-1}(x^\beta \partial^\alpha F(u)).$$

To obtain (1.12) it is sufficient to prove that the three terms in the right-hand side of (3.2) are in $L_{v_{d-\varepsilon}}^\infty$ for every $\varepsilon > 0$. We first consider $q_1(D)u$. By direct computation we can write

$$[(\chi p)(D), x^\beta] \partial^\alpha u = \sum_{0 \neq \gamma \leq \beta} (-i)^{|\gamma|} \binom{\beta}{\gamma} (\partial_\xi^\gamma (\chi p))(D) (x^{\beta-\gamma} \partial^\alpha u),$$

where the derivatives of χp in the right-hand side are meant in the sense of distributions. By the transposed Leibniz formula² we obtain (since $|\beta| \leq |\alpha|$ and $\gamma \neq 0$)

$$(3.3) \quad [(\chi p)(D), x^\beta] \partial^\alpha u = \sum_{0 \neq \gamma \leq \beta} \sum_{\tilde{\alpha}, \tilde{\beta}: |\tilde{\beta}| \leq |\tilde{\alpha}| < |\alpha|} C_{\alpha, \beta, \tilde{\alpha}, \tilde{\beta}} (\partial_\xi^\gamma (\chi p))(D) \partial^{\tilde{\gamma}} (x^{\tilde{\beta}} \partial^{\tilde{\alpha}} u).$$

where $\tilde{\gamma}$ is a suitable multi-index depending on $\alpha, \beta, \tilde{\alpha}, \tilde{\beta}, \gamma$, with $|\tilde{\gamma}| = |\gamma|$, and $C_{\alpha, \beta, \tilde{\alpha}, \tilde{\beta}}$ are suitable constants. Moreover we have $x^{\tilde{\beta}} \partial^{\tilde{\alpha}} u \in L_{v_{d-\varepsilon}}^\infty$ by the inductive hypothesis. Now we observe that the operator $\partial_\xi^\gamma (\chi p)(D) \partial^{\tilde{\gamma}}$ has a symbol given by a sum of homogeneous functions of nonnegative degree multiplied by cut-off functions, hence by Proposition 2.3 we deduce that its integral kernel belongs to $L_{v_d}^\infty$, so that it maps $L_{v_{d-\varepsilon}}^\infty \rightarrow L_{w_{d-\varepsilon}}^\infty \hookrightarrow L_{v_{d-2\varepsilon}}^\infty$ by Proposition 2.5. By Proposition 2.6 the same happens for $p(D)^{-1} : L_{v_{d-2\varepsilon}}^\infty \rightarrow L_{v_{d-3\varepsilon}}^\infty$, and therefore we have $q_1(D) : L_{v_{d-\varepsilon}}^\infty \rightarrow L_{v_{d-3\varepsilon}}^\infty$ for every $\varepsilon > 0$, and therefore $q_1(D)u \in L_{v_{d-\varepsilon}}^\infty$ for every $\varepsilon > 0$.

Concerning $q_2(D)u$, by the symbolic calculus we have

$$(3.4) \quad \begin{aligned} q_2(D)u &= p(D)^{-1}[(1-\chi)p(D), x^\beta] \partial^\alpha u \\ &= \sum_{0 \neq \gamma \leq \beta} (-i)^{|\gamma|} \binom{\beta}{\gamma} p(D)^{-1} (\partial_\xi^\gamma ((1-\chi)p))(D) (x^{\beta-\gamma} \partial^\alpha u). \end{aligned}$$

Using (3.1) and $|\beta - \gamma| \leq N - 1$ we see that $x^{\beta-\gamma} \partial^\alpha u \in L_{d-\varepsilon}^\infty$. On the other hand, the multiplier $p(D)^{-1} (\partial_\xi^\gamma ((1-\chi)p))(D)$ has a smooth symbol of negative order and

²Namely,

$$x^\beta \partial^\alpha u(x) = \sum_{\gamma \leq \beta, \gamma \leq \alpha} \frac{(-1)^{|\gamma|} \beta!}{(\beta - \gamma)!} \binom{\alpha}{\gamma} \partial^{\alpha-\gamma} (x^{\beta-\gamma} u(x)).$$

therefore by Proposition 2.2 it maps $L_{v_r}^\infty$ into itself continuously if $0 < r \leq d$. Hence $q_2(D)u \in L_{d-\varepsilon}^\infty$.

Finally, concerning the nonlinear term, by the Faà di Bruno formula and the interpolation inequalities we have

$$(3.5) \quad \|D^N F(u)\|_{L^\infty(I)} \leq C \sum_{1 \leq \nu \leq N} \|F'\|_{C^{\nu-1}} \|u\|_{L^\infty(I)}^{\nu-1} \|D^\nu u\|_{L^\infty(I)}.$$

cf. [17, Formula (3.1.9)]. In this formula, we use the same notation as in Proposition 2.8, and the norm of F' is meant on the range of u .

We now estimate each term by observing that, thanks to Theorem 1.1 for $\nu > 1$ (integer) we have $|u|^{\nu-1} \in L_{v_{(\nu-1)d}}^\infty$, which implies $\langle x \rangle |u|^{\nu-1} \in L^\infty$; when $\nu = 1$ instead $|F'(u)| \leq C|u| \in L_{v_d}^\infty$, so that $\langle x \rangle F'(u) \in L^\infty$. On the other hand, we also know from (3.1) that $\partial^\alpha u \in L_{v_{d+N-1-\varepsilon}}^\infty$ if $|\alpha| = N$. We deduce by (3.5) that $D^N F(u) \in L_{v_{d+N-\varepsilon}}^\infty$. By Proposition 2.6 we get $p(D)^{-1}(x^\beta \partial^\alpha F(u)) \in L_{v_{d-\varepsilon}}^\infty \subset L_{v_{d-2\varepsilon}}^\infty$ for every $\varepsilon > 0$. The estimate (1.12) is then proved.

Let us now prove (1.13) in the case $d = 1$. We argue as in the first part of the present proof, assuming (1.13) for $|\alpha| \leq N - 1$ and prove it for $|\alpha| \leq N$. Moreover by the first part of the present proof we already know that

$$(3.6) \quad D^N u \in L_{v_{1+N-\varepsilon}}^\infty.$$

We will exploit the particularly simple structure of positively homogeneous functions in dimension $d = 1$, which allows us to eliminate the above ε -loss in the decay.

Let us estimate $q_1(D)u$ in (3.2). We observe that in (3.3) we have $x^{\tilde{\beta}} \partial^{\tilde{\alpha}} u \in L_{v_1}^\infty$ by the inductive hypothesis. Moreover we can split further the right-hand side of (3.3): by the Leibniz formula and using $p(\xi) = \sum_{j=0}^h p_j(\xi)$ we see that the operator $p(D)^{-1} \partial_\xi^\gamma (\chi p)(D) \partial^{\tilde{\gamma}}$ (now with $\tilde{\gamma} = \gamma$ because $|\tilde{\gamma}| = |\gamma|$ and $d = 1$) has as symbol i^γ times the function

$$\sum_{0 \leq \mu < \gamma} \binom{\gamma}{\mu} p(\xi)^{-1} \partial^{\gamma-\mu} \chi(\xi) \xi^\gamma \partial^\mu p(\xi) + \sum_{j=1}^h \chi(\xi) p(\xi)^{-1} \xi^\gamma \partial^\gamma p_j(\xi) + \chi(\xi) p(\xi)^{-1} \xi^\gamma \partial^\gamma p_0(\xi).$$

The first sum is a smooth compactly supported symbol and therefore by Proposition 2.2 gives rise to a bounded operator $L_{v_1}^\infty \rightarrow L_{v_1}^\infty$.

For the second term we write

$$\chi(\xi) p(\xi)^{-1} \xi^\gamma \partial^\gamma p_j(\xi) = \frac{\chi(\xi) \xi^\gamma \partial^\gamma p_j(\xi)}{p_0(\xi)} \cdot \frac{1}{1 + \sum_{j=1}^h p_{m_j}(\xi)/p_0(\xi)}$$

which therefore gives rise to a bounded operator $L_{v_1}^\infty \rightarrow L_{v_1}^\infty$ by Propositions 2.3, 2.4 and 2.1, since $\xi^\gamma \partial^\gamma p_j(\xi)/p_0(\xi)$ has degree $m_j > 0$.

Finally the last term

$$\chi(\xi)p(\xi)^{-1}\xi^\gamma\partial^\gamma p_0(\xi)$$

vanishes identically, because in dimension 1 we have $p_0(\xi) = p_0^+$ if $\xi > 0$ and $p_0(\xi) = p_0^-$ for $\xi < 0$ ($p_0^+, p_0^- \in \mathbb{C}$), so that $\partial^\gamma p_0(\xi) = (p_0^+ - p_0^-)\partial^{\gamma-1}\delta$ and $\xi^\gamma\partial^{\gamma-1}\delta = 0$. This is in fact the only point where the assumption $d = 1$ plays an essential role.

Concerning the term $q_2(D)u$ in (3.4), we can use the information (3.6) which implies $x^{\beta-\gamma}\partial^\alpha u \in L_{v_2-\varepsilon}^\infty \subset L_{v_1}^\infty$ (if $\varepsilon < 1$) because $|\beta - \gamma| \leq N - 1$, and we conclude that $q_2(D)u \in L_{v_1}^\infty$, since $p(D)^{-1}(\partial_\xi^\gamma((1 - \chi)p))(D)$ is bounded on $L_{v_1}^\infty$, as already observed.

Finally, for the nonlinear term $p(D)^{-1}(x^\beta\partial^\alpha F(u))$ in (3.2) we can argue as above, but now we use again the information (3.6) in place of (3.1) (with $d = 1$). By (3.5) we now obtain $D^N F(u) \in L_{v_2+N-\varepsilon}^\infty$. By Proposition 2.6 we have $p(D)^{-1} : L_{v_2-\varepsilon}^\infty \rightarrow L_{w_2-\varepsilon}^\infty = L_{v_1}^\infty$ (if $\varepsilon < 1$), so that $p(D)^{-1}(x^\beta\partial^\alpha F(u)) \in L_{v_1}^\infty$. This concludes the proof of (1.13). □

REFERENCES

- [1] C.J. Amick and J.F. Toland, *Uniqueness and related analytic properties for the Benjamin-Ono equation - a nonlinear Neumann problem in the plane*, Acta Math. **167** (1991), 107–126.
- [2] T.B. Benjamin, *Internal waves of permanent form in fluids of great depth*, J. Fluid Mech. **29** (1967), 559–592.
- [3] H.A. Biagioni and T. Gramchev, *Fractional derivative estimates in Gevrey classes, global regularity and decay for solutions to semilinear equations in \mathbb{R}^n* , J. Differential Equations **194** (2003), 140–165.
- [4] J. Bona and Y. Li, *Analyticity of solitary-wave solutions of model equations for long waves*. SIAM J. Math. Anal. **27** (1996), n. 3, 725–737.
- [5] J. Bona and Y. Li, *Decay and analyticity of solitary waves*, J. Math. Pures Appl. **76** (1997), 377–430.
- [6] M. Cappelletto, T. Gramchev and L. Rodino, *Semilinear pseudo-differential equations and travelling waves*, Fields Institute Communications, **52** (2007), 213–238.
- [7] M. Cappelletto, T. Gramchev and L. Rodino, *Sub-exponential decay and uniform holomorphic extensions for semilinear pseudodifferential equations*. Comm. Partial Differential Equations **35** (2010) n. 5, 846–877.
- [8] M. Cappelletto, T. Gramchev and L. Rodino, *Decay estimates for solutions of nonlocal semilinear equations*, Nagoya Math. J. **218** (2015), 175198.
- [9] M. Cappelletto and F. Nicola, *Holomorphic extension of solutions of semilinear elliptic equations*, Nonlinear Anal. **74** (2011), 2663–2681.
- [10] M. Cappelletto and F. Nicola, *Regularity and decay of solutions of nonlinear harmonic oscillators*, Adv. Math. **229** (2012), 1266–1299.
- [11] M. Cappelletto and F. Nicola, *Sharp decay estimates and smoothness for solutions to nonlocal semilinear equations*, Discrete Contin. Dyn. Syst. **36** (2016), 1869–1880.
- [12] K. Gröchenig, *Foundations of Time-Frequency Analysis*, Birkäuser, Boston, MA., 2001.

- [13] Y.C. Lee and H.H. Chen, *Nonlinear dynamical models of plasma turbulence*, Phys. Scr. **T2/1** (1982), 44-47.
- [14] M. Maris, *On the existence, regularity and decay of solitary waves to a generalized Benjamin-Ono equation*, Nonlinear Anal. **51** (2002), n. 6, 1073-1085.
- [15] H. Ono, *Algebraic solitary waves in stratified fluids*, J. Phys. Soc. Japan **39** (1975), 1082-1091.
- [16] G.I. Sivashinsky, *Nonlinear analysis of hydrodynamic instability in laminar flames. I. Derivation of basic equations*, Acta Astronaut. **4** (1977), no. 11-12, 1177-1206.
- [17] M. Taylor, *Pseudodifferential operators and nonlinear PDE*. Birkhäuser, 1991.
- [18] O. Thual, U. Frisch and M. Hénon, *Application of pole decomposition to an equation governing the dynamics of wrinkled flame fronts*, J. Physique **46** (1985) n. 9, 1485-1494.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY

E-mail address: marco.cappiello@unito.it

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

E-mail address: fabio.nicola@polito.it