

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Corrigendum to 'Population structure analysis provides insights into the infection biology and invasion strategies of *Kretzschmaria deusta* in trees'

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1638548> since 2017-05-25T16:51:14Z

Published version:

DOI:10.1016/j.funeco.2016.05.003

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution:

*Questa è la versione dell'autore dell'opera:
[Guglielmo F. et al. 2016, Fungal Ecology, 23, 172]*

The definitive version is available at:

*La versione definitiva è disponibile alla URL:
[<http://www.sciencedirect.com/science/article/pii/S1754504816300460>]*

Corrigendum to “Population structure analysis provides insights into the infection biology and invasion strategies of *Kretzschmaria deusta* in trees”

Fabio Guglielmo, Serena Michelotti, Giovanni Nicolotti ¹, Paolo Gonthier

The December 2012 edition of *Fungal Ecology* included a research article on the population structure of the wood decay fungus *Kretzschmaria deusta* in the stem bases of *Acer platanoides*, *Acer pseudoplatanus*, *Fagus sylvatica* and *Platanus x acerifolia* in northern Italy (Guglielmo et al., 2012). The Introduction of the article referred to the significance of this fungus in tree hazard assessment. In this context, it was incorrectly indicated that Lonsdale (1999) stated that “Sudden limb failures are also often associated with *Kretzschmaria deusta*”. *K. deusta* (syn. *Ustulina deusta*) is found typically in the stem bases and central roots of its host trees (Schwarze et al., 2000). It is not generally known to cause branch failures but colonisation initiated via basal wounds occasionally extends 4 m or more up the stem (Lonsdale, 1999). The above citation error came to light as a result of enquiries that followed a rare occurrence of this fungus in association with branch failure in *P. x acerifolia*. The authors would like to apologise for the misinterpretation and would like to thank Dr David Lonsdale for the clarification above.

References

- Guglielmo, F., Michelotti, S., Nicolotti, G., Gonthier, P., 2012. Population structure analysis provides insight into the infection biology and invasion strategies of *Kretzschmaria deusta* in trees. *Fungal Ecol.* 5, 714–725. <http://dx.doi.org/10.1016/j.funeco.2012.06.001>.
- Lonsdale, D., 1999. Principles of Tree Hazard Assessment and Management. Research for Amenity Trees 7. The Stationery Office, London.
- Schwarze, F.W.M.R., Engels,