

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Cross-sectional investigation on sheep sarcosporidiosis in Sardinia, Italy

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1634514> since 2017-05-16T11:05:24Z

Published version:

DOI:10.1016/j.vprsr.2016.05.004

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in VETERINARY PARASITOLOGY, 3-4, 2016, 10.1016/j.vprsr.2016.05.004.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.vprsr.2016.05.004

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S2405939015300125>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

Anna Paola Pipia a, Antonio Varcasia a, Antonella Zidda a, Giorgia Dessì a, Romina Panzalis a, Claudia Tamponi a, Raffaele Marrosu b, Gabriele Tosciri a, Giuliana Sanna a, Francesco Dore a, Francesco Chiesa c, Antonio Scala a

a Laboratory of Parasitology, Veterinary Teaching Hospital, Veterinary Department, University of Sassari, Sassari, Italy

b Azienda Sanitaria Locale (ASL) N.1, Sassari, Italy

c Department of Veterinary Science, University of Torino, Torino, Italy

Abstract

An epidemiological survey on sarcosporidiosis was carried out with a cross sectional investigation on macroscopic and microscopic *Sarcocystis* spp. in Sarda breed sheep slaughtered in different abbatoirs of Sardinia, Italy. For the macroscopic survey, muscular samples (diaphragm, abdominal and intercostals muscles, cutaneous muscles and muscles of the thigh) from 769 slaughtered Sarda sheep, oesophagus (n = 365) and laryngeal and pterygoid muscles (n = 521) were macroscopically investigated and Polymerase Chain Reaction (PCR) on selected macroscopic cysts was performed for a molecular identification of macroscopic *Sarcocystis* species. For the microscopic investigation 112 heart samples from slaughtered Sarda sheep were collected and investigated with two different protocols: unstained (compression) examination and a molecular technique. The overall prevalence of infection for macroscopic forms of sarcocysts was of 23.3% (179/769) with prevalences higher in the oesophagus (31.6%; 125/395) compared with the other investigated tissue type; two different morphotypes, classified as large oval (LO) macroscopic cysts, identified as *Sarcocystis gigantea*, and slender fusiform (SF) sarcocysts, were identified. The examination of heart samples revealed an overall prevalence of 77.7% (87/112) for *Sarcocystis* spp.; the nested-PCR analysis of heart samples allowed to identify the microscopic species, *Sarcocystis tenella* and *Sarcocystis arieticanis*, with prevalences of 95.5% (107/112) and 17.8% (5/112) respectively. Reported results highlight the high prevalence of *Sarcocystis* infection in the island and suggests the need of an improvement of control and prevention strategies for this parasitosis.

1. Introduction

Members of the genus *Sarcocystis* (Apicomplexa, Eimerinae, Coccidea, Sarcocystidae) (Dubey, 1988) are among the most common protozoan parasites of striated muscles of livestock such as cattle, sheep and goats (Mirzaei Dehaghi et al., 2013). These parasites have an obligatory two-host life-cycle with carnivores as definitive hosts and herbivores and omnivores as intermediate hosts (Tenter, 1995).

Four species of *Sarcocystis* (*Sarcocystis gigantea*, *Sarcocystis medusiformis*, *Sarcocystis tenella* and *Sarcocystis arieticanis*) have been described in sheep (Dubey et al., 1989), although two other species, *Sarcocystis mihoensis* and *Sarcocystis gracilis*-like, originally isolated in roe deer, were described also in this domestic small ruminant (Saito et al., 1997; Giannetto et al., 2005).

S. gigantea and *S. medusiformis* transmitted by felids, are generally considered non-pathogenic and produce macroscopically visible cysts (Bahari et al., 2014); while *S. tenella* and *S. arieticanis*, producing microscopic cysts (Bahari et al., 2014) are transmitted by canids and considered pathogenic (Heckerroth and Tenter, 1999). Dogs are also recognized as the definitive hosts of *S. mihoensis* and together with foxes of *S. gracilis*-like (Giannetto et al., 2005).

S. tenella, *S. gigantea* and *S. arieticanis* are distributed worldwide, while infections with *S. medusiformis* have been reported only from Australia, New Zealand, Iran and Italy (Heckerroth and Tenter, 1999; Scala et al., 2008).

Sheep become infected with *Sarcocystis* spp. by ingesting sporocysts with contaminated food or water. The presence of macroscopic *Sarcocystis* spp. in sheep, causes great concern to the meat industry as part or even the whole infected carcasses may be rejected for human consumption, resulting in serious economic losses (Dubey et al., 1988; Oryan et al., 1996). The microscopic species, on the other hand, may cause serious pathological condition in the infected animals, especially during acute forms and also result in heavy production losses (Fayer, 1976; Munday, 1979, 1986). The severity of the disease in sheep caused by *S. tenella* and *S. arieticanis* seems to be related with the dose of ingested sporocysts and of the immune status of the host (Heckerroth and Tenter, 1999). During the early multiplication of the parasites by endopolygeny, a primary infection with one of the pathogenic *Sarcocystis* species may lead to acute sarcosporidiosis with encephalitis, encephalomyelitis, and haemorrhagic diathesis which can cause the death of the animal (Heckerroth and Tenter, 1999). In pregnant sheep, acute *Sarcocystis* infection sometimes results in foetal death, abortion, or premature birth of the lamb (Munday, 1981; Fayer and Dubey, 1988). Sheep may be infected with different pathogenic and non-pathogenic *Sarcocystis* species at the same time (Gjerde, 2013).

The most common taxonomic criteria for the identification of *Sarcocystis* species, is the ultrastructure of the sarcocysts wall (Giannetto et al., 2005). Several authors have reported that the sarcocysts wall varies from being relatively simple to highly complex (Dubey et al., 1988; Obendorf and Munday, 1987; O'Toole, 1987; Mehlhorn et al., 1975).

Nevertheless in recent years molecular methods have been used for the identification of *Sarcocystis* spp., particularly those infecting domestic animals (Gjerde, 2013; Tenter, 1995). Several molecular studies on *Sarcocystis* spp., have been carried using nuclear ribosomal DNA unit, particularly the small subunit (18S) rRNA gene (Gjerde, 2013).

Several reports on the prevalence of ovine *Sarcocystis* infection have revealed that this parasitosis is still common even in developed countries (Mirzaei and Rezaei, 2014; Mirzaei Dehaghi et al., 2013; Savini et al., 1992) particularly in regions where sheep breeding is still carried out with extensive methods, like Mediterranean countries. Within this area, Sardinia plays an important role as epidemiological observatory as N 3,300,000 (Ministero della Salute, 2013) sheep are raised with traditional extensive methods in the island. Many parasitological diseases, including zoonosis, are still to date widespread in Sardinia due the isolation of animals and parasitic population due to insularity but also for political, cultural and also breeding methods (Varcasia et al., 2011).

In Sardinia, (Italy) Scala and Nieddu, 1990, reported a prevalence of 66% for *S. gigantea*. This specie is commonly found in the oesophagus and other localization (as skeletal muscles, diaphragm, heart, tongue and larynx) of slaughtered sheep (Bahari et al., 2014). Furthermore the same authors found microscopic cysts with prevalences ranging between 36% to 81% in the oesophagus and heart respectively (Scala and Nieddu, 1990).

Despite this, data on epidemiology and molecular characterization of sheep sarcosporidiosis are quite outdated and mainly present in grey scientific literature (regional papers in Italian).

Hence, the goal of this study was to fill this gap of knowledge on sheep sarcosporidiosis with a cross-sectional investigation on macroscopic and microscopic species.

2. Material and methods

2.1. Macroscopic sarcocysts

During 2013 a total of 769 Sarda breed sheep, females, aged between 3–7 years slaughtered in 4 abattoirs (Thiesi (SS) n = 282; Tula (SS) n = 157; Settimo S. Pietro (CA) n = 103; Nule (NU) n = 227) of Sardinia Island, Italy, were investigated for *Sarcocystis* infection. During the postmortem inspection, diaphragm, abdominal and intercostal muscles, cutaneous muscles and muscles of the thigh were examined for the detection of macroscopic cysts of *Sarcocystis* spp. In addition, the oesophagus and laryngeal/pterygoid muscles were investigated in 365 and 521 of these 769 sheep, respectively. Macroscopic cysts were classified according to Dubey et al. (1989).

In order to confirm the taxonomy of macroscopic cysts, a molecular study was carried out on 30 individual cysts samples isolated during the macroscopic examination. Individually isolated sarcocysts were washed twice with distilled water, placed in 1.5 ml Eppendorf tubes with ethanol 70% and stored at – 20 °C until DNA extraction with a commercial kit (PureLink® Genomic DNA Mini Kit – Invitrogen, USA), according to the manufacturer's instructions. Only the partial 28S and 18S rRNA genes were amplified using the primers pairs KL5a (5' GAC CCT GTT GAG CTT GAC 30) and KL2 (5' ACT TAG AGG CGT TCA GTC 3') and 1L (5' CCATGCATGTCTAAGTATAAGC-3') and 1H (5' -TATCCCCATCACGATGCATAC-3') as described by Mugridge et al. (1999) and Yang et al. (2001). Polymerase chain reaction (PCR) was performed in a 25 µl total volume containing 2.5 µl 10 × PCR buffer, 2.5 µl 2 mM dNTP mix, 0.1 µM each primer, 0.5 µl *Thermus aquaticus* DNA Polymerase (Thermo Scientific), 2.5 µl MgCl₂, 0.2 µg genomic DNA and the remaining volume of water. Following the initial 3 min denaturation step at 94 °C, 30 amplification cycles were carried out at 94 °C for 45 s and at 65 °C (62 °C using primer pairs 1L/1H) for 45 s, extension at 72 °C for 1 min and a final extension of 5 min at 72 °C. PCR products were purified with a commercial kit (High Pure PCR Product Purification Kit, Roche) and sequenced through an external service (MWG Eurofins). Obtained sequences were then first compared with BLAST databases (<http://blast.ncbi.nlm.nih.gov/Blast>).

2.2. Microscopic sarcocysts

During the same period, a further investigation was carried out on other 112 Sarda breed sheep coming from 4 abattoirs of Sardinia (Thiesi (SS) n = 40; Tula (SS) n = 20; Settimo S. Pietro (CA) n = 18; Nule (NU) n = 34) for the identification of microscopic *Sarcocystis* spp., with two different protocols, light microscopy of unstained samples and molecular techniques. Heart was chosen as target organ according to previous studies (Pérez-Creo et al., 2013; Wheeler et al., 1987). All samples were examined immediately after slaughtering at light microscopy for the presence of microscopic species of *Sarcocystis* as described by Fukuyo et al. (2002). For each heart samples two samples of muscle tissue were obtained; from the atrioventricular septum (AVS) and from the left ventricle (LV). About 0.5 g of muscles (2 mm × 8 mm) were cut and squashed between two glass slides and examined by light microscopy (× 100) and other 10 g of each samples were

stored for the further molecular investigations. In this step of the survey, 10 g of tissue were processed from all 112 heart samples. After homogenization, 0.05 g of each sample was processed for the DNA extraction with a commercial kit (PureLink® Genomic DNA Mini Kit – Invitrogen, USA). A nested PCR targeting the multicopy 18S rRNA, was performed on all samples as previously described by Heckeroth and Tenter (1999). The nested PCR (ST-nested-PCR) for *S. tenella* was performed using the external primer pair ST1(5′ GGA TCG CAT TAT GGT CAT-3′) AP2 (5′ CCC GGG ATC CAA GCT TGA TCC TTC TGC AGG TTC ACC TAC-3′) and the nested primer pair 8 (5′ -TTT GAC TCA ACA CGG G-3′) and ST3 (5′ CGT TGCCGC GCG TTA A-3′). For *S. arieticanis* the nested PCR was performed using the external primer pair STA (5′ -TTT CGC AAG GAA GAG GA -3′) and SA2 (5′ TGA AAC GGC GCG TAG A-3′) with the internal primers 2 (5′ AGG GTT CGA TTC CGG AG -3′) and SA1(5′ GCG GGA AGA GGAGAA T-3′). The PCR was performed in a 100 μl reaction volumes containing 10 mM Tris-HCl, 50 mM potassium chloride, 0.1% Triton X-100, 1.75 mM magnesium chloride, 0.1 mM each of deoxynucleotide triphosphate (dNTP), 100 pmol of each primer and 1.5 U of *T. aquaticus* DNA Polymerase (Thermo Scientific). Amplifications were carried out in an GeneAmp PCR System 9700 Thermal Cycler (Applied Biosystem, USA) using the following PCR protocol conditions: initial denaturation at 94 °C for 4 min, 26 cycles at 93 °C for 2 min, 57 °C for 2 min, 72 °C for 2 min, with a final extension step of 5 min to complete the process.

PCR products were separated on 1.5% agarose gel and visualized after staining with Gel Red to check for appropriately sized product. PCR products were then purified using the NucleoSpin Gel and PCR clean up (Machery-Nagel, Germany) and sent to an external sequencing service (Eurofins MWG Operon, Germany). Sequences analysis was performed as described above for macroscopic *Sarcocystis* spp.

2.3. Statistical analysis

Data were analysed using the statistical package Epi-Info (version 7.0, CDC/WHO, Atlanta, GA, USA). The Chi-squared test (χ^2) was used to determine significant differences on prevalences between the different examined tissues. Differences were considered with statistical significance when the P value was ≤ 0.05 . Diagnostic tests agreement (compression method vs PCR) was evaluated using K statistic test.

3 Results

3.1. Macroscopic sarcocysts

The overall prevalence of infection for macroscopic *Sarcocystis* spp., in examined sheep was of 23.3% (179/769). Prevalence of *Sarcocystis* spp., infection between the 4 abattoirs across the island are summarized in Table 1. In particular, it was possible to isolate two different morphotypes, identified as large oval (LO) macroscopic cysts (Fig. 1) and recognized as *S. gigantea* with a mean length of $7308.5 \pm 1686.8 \mu\text{m}$ and a mean width of $5421.25 \pm 1274.6 \mu\text{m}$ and slender fusiform (SF) sarcocysts with an average length of $7704.75 \pm 549.9 \mu\text{m}$ and a mean width of $170.25 \pm 30 \mu\text{m}$ (Fig. 2) localized especially in skeletal muscles and being similar to *S. medusiformis*.

Detailed prevalences of infection and *Sarcocystis* spp., found in the examined tissues are shown in Table 2 with the respectively Odds Ratio values. Differences were statistically significant (χ^2 with 6 degrees of freedom = 336.11; $P \leq 0.05$). Oesophagus was the most infected tissue (31.6%; 125/395), followed by abdominal muscles (20.1%; 157/ 769). Macroscopic cysts classified as LO were mostly observed in the oesophagus (31.6%; 125/395) while SF cysts were mainly found in the abdominal muscles (12.3%; 95/769). Difference was statistically significant ($\chi^2 = 63.36$; $P \leq 0.05$).

The DNA extracted from the LO (N = 5) and SF (N = 4) macroscopic cysts was used as a template for genetic characterization of the two different macroscopic sarcocyst types: primers targeting the large ribosomal subunit (*lsu*) and the small ribosomal subunit of the rRNA gene yielded an 800 bp and 500 bp fragment, respectively, for all the samples. Type LO sarcocysts were identified as *S. gigantea* (showing a 99% nucleotide sequence identity for both genes with the sequences deposited in GenBank by Mugridge et al. (1999) and Gjerde (2013) (GenBank accession number: U85706.1 and KC209733.1) respectively. Type SF sarcocysts showed a 94% nucleotide sequence similarity with *S. gigantea* and *S. moulei* for both genes. The SF *Sarcocystis* could be considered consistent with *S. medusiformis* as described by Dubey et al. (1989) and Tenter (1995) though the sequences could not be compared with *S. medusiformis*, as sequences derived from this parasite are not present on GenBank. All sequences obtained from *S. gigantea* and *S. medusiformis* were submitted to GenBank and given the accession numbers (Genbank ID: KX223753; KX223754).

3.2. Microscopic sarcocysts

Sarcocysts were found in the 77.7% (87/112) of examined heart samples; no statistical difference in the frequency of sarcocysts between the two examined sites, AVS (63.4%; 71/112) and LV (51.8%; 58/112) was observed ($\chi^2 = 3.09$; $P \geq 0.05$). In Table 2 are detailed prevalences of infection found in the different abattoirs.

The nested-PCR analysis of heart samples allowed to identify the microscopic species *S. tenella* and *S. arieticanis*, with prevalences of 95.5% (107/112) and 17.8% (5/112) respectively; the difference between the prevalences of the two species was statistically significant ($\chi^2 = 137.63$; $P < 0.05$).

The alignment of our sequences with BLAST showed an homology of 100% and 99% within our samples sequences (Genbank ID KX223751; KX223752) and those of *S. tenella* (KP263759.1) with *S. arieticanis* (L24382.1) respectively, available in GenBank.

The 79.5% (83/112) of the examined heart samples tested positive both at the light microscopy method and PCR. The 21.4% (24/112) of heart samples were positive to PCR and negative to light microscopy. Conversely, the 3.6% (4/112) of heart samples were positive at light microscopy and at the same time PCR negative. No agreement between the two test was observed ($K < 0.01$).

4. Discussion

The results herein reported indicate the high distribution of macroscopic *Sarcocystis* spp., infection in slaughtered sheep of Sardinia, Italy. The morphological classification of the cysts allowed us to identify two macroscopic forms identified as LO and SF with morphological features comparable with those described by Dubey et al. (1989) and Tenter (1995) for *S. gigantea* and *S. medusififormis*, respectively. LO cysts were mainly localized in the oesophagus (57%) while SF cysts in the abdominal muscles (12.3%). This finding is in accordance with those reported by other authors (Oryan et al., 1996) and suggests that during the official inspection of the carcasses at slaughterhouses, abdominal muscles should be primarily inspected by the veterinary for *Sarcocystis* spp., infection. Molecular methods applied on the two morphotypes enabled to characterize for the first time in the island LO cysts as *S. gigantea* while it was not possible to confirm the morphological description of SF cysts as *S. medusififormis* due to the lack of sequences in the databases. Furthermore with the present survey it was possible to report and characterized by the molecular biology, for the first time in Sardinia the presence of the microscopic species, *S. tenella* and *S. arieticanis*.

On the evidence of the good performances obtained by the use of the light microscopy unstained (77.7% of prevalences vs 95.5% for the nestedPCR; χ^2 Yates corrected = 13.89; $P < 0.05$) this technique simple, inexpensive and rapid could be considered as a useful tool suitable in the routine laboratory practice. On the other hand PCR assays may increase the detection sensitivity of *Sarcocystis* spp. and contribute to diagnostic precision (Mirzaei Dehaghi et al., 2013). The specificity of the 18S rRNA was previous confirmed by many authors that considered as a powerful tool for species-specific differentiation of the ovine *Sarcocystis* species (Heckerroth and Tenter, 1999). In our study *S. tenella*, one of the most pathogenic species in sheep (Dubey, 1988), was reported with a high prevalence (95.5%). Similar results had previously reported in other countries: 91.7% in Romania (Titilincu et al., 2008); 86.5% in Turkey (Beyazit et al., 2007), France 94.8% (Diéz-Baños, 1978), 87.6% in Slovaki (Mala and Baranova, 1995) 96.9% in Mongolia (Fukuyo et al., 2002); 97.0% in Iraq (Latif et al., 1999); and 76% in Italy (Giannetto et al., 2005).

5. Conclusion

The presence of macroscopic and microscopic sarcocysts in Sarda sheep is an important epidemiological data furthermore their presence seems to be evenly spread across the island. The lower frequency of macroscopic sarcocysts infection herein reported (23.3%) compared with microscopic sarcocysts (95.5%) may be due to the fact that Sardinian sheep for breeding reasons have more contact with dogs (shepherd dogs) than with cats (Varcasia et al., 2011). On the other hand *Sarcocystis* transmitted by felids play an important role for the monitoring of another important zoonosis like *Toxoplasma gondii*, that have the same lifecycle.

Although the macroscopic species seems to play a marginal pathogenic role, their presence in various organs causes important economic losses in the meat industry (Oryan et al., 1996) because during the official veterinary control at slaughterhouses, carcasses with a massive infection with macroscopic cysts must be condemned by veterinarians (Bahari et al., 2014; Reg CE 854/2004). On the other hand the high prevalences of the microscopic species could have an important impact on sheep production (Oryan et al., 1996) due to the negative effects on the growth and weight gain of the animals with heavy production losses and their role as a cause of abortion (Munday, 1979, 1986; Fayer and Dubey, 1988).

For these reasons, *Sarcocystis* spp., in sheep should be routinely investigated during clinical practice by practitioners, considering also that 47% of sheep abortions in Sardinia remains without a specific diagnosis (Firinu, 1989).

Molecular diagnosis has proved to be as a sensitive technique for specie-specific differentiation of *Sarcocystis* species and also other sheep protozoa like *Toxoplasma* and *Neospora* (Tamponi et al., 2015) confirming the importance of the molecular techniques for epidemiological and diagnostic surveys. Reported results highlight the high prevalence of *Sarcocystis* spp., infection in the island and suggests the need of an improvement of control and prevention strategies for these parasites.

Conflict of interest

The authors declare there is no conflict of interest which affects the outcome of this paper.

Acknowledgements

The research was partially funded by Fondazione Banco di Sardegna, Prot. U140.2015/AI.113.MGB.

References

1. Bahari, P., Salehi, M., Seyedabadi, M., Mohammadi, A., 2014. Molecular identification of macroscopic and microscopic cysts of *Sarcocystis* in sheep in North Khorasan Province, Iran. *Int. J. Mol. Cell. Med.* Winter 3 (1), 51–56.
2. Beyazit, A., Yaziciog˘lu, O.˘., Karaer, Z., 2007. The prevalence of ovine *Sarcocystis* species in Izmir province. *Ank. Univ. Vet. Fak. Derg.* 54, 111–116.
3. Diéz-Baños, P., 1978. Sobre la prevalencia de la *Sarcosporidiosis* ovina en la provincia de Leon, con un estudio comparative de diversos metode diagnosticos. *An. Fac. Vet. Leon.* 24, 195–199.
4. Dubey, J.P., 1988. Lesions in sheep inoculated with *Sarcocystis tenella* sporocysts from canine feces. *Vet. Parasitol.* 26, 237–252.
5. Dubey, J.P., Lindsay, D.S., Speer, C.A., Fayer, R., Livingstone Jr., C.W., 1988. *Sarcocystis arieticanis* and other *Sarcocystis* species in sheep in the United States. *J. Parasitol.* 74, 1033–1038.
6. Dubey, J.P., Speer, C.A., Fayer, R., 1989. *Sarcocystosis of Animals and Man.* CRC Press, Boca Raton.
7. Fayer, R., 1976. Economic losses to sarcocystosis. *Natl. Wool Grow.* 66, 22–28.
8. Fayer, R., Dubey, J.P., 1988. *Sarcocystis* induced abortion and fetal death. *Prog. Clin. Biol. Res.* 281, 153–164.
9. Firinu, A., 1989. Gli aborti ovi-caprini in Europa e Nord Africa. *Atti del Seminario di studio su: “Gli aborti degli ovini e dei caprini”.* Quaderni dell'Istituto Zooprofilattico Sperimentale della Sardegna.
10. Fukuyo, M., Battsetseg, G., Byamba, B., 2002. Prevalence of *Sarcocystis* infection in meatproducing animals in Mongolia. *Southeast Asian J. Trop. Med. Public Health* 33, 490–495.
11. Giannetto, S., Poglayen, G., Brianti, E., Gaglio, G., Scala, A., 2005. *Sarcocystis gracilis*-like sarcocysts in a sheep. *Vet. Rec.* 156, 322–323.
12. Gjerde, B., 2013. Phylogenetic relationships among *Sarcocystis* species in cervids, cattle, and sheep inferred from the mitochondrial cytochrome c oxidase subunit I gene. *Int. J. Parasitol.* 43, 579–591.
13. Heckerth, A.R., Tenter, A.M., 1999. Development and validation of species-specific nested PCRs for diagnosis of acute sarcocystosis in sheep. *Int. J. Parasitol.* 29, 1331–1349.
14. Latif, B., Al-Delemi, J., Mohammed, B., Al-Bayati, S., Al-Amiry, A., 1999. Prevalence of *Sarcocystis* spp. in meat producing animals in Iraq. *Vet. Parasitol.* 84, 599–608.
15. Mala, P., Baranova, M., 1995. Diagnosis of *Sarcocystis* infection in slaughter animals at veterinary meat inspection. *Vet. Med.* 40, 97–100.
16. Mehlhorn, H., Heydorn, A.O., Gestrich, R., 1975. Licht und elektronenmikroskopische Untersuchungen an Cysten von *Sarcocystis ovis* Heydorn et al. (1975) in der Musculatur von Schafen. *Z. Parasitenkd.* 48, 83–93.
17. Ministero della Salute, 2013. Anagrafe Zootecnica istituita presso il CSN dell’Istituto “G.Caporale” di Teramo.
18. Mirzaei, M., Rezaei, H., 2014. The role of sheep in the epidemiology of *Sarcocystis* spp. in Tabriz area northwest of Iran. *J. Parasit. Dis.* <http://dx.doi.org/10.1007/s12639-014-0495-6>.
19. Mirzaei Dehaghi, M., Fallahi, M., Sami, M., Radfar, M.H., 2013. Survey of *Sarcocystis* infection in slaughtered sheep in Kerman Abattoir, Kerman, Iran. *Comp. Clin. Pathol.* 22, 343–346.
20. Mugridge, N.B., Morrison, D.A., Johnson, A.M., Luton, K., Dubey, J.P., Votýpka, J., Tenter, A.M., 1999. Phylogenetic relationships of the genus *Frenkelia*: a review of its history and new knowledge gained from comparison of large subunit ribosomal ribonucleic acid gene sequences. *Int. J. Parasitol.* 29, 957–972.
21. Munday, B.L., 1979. The effect of *Sarcocystis ovis* on growth rate and haematocrit in lambs. *Vet. Parasitol.* 5, 129–135.
22. Munday, B.L., 1981. Premature parturition in ewes inoculated with *Sarcocystis ovis*. *Vet. Parasitol.* 9, 17–26.
23. Munday, B.L., 1986. Effects of different doses of dog derived *Sarcocystis* sporocysts on growth rate and haematocrit in lambs. *Vet. Parasitol.* 21, 21–24.
24. Obendorf, D.L., Munday, B.L., 1987. Experimental infection with *Sarcocystis medusiformis* in sheep. *Vet. Parasitol.* 24, 59–65.

25. Oryan, A., Moghaddar, N., Gaur, S.N.S., 1996. The distribution pattern of *Sarcocystis* species, their transmission and pathogenesis in sheep in Fars province of Iran. *Vet. Res. Commun.* 20, 243–253.
26. O'Toole, D., 1987. Experimental ovine sarcocystosis: sequential ultrastructural pathology in skeletal muscle. *J. Comp. Pathol.* 97, 51–60.
27. Pérez-Creo, A., Panadero, R., López, C., Díaz, P., Vázquez, L., Díez-Baños, P., Morrondo, P., 2013. Prevalence and identity of *Sarcocystis* spp. in roe deer (*Capreolus capreolus*) in Spain: a morphological study. *Res. Vet. Sci.* 95, 1036–1040.
28. Saito, M., Shibata, Y., Kubo, M., Itagaki, H., 1997. *Sarcocystis mihoensis* n. sp. from sheep in Japan. *J. Vet. Med. Sci. Jpn.* 59, 103–106.
29. Savini, G., Dunsmore, J., Robertson, I., Seneviratna, P., 1992. The epidemiology of *Sarcocystis* spp. in cattle of Western Australia. *Epidemiol. Infect.* 108, 107–113.
30. Scala, A., Nieddu, A.M., 1990. La sarcosporidiosi dell'ovino in Sardegna. *Parassitologia* 32, 249–250.
31. Scala, A., Cadeddu, F., Sechi, F., Sedda, G., Ripoché, M., Varcasia, A., Mula, P., Giobbe, M., Giannetto, S., Mulas, D., 2008. Le sarcosporidiosi ovine in Sardegna: note epidemiologiche sulle forme macroscopiche trasmesse dal gatto. *Large Anim. Rev.* 14, 223.
32. Tamponi, C., Varcasia, A., Pipia, A.P., Zidda, A., Panzalis, R., Dore, F., Dessì, G., Sanna, G., Salis, F., Bjorkmann, C., Scala, A., 2015. ISCOM ELISA in milk as screening for *Neospora caninum* in dairy sheep. *Large Anim. Rev.* 5, 213–216.
33. Tenter, A.M., 1995. Current research on *Sarcocystis* species of domestic animals. *Int. J. Parasitol.* 25, 1311–1330.
34. Titilincu, A., Mircean, V., Blaga, R., Bratu, C.N., Cozma, V., 2008. Epidemiology and etiology in sheep sarcocystosis. *Bull. UASVM Vet. Med.* 65 (2), 50–54.
35. Varcasia, A., Tanda, B., Giobbe, M., Solinas, C., Pipia, A.P., Malgor, R., Carmona, C., Garippa, G., Scala, A., 2011. Cystic echinococcosis in Sardinia: farmers' knowledge and dog infection in sheep farms. *Vet. Parasitol.* 181, 335–340.
36. Wheeler, P.R., Burkitt, H.G., Daniels, V.G., 1987. *Texto y Atlas en color de Histología Funcional*. Ed. JIMS, Barcelona (348 pp.).
37. Yang, Z.Q., Zuo, Y.X., Yao, Y.G., Chen, X.W., Yang, G.C., Zhang, Y.P., 2001. Analysis of the 18S rRNA genes of *Sarcocystis* species suggests that the morphologically similar organisms from cattle and water buffalo should be considered the same species. *Mol. Biochem. Parasitol.* 115, 283–288.




Fig. 1. Large oval (LO) macroscopic cysts.



Fig. 2. Slender fusiform (SF) sarcocysts localized in skeletal muscles.

Table 1

Prevalences of infection and *Sarcocystis* spp., found in the examined tissues.

Abattoirs	Macroscopic survey			Microscopic survey		
	Number of sheep	Number of positive sheep	P (%)	Number of sheep	Number of positive sheep	Prevalence of infection
Thiesi	282	55	19.5	40	29	72.5
Tula	157	34	21.6	20	16	80.0
Sinnai	103	26	25.2	18	14	77.8
Nule	227	64	28.2	34	28	82.3
Statistical analysis	χ^2 with 3 degrees of freedom = 5.78; P N 0.05			χ^2 with 3 degrees of freedom = 1.11; P N 0.05		

Table 2

Prevalences of macroscopic and microscopic *Sarcocystis* spp. between abattoirs across the island.

Muscle tissue	Examined sheep (n)	Positive (n)	(P) %	Odds ratio	LO cysts (<i>S. gigantea</i> -like)		SF cyst (<i>S. medusiformis</i> -like)		P (%)
					n of positive sheep	P (%)	n of positive sheep	P (%)	
Superficial muscles of the thigh	769	15	2.0%	0.33	6	0.8%	13	1.7%	
Diaphragm	769	44	5.7%	1.00	17	2.2%	31	4%	
Cutaneous muscles	769	63	8.2%	1.47	0		63	8.2%	
Intercostal muscles	769	64	8.3%	1.50	28	3.6%	47	6.1%	
Laryngeal ^a and pterygoid muscles	521	84	16.1%	3.17	84	16.1%	6	1.5%	
Abdominal muscles	769	157	20.4%	4.23	90	11.7%	95	12.3%	
Oesophagus ^a	395	125	31.6%	7.63	225	57%	5	1.3%	

^a Tissues not always available during the slaughterhouse inspection; LO cyst = large oval; SF cyst = slender fusiform (Dubey et al., 1989).