IRIS AperTO

UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A Prospective International Aspergillus terreus Survey: An EFISG, ISHAM and ECMM Joint Study

This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/1638082 since 2017-10-09T18:02:32Z

Published version:

DOI:doi.org/10.1016/j.cmi.2017.04.012

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

10 September 2026

Accepted Manuscript



A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study

Brigitte Risslegger, Tamara Zoran, Michaela Lackner, Maria Aigner, Ferran Sánchez-Reus, Antonio Rezusta, Anuradha Chowdhary, Saad Jaber Taj-Aldeen, Maiken C. Arendrup, Salvatore Oliveri, Dimitrios P. Kontoyiannis, Ana Alastruey-Izquierdo, Katrien Lagrou, Giuliana Lo Cascio, Jacques F. Meis, Walter Buzina, Claudio Farina, Miranda Drogari-Apiranthitou, Anna Grancini, Anna Maria Tortorano, Birgit Willinger, Axel Hamprecht, Elizabeth Johnson, Lena Klingspor, Valentina Arsic-Arsenijevic, Oliver A. Cornely, Joseph Meletiadis, Wolfgang Prammer, Vivian Tullio, Jörg-Janne Vehreschild, Laura Trovato, Russell E. Lewis, Esther Segal, Peter-Michael Rath, Petr Hamal, Manuel Rodriguez-Iglesias, Emmanuel Roilides, Sevtap Arikan-Akdagli, Arunaloke Chakrabarti, Arnaldo L. Colombo, Mariana S. Fernández, M. Teresa Martin-Gomez, Hamid Badali, Georgios Petrikkos, Nikolai Klimko, Sebastian M. Heimann, Jos Houbraken, Omrum Uzun, Michael Edlinger, Sonia de la Fuente, Cornelia Lass-Flörl

PII: S1198-743X(17)30220-3

DOI: 10.1016/j.cmi.2017.04.012

Reference: CMI 924

To appear in: *Clinical Microbiology and Infection*

Received Date: 24 February 2017

Revised Date: 7 April 2017

Accepted Date: 9 April 2017

Please cite this article as: Risslegger B, Zoran T, Lackner M, Aigner M, Sánchez-Reus F, Rezusta A, Chowdhary A, Taj-Aldeen SJ, Arendrup MC, Oliveri S, Kontoyiannis DP, Alastruey-Izquierdo A, Lagrou K, Lo Cascio G, Meis JF, Buzina W, Farina C, Drogari-Apiranthitou M, Grancini A, Tortorano AM, Willinger B, Hamprecht A, Johnson E, Klingspor L, Arsic-Arsenijevic V, Cornely OA, Meletiadis J, Prammer W, Tullio V, Vehreschild J-J, Trovato L, Lewis RE, Segal E, Rath P-M, Hamal P, Rodriguez-Iglesias M, Roilides E, Arikan-Akdagli S, Chakrabarti A, Colombo AL, Fernández MS, Martin-Gomez MT, Badali H, Petrikkos G, Klimko N, Heimann SM, Houbraken J, Uzun O, Edlinger M, Fuente Sdl, Lass-Flörl C, A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study, *Clinical Microbiology and Infection* (2017), doi: 10.1016/j.cmi.2017.04.012.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

RESEARCH NOTE

A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study

Brigitte Risslegger[1]*, Tamara Zoran[1]*, Michaela Lackner[1], Maria Aigner[1], Ferran Sánchez-Reus[2], Antonio Rezusta[3], Anuradha Chowdhary[4], Saad Jaber Taj-Aldeen[5], Maiken C. Arendrup[6], Salvatore Oliveri[7], Dimitrios P. Kontoyiannis[8], Ana Alastruey-Izquierdo[9], Katrien Lagrou[10], Giuliana Lo Cascio[11], Jacques F. Meis[12], Walter Buzina[13], Claudio Farina[14], Miranda Drogari-Apiranthitou[15], Anna Grancini[16], Anna Maria Tortorano[17], Birgit Willinger[18], Axel Hamprecht[19], Elizabeth Johnson[20], Lena Klingspor[21], Valentina Arsic-Arsenijevic[22], Oliver A. Cornely[23], Joseph Meletiadis[24], Wolfgang Prammer[25], Vivian Tullio[26], Jörg-Janne Vehreschild[27], Laura Trovato[28], Russell E. Lewis[29], Esther Segal[30], Peter-Michael Rath[31], Petr Hamal[32], Manuel Rodriguez-Iglesias[33], Emmanuel Roilides[34], Sevtap Arikan-Akdagli[35], Arunaloke Chakrabarti[36], Arnaldo L. Colombo[37], Mariana S. Fernández[38], M. Teresa Martin-Gomez[39], Hamid Badali[40], Georgios Petrikkos[41], Nikolai Klimko[42], Sebastian M. Heimann[43], Jos Houbraken[44], Omrum

Uzun[45], Michael Edlinger[46], Sonia de la Fuente[47], Cornelia Lass-Flörl[1]

* contributed equally

Author affiliations:

[1]Division of Hygiene and Medical Microbiology, Medical University of Innsbruck, Innsbruck/Austria, [2]Servei de Microbiologia, Hospital de la Santa Creu I Sant Pau, Barcelona/Spain, [3]Microbiologia, Hospital Universitario Miguel Servet, IIS Aragon, Universidad de Zaragoza, Zaragoza/Spain, [4]Department of Medical Mycology, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi/India, [5]Microbiology Division, Department of Laboratory Medicine and Pathology, Hamad Medical Corporation, Doha/Qatar, [6]Statens Serum Institute, Unit of Mycology, Copenhagen/Denmark & Department of Clinical Microbiology, Copenhagen University, Rigshospitalet, Copenhagen/Denmark, [7]Department of Biomedical and Biotechnological Sciences, University of Catania, Catania/Italy, [8]The University of Texas MD Anderson Cancer Center, Houston, Texas/USA, [9]National Centre for Microbiology, Instituto de Salud Carlos III, Madrid/Spain, [10]Department of Microbiology and Immunology, KU Leuven, Leuven/Belgium, [11]Unità Operativa Complessa di Microbiologia

e virologia, Dipartimento di Patologia e diagnostica, Azienda Ospedaliera Universitaria Integrata, Verona/Italy, [12]Department of Medical Microbiology and Infectious Diseases, Canisius Wilhelmina Hospital, Nijmegen/The Netherlands, [13]Institute of Hygiene, Microbiology and Environmental Medicine, Medical University of Graz, Graz/Austria, [14]Microbiology Institute, ASST Papa Giovanni XXIII, Bergamo/Italy, [15]Infectious Diseases Research Laboratory/4th Department of Internal Medicine, ATTIKON University Hospital, National and Kapodistrian University of Athens, Athens/Greece, [16]Laboratorio Centrale di Analisi Chimico Cliniche e Microbiologia, IRCCS Foundation, Cà Granda Ospedale Maggiore Policlinico, Milan/Italy, [17]Department of Biomedical Sciences for Health, Università degli Studi di Milano, Milan/Italy, [18]Department of Laboratory Medicine, Division of Clinical Microbiology, Medical University of Vienna, Vienna/Austria, [19]Institute for Medical Microbiology, Immunology and Hygiene, University of Cologne, Cologne/Germany, [20]Mycology Reference Laboratory, Public Health England, Bristol/United Kingdom, [21]Karolinska Institutet, Department of Laboratory Medicine, F 68, Karolinska University Hospital, Huddinge, Stockholm/Sweden, [22]National Reference Medical Mycology Laboratory, Institute of Microbiology and Immunology, Faculty of Medicine,

University of Belgrade, Belgrade/Serbia, [23]Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), Department I of Internal Medicine, Clinical Trials Centre Cologne (ZKS Köln), Center for Integrated Oncology (CIO Köln-Bonn), German Centre for Infection Research (DZIF), University of Cologne, Cologne/Germany, [24]Clinical Microbiology Laboratory, National Kapodistrian University of Athens, ATTIKON University Hospital Athens, Athens/Greece, [25]Department of Hygiene and Medical Microbiology, Klinikum Wels-Grieskirchen, Wels/Austria, [26]Department of Public Health and Pediatrics, Microbiology Division, Turin/Italy, [27]Department I for Internal Medicine, University Hospital of Cologne, Cologne/Germany and German Centre for Infection Research, partner site Bonn-Cologne, Germany, [28]A.O.U. Policlinico Vittorio Emanuele Catania, Biometec – University of Catania/Italy, [29]Infectious Diseases Unit, S. Orsola-Malpighi, Department of Medical and Surgical Sciences, University of Bologna, Bologna/Italy, [30]Department of Clinical Microbiology and Immunology, Sackler School of Medicine, Tel Aviv University, Tel Aviv/Israel, [31]Institute of Medical Microbiology, University Hospital Essen, University of Duisburg- Essen, Essen/Germany, [32]Department of of Microbiology, Faculty of Medicine and Dentistry, Palacky University Olomouc and University Hospital Olomouc/Czech

Republic, [33]Clinical Microbiology, Puerta del Mar University Hospital, University of Cádiz, Cádiz/Spain, [34]Infectious Diseases Unit, 3rd Department of Pediatrics, Faculty of Medicine, Aristotle University School of Health Sciences, Hippokration General Hospital, Thessaloniki/Greece, [35]Department of Medical Microbiology, Hacettepe University Medical School, Ankara/Turkey, [36]Division of Mycology, Department of Medial Microbiology, Chandigarh/India, [37]Escola Paulista de Medicina, Federal University of São Paulo, São Paulo/Brazil, [38]Departmento de Micología, Instituto de Medicina Regional, Universidad Nacional del Nordeste, CONICET, Resistencia/Argentina,[39] Division of Clinical Mycology, Department of Microbiology, Vall d'Hebron University Hospital, Barcelona, [40]Department of Medical Mycology and Parasitology/Invasive Fungi Research Center, Mazandaran University of Medical Sciences, Sari/Iran, [41]School of Medicine, European University Cyprus, Nicosia/Cyprus, [42]Department of Clinical Mycology, Allergy and Immunology, North Western State Medical University, Saint Petersburg/Russia, [43]Department I for Internal Medicine, University Hospital of Cologne, Cologne/Germany, [44]CBS-KNAW Fungal Biodiversity Centre, Utrecht/The Netherlands, [45]Hacettepe University Medical School, Department of Infectious Diseases and Clinical Microbiology, Ankara/Turkey, [46]Department of Medical

Statistics, Informatics and Health Economics, Medical University of Innsbruck, Innsbruck/Austria, [47]Department of Dermatology, Hospital Ernest Lluch Martin, Calatayud, Zaragoza/Spain

Running title: TerrNet Study

Corresponding author: Cornelia Lass-Flörl, Division of Hygiene and Medical Microbiology. University of Innsbruck, Schöpfstraße 41, 6020 Innsbruck, Austria
Tel.No. 0043 512 9003-70703, Fax No. 0043 512 9003-73700
email: cornelia.lass-floerl@i-med.ac.at

Length of abstract : 87 words
Length of paper (excluding abstract and ref.) : 1172 words

ACCEPTED MANUSCRIPT

Abstract

Objectives: A prospective international multicentre surveillance study was conducted to investigate the prevalence and amphotericin B (AMB) susceptibility of *Aspergillus terreus* species complex infections.

Methods: Three hundred seventy cases from 21 countries were evaluated.

Results: The overall prevalence of *A. terreus* species complex among patients investigated and with mold positive cultures was 5.2% (370/7116). AMB MICs were ranging from 0.125 to 32 mg/L, (median 8mg/L).

Conclusions: *A. terreus* species complex infections cause a wide spectrum of aspergillosis and the majority of cryptic species display high AMB MICs.

ACCEPTED MANUSCRIPT

Introduction:

Aspergillus terreus species complex holds an exceptional position within the aspergilli, as it appears to be a rare pathogen of infection and displays polyene resistance [1,2,3]. *A. terreus* is a common cause of invasive aspergillosis (IA) at the M. D. Anderson Cancer Center in Houston, USA, and the University Hospital of Innsbruck, Austria [3,4,5]. Almost no data are available on how frequently this species occurs elsewhere and whether differences within amphotericin B (AMB) susceptibility exist. Our objective was to investigate the global prevalence of *A. terreus* species complex in fungal diseases and to survey AMB susceptibility.

Methods:

An international surveillance network was established on behalf of the European Fungal Infection Study Group, the International Society for Human and Animal Mycology *Aspergillus terreus* working group, and the European Confederation of Medical Mycology. 38 centres from 21 countries participated. Each centre collected isolates and reported the number of *A. terreus* and fungal pathogens detected for 12 consecutive months (2014-2015). Patient characteristics', epidemiological data, and antifungal treatment were documented through an online questionnaire

using www.clinicalsurveys.net online platform. Patients were classified according to the European Organisation for the Research and Treatment of Cancer/Mycoses Study Group consensus definitions [6] by the participating centres. Unless otherwise noted, the isolation of *A. terreus* from sputa of non-neutropenic patients was categorized as colonisation. Isolates were sent to the Division of Hygiene and Medical Microbiology for molecular species identification [7,8] and susceptibility testing according to EUCAST (European Committee on Antimicrobial Susceptibility Testing) method [2]. A. *terreus* strains were identified to the cryptic species level by sequencing partial beta-tubulin and applying a validated in-house database owned by Jos Houbraken, CBS Fungal Biodiversity Center, Utrecht, The Netherlands. An AMB epidemiological cut-off value of 4mg/L was set for *A. terreus* [2].

This study was approved by the Ethics Commission of the Medical University of Innsbruck (UN4926).

Results:

461 cases were enrolled of which 91 were excluded because of insufficient patient documentation (n=45) or lack of fungal isolates (n=46) being available. Consequently, this survey comprises 370 eligible cases with an equal number of corresponding *A. terreus* isolates. Cases derived from Europe

(n=261), followed by Middle East (n=70), India (n=19), South America (n=10), and North America (n=10) (Figure 1). *A. terreus* sensu stricto (n=315), *A. citrinoterreus* (n=36), *A. alabamensis* (n=6), *A. hortai* (n=10), *A. floccosus* (n=1), and *A. neoafricanus* (n=1) were identified. One isolate (*A. terreus 1214*) was most close to *A. alabamensis* and might represent a new species. Thus, cryptic species accounted for 14.9% (55/370) with *A. citrinoterreus* (36/55, 65.5%) being dominant. AMB MICs ranged from 0.125 to 32 mg/L for *A. terreus* sensu stricto; MICs for all cryptic species were consistently higher, ranging from 2 to 32 mg/L, see Table 1. According to the EUCAST cut-off values, 194 isolates (52.4%) were classified as non-wild types. A proportion of 6.3% (n=20) of the *A. terreus* sensu stricto isolates displayed lower MICs, ranging from 0.25 – 0.5 mg/L. Isolates were predominantly acquired from Spain (n=85) and Austria (n=49), see Figure 1.

Underlying diseases e described in Table 2. Species distribution did not differ per underlying disease and specimen investigated (Table 2). Diseases comprised IA (25.1%), allergic broncho-pulmonary aspergillosis (12.4%), chronic aspergillosis (11.4%), COPD exacerbation (5.5%), aspergilloma (3.7%), otitis externa (2.5%), and wound infections (0.7%). 25.1% and 27.3% of the patients suffered from proven and probable IA, 28.6% were colonized, 10.1% had onychomycosis, and 8.9% had

mycological documented diseases such as otitis externa, aspergilloma and others.

Using a random effects model the pooled estimated proportion was 5.6% (95% CI 3.8 to 7.7) with I^2 = 92% ($p<0.0001$) and the proportions ranged from 0.0% to 58.3%. These calculations were done with MedCalc 16.8.4. Four reference centres and one centre dealing with onychomycosis only were excluded from the analysis.

A total of 68 patients received antifungal treatment at the time of fungal diagnosis, 12 were treated with AMB or liposomal - AMB. The remaining 56 received combinations of azoles and echinocandins and improved. Only one patient died due to the *A. terreus* infection. No information on outcome was available in 13 patients.

Discussion:

Infections due to *A. terreus* species complex were detected in 21 countries and 38 centres with an overall prevalence of 5.2% among mold infections. High AMB MICs were frequently observed and crossed all cryptic species. Infections were reported from all over the world with three main specific findings. Firstly, Spain and Austria were the countries with the highest density of *A. terreus* isolates collected. Secondly, the number of *A. terreus* cases enrolled varied from centre to

centre, and displayed a broad range from zero to several cases per country. Thirdly, it seems that few susceptible AMB variants exist within *A. terreus* sensu stricto.

Taking into account the differences on the environmental conditions, host related characteristics, and the use of antifungal agents, it is not possible to conclude on the particular biogeography of *A. terreus* species complex. In addition, one has to be aware that data collected may depend on the quality of care, patient demographics, infection control practices, frequency of specimen collection, and laboratory methodology. Hence, further studies are needed to determine whether specific risk and/or environmental factors are associated with infections by *A. terreus.*

Notable was the fact that *Aspergillus* section *Terrei* was most commonly isolated from patients suffering from chronic lung diseases (39.2%). No similar data have been reported [10] and it remains to be seen whether *A. terreus* reflects an emerging pathogen of this disease entity.

A. terreus is a poor target for AMB and hence is reported as resistant [2]. The role of isolates with MICs <0.5 mg/L needs further evaluation. The pharmacodynamic target may be attained with the standard AMB dose for isolates with MICs ≤0.25 mg/l [10] and infections were successfully treated with high dose liposomal-AMB [11].

Cryptic species accounted for 14.8%, with *A. citrinoterreus* being the most prevalent. Although the clinical implications of sibling species of *A. terreus* are less well understood, our study confirms that these species are generally resistant to AMB and are causing a wide spectrum of invasive and non-invasive aspergillosis. Guinea et al. [12] observed *A. citrinoterreus* acting mainly as a co-pathogen with *A. fumigatus*.

Our study has some limitations. We do not have a comprehensive worldwide *A. terreus* survey network and some countries are missing for a variety of reasons. Also, generally, the diagnosis of fungal infections is difficult to obtain and may often be based on detection of biomarkers rather than on isolation of the infecting organism. Hence, some cases may have been missed and chronic lung diseases were not specified in more detail. Further, we have no data available on co-infections which may complicate diseases. The centres included represent a convenience sample. However, this is the largest and geographically most diverse study on the contemporary epidemiology of *A. terreus* species complex infections worldwide.

Our study shows that *A. terreus* sensu stricto is widely distributed in climatically divergent countries, and that cryptic species display high AMB MICs. *A. terreus* species complex was

292 most commonly isolated from patients suffering from chronic

293 lung diseases (39.2%).

ACCEPTED MANUSCRIPT

Funding

This work was supported by ECMM, ISHAM and EFISG and in part by an unrestricted research grant through the Investigator Initiated Studies Program of Astellas and Pfizer as well as by the Austrian Science Fund (W1253-B24 doctoral program HOROS).

Transparency Declaration

We declare that we have no conflicts of interest related to this study.

CLF received research grants, travel grants or honorarium as a speaker or advisor from Gilead Sciences, Pfizer, Schering Plough, MSD and Basilea, VAA received research grants or honorarium as a speaker or advisor from Astellas, Pfizer and Schering Plough, MSD. WB received honorarium as a speaker from Pfizer, Schering Plough and MSD. CF received research grants or honorarium as a speaker or advisor from Astellas, Gilead Sciences, MSD and Basilea. AH received travel grants from Astellas and served on the advisory board of Gilead. EJ received travel grants or honorarium as a speaker or advisor from Bio-Rad, Gilead Sciences, Pfizer, Schering Plough, and MSD. DPK has received research support from Pfizer, Astellas Pharma, honoraria from Astellas, Merck, Pfizer and he serves

on the advisory boards of Merck, Amplyx, Cidara and F2G. JFM received grants from Astellas, Basilea and Merck. He has been a consultant to Astellas, Basilea and Merck and received speaker's fees from Merck, United Medical and Gilead. ALC received educational grants from Pfizer, Gilead and United Medical and research grant from Pfizer, Astellas. MCA received research grants/contract work (Paid to SSI) from Amplyx, Basilea, Cidara, F2G, Gilead, Pfizer & T2Candida, speaker honoraria from Astellas, Basilea, Gilead, MSD, Novartis, Pfizer & T2Candida. NK has received research support from Pfizer, honoraria from Astellas, Merck and Pfizer, and he serves on the advisory boards of Merck and Pfizer. OAC is supported by the German Federal Ministry of Research and Education, has received research grants from Actelion, Aranis, Astellas, AstraZeneca, Basilea, Bayer, Cidara, Duke University (NIH UM1AI104681), F2G, Gilead, GSK, Leeds University, MedPace, Melinta Therapeutics, Merck/MSD, Miltenyi, Pfizer, Rempex, Roche, Sanofi Pasteur, Scynexis, Seres Therapeutics, The Medicine Company, is a consultant to Anacor, Amplyx, Actelion, Astellas, Basilea, Cidara, Da Volterra, F2G, Gilead, Inositec AG, Janssen Pharmaceuticals, Matinas, Menarini, Merck/MSD, Paratek Pharmaceuticals, Scynexis, Seres, Summit, Vical, Vifor, and received lecture honoraria from Astellas, Basilea, Gilead and Merck/MSD. BW received

research support from Pfizer, travel grants or honorarium as a speaker or advisor from Gilead Sciences, Pfizer, Astellas, MSD and Basilea. LK grant from Gilead and has been an adviser to Astellas, Gilead, Schering-Plough, has received research grants from Gilead, Merck, Sharpe&Dohme, Schering-Plough and has received honoraria for educational lectures from Gilead, Pfizer, Merck, Sharpe & Dohme, Schering-Plough and Janssen. AMT received research support from Gilead Sciences, travel grants or honorarium as speaker from Gilead Sciences, Pfizer, Astellas. KL has received research grants from Gilead, MSD and Pfizer, received travel support from MSD, Pfizer and Gilead and received lecture honoraria from Gilead, MSD, and Pfizer. SAA received Investigator Initiated Research grant support from Pfizer and speaker honoraria or travel grants from Astellas, Gilead, Merck and Pfizer. SO received research grants from Pfizer and Astellas, honorarium as advisor from MSD, Astellas Pharma and Gilead Sciences. ER reports grants, and non-financial support from Astellas, Pfizer, Gilead, and Merck, outside the submitted work. AAI received research and/or travel grants from Gilead Sciences, F2G and Pfizer. OU was consultant for and has received travel grants from Merck, Astellas, Pfizer, and Gilead. S.M.H. has received research and travel grants from Astellas, Gilead, and MSD, research grants from Basilea, and travel grants from Pfizer; was consultant to

366 Basilea; and received honoraria as a speaker from Astellas and

367 Merck.

ACCEPTED MANUSCRIPT

References

1. Perkhofer S, Lass-Flörl C, Hell M, Russ G, Krause R, Hönigl M, et al. The Nationwide Austrian *Aspergillus* Registry: a prospective data collection on epidemiology, therapy and outcome of invasive mould infections in immunocompromised and/or immunosuppressed patients. Int J Antimicrob Agents 2010;36:531-6.
2. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope WW; European Committee on Antimicrobial Susceptibility Testing Subcommittee on Antifungal Susceptibility Testing (EUCAST-AFST). EUCAST technical note on *Aspergillus* and amphotericin B, itraconazole, and posaconazole. Clin Microbiol Infect. 2012;18:E248-50.
3. Lass-Flörl C, Grif K, Mayr A, Petzer A, Gastl G, Bonatti H et al. Epidemiology and outcome of infections due to *Aspergillus terreus*: 10-year single center experience. Br J Haematol. 2005;131:201-207.
4. Lass-Florl C, Grif K, Kontoyiannis DP. Molecular typing of *A. terreus* isolates collected in Houston, Texas and Innsbruck, Austria: evidence of great genetic diversity. J Clin Microbiol. 2007;45:2686-90.
5. Lackner M, Coassin S, Haun M, Binder U, Kronenberg F, Haas H et al. Geographically predominant genotypes of *Aspergillus terreus* species complex in Austria: a

microsatellite typing study. Clin Microbiol Infect. 2016;22:270-276.

6. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T et al. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. Clin Infect Dis 2008;46:1813-21.
7. Balajee S A, Baddley JW, Peterson SW, Nickle D, Varga J, Boey A et al., ISHAM Working Group on *A. terreus. Aspergillus alabamensis*, a new clinically relevant species in the section *Terrei*. Eukaryotic Cell 2009;8:713–22.
8. Kathuria S, Sharma C, Singh PK, Agarwal P, Agarwal K, Hagen F et al. Molecular epidemiology and in-vitro antifungal susceptibility of *Aspergillus terreus* species complex isolates in Delhi, India: Evidence of genetic diversity by amplified fragment length polymorphism and microsatellite typing. PLoS ONE 2015;10:e0118997.
9. Denning DW, Cadranel J, Beigelman-Aubry C, Ader F, Chakrabarti A, Blot S et al. Chronic pulmonary aspergillosis:

rationale and clinical guidelines for diagnosis and management. Eur Respir J. 2016;47:45-68.

10. Elefanti A, Mouton JW, Verweij, PE, Zerva L, Meletiadis J. Susceptibility breakpoints for amphotericin B and *Aspergillus* species in an in vitro pharmacokinetic-pharmacodynamic model simulating free-drug concentrations in human serum. Antimicrob Agents Chemother. 2014;58:2356-2362.
11. Lewis RE, Albert NP, Liao G, Wang W, Prince RA, Kontoyiannis DP. High-dose induction liposomal amphotericin B followed by de-escalation is effective in experimental *Aspergillus terreus* pneumonia. J Antimicrob Chemother. 2013;68:1148-1151.
12. Guinea J, Sandoval-Denis M, Escribano P, Peláez T, Guarro J, Bouza E. *Aspergillus citrinoterreus*, a new species of section Terrei isolated from samples of patients with non-hematological predisposing conditions. J Clin Microbiol. 2015;53:611-617.

ACCEPTED MANUSCRIPT

Table 1. Distribution of amphotericin B MICs against *Aspergillus terreus* species complex isolates collected during the study period and tested according to EUCAST methodology

Species	Amphotericin B MICs, mg/L								
	0.125	0.25	0.5	1	2	4	8	16	32
A. terreus sensu stricto	3	7	10	14	36	81	86	55	23
A. citrinoterreus					3	13	8	7	5
A. hortai					1	2	5	2	
A. alabamensis					2	3	1		
A. floccosus						1			
A. neoafricanus									1
Potential new species							1		

Table 2. Species distribution of *Aspergillus terreus* species complex isolated from the various human specimens

Species	Specimens, total numbers						
	Sputa	Bronchoalveolar lavages and tracheal secretions	Body-fluids	Biopsies	Swabs	Others	Total
A. terreus sensu stricto	126	65	53	33	17	21	315
A. citrinoterreus	14	7	3	5	3	4	36
A. hortai	4	2			1	3	10
A. alabamensis	3	2			1		6
A. floccosus					1		1
A. neoafricanus						1	1
Potential new species				1			1
Total	147	76	56	39	23	29	370

* aspirates, wound secretions, nails

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

Fig. 1a - c. Overview of countries and *Aspergillus terreus* species complex isolated numbers collected during the study period:

a) *Aspergillus citrinoterreus*

b) *Aspergillus terreus* sensu stricto

c) *Aspergillus hortai*

ACCEPTED MANUSCRIPT

Aspergillus citrinoterreus



Total=36

ACCEPTED

Aspergillus terreus sensu stricto



Total=315

- **Austria**: 48 (15.24%)
- **Belgium**: 3 (0.95%)
- **Brazil**: 6 (1.90%)
- **Czech republic**: 10 (3.17%)
- **Denmark**: 17 (5.40%)
- **France**: 11 (3.49%)
- **Germany**: 24 (7.62%)
- **Greece**: 5 (1.59%)
- **India**: 20 (6.35%)
- **Iran**: 7 (2.22%)
- **Israel**: 28 (8.89%)
- **Italy**: 22 (6.98%)
- **Netherland**: 8 (2.54%)
- **Portugal**: 2 (0.63%)
- **Quatar**: 15 (4.76%)
- **Serbia**: 2 (0.63%)
- **Spain**: 65 (20.63%)
- **Sweden**: 2 (0.63%)
- **USA**: 8 (2.54%)
- **Turkey**: 6 (1.90%)
- **UK**: 4 (1.27%)
- **Unknown**: 2 (0.63%)

ACCEPTED


Aspergillus hortai
Czech republic: 3 (30 %)
Italy: 2 (20 %)
Brazil: 1 (10 %)
Israel: 1 (10 %)
Texas: 1 (10 %)
Spain: 2 (20 %)
Total=10

ACCEPTED