

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

### Short-term intensive family therapy for adolescent eating disorders: 30-month outcome

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1653608> since 2021-03-11T12:17:49Z

*Published version:*

DOI:10.1002/erv.2353

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## Short-Term Intensive Family Therapy for Adolescent Eating Disorders: Thirty-month Outcome

Journal:	<i>European Eating Disorders Review</i>
Manuscript ID:	ERV-RA-2014-12-1063.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Marzola, Enrica; University of Turin, Neuroscience, Section of Psychiatry Knatz, Stephanie; University of California, San Diego, Psychiatry Murray, Stuart; University of California San Diego, Psychiatry Rockwell, Roxanne; University of California San Diego, Psychiatry Boutelle, Kerri; University of California San Diego, Psychiatry Eisler, Ivan; King's College London, Institute of Psychiatry, Child Psychiatry Kaye, Walter; University of California San Diego, Psychiatry
Keywords:	anorexia nervosa, family therapy
Abstract:	<p>Family therapy approaches have generated impressive empirical evidence in the treatment of adolescent eating disorders (ED's). However, the paucity of specialist treatment providers limits treatment uptake; therefore, our group developed the Intensive Family Therapy (IFT) – a 5-day treatment based on the principles of family-based therapy for ED's. We retrospectively examined the long-term efficacy of IFT in both single-family (S-IFT) and multi-family (M-IFT) settings evaluating 74 eating disordered adolescents who underwent IFT at the University of California, San Diego, between 2006 and 2013. Full remission was defined as normal weight (<math>\geq 95\%</math> of expected for sex, age, and height), EDE-Q global score within 1 SD of norms, and absence of binge-purging behaviors. Partial remission was defined as weight <math>\geq 85\%</math> of expected or <math>\geq 95\%</math> but with elevated EDE-Q global score and presence of binge-purging symptoms (<math>&lt; 1/\text{week}</math>). Over a mean follow-up period of 30 months, 87.8% of participants achieved either full (60.8%) or partial remission (27%), whilst 12.2% reported a poor outcome, with both S-IFT and M-IFT showing comparable outcomes. Short-term, intensive treatments may be cost-effective and clinically useful where access to regular specialist treatment is limited.</p>

**Short-Term Intensive Family Therapy for Adolescent Eating Disorders: Thirty-month Outcome**

Enrica Marzola<sup>1</sup>, MD, Stephanie Knatz<sup>2</sup>, PhD, Stuart B. Murray<sup>2</sup>, PhD, Roxanne Rockwell<sup>2</sup>, PhD, Kerri Boutelle<sup>2</sup>, PhD, Ivan Eisler<sup>3</sup>, PhD, Walter H. Kaye<sup>2</sup>, MD

<sup>1</sup>Department of Neuroscience, University of Turin, Turin, Italy.

<sup>2</sup>Department of Psychiatry, University of California, San Diego, CA, USA.

<sup>3</sup>Institute of Psychiatry, Kings College, London, UK.

Corresponding Author:  
Walter H Kaye, MD  
Professor of Psychiatry  
Director, UCSD Eating Disorder Research and Treatment Program  
UCSD Department of Psychiatry  
Chancellor Park  
4510 Executive Dr., Suite 315  
San Diego, CA 92121  
<http://eatingdisorders.ucsd.edu>  
[wkaye@ucsd.edu](mailto:wkaye@ucsd.edu)  
[whkaye@gmail.com](mailto:whkaye@gmail.com)  
858 205 7293 cell  
858 534-3951 office  
858 534 6727 fax

The study took place at the University of California San Diego, Eating Disorders Research and Treatment Program, Chancellor Park, 4510 Executive Drive, Suite 315, San Diego CA 92121

Funding provided by the Global Foundation for Eating Disorders, the Price Foundation, the Davis Foundation, and the Peterson Family.

RUNNING HEAD: SHORT-TERM INTENSIVE FAMILY THERAPY

### Abstract

Family therapy approaches have generated impressive empirical evidence in the treatment of adolescent eating disorders (ED's). However, the paucity of specialist treatment providers limits treatment uptake; therefore, our group developed the Intensive Family Therapy (IFT) – a 5-day treatment based on the principles of family-based therapy for ED's. We retrospectively examined the long-term efficacy of IFT in both single-family (S-IFT) and multi-family (M-IFT) settings evaluating 74 eating disordered adolescents who underwent IFT at the University of California, San Diego, between 2006 and 2013. Full remission was defined as normal weight ( $\geq 95\%$  of expected for sex, age, and height), Eating Disorder Examination-Questionnaire (EDE-Q) global score within 1 SD of norms, and absence of binge-purging behaviors. Partial remission was defined as weight  $\geq 85\%$  of expected or  $\geq 95\%$  but with elevated EDE-Q global score and presence of binge-purging symptoms ( $< 1/\text{week}$ ). Over a mean follow-up period of 30 months, 87.8% of participants achieved either full (60.8%) or partial remission (27%), whilst 12.2% reported a poor outcome, with both S-IFT and M-IFT showing comparable outcomes. Short-term, intensive treatments may be cost-effective and clinically useful where access to regular specialist treatment is limited.

**Keywords:** intensive family therapy, family therapy, eating disorders, anorexia nervosa, multi-family therapy

## Introduction

Eating disorders (ED's) rank amongst the most pernicious of all psychiatric disorders, demonstrating marked medical complications alongside poor rates of treatment outcome and high rates of relapse.(Bulik, Berkman, Brownley, Sedway, & Lohr, 2007; Steinhausen, 2002) However, despite adult populations being plagued by 'unacceptable treatment outcomes'(Bulik *et al.*, 2007), a burgeoning body of evidence suggests that early intervention may afford more promising outcomes for those with ED's.(Steinhausen, 2002) As such, much contemporary research and treatment interventions have been oriented towards enhancing outcomes in adolescent presentations of ED's; the most common time of onset.(Neubauer, Weigel, Daubmann, Wendt, Rossi, Lowe, & Gumz, 2014)

With respect to adolescent presentations of AN, specifically, **eating disorder focused family therapy (FT-AN; sometimes referred to as family based-treatment or FBT or Maudsley therapy)** has garnered a particularly impressive evidence-base over the last decade,(Eisler, Wallis, & Dodge, 2015) with an array of empirical evidence documenting that between 50-70% of adolescents are typically weight restored within 12 months of commencing treatment,(Downs & Blow, 2013; le Grange & Eisler, 2009) with swifter and more robust rates of weight restoration when compared with other adolescent-focused treatments.(Couturier, Kimber, & Szatmari, 2013b; Lock, Le Grange, Agras, Moye, Bryson, & Jo, 2010) Furthermore, **FT-AN** also demonstrates favorable cognitive remission throughout treatment, with up to 40% being remitted of cognitive symptomatology within 12 months of commencing treatment.(Lock *et al.*, 2010) Accordingly, **FT-AN** has been found to be effective in reducing the frequency and duration of costly hospital admissions,(Madden, Miskovic-Wheatley, Wallis, Kohn, Lock, Le Grange, Jo, Clarke, Rhodes, & Touyz, 2014) limiting the overall cost of treatment, with long-term follow-up further suggesting that rates of relapse and readmission are curtailed significantly.(Couturier *et al.*, 2013b; le Grange, Lock, Accurso, Agras, Darcy, Forsberg, & Bryson, 2014) Specialist

outpatient services using FT-AN have been shown to reduce the need for hospital treatment 2-3 fold.(House, Craig, Simic, Nicholls, Hugo, & Eisler, 2012)

One of the most significant barriers to the uptake and efficacy of these specialty treatments lies in the limited number of trained providers beyond the academic institutions in which these treatments were developed.(Murray & Le Grange, 2014) For instance, many families report that access to specialized family therapy treatments is made significantly more difficult by the limited number of specialist treatment centers in particular geographical regions.(Brown, 2010; Parent & Parent, 2008) Similarly, clinical practitioners report that geographical location and limited access to specialist supervision restricts their adoption of specialized family therapy, with clinicians feeling less equipped to conduct specialist family treatments in remote areas without regular access to training and supervision.(Couturier *et al.*, 2013b) Indeed, the challenges facing families in accessing specialized family therapy in remote regions has been noted in the United States(Brown, 2010), Canada(Couturier *et al.*, 2013b), and Australia(Wallis, Alford, Hanson, Titterton, Madden, & Kohn, 2012) alike.

In light of this limited access to treatment for some families, recent clinical endeavors have sought to develop short-term and intensive treatment interventions, allowing for temporary immersion into brief treatment programs with a view to swiftly mobilize treatment processes.(Rockwell, Boutelle, Trunko, Jacobs, & Kaye, 2011; Wallis *et al.*, 2012) In order to explore a creative solution to this gap in service provision, our group adopted the principles of FT-AN (Eisler, Le Grange, & Lock, In Press) and constructed a week-long Intensive Family Therapy (IFT) outpatient program, allowing families from remote regions to attend the otherwise unavailable treatment programs in the United States, albeit over a shorter period than typically recommended if local treatment were available.(Rockwell *et al.*, 2011). In fact, IFT originally started with the overarching goal to fill not only the need to extend treatment to families in areas where there was no access to specialist family therapy services but also to offer ancillary services for treatment-resistant cases (Knatz, Kaye, Marzola, & Boutelle, 2015).

In keeping with recent Cochrane Review guidelines,(Fisher, Hetrick, & Rushford, 2010) the escalating costs of specialist treatment, and the development of Multi-Family Therapy (MFT) as a clinically viable alternative to FBT (Dare & Eisler, 2000; Eisler *et al.*, 2015) our intensive program evolved to incorporate MFT principles. Whilst building on established family therapy principles, MFT also leverages alternate mechanisms of therapeutic change in ensuring symptom remission, relying more centrally on group processes that maximize families' own resources, while reducing feelings of helplessness, isolation and shame.(Dare & Eisler, 2000; Eisler, 2005; MRM Scholz, Scholz, Gantchev, & Thömke, 2005; Simic & Eisler, 2015) Empirical evidence suggests that MFT is helpful for families and features lower rates of dropout than other forms of family therapies,(Eisler *et al.*, 2015; Hollesen, Clausen, & Rokkedal, 2013; Salaminou, Campbell, Kuipers, & I, In Press) although the long-term efficacy of MFT remains to be fully explicated.

To date, preliminary data suggests that short-term family intervention programs may result in weight gain between admission and follow-up, although little evidence has tracked the trajectory of eating disorder symptomatology throughout these intensive treatment adaptations.(Dare & Eisler, 2000; Rockwell *et al.*, 2011) In addition, the absence of long-term follow-up data has precluded a thorough assessment of the efficacy of short-term intensive family therapy, suggesting that despite promising preliminary findings, more rigorous investigations ought to document the efficacy of the short-term intensive family therapy of adolescent ED's.

Thus, we sought to investigate the long-term efficacy of our short-term IFT program developed at the University of California, San Diego. Our short-term intensive treatment program consisted of two iterations of a one-week intensive family-based treatment program; one for single families (S-IFT) (Rockwell *et al.*, 2011) and another for multiple families (M-IFT).(Knatz *et al.*, 2015) This study represents the first long-term follow-up of short-term intensive treatments, and also represents the first comparison of single- versus multi-family

therapy of adolescent ED's. Thus, the present study sought to a) evaluate the outcome of adolescent patients who participated in the IFT programs, as well as the acceptability and feasibility; b) to compare the S-IFT/M-IFT models; and c) to compare the outcome of adolescent patients with Anorexia Nervosa (AN) and adolescent patients with Eating Disorder Not Otherwise Specified (EDNOS) - restricting subtype. We hypothesized that both S-IFT and M-IFT would result in sustained physiological and psychological symptom remission, and would be deemed acceptable to families undergoing treatment.

For Review Only



## Method

### ***Participants***

One hundred and eighteen families that participated in IFT (S-IFT+M-IFT) between November 2006 and June 2013 at the University of California, San Diego were considered for inclusion in this report. All families completed the full treatment although because of medical instability one patient had to be briefly hospitalized and then completed IFT. We used the following criteria to exclude participants for this analyses: a) meeting DSM-IV-TR(American Psychiatric Association, 2000) criteria for Bulimia nervosa (BN), b) sub-threshold variants of BN (i.e., EDNOS with binge-purging behaviors) and c) lack of written informed consent. Fourteen families were excluded due to diagnostic criteria and 12 were excluded for consent reasons. Thus, 92 families were contacted to participate in this study. Of the 92, 11 (11.9%) declined to participate and 7 (7.6%) were unable to be contacted. We collected data on 74 (80.5%) of the families contacted. The study was conducted according to the Institutional Review Board regulations of the University of California, San Diego.

### ***Treatment***

#### ***Single-Family Intensive Family Therapy (S-IFT)***

The single-family intensive family therapy (S-IFT) previously described by our team(Rockwell *et al.*, 2011) is a one-week brief intervention for individual families with an adolescent with AN (Table 1). The S-IFT treatment team comprised psychiatrists, psychologists, nurses, and social workers, with a total of 40 hours of intensive treatment delivered over the course of the week. In terms of the clinical content of S-IFT, this treatment may be most centrally characterized by an array of intensified FT-AN practices, including repeated anxiety-raising sessions, repeated family meal sessions with coaching, and intensified attempts to mobilize parental authority. However, S-IFT also included behavioral contracting and psychoeducation. The contracting sessions were focused on developing a behavioral contract to

reinforce adolescents' positive behaviors related to food and eating-related challenges. Additionally, families attended three one-hour psychoeducation sessions lead by a psychiatrist with extensive experience in the treatment of ED, with the goal of empowering parents and adolescents providing them with a better understanding of ED symptoms. Treatment providers were trained in FBT (Lock & le Grange, 2005) and received ongoing supervision from a certified FBT supervisor (K.B.). All families received psychiatric and medical consultations.

### *Multi-Family Intensive Family Therapy (M-IFT)*

Our multi-family intensive family therapy program (M-IFT) has previously been described by our team (Knatz *et al.*, 2015) and features up to 6 families with adolescents with heterogeneous ED presentations. Differences and similarities with S-IFT are presented in Table 1. The treatment centers on the principles of parent-led symptom reduction, and includes three sequential phases: observation, intervention, and reinforcement/planning. In keeping with the multi-family therapy literature, (Eisler, 2005; M Scholz, Asen, Gantchev, Schell, & Süß, 2002; MRM Scholz *et al.*, 2005; Simic & Eisler, 2015) a less direct therapeutic stance was taken where therapists acted as facilitators and families were encouraged to connect with one another to share feedback. To this end, our M-IFT program consisted of a reflexive blend of FT-AN, structural, systemic, strategic, narrative and psychodrama-based family therapy practices. Additionally, the group format was designed in order to enhance families' opportunities for learning, allowing parent-to-parent consultation, and promoting family cohesion in the context of a supportive and reflexive environment, which also help reduce the families' sense of isolation and stigma. Behavioral contracting and psychoeducation sessions were structured as described earlier. Treatment providers were trained in FBT (Lock & le Grange, 2005) and multi-family therapy for AN and received ongoing supervision from a certified FBT supervisor (K.B.) as well as less frequent supervision from an expert in MFT (I.E.) (Voriadaki, Simic, Espie, & Eisler, 2015) All families received psychiatric and medical consultations.

## **Measures**

### *Eating Disorder symptoms*

A semi-structured interview adapted from the Eating Disorders Examination (C. Fairburn, Cooper, & O'Connor, 2008) was used to collect details about past participants' self-reported height and weight, eating disorder symptomatology, treatments, and medications. We inquired about illness trajectory since time of discharge from the IFT program as well as symptoms within the 3 months prior to the time of interview. Additionally, the interview asked participants about their opinion on the usefulness of IFT. In case of positive answer, the interviewees were required to specify what components of IFT they considered particularly helpful choosing between: a) psycho-education; b) group format (in case of M-IFT); c) behavioral contracts and their combination.

All interviews were conducted over the phone from July to November 2013 by the same interviewer. The interviews were conducted with parents for individuals who were still adolescents (less than 18 years old at the time of the follow up, N=39) and either parents (N=22) or past participants for individuals over the age of 18 (N=13).

Additionally, parents and past participants of all ages were also asked to complete an online self-report Eating Disorder Examination Questionnaire (EDE-Q). The EDE-Q (C. G. Fairburn & Beglin, 1994) is a 28-item self-report questionnaire with high internal consistency (Cronbach's alpha global scale = 0.93; (Mond, Hay, Owen, & Beumont, 2004)) that provides a measure of the range and severity of eating disorder features in a total score and four subscales: Restraint, Eating Concern, Shape Concern, and Weight Concern. A parent version (P-EDEQ), (Loeb, 2008) was also used depending on participants' age. In case of overlap or disagreement between adults and parents' reports the former were included in the analysis.

To allow a comparison of our findings with those in the existing literature, we adopted outcome criteria commonly used for FBT studies (Eisler & Dare, 2000; le Grange, Crosby,

Rathouz, & Leventhal, 2007; Le Grange, Lock, Agras, Moye, Bryson, Jo, & Kraemer, 2012b; Lock *et al.*, 2010) thus categorizing outcome as follows: 1) Full remission: a minimum of 95% expected body weight (%EBW) for sex, age, and height as calculated using the Centers for Disease Control and Prevention weight charts, (Kuczmarski, Ogden, Guo, Grummer-Strawn, Flegal, Mei, Wei, Curtin, Roche, & Johnson, 2002) EDE-Q global score within 1 SD of published community norms (1.59; (Mond, Hay, & Owen, 2006)), and absence of binge-purging behaviors (as assessed per EDE-Q items and clinical interview) in the previous 28 days. 2) Partial remission: weight either greater than 85% or higher than 95% EBW but with elevated EDE-Q global mean score and binge-purging symptoms occurring less than once per week. 3) Poor outcome: failure to achieve a minimum of 85% EBW or presence of binge-purging episodes during the previous 28 days with a frequency of once a week or more. Remission was evaluated in our sample using interview data (height, weight and binge-purge behaviors) and the EDE-Q global score. If the EDE-Q was not completed (N=19), we utilized the interview data to determine outcome based on weight, height and binge/purge behaviors.

### *Demographics*

From chart review we collected past participants' gender, ethnicity, place of origin, diagnosis, treatment setting, age, age of onset, duration of illness, medication use, number of previous hospitalizations and residential treatments, and family history of both ED's and psychiatric disorders. Weight and height as measured by a nurse upon admission were also garnered from chart review to calculate %EBW. (le Grange, Doyle, Swanson, Ludwig, Glunz, & Kreipe, 2012a)

### *Statistical analysis*

Preliminary lost-case analysis evaluated any differences in baseline variables between those who agreed to participate and those who declined using Mann-Whitney-Wilcoxon test and Fisher's exact test to analyze continuous and categorical variables, respectively. Moreover, a sensitivity analysis evaluated whether those who were lost at follow-up could significantly

influence outcome. We evaluated the relationship between diagnosis, age at intervention, duration of illness, age of onset, and length of follow-up with outcome diagnosis using binary logistic regression. Changes in %EBW over time in the overall sample were analyzed with a paired-sample t-test. Furthermore, we evaluated changes in %EBW over time, and outcome differences between diagnosis (AN versus EDNOS) and programs (S-IFT versus M-IFT) using Mann-Whitney-Wilcoxon test and Fisher's exact test. Differences in treatment received after discharge across outcome categories were investigated using the Fisher's exact test.

For Review Only

## Results

### ***Participant characteristics at baseline***

As shown in **Table 2**, patients' mean age at entry into treatment was 14.8 (SD=2.7) years, with 42.4% residing in California (N=39), 52.2% residing in other states within the United States (N=48), and 5.4% residing outside of the United States (N=5).

Of the overall sample who participated in the follow-up (N=74), 59.5% (N=44) met DSM-IV diagnostic criteria for AN; of these, 88.6% met criteria for AN-restricting subtype (ANR; N=39), whilst 11.4% for AN-binge/purge subtype (ANBN; N=5; see **Table 2**). The remaining 40.5% of the sample met DSM-IV criteria for EDNOS-restricting type (N=30). The overall mean %EBW at entry into the program was 86.36 (SD=8.74) and mean illness duration was 1.86 years (SD=1.97 years), with 31.1% of participants reporting previous ED-related hospitalizations before coming to UCSD (N=23). **The EDNOS subgroup was mainly composed of females (83.3%, N=25) with mean duration of illness of 1.7 (SD= 1.49) years, mean age of 14.5 (SD=3) years, and mean age of onset of 12.9 (SD=2.2) years. The majority of those affected by EDNOS underwent M-IFT (73.3%, N=22).**

### ***Lost case analysis***

As shown in **Table 2**, the lost case analyses revealed no significant differences between those individuals who agreed to participate (N=74) and those who did not (N=18), with respect to any of the variables measured at baseline (age, duration of illness, treatment structure, %EBW, age of onset and place of origin), except number of hospitalizations before IFT.

### ***Study participation and attrition***

A total of 74 families participated in the outcome study (see Figure 1), of which 27% (N=20) completed S-IFT and 73% (N=54) completed M-IFT. Of this sample, at time of follow-up 47.3% (N=35) were young adults and 52.7% (N=39) were adolescents. The mean length of

follow-up was 30.85 months (SD=20.2 months; range=4-83 months), and it should be noted the M-IFT group featured a significantly shorter follow-up period (Mann-Whitney-Wilcoxon test  $p<0.001$ ), due to the fact that S-IFT was implemented in November 2006, and M-IFT was implemented in March 2010.

### ***Characteristics of participants at follow-up***

At follow-up, mean %EBW of the overall sample was 99% (SD=12.6; see Table 3), representing a statistically significant increase in weight ( $t=7.99$ ,  $p<0.001$ ) with a very large effect size ( $d=1.18$ ). A significant increase in %EBW emerged in both diagnosis subgroups (Figure 2a). Since discharge from UCSD 24.2% (N=18) received no other treatments, 50% (N=37) individual psychotherapy, 6.8% (N=5) family counselling, 2.7% (N=2) intensive outpatient program, 9.5% (N=7) a combination of individual psychotherapy and psychiatric consultations, 2.7% (N=2) psychiatric visits only, and 4.1% (N=3) other kinds of treatments. Taken cumulatively across both treatment groups, 87.8% (N=65) of our overall sample achieved a positive outcome, with significantly less hospitalizations following IFT when compared to those required before IFT. More specifically, 87.8% (N=65) of participants achieved either full (60.8%, N=45) or partial remission (27%; N=20), whilst 12.2% (N=9) reported a poor outcome (See Figure 3).

### ***Outcome sensitivity analyses***

The sensitivity analysis revealed that even if those who were lost at follow-up had all positive outcomes the findings of this study would not be affected. However, the overall findings could be modified if those lost at follow-up reported all poor outcomes or in case of all positive outcomes in one condition and all negative in the other.

### ***Treatment after discharge and outcome***

No significant differences emerged across outcome groups with respect to kinds of treatment received after discharge from UCSD (Fisher's exact test  $p=0.69$ ).

### ***Relationship between baseline variables and outcome***

Binary logistic regressions failed to show an association between outcome and the following baseline variables: diagnosis (Wald test= 2.92,  $p=0.09$ ), age at intervention (Wald test= 0.45,  $p=0.5$ ), age of onset (Wald test= 0.24,  $p=0.62$ ), and duration of illness (Wald test= 2.43,  $p=0.12$ ).

### ***Comparison of diagnosis; AN and EDNOS-restricting subtype***

Among baseline characteristics, individuals affected by AN versus EDNOS-restricting subtype differed only with respect to %EBW (see [Table 2](#)). No differences emerged between AN and EDNOS groups with respect to outcome (Fisher's exact test  $p=0.16$ ), medications (Fisher's exact test  $p=0.16$ ), or higher levels of care in the interval since IFT treatment (Fisher's exact test hospitalizations  $p=0.7$ ; residential treatments  $p=0.18$ ).

### ***Comparison of different models of delivery; S-IFT and M-IFT***

No differences emerged between S-IFT and M-IFT on baseline characteristics (data not shown). Furthermore, no differences between S-IFT and M-IFT groups emerged with respect to %EBW at follow-up (Mann-Whitney-Wilcoxon test  $p=0.77$ ; see Figure 2b) outcome (Fisher's exact test  $p=1$ ), medications (Fisher's exact test  $p=0.61$ ), or higher levels of care in the interval since IFT treatment (Fisher's exact test hospitalizations  $p=0.659$ ; residential treatments  $p=0.72$ ).

### ***Acceptability of IFT***

The vast majority of the interviewees reported IFT to be useful ( $N=68$ , 91.9%). In particular, 60% of those who underwent S-IFT evaluated as particularly helpful the combination



of psycho-educational activities and behavioral contracting whilst 87% of those who participated in M-IFT valued the combination of all three components (i.e., psycho-educational activities, behavioral contracts, and group setting).

For Review Only

## Discussion

The main aim of this study was to assess long-term outcome of two forms of short-term, intensive family therapy in the context of adolescents affected by AN and EDNOS restricting subtype. When considered together, both S-IFT and M-IFT showed similarly significant weight- and behavioral symptom remission, with approximately 88% demonstrating full (61%) or partial remission (27%), and only 12% reporting a poor outcome at a mean of 30 months follow-up. Furthermore, both forms of intensive family therapy were deemed acceptable to families and adolescents. Cumulatively these data represent novel findings, and may provide preliminary support for the clinical utility of short-term intensive treatments in adolescents with ED's.

In keeping with outcome criteria widely endorsed in the extant outcome literature,(Eisler, Simic, Russell, & Dare, 2007; Le Grange *et al.*, 2012b; Lock *et al.*, 2010) the present findings demonstrate comparable outcomes to those reported in controlled trials of **FT-AN**, which typically illustrate that up to 49% report full and 77% report partial remission at 12-month follow-up, with 23% reporting poor outcome.(Le Grange *et al.*, 2012b; Lock *et al.*, 2010) In this study, we found comparable outcomes for S-IFT and M-IFT, although there were differences in follow-up lengths. Other studies presenting longer term follow-ups indicate that there is an ongoing improvement following family therapy and that those who do well maintain their achievements and have low relapse rates.(Eisler, Dare, Russell, Szmulker, Le Grange, & Dodge, 1997; Eisler *et al.*, 2007; le Grange *et al.*, 2014; Lock, Couturier, & Agras, 2006)

This study is the first to present data on long-term outcomes using intensive treatments for adolescents with AN and EDNOS-restricting subtype. This model provides a treatment alternative that may be able to reach a greater proportion of the families who have a child with an ED. It is important to note the scarcity of expert **FT-AN** therapists (Murray & Le Grange, 2014) alongside the propensity for lack of treatment fidelity amongst those in unsupported settings (Couturier, Kimber, Jack, Niccols, Van Blyderveen, & McVey, 2013a, 2014; Wallace & von Ranson, 2012) which could impinge upon both the uptake and outcome of specialist

treatments.(House *et al.*, 2012) This intensive therapy model could provide an empirically supported treatment for families whose geographical location restricts their access to specialist outpatient ED services that are most likely to provide evidence-based treatments.(Couturier *et al.*, 2013a, 2014)

Furthermore, our findings add to a growing body of literature suggesting the efficacy of the multi-family intensive treatment for families with a child with an ED.(Gabel, Pinhas, Eisler, Katzman, & Heinmaa, 2014; Salaminou *et al.*, In Press) Indeed, in light of the similar treatment outcomes between our S-IFT and M-IFT, and the empirically supported individual FT-AN,(Lock, 2011) the simultaneous treatment of multiple families in intensive treatment programs without any loss of efficacy may represent a particularly cost-effective avenue for further empirical and clinical endeavors, although these treatments were not directly compared in this study. This assertion may be further underscored in light of the present findings drawn from adolescents engaged in costly hospital and partial hospital programs,(Madden *et al.*, 2014) with reduced rates of readmissions to hospital settings further underscoring the cost-effectiveness of this treatment.

In addition to our overall findings, delineating the differential effects of short-term intensive treatments across diagnoses further supported the efficacy of treatment, with both those with AN and those with EDNOS-restricting type presentations reporting comparable treatment outcomes. This may be particularly important to consider when considering the heterogeneity and complexity of presentation (i.e., high levels of Obsessive Compulsive Disorder, complex family presentations) and the noted ambiguity surrounding the treatment of atypical presentations of disordered eating.(Keel & Brown, 2010) There is evidence that more complex cases require a greater degree of flexibility and tailoring to the specific needs of a particular family(Webb, Thuras, Peterson, Lampert, & Crow, 2011) and it is possible that the intensive and immersive nature of short-term treatment allows for more observation and

analyses of the subtle nuances of atypical presentations, which may not emerge in less intensive and time-limited outpatient settings.

Alongside the notable clinical implications, the present findings, if replicated in larger and more controlled trials, may likely impact the development of treatment policies. Firstly, providing centralized access to leading evidence-based treatments mitigates the challenges to treatment uptake brought about by the scarcity of specialized treatment providers. This may be particularly important when considering access to treatments most typically delivered on an outpatient basis, as long distance commutes may likely impinge upon regular treatment schedules. It is of note that in the present study, 57.6% of those seeking intensive specialist treatment came from other states or outside the United States, confirming the difficulty for patients and their families in accessing specialist treatment programs. Furthermore, grouping several families together throughout treatment allows for the division of treatment costs between families, providing more financially viable and equally efficacious treatment options for families and insurance providers alike. This is particularly noteworthy in the context of costly treatment programs for AN, which typically approximate up to \$3,979 per day in the United States. (Madden *et al.*, 2014)

However, despite these encouraging findings, limitations should also be noted. This study relies on self-report data and the semi-structured interview used is not validated and was not administered to all patients at the different time-points considered. In fact, in some cases either the EDE-Q or the interview were used to evaluate outcome. Moreover, a number of participants were lost at follow-up assessment, and our sensitivity analyses suggested that if all the participants who were lost to follow up did poorly, the results of this study may not stand. Furthermore, because of the evolution of this clinical treatment program, the length of follow-up varied between individual and multi-family intensive treatments. Moreover, in keeping with our clinical policy to exercise no exclusion criteria for treatment, there were both AN and EDNOS-restricting subtype included in the present study, although our analyses were not powered to look at any differential effects of treatment by diagnosis. Finally, it is possible that there was a

selection bias, in that families who participated in our intensive program were particularly motivated to engage in treatment, given that they had to travel to San Diego and often pay out of pocket for treatment. Furthermore, those who completed the follow-up assessment reported less frequently a history of hospitalizations than those who declined to participate in this study. Lastly, a significant proportion of participants received follow-up treatments after attending the program. This confounds the reported outcome data as it is possible that changes in reported outcomes may be partially attributed to follow up services. However, a quarter of individuals did not undergo any other treatments after receiving IFT, and of those who did, the majority did not receive specialty family therapy services. Given these limitations, it remains to be proven whether these results will generalize to all patients with AN and EDNOS-restricting subtype and their families.

The present results provide novel data as to the preliminary efficacy of one week, intensive treatments for adolescent ED's. Whilst preliminary at this stage, these promising findings suggest that further research should replicate the present findings in larger and more rigorously controlled trials.

### **Acknowledgements**

We thank the families who participated in this study, the treatment staff, Anthony Gamst, PhD, Professor of Biostatistics and Bioinformatics, University of California San Diego, who assisted with statistical review, and the Price Foundation and Davis/Wismer Foundation. Dr. Marzola would like to thank Secondo Fassino, MD, and Giovanni Abbate-Daga, MD, Professors of Psychiatry at the University of Turin, Italy, for their kind support.

For Review Only

## References

- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders - Fourth Edition - Text Revision (DSM-IV-TR)*.
- Brown, H. (2010). *Brave Girl Eating*. HarperCollins; New York.
- Bulik, C., Berkman, N., Brownley, K., Sedway, J., & Lohr, K. (2007). Anorexia nervosa treatment: a systematic review of randomized controlled trials. *Int J Eat Disord, 40*(4), 310-320.
- Couturier, J., Kimber, M., Jack, S., Niccols, A., Van Blyderveen, S., & McVey, G. (2013a). Understanding the uptake of family-based treatment for adolescents with anorexia nervosa: therapist perspectives. *Int J Eat Disord, 46*(2), 177-188.
- Couturier, J., Kimber, M., Jack, S., Niccols, A., Van Blyderveen, S., & McVey, G. (2014). Using a knowledge transfer framework to identify factors facilitating implementation of family-based treatment. *Int J Eat Disord, 47*(4), 410-417. doi: doi: 10.1002/eat.22225
- Couturier, J., Kimber, M., & Szatmari, P. (2013b). Efficacy of family-based treatment for adolescents with eating disorders: a systematic review and meta-analysis. *Int J Eat Disord, 46*(1), 3-11.
- Dare, C., & Eisler, I. (2000). A multi-family group day treatment programme for adolescent eating disorder. *Eur Eat Disord Rev, 8*(1), 4-18.
- Downs, K., & Blow, A. (2013). A substantive and methodological review of family-based treatment for eating disorders: the last 25 years of research. *J Fam Therapy, 35*(S1), 3-28.
- Eisler, I. (2005). The empirical and theoretical base of family therapy and multiple family day therapy for adolescent anorexia nervosa. *J Fam Therapy, 27*(2), 104-131. doi: 10.1111/j.1467-6427.2005.00303.x
- Eisler, I., & Dare, C. (2000). Family therapy for adolescent anorexia nervosa: the results of a controlled comparison of two family interventions. *J Child Psychol Psychiat, 41*(6), 727-736.

- Eisler, I., Dare, C., Russell, G. F., Szumukler, G., Le Grange, D., & Dodge, E. (1997). Family and individual therapy in anorexia nervosa. A 5-year follow-up. *Arch Gen Psychiatry*, *54*, 1025-1030.
- Eisler, I., Le Grange, D., & Lock, J. (In Press). Family therapy for adolescent eating disorders. In Sexton T & Lebow J (eds.) *Handbook of Family Therapy 4th Edition*. Routledge.
- Eisler, I., Simic, M., Russell, G., & Dare, C. (2007). A randomised controlled treatment trial of two forms of family therapy in adolescent anorexia nervosa: a five year follow-up. *J Child Psychol Psychiatry*, *48*(6), 552-560.
- Eisler, I., Wallis, A., & Dodge, E. (2015). What's New is Old and What's Old is New: The Origins and evolution of Family Therapy for Eating Disorders! In K. Loeb, D. Le Grange & J. Lock (Eds.), *Family Therapy for Adolescent Eating and Weight Disorder: New Applications*: Routledge.
- Fairburn, C., Cooper, Z., & O'Connor, M. (2008). Eating disorder examination. In : Fairburn CG, editor. *Cognitive Behavior Therapy and Eating Disorders*, 16th edition. New York: Guilford Press, pages 265-308.
- Fairburn, C. G., & Beglin, S. (1994). Assessment of eating disorders: interview or self-report questionnaire? *Int J Eat Disord*, *16*, 363-370.
- Fisher, C., Hetrick, S., & Rushford, N. (2010). Family therapy for anorexia nervosa. *Cochrane Database of Systematic Reviews* (4):CD004780. doi: doi: 10.1002/14651858.CD004780.pub2
- Gabel, K., Pinhas, L., Eisler, I., Katzman, D., & Heinmaa, M. (2014). The effect of multiple family therapy on weight gain in adolescents with anorexia nervosa: Pilot data. *J Can Acad Child Adolesc Psychiatry*, *23*(3), 196-199.
- Hollesen, A., Clausen, L., & Rokkedal, K. (2013). Multiple family therapy for adolescents with anorexia nervosa: a pilot study of eating disorder symptoms and interpersonal functioning. *J Fam Therapy*, *35*, 53-67.



- House, J. S., U, Craig, M. L., S, Simic, M., Nicholls, D., Hugo, P. B., M, & Eisler, I. (2012). Comparison of specialist and nonspecialist care pathways for adolescents with anorexia nervosa and related eating disorders. *Int J Eat Disord*, *45*(8), 949-956. doi: doi: 10.1002/eat.22065
- Keel, P., & Brown, T. (2010). Update on course and outcome in eating disorders. *Int J Eat Disord*, *43*, 195-204.
- Knatz, S., Kaye, W., Marzola, E., & Boutelle, K. (2015). A brief, intensive application of family-based treatment for eating disorders. In K. Loeb, D. Le Grange & J. Lock (Eds.), *Family Therapy for Adolescent Eating and Weight Disorder: New Applications*: Routledge.
- Kuczmarski, R., Ogden, C., Guo, S., Grummer-Strawn, L., Flegal, K., Mei, Z., Wei, R., Curtin, L., Roche, A., & Johnson, C. (2002). 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat*, *246*, 1-190.
- le Grange, D., Crosby, R., Rathouz, P., & Leventhal, B. (2007). A randomized controlled comparison of family-based treatment and supportive psychotherapy for adolescent bulimia nervosa. *Arch Gen Psychiatry*, *64*(9), 1049-1056.
- le Grange, D., Doyle, P., Swanson, S., Ludwig, K., Glunz, C., & Kreipe, R. (2012a). Calculation of expected body weight in adolescents with eating disorders. *Pediatrics*, *129*(2), e438-e446.
- le Grange, D., & Eisler, I. (2009). Family interventions in adolescent anorexia nervosa. *Child Adolesc Psychiatr Clin N Am*, *18*(1), 159-173. doi: 10.1016/j.chc.2008.07.004
- le Grange, D., Lock, J., Accurso, E., Agras, W., Darcy, A., Forsberg, S., & Bryson, S. (2014). Relapse from remission at two- to four-year follow-up in two treatments for adolescent anorexia nervosa. *J Am Acad Child Adolesc Psychiatry*, *53*(11), 1162-1167. doi: 10.1016/j.jaac.2014.07.014
- Le Grange, D., Lock, J., Agras, W., Moye, A., Bryson, S., Jo, B., & Kraemer, H. (2012b). Moderators and mediators of remission in family-based treatment and adolescent focused therapy for anorexia nervosa. *Behaviour Research and Therapy*, *50*, 85-92.

- Lock, J. (2011). Evaluation of family treatment models for eating disorders. *Curr Opin Psychiatry*, 24(4), 274-279.
- Lock, J., Couturier, J., & Agras, W. (2006). Comparison of long-term outcomes in adolescents with anorexia nervosa treated with family therapy. *J Am Acad Child Adoles Psychiatry*, 45(6), 666-672.
- Lock, J., & le Grange, D. (2005). Family-based treatment of eating disorders. *Int J Eat Disord*, 37, Suppl: S64-S67.
- Lock, J., Le Grange, D., Agras, W., Moye, A., Bryson, S., & Jo, B. (2010). Randomized clinical trial comparing family-based treatment with adolescent-focused individual therapy for adolescents with anorexia nervosa. *Arch Gen Psychiatry*, 67, 1025-1032.
- Loeb, K. (2008). Eating Disorder Examination Questionnaire – Parent Version (P-EDEQ), Version 1.4. Unpublished measure based on Fairburn, C.G. & Beglin, S.J. The Eating Disorder Examination Questionnaire (EDE-Q). In C. G. Fairburn (Ed.), *Cognitive behavior therapy and eating disorders*. New York: Guilford Press. .
- Madden, S., Miskovic-Wheatley, J., Wallis, A., Kohn, M., Lock, J., Le Grange, D., Jo, B., Clarke, S., Rhodes, P., & Touyz, S. (2014). A randomized controlled trial of in-patient treatment for anorexia nervosa in medically unstable adolescents. *Psychol Med*, 14, 1-13. doi: <http://dx.doi.org/10.1017/S0033291714001573>
- Mond, J., Hay, P. R., B, & Owen, C. (2006). Eating Disorder Examination Questionnaire (EDE-Q): norms for young adult women. *Behav Res Ther*, 44(1), 53-62.
- Mond, J., Hay, P. R., B, Owen, C., & Beumont, P. (2004). Validity of the Eating Disorder Examination Questionnaire (EDE-Q) in screening for eating disorders in community samples. *Behav Res Ther*, 42, 551-567.
- Murray, S., & Le Grange, D. (2014). Family therapy for adolescent eating disorders: an update. *Curr Psychiatr Rep*, 16(5), 447. doi: doi: 10.1007/s11920-014-0447-y

- Neubauer, K., Weigel, A., Daubmann, A., Wendt, H., Rossi, M., Lowe, B., & Gumz, A. (2014). Paths to first treatment and duration of untreated illness in anorexia nervosa: Are there differences according to age of onset? *Eur Eat Disord Rev*, *22*(4), 292-298. doi: 10.1002/erv.2300
- Parent, B., & Parent, T. (2008). Anorexia, Maudsley and an impressive recovery: One family's story. *J Pediatr Child Hlth*, *44*, 70-73.
- Rockwell, R., Boutelle, K., Trunko, M., Jacobs, M., & Kaye, W. (2011). An innovative short-term, intensive, family-based treatment for adolescent anorexia nervosa: case series. *Eur Eat Disord Rev*, *19*(4), 362-367.
- Salaminou, E., Campbell, M., Kuipers, E., & I, E. (In Press). Intensive Multi Family Therapy for adolescent anorexia nervosa: An open study of 30 families. *J Fam Therapy*.
- Scholz, M., Asen, E., Gantchev, K., Schell, B., & Süß, U. (2002). A family day clinic in child psychiatry. The Dresden model--concepts and first experiences. [Article in German]. *Psychiatr Prax*, *29*(3), 125-129.
- Scholz, M., Scholz, K., Gantchev, K., & Thömke, V. (2005). Multiple family therapy for anorexia nervosa: concepts, experiences and results. *J Fam Therapy*, *27*, 132-141.
- Simic, M., & Eisler, I. (2015). Multifamily therapy for eating disorders. In Loeb, K., Le Grange, D., & Lock, J. (Eds). *Family Therapy for Adolescent Eating and Weight Disorders*. New York: Guilford Press.
- Steinhausen, H. C. (2002). The outcome of anorexia nervosa in the 20th century. *Am J Psychiatry*, *159*(8), 1284-1293.
- Voriadaki, T., Simic, M., Espie, J., & Eisler, I. (2015). Intensive multi-family therapy for adolescent anorexia nervosa: Adolescents' and parents' day to day experiences. *J Fam Therapy*, *37*, 5-23.
- Wallace, L., & von Ranson, K. (2012). Perceptions and use of empirically-supported psychotherapies among eating disorder professionals. *Behav Res Ther*, *50*, 215-222.

- Wallis, A., Alford, C., Hanson, A., Titterton, J., Madden, S., & Kohn, M. (2012). Innovations in Maudsley family-based treatment for anorexia nervosa at the Children's Hospital at Westmead: a family admission programme. *J Fam Therapy Special Issue: Innovations in the Treatment of Adolescent Eating Disorders*, 35(S1), 68-81. doi: doi: 10.1111/j.1467-6427.2012.00604
- Webb, C., Thuras, P., Peterson, C., Lampert, J. M., D, & Crow, S. (2011). Eating-related anxiety in individuals with eating disorders. *Eat Weight Disord*, 16(4), e236-e241.

For Review Only

**Table 1. Differences and similarities of single-family (S-IFT) and multi-family Intensive Family Therapy (M-IFT).**

	Single-family intensive family therapy (S-IFT)	Multi-family intensive family therapy (M-IFT)
Duration/Weekly hours	5 days/40	5 days/40
Families involved	1	Up to 6
Group setting	-	+
Multiple family meals	-	+
Psychiatrists, psychologists, nurses, and social workers	+	+
Anxiety-raising sessions, family meal sessions with coaching, intensified attempts to mobilize parental authority	+	+
Behavioral contracting	+	+
Psychoeducation	+	+
Parent-to-parent consultation	-	+
Medical consultation	+	+
Broader systemic work (e.g., family sculpt, inter-family role plays)	-	+

**Table 2.** Participant characteristics at time of entry into IFT and lost case analysis.

	Total sample (N=92, 100%)	Participating in FU study (N=74, 80.5%)	Not participating in FU study (N=18, 19.5%)	p-value
Gender, female, N(%)	85(92.4)	68(91.9)	17(94.4)	1
Ethnicity, Caucasian, N(%)	83(91.2)	68(91.9)	15(88.2)	1
Place of origin, N(%)				0.06
Local area	39(42.4)	31(41.9)	8(44.4)	
USA	48(52.2)	41(55.4)	7(38.9)	
Out of the USA	5(5.4)	2(2.7)	3(16.7)	
Diagnosis, N(%)				0.42
AN	57(62)	44(59.5)	13(72.2)	
EDNOS	35(38)	30(40.5)	5(27.8)	
Treatment setting, N(%)				0.16
S-IFT	28(30.4)	20(27)	8(44.4)	
M-IFT	64(69.6)	54(73)	10(55.6)	
Age at entry, years, Mean(SD)	14.84(2.74)	14.74(2.81)	15.27(2.44)	0.65
Age of onset, years, Mean(SD)	13.11(2.42)	13.05(2.52)	13.38(2)	0.63

Duration of illness, years, Mean(SD)	1.9(1.95)	1.86(1.97)	2.1(1.93)	0.79
%EBW, Mean(SD)	86.14(8.65)	86.36(8.74)	84.62(8.33)	0.41
AN	80.44(4.67)	80.52(4.84)	80(3.85)	
EDNOS	94.8(5.55)	94.93(5.46)	93.87(7)	
Medication use, N(%)	39(42.4)	31(41.9)	8(44.4)	1
Previous hospitalizations, individuals, N(%)	35(38)	23(31.1)	12(66.7)	<b>0.007</b>
Previous residential treatments, individuals, N(%)	5(5.4)	5(6.8)	0(0)	0.58
Family history of psychiatric disorders, N(%)	63(68.5)	53(71.6)	10(55.6)	0.33
Family history of eating disorders, N(%)	30(32.6)	25(33.8)	5(27.8)	0.82

Fisher's Exact test and Mann-Whitney-Wilcoxon test were used to assess statistical significance for categorical and continuous variables, respectively.

Legend: S-IFT: single-family Intensive Family Therapy; M-IFT: multi-family Intensive Family Therapy; AN: anorexia nervosa; EDNOS: eating disorder not otherwise specified, restricting type; %EBW: % expected body weight.

**Table 3.** Participant characteristics at follow-up and by treatment structure in those who completed the follow-up study.

	Total sample participating in FU study (N=74)		S-IFT (N=20, 27%)		M-IFT (N=54, 73%)		p-value
Length of follow-up, months, Mean(SD)	30.85	(20.2)	53.4	(16.07)	22.5	(14.37)	<0.001
Age at follow-up, years, Mean(SD)	17.18	(3.4)	19.2	(3.03)	16.44	(3.25)	0.002
%EBW, Mean(SD)	99	(12.57)	102.17	(17.47)	97.83	(10.14)	0.57
AN	96.89	(14.42)	104.16	(21.92)	94.17	(9.47)	0.27
EDNOS	102.11	(8.52)	99.18	(7.52)	103.17	(8.78)	0.26
Outcome, N(%)							1
Full remission	45	(60.8)	13	(65)	32	(59.3)	
Partial remission	20	(27)	5	(25)	15	(27.7)	
No remission	9	(12.2)	2	(10)	7	(13)	
Undergoing treatment, N(%)	36	(50.7)	6	(31.6)	30	(57.7)	0.06




On medications, N(%)	34	(45.9)	8	(40)	26	(48.1)	0.61
How long on medications, N(%)							0.9
<6 months	9	(26.5)	3	(37.5)	6	(23.1)	
6-12 months	8	(23.5)	1	(12.5)	7	(26.9)	
>12 months	12	(35.3)	3	(37.5)	9	(34.6)	
n.a.	5	(14.7)	1	(12.5)	4	(15.4)	
Hospitalization since IFT, individuals, N(%)	6	(8.1)	2	(10)	4	(7.4)	0.66
Residential treatments since IFT, individuals, N(%)	11	(14.9)	2	(10)	9	(16.7)	0.72

Fisher's Exact test and Mann-Whitney-Wilcoxon test were used to assess statistical significance for categorical and continuous variables, respectively.

Legend: S-IFT: single-family Intensive Family Therapy; M-IFT: multi-family Intensive Family Therapy; %EBW: % expected body weight


Figure 1. Outline of family enrollment in this study.




32x41mm (600 x 600 DPI)

**Figure 2. Comparison of % Expected Body Weight (EBW%) at baseline and follow-up between (a) individuals with Anorexia Nervosa (AN) and (b) Eating Disorder Not Otherwise Specified – restricting subtype (EDNOS) and Single-Family Intensive Family Therapy (S-IFT) and Multi-Family Intensive Family Therapy (M-IFT).**

**Figure 2a.**




**Figure 2b.**



Comparison of % Expected Body Weight (EBW%) at baseline and follow-up between (a) individuals with Anorexia Nervosa (AN) and (b) Eating Disorder Not Otherwise Specified – restricting subtype (EDNOS) and Single-Family Intensive Family Therapy (S-IFT) and Multi-Family Intensive Family Therapy (M-IFT)  
 33x43mm (600 x 600 DPI)

Figure 3. Outcomes at follow-up for total sample.



Outcomes at follow-up for total sample  
15x8mm (600 x 600 DPI)

Review Only