

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A SNP in the goat CSN2 promoter region is associated with the absence of β -casein in the milk

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1507156> since 2018-03-17T11:54:58Z

Published version:

DOI:10.1111/j.1365-2052.2007.01649.x

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

G. Cosenza, A. Pauciullo, L. Colimoro, A. Mancusi, A. Rando, D. Di Bernardino and L. Ramunno. A SNP in the goat CSN2 promoter region is associated with the absence of β -casein in milk *Animal Genetics*, 2007, 38, 655-658

DOI 10.1111/j.1365-2052.2007.01649.x

The publisher's version is available at:

<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2052.2007.01649.x/abstract;jsessionid=2B0392FDF47CA92BD21812A24B626335.f04t01?systemMessage=Wiley+Online+Library+is+migrating+to+a+new+platform+powered+by+Atypon%2C+the+leading+provider+of+scholarly+publishing+platforms.+The+new+Wiley+Online+Library+will+be+migrated+over+the+weekend+of+March+17+and+18.You+should+not+experience+any+issues+or+loss+of+access+during+this+time.+For+more+information%2C+please+visit+our+migration+page%3A++http%3A%2F%2Fwww.wileyactual.com%2FWOLMigration%2F>

When citing, please refer to the published version.

Link to this full text:

<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2052.2007.01649.x/epdf>

1 **Short communication**

2 **An SNP in the goat CSN2 promoter region is associated with the absence of b-casein in milk**

3

4 G. Cosenza*, A. Pauciullo*, L. Colimoro*, A. Mancusi*, A. Rando†, D. Di Bernardino* and L.
5 Ramunno*

6

7

8 *Dipartimento di Scienze del Suolo, della Pianta, dell'Ambiente e delle Produzioni Animali, Facoltà
9 di Agraria, Università degli Studi di Napoli Federico II, Portici (Na) 80055, Italy.

10

11 †Dipartimento di Scienze delle Produzioni Animali, Facoltà di Agraria, Università degli Studi della
12 Basilicata, Potenza 85100, Italy

13

14

1 **Summary**

2 So far, at least eight alleles in the goat CSN2 locus have been associated with the level of b-casein
3 expression in milk. Alleles CSN2 A, CSN2 A1, CSN2 B, CSN2 C, CSN2 D and CSN2 E have been
4 associated with normal content (allele effects of about 5 g of β -casein per litre), whereas the CSN2 0
5 and CSN2 01 alleles have been associated with non-detectable levels of β -casein. Most of these
6 alleles have been characterized genetically. Herein, we report the identification of a previously
7 unreported SNP in the goat CSN2 promoter region (AJ011018:g.1311T>C), which is associated with
8 the absence of β -casein in the milk. Furthermore, we developed a PCR-based method that allows
9 detection of this mutation.

10

11

12 **Keywords** β -casein, goat, null allele, promoter, SNP.

13

1 β -casein is the most abundant protein component (about 10 g/l) in goat milk (Grosclaude et al. 1987).
2 The goat b-casein (CSN2) gene maps to chromosome 6 and its organization is similar to that observed
3 in other species (Rijnkels 2002). To date, eight alleles identified at this locus are associated with two
4 expression levels: the CSN2A, CSN2A1, CSN2B, CSN2C, CSN2D and CSN2E alleles are associated
5 with normal content (Roberts et al. 1992; Mahe´ & Grosclaude 1993; Neveu et al. 2002; Galliano et
6 al. 2004; Cosenza et al. 2005; Caroli et al. 2006), whereas the CSN20 and CSN201 alleles are
7 associated with non-detectable amounts of this protein (Ramunno et al. 1995; Persuy et al. 1999).
8 Analysis of the coagulation properties of milk with and without b-casein demonstrates that milk
9 without b-casein shows longer coagulation times (about three times the normal value, 15–25 min vs.
10 4–7 min) and low curd firmness (so low that it is impossible to measure). Furthermore, cheese yield
11 (caciotta at 30 days ageing) from milk without b-casein is about 80% of that obtained from milk with
12 a normal content of this casein fraction (Chianese et al. 1993). With the exception of the CSN2B
13 allele, the DNA and protein sequences of these alleles are known (Roberts et al. 1992; Mahe´ &
14 Grosclaude 1993; Ramunno et al. 1995; Persuy et al. 1999; Neveu et al. 2002; Galliano et al. 2004;
15 Cosenza et al. 2005; Caroli et al. 2006). The CSN201 allele is characterized by a single-nucleotide
16 substitution at position 373 of the seventh exon (Ramunno et al. 1995) (AJ011018:g.8915C>T), while
17 the CSN20 has a single nucleotide deletion (adenine) in a row of four adenines between nucleotide
18 16 and 19 of exon 7 (Persuy et al. 1999) (AJ011018:g.8561delA). These mutations result in a
19 truncated protein at amino acids 57 (Persuy et al. 1999) (p.Ile49SerfsX10) and 181 (Ramunno et al.
20 1995) (p.Gln182X) respectively.
21 Levels of mRNA product transcribed by the CSN20 and CSN201 alleles are about 100 (Persuy et al.
22 1999) and 10 (Ramunno et al. 1995) times lower than normal expression respectively. More recently,
23 Cunsolo et al. (2005) reported on the identification and characterization of a truncated b-casein in the
24 milk of goats homozygous for the CSN201 allele. The truncated protein contained 1–166 amino acid
25 residues of the mature b-casein variant A. So far, research on the polymorphisms associated with
26 differences in the expression of b-casein has been mainly limited to exonic regions of CSN2.

1 However, mutations in the promoter region might be responsible for differences in the level of gene
2 expression by modifying either the level of transcription or the mRNA stability and, consequently,
3 the content of a particular protein in the milk (Martin et al. 2002; Prinzenberg et al. 2003;
4 Szymanowska et al. 2004; Kuss et al. 2005). Moreover, it has been suggested that differential
5 expression of various milk-protein alleles is a possible result of linkage between variants of coding
6 and regulatory regions of their genes (van Eenennaam & Medrano 1991; Ramunno et al. 2005).
7 Therefore, the objective of this study was to examine associations between polymorphisms in the 5'-
8 flanking region of the goat CSN2 gene and the b-casein content in the milk.

9 Comparisons between sequences of goat CSN2A (AJ011018) and CSN201 (AJ011019) alleles
10 showed two transitions located in the 5' gene-flanking region: g.1538A>G, which was already
11 reported by Pappalardo et al. (1997), and g.1311T>C. Because nucleotides T in position 1311 and A
12 in position 1538 have been found not only in *Capra hircus* promoter sequences (DQ673920,
13 DQ673919, AY834229, AY398686, M90559, AY311384) but also in other ruminant species such as
14 *Ovis aries* (X79703), *Bubalus bubalis* (AY352050), *Bos taurus* (AJ973327, U47012, M55158,
15 U47013, X14711, M75888) and *Bos grunniens* (AF194986), these sequences might represent the
16 ancestral state of the gene.

17 To assess the effect of these two mutations in the promoter region on b-casein level in goat milk, we
18 analysed the genomic DNA of 854 goats in five breeds: 74 Garganica, 115 Malta, 90 Alpine, 95
19 Saanen and 480 Southern Italy population of undefined genetic type. Genomic DNA was extracted
20 from leucocytes (Goossens & Kan 1981) obtained from blood samples collected using Na₂EDTA as
21 anticoagulant.

22 Milk from the same individuals was previously analysed by SDS-PAGE according to Grosclaude et
23 al. (1987). Moreover, each DNA sample was genotyped for the presence of the mutation g.8915C>T
24 by allele-specific amplification (AS-PCR) (Ramunno et al. 1995). Genotyping for the transition
25 g.1538A>G was performed by MseI PCRRFLP (Pappalardo et al. 1997) and an AS-PCR was set up
26 for the transition g.1311T>C. Sequence primers used for the AS-PCR for the transition g.1311T>C

1 (184 bp in length) and amplification conditions are listed in Table S1. Primers for the amplification
2 of goat CSN2 exon 9 were also included in the PCR reactions as a positive control (360 bp in length).
3 Amplifications were performed in a 50- μ l volume containing from 100 to 200 ng of genomic DNA,
4 1x PCR Buffer, 1.5 mM MgCl₂, 200 μ M of each dNTP, 0.04% BSA, 10 pmol of each primer, 2.5 U
5 Taq DNA polymerase (Promega). Amplified fragments were analysed by means of electrophoresis
6 in 2% agarose gels stained with ethidium bromide. In the present study, the lack of b-casein in the
7 milk was always in cis with the g.1538G allele, but this allele has been associated with normal b-
8 casein levels in other studies. Therefore, this mutation is believed not to be involved in the lack of
9 expression of CSN2. On the contrary, all goats whose milk produced an SDS-PAGE pattern apparently
10 without the b-casein fraction (Fig. 1, lane 1) (15 Southern Italy and three Malta samples) and those
11 goats with a band less intense than normal (about 50%, Fig. 1, lane 2) (11 Garganica, 15 Malta and
12 81 Southern Italy animals) were homozygous or heterozygous respectively for the g.1311C and
13 g.8915T alleles (Table 1). All samples that produced a normal SDS-PAGE pattern (Fig. 1, lane 3) (63
14 Garganica, 97 Malta, 90 Alpine, 95 Saanen, 384 Southern Italy samples) were homozygous for the
15 g.1311T and g.8915C alleles (Table 1). The frequencies of the g.1311C and g.8915T alleles were
16 0.074, 0.091 and 0.115 for the Garganica and Malta breeds and Southern Italy population
17 respectively. The overall frequencies of g.1311C and g.1538G were 0.084 and 0.152 respectively.
18 Sequence analyses of the CSN2 promoter region based on TRANSFAC 7.0 database
19 (<http://www.gene-regulation.com/pub/databases.html>) indicate that the presence of g.1311C does not
20 alter or create any regulatory sites (Fig. 2). However, given the absence of the b-casein in animals
21 with this allele, it is possible that this mutation is involved in gene regulation processes. It is also
22 possible this allele is in linkage disequilibrium with a causative mutation for lack of expression.
23 Sequencing of the complete genomic sequence of the CSN2 gene in these animals is needed.

24

25 **Acknowledgement**

1 This is contribution no. 149 from the Department of Soil, Plant, Environment and Animal Production
2 Sciences, Agriculture Faculty, University of Naples Federico II.

3

4 **References**

5 Caroli A., Chiatti F., Chessa S., Rignanese D., Bolla P. & Pagnacco G. (2006) Focusing on the goat
6 casein complex. *Journal of Dairy Science* 89, 3178–87.

7

8 Chianese L., Garro G., Nicolai M.A. et al. (1993) The nature of b-casein heterogeneity in caprine
9 milk. *Lait* 73, 533–47.

10

11 Cosenza G., Pauciullo A., Gallo D., Di Berardino D. & Ramunno L. (2005) A SspI PCR-RFLP
12 detecting a silent allele at the goat CSN2 locus. *Journal of Dairy Research* 72, 456–9.

13

14 Cunsolo V., Galliano F., Muccilli V., Saletti R., Marletta D., Bordonaro S. & Foti S. (2005) Detection
15 and characterization by high-performance liquid chromatography and mass spectrometry of a goat b-
16 casein associated with a CSN2 null allele. *Rapid Communications in Mass Spectrometry: RCM* 19,
17 2943–9.

18


19 van Eenennaam A.L. & Medrano J.F. (1991) Differences in allelic protein expression in the milk of
20 heterozygous j-casein cows. *Journal of Dairy Science* 74, 1491–6.

21

22 Galliano F., Saletti R., Cunsolo V., Foti S., Marletta D., Bordonaro S. & D'Urso G. (2004)
23 Identification and characterization of a new b-casein variant in goat milk by high-performance liquid
24 chromatography with electrospray ionization mass spectrometry and matrix-assisted laser
25 desorption/ionization mass spectrometry. *Rapid Communications in Mass Spectrometry: RCM* 18,
26 1972–82.

- 1
- 2 Goossens M. & Kan Y.W. (1981) DNA analysis in the diagnosis of hemoglobin disorders. *Methods*
- 3 *in Enzymology* 76, 805–17.
- 4
- 5 Grosclaude F., Mahe´ M.F., Brignon G., Di Stasio L. & Jeunet R. (1987) A Mendelian polymorphism
- 6 underlying quantitative variations of goat α 1-casein. *Genetics Selection Evolution* 19, 399–412.
- 7
- 8 Kuss A.W., Gogol J., Bartenschlager H. & Geldermann H. (2005) Polymorphic AP-1 binding site in
- 9 bovine CSN1S1 shows quantitative differences in protein binding associated with milk protein
- 10 expression. *Journal of Dairy Science* 88, 2246–52.
- 11
- 12 Mahe´ M.F. & Grosclaude F. (1993) Polymorphism of b-casein in the Creole goat of Guadeloupe:
- 13 evidence for a null allele. *Genetics Selection Evolution* 21, 127–9.
- 14
- 15 Martin P., Szymanowska M., Zwierzchowski L. & Leroux C. (2002) The impact of genetic
- 16 polymorphisms on the protein composition of ruminant milks. *Reproduction, Nutrition, Development*
- 17 42, 433–59.
- 18
- 19 Neveu C., Molle´ D., Moreno J., Martin P. & Le´onil J. (2002) Heterogeneity of Caprine beta-casein
- 20 elucidated by RP-HPLC/MS: genetic variants and phosphorylations. *Journal of Protein Chemistry* 21,
- 21 557–67.
- 22
- 23 Pappalardo M., Rando A., Di Gregorio P., Masina P. & Ramunno L. (1997) A MseI RFLP in the 5¢
- 24 DNA region of the goat b-casein gene. *Animal Genetics* 28, 242.
- 25

- 1 Persuy M.A., Printz C., Medrano J.F. & Mercier J.C. (1999) A single nucleotide deletion resulting in
2 a premature stop codon is associated with marked reduction of transcripts from a goat b-casein null
3 allele. *Animal Genetics* 30, 444–51.
- 4
- 5 Prinzenberg E.M., Weimann C., Brandt H., Bennewitz J., Kalm E., Schwerin M. & Erhardt G. (2003)
6 Polymorphism of the bovine CSN1S1 promoter: linkage mapping, intragenic haplotypes, and effects
7 on milk production traits. *Journal of Dairy Science* 86, 2696–705.
- 8
- 9 Ramunno L., Mariani P., Pappalardo M., Rando A., Capuano M., Di Gregorio P. & Cosenza G. (1995)
10 Un gene ad effetto maggiore sul contenuto di caseina b nel latte di capra. *Proceedings XI Congress*
11 *National Scientific Association of Animal Production (ASPA)*, 185–6.
- 12
- 13 Ramunno L., Cosenza G., Rando A., Pauciullo A., Illario R., Gallo D., Di Berardino D. & Masina P.
14 (2005) Comparative analysis of gene sequence of goat CSN1S1 F and N alleles and characterization
15 of CSN1S1 transcript variants in mammary gland. *Gene* 345, 289–99.
- 16
- 17 Rijnkels M. (2002) Multispecies comparison of the casein gene loci and evolution of casein gene
18 family. *Journal of Mammary Gland Biology and Neoplasia* 7, 327–45.
- 19
- 20 Roberts B., Di Tullio P., Vitale J., Hehir K. & Gordon K. (1992) Cloning of the goat b-casein-
21 encoding gene and expression in transgenic mice. *Gene* 121, 255–62.
- 22
- 23 Szymanowska M., Malewski T. & Zwierzchowski L. (2004) Transcription factor binding to variable
24 nucleotide sequences in 5' flanking regions of bovine casein genes. *International Dairy Journal* 14,
25 103–15.
- 26



1

2 **Figure 1** Electrophoretic pattern in SDS-PAGE of individual goat samples. Lane 1, CSN201/01; lane
 3 2, CSN2A*/01 and lane 3, CSN2A*/A*, where A* $\frac{1}{4}$ CSN2A, CSN2A1, CSN2B, CSN2C, CSN2D
 4 or CSN2E (indistinguishable by SDS-PAGE).

5

tccaattggtgagagacagtcacttaggaatgctgtgtttattgcaacaaat**tgtaaagcatcttctctgagaaaaataaag**
 STAT5
 ggaaatggtgaatgggaaggatatgctttcttttgtatt**ctctttctgagaaatcag**actttttcaactgtggccttggc
 GR
 caccaaaagctaacaataaaggcatatgaagtagccaaggcctttctagtatat**ctatgacactgagttcatt**cat
 ETS
 cttttatttt**ctctgactctctctggg**tccatatgagcagctcttagaatgaatattagctgaataatccaaatacatagt
 agatggtgatttgggttttctaagcaatccaagacttgatgacagtaagatgtattaccatccaacacacatctcagca
 tgatataaatgcaaggatatttgaagaaaaatttttaattatgtcaaagtgttactttagaaggctcatctatctgtc
 ccaaagctgtgaatatatatattgaaggtaatgaatagatgaagctaaccttgtaaaaatgagtgtgaaatacaact
 acaattatgaacatctgtcactaaagaggcaaagaaacttgaagattgcttttgcaaagggtcctattaataaaaagt
 NFKB
 acttttgaggctgtggctcagactctattgttagtact**tagggtaaagacctc**ctccgtagggttccattttctttctgtc
 STAT5
 ttccctcatt**tgccctccatgaatact**agctgataaacattgactataaaaagatatgaggccaaacttgagctgtcca
 GR, PR/ER
 ttttaataaatctgtataaataatattg**ttctac**aaaagtattatctaataaatgttactttctgtcttaaaatccct
 caacaaatccccactatctagagaataagattgacattccctggaatcacagcatgctttgtctgccattatctgacccc
 tttctctttctctctctccctccatctactcctttttctctgcaattcatgaccagattcactgtttgatttggctt
 E2F
 gcatgtgtgtgtgctgagttgctgtgactggtatcaacccccatgaatgatagtcaccaggtctactgtccatgaat
 MEF2
tttcagtcaagaataactggagtggttgcatttctactccatttgattaatttagtgacttttaatttcttttcca
 tttcggggagcctattcttcttttagtctatactctcttccactcttcagggtcaaggtatcatcgtgtgcttgttagc
 HOXF 1311 PR GR, PR/ER
 ttgttactttctccattatagcttaagcac^taacaactggtcaggttgcatgaaattggttctttgtgtggcctgtat
 c
 atttctgtgtgtattagaatttaccccaagatctcaaagaccactgaataactaaagagacctcattgtggttacaata
 SP1 EF2
 atttg**gggactgggccaaaactt**ccgtgcatcccagccaagatctgtagctactggacaatttcatttcccttatcagat
 1538 AP-2alpha MGF milk box element C/EBPalpha
 tgtgagttattcctgtt^aaaatgctccccagaatttctggggacagaaaaataggaagaatt**catttccctaatcat**gag
 g
 MGF OCT-1 SV40 core enh.seq. OCT-1 TATABOX
 atttctaggaatt**caaat**ccactattggtttttatttcaaccagaaaaattagcatgcatttaaatactatataaaacag

 Exon 1
 ccactaaatcagatcattATCCATTCAGCTTCTCCTTCACTTCTTCTCCTCTACTTTGGAAAAAAG.....

1
2
3
4
5
6
7
8
9
10
11
12

Figure 2 Nucleotide sequence of the 5' flanking region and partial exon 1 of goat CSN2 gene (numbering is according to AJ011018). Congruent and putative factors are double underlined, bold letters, or boxed. Transcription factor abbreviations: C/EBP, CCAAT/enhancer-binding protein; OCT-1, nuclear factor octamer-1; PR, progesterone receptor; MEF2, myocyte-specific enhancer-binding factor; EF2, E2F-myc activator/cell cycle regulator; SP1, GC-box factors; STAT5, signal transducer and activator of transcription 5; MGF, mammary gland factor recognition sequence; AP-2, alpha, activating enhancer binding protein 2 alpha; GR, glucocorticoid response element; NFKB, nuclear factor kappa B; HOXF, factors with moderate activity to homeo domain consensus sequence, ETS, E26 transformation-specific family of transcription factors.

1

Table 1 Distribution of genotypes at the goat CSN2 locus.

SDS-PAGE	g.1311T>C			g.8915C>T			g.1538A>G			Total
	TT	TC	CC	CC	CT	TT	AA	AG	GG	
<i>CSN2</i> ^{A*/A*}	729	-	-	729	-	-	648	66	15	729
<i>CSN2</i> ^{A*/0}	-	107	-	-	107	-	-	86	21	107
<i>CSN2</i> ^{0/0}	-	-	18	-	-	18	-	-	18	18
Total	729	107	18	729	107	18	648	152	54	854

2

3 A*, alleles associated with a normal content of b-casein (*CSN2A*, *CSN2A1*, *CSN2B*, *CSN2C*,
4 *CSN2D* and *CSN2E*).