

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Mitotic index and aneuploidy variation in growing day 2- to day 7- IVP bovine embryos of the Agerolese breed of cattle

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1507152> since 2018-03-22T18:03:32Z

Published version:

DOI:10.1007/s10577-012-9313-0

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Rubessa M., Iannuzzi A., Peretti V., Pauciullo A., Cosenza G., Ramunno L., Iannuzzi L., Rubes J., Di Berardino D. Mitotic index and aneuploidy variation in growing day 2- to day 7- IVP bovine embryos of the Agerolese breed of cattle. *Chromosome Research* (2012) 20: 795-796

DOI: 10.1007/s10577-012-9313-0

The publisher's version is available at:

<https://link.springer.com/article/10.1007%2Fs10577-012-9313-0>

When citing, please refer to the published version.

Link to this full text:

<https://link.springer.com/article/10.1007%2Fs10577-012-9313-0>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

Mitotic index and aneuploidy variation in growing day 2- to day 7- IVP bovine embryos of the Agerolese breed of cattle

Rubessa M.¹, Iannuzzi A.², Peretti V.³, Pauciullo A.¹, Cosenza G.¹, Ramunno L.¹, Iannuzzi L.², Rubes J.⁴, Di Berardino D.¹

¹DISSPAPA Dept., University of Naples “Federico II”, Portici, Italy

²National Research Council (CNR), ISPAAM, Lab. of Animal Cytogenetics and Gene Mapping, Naples, Italy

³DISCIZIA Dept., University of Naples “Federico II”, Naples, Italy.

⁴Veterinary Research Institute, Brno (Czech Republic).

The present study reports on the variation of the mitotic index in growing day 2- to day 7- IVP bovine embryos of the Agerolese breed of cattle. After IVM (24 h), COCs were transferred into 25/well with 300 µl IVF-TALP and covered with mineral oil. Frozen sperm from a bull were selected by centrifugation on a Percoll discontinuous gradient (45–80 %). The pellet was diluted in IVF medium and added to the COCs at the concentration of 1×10^6 sperm/mL. Gametes were co-incubated for 20–22 h at 39 °C, in 5 % CO₂ in air. After co-incubation, presumptive zygotes were vortexed to remove cumulus cells and randomly allocated in six groups, each into 400 µl of SOF medium, with 30 µl/ml essential amino acids, 10 µl/ml non-essential amino acids, 0.34 mM tri-sodium citrate, 2.77 mM myo-inositol, and 5 % BS. Zygotes were incubated in a humidified mixture of 5 % CO₂, 7 % O₂ and 88% N₂ in air at 39 °C for 20–22 h. Starting on Day 0 (IVF day), embryos were taken out of the incubator at day 2(48 h), 3 (72 h), 4 (96 h), 5 (120 h), 6(144 h) and 7(168 h), examined under a stereomicroscope, treated in a lysing buffer (0.01 N HCl, 0.1 % Tween 20) for 30 s, transferred in a small droplet to a precleaned slide and fixed with methanolacetic acid (3:1). Out of 178 embryos (3,100 cells) analyzed, the mitotic index was 19.6 % (28/143 cells) at day 2, 18.6 % (41/221 cells) at day 3, 10.3 % (27/263 cells) at day 4, 7.1 % (26/367 cells) at day 5, 0.7 % (5/681 cells) at day 6 and 0.6 (9/1425 cells) at day 7. FISH analysis is undergoing and will be reported elsewhere.

Acknowledgements: This work was supported by the RARECA- PSR 214 project of the Campania Region (Italy).