

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## A cytogenetic investigation on the yak (*Bos grunniens*) reared in central Italy

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1507158> since 2018-03-18T15:54:03Z

*Published version:*

DOI:10.1007/s10577-008-1922-2

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

D. Nicodemo, A. Pauciullo, A. Castello, G. Cosenza, V. Peretti, A. Perucatt, G.P. Di Meo, G. Ficco, L. Ramunno, L. Iannuzzi, J. Rubes and D. Di Bernardino. A cytogenetic investigation on the yak (*Bos grunniens*) reared in central Italy. *Chromosome Research* (2008) 16:1027-1071  
DOI: 10.1007/s10577-008-1922-2

**The publisher's version is available at:**

<https://link.springer.com/article/10.1007%2Fs10577-008-1922-2>

**When citing, please refer to the published version.**

**Link to this full text:**

<https://link.springer.com/article/10.1007%2Fs10577-008-1922-2>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

## **A cytogenetic investigation on the yak (*Bos grunniens*) reared in central Italy**

D. Nicodemo<sup>1</sup>, A. Pauciullo<sup>1</sup>, A. Castello<sup>2</sup>, G. Cosenza<sup>1</sup>, V. Peretti<sup>3</sup>, A. Perucatti<sup>4</sup>, G.P. Di Meo<sup>4</sup>, G. Ficco<sup>5</sup>, L. Ramunno<sup>1</sup>, L. Iannuzzi<sup>4</sup>, J. Rubes<sup>6</sup> and D. Di Berardino<sup>1</sup>

<sup>1</sup>Department of Soil, Plant, Environment and Animal Production Sciences, University of Naples Federico II, Portici, Italy

<sup>2</sup>Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Cientificas, Madrid, Espana

<sup>3</sup>Department of Animal Science and Food Inspection, University of Naples Federico II, Naples, Italy

<sup>4</sup>Research National Council (CNR), ISPAAM, Laboratory of Animal Cytogenetics and Gene mapping, Naples, Italy

<sup>5</sup>CRA-PCM, Research Center for the production of meat and Genetic improvement, Monterotondo, Italy

<sup>6</sup>Veterinary Research Institute, Brno, Czech Republic

The domestic yak (*Bos grunniens*) is an important Bovidae species which in Asia plays a remarkable role for the economy of the Tibetan highlands (over 4000 m above sea level.). Despite its strategic and economical importance, this species has been almost neglected, from a cytogenetic point of view, as demonstrated by the few papers present in the literature. To fill this gap, we undertook a cytogenetic investigation by using blood samples of eight yak bulls recently imported into central Italy. CGR-banding, silver staining and FISH-techniques were applied. The preliminary results were as follows: (a) the chromosomal make-up of the yak was  $2n=60,XY$ , (b) the animals investigated showed normal karyotype, (c) the frequency of chromosome/chromatid breaks was similar to that of cattle (3.7 vs 3.0 %, respectively), (d) the mean rate of SCE/cell at 10 2g/ml (f.c.) was significantly ( $P<0.001$ ) lower compared to that of cattle (5.2 vs 8.3, respectively), (e) the GTG-RBA-RBG banding patterns Y at the 400 bands level of resolution Y did not reveal structural differences compared to cattle, (f) the CBA-banding pattern was also similar to cattle, i.e. no C-bands on the subcentromeric X, while the Y- chromosome showed heterochromatic tips in the short arms.

Zoo-FISH with bovine painting probes derived from microdissected chromosomes 5-X-Xcen and Y-upon yak metaphase chromosomes showed complete hybridization, thus confirming the close homology between the two species. In addition, FISH-mapping with bovine BAC-clones containing ZFY and SRY genes to the yak Y-chromosome also revealed the same localization as reported in cattle (Yp12.2 and Yq23dist, respectively).

However, the fact that *Bos taurus* x *Bos grunniens* F1 male hybrids are sterile, while the females are normally fertile, suggests that the main difference between yak and cattle might be in the X-Y pseudoautosomal (PAR) region of the two species. Further investigation, therefore, is necessary with more effective cytogenetic techniques, such as high resolution banding, fine gene mapping analysis and DNA sequencing.