



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

# Tannery mixed liquors from an ecotoxicological and mycological point of view: Risks vs potential biodegradation application

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1660424 since 2018-02-21T14:56:09Z
Published version:
DOI:10.1016/j.scitotenv.2018.01.240
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)





## This is the author's final version of the contribution published as:

Valeria Tigini, Federico Bevione, Valeria Prigione, Anna Poli, Lucrezia Ranieri, Francesco Spennati, Giulio Munz, Giovanna Cristina Varese, Tannery mixed liquors from an ecotoxicological and mycological point of view: Risks vs potential biodegradation application, Science of the Total Environment, 627, 2018, pagg. 835-843, DOI: 10.1016/j.scitotenv.2018.01.240

### The publisher's version is available at:

https://doi.org/10.1016/j.scitotenv.2018.01.240

When citing, please refer to the published version.

## Link to this full text:

https://doi.org/10.1016/j.scitotenv.2018.01.240

This full text was downloaded from iris-Aperto: https://iris.unito.it/

University of Turin's Institutional Research Information System and Open Access Institutional Repository

1 TITLE: Tannery mixed liquors from an ecotoxicological and mycological point of view: risks vs potential biodegradation application

AUTHORS: Valeria Tigini<sup>1\*</sup>, Federico Bevione<sup>1</sup>, Valeria Prigione<sup>1</sup>, Anna Poli<sup>1</sup>, Lucrezia Ranieri<sup>1</sup>, Francesco
 Spennati<sup>2</sup>, Giulio Munz<sup>2</sup>, Giovanna Cristina Varese<sup>1</sup>

AFFILIATION: <sup>1</sup> Department of Life Sciences and Systems Biology, University of Turin, viale Mattioli 25,
10125 Turin, Italy. <sup>2</sup> Department of Environmental and Civil Engineering, University or Florence, via Santa
Marta 3, 50139 Firenze, Italy.

8

\*CORRESPONDING AUTHOR: Mailing address: Department of Life Sciences and Systems Biology,
University of Turin, viale Mattioli, 25, 10125 Turin, Italy. Phone: +39-011-6705964. Fax: +39-011-6705962. Email: valeria.tigini@unito.it

12

#### 13 Abstract

14 Fungi are known to be present in the activated sludge of wastewater treatment plants (WWTP). Their 15 study should be at the base of an overall vision of the plant effectiveness and of effluents sanitary impact. 16 Moreover, it could be fundamental for the implementation of successful bioaugmentation strategies aimed at the 17 removal of recalcitrant or toxic compounds. This is one of the first studies on the cultivable autochthonous 18 mycoflora present in the mixed liquors of two WWTP treating either vegetable or chromium tannery effluents. 19 All samples showed a risk associated with potential pathogens or toxigenic species and high ecotoxicity 20 (Lepidium sativum and Raphidocelis subcapitata were the most sensitive organisms). Diverse fungal populations 21 developed, depending on the origin of the samples (63% of the 102 identified taxa were sample-specific). The 22 use of a fungistatic was determinant for the isolation and, thus, for the identification of sample-specific species 23 with a lower growth rate. The incubation temperature also affected the mycoflora composition, even though at 24 lower extent. A selective medium, consisting of agarised wastewater, allowed isolating fungi with a 25 biodegradation potential. Pseudallescheria boydii/Scedosporium apiospermum species complex was 26 ubiquitously dominant, indicating a possible role in the degradation of pollutants in both WWTP. Other species, 27 i.e. Trichoderma spp., Trematosphaeria grisea, Geotrichum candidum, Lichtheimia corvmbifera, Acremonium 28 furcatum, Penicillium simplicissimum, Penicillium dangeardii, Fusarium solani, Scopulariopsis brevicaulis 29 potentially could be involved in the degradation of specific pollutants of vegetable or chromium tannery wastewaters. However, several of these fungi are potential pathogens and their application, for an *in situ*treatment, must be carfully evaluated.

32

33 Key Words: Tannery wastewater, activated sludge, fungi, autochthonous mycoflora, ecotoxicity.

34

#### 35 1 Introduction

36 The secondary treatments, which consist of biological processes operated with either suspended or 37 attached biomasses, are generally the core of wastewater treatment plants (WWTPs) (Leyva-Diaz et al., 2017). 38 The knowledge on composition and dynamics of microbial community has been fundamental for the 39 development of biological treatment technologies (Orhon, 2015). However, studies were mainly focused on the 40 bacterial fraction of these complex ecosystems, as well as mathematical models applied to describe the processes 41 (Orhon, 2015). Recently, the literature has pointed out a consistent presence of autochthonous mycoflora in the 42 activated sludge of WWTPs treating paper mill, vinery or municipal wastewaters (Grabinskaloniewska et al., 43 1993; Awad and Kraume, 2011; Evans and Seviour, 2012). Its characterisation should be included in an overall 44 vision of the effectiveness and impact of a treatment plant, considering functional, ecological and sanitary 45 aspects (More et al., 2010; Awad and Kraume, 2011; Korzeniewska, 2011).

The role of fungal organisms in the depuration process has been recently demonstrated. In particular,
fungi seem to perform a complementary action with respect to bacteria in the removal of pollutants from
wastewaters and can actively interact with them (Liu et al., 2017; Svobodová et al., 2016; Anastasi et al., 2012).

The study of autochthonous fungi in WWTPs may acquire particular importance when recalcitrant or toxic pollutants are present in wastewaters, in order to improve depuration performance by bioaugmentation or biostimulation of naturally present fungi (More et al., 2010; Djelal and Amrane, 2013; Herrero and Stukey, 2015).

Tannery wastewaters, regardless of the type of industrial process (chromium or vegetable), are among the most difficult to treat, basically on account of their recalcitrance and/or their toxicity towards bacteria (Lofrano et al., 2013). On the contrary, the ability of fungi in the degradation of tannery pollutants has been already demonstrated (Sharma and Malaviya, 2016; Zhang et al., 2015).

57 Besides the study of diversity and ecological function of the microbial community, a successful 58 bioaugmentation strategy should contemplate the assessment of possible toxicity effects towards the biota 59 (Herrero and Stukey, 2015). Ecotoxicity tests, associated with chemical and biological characterisations, can be a informative tool for the efficiency evaluation and the management of the secondary treatment in WWTPs
(Chapman, 2000). Nevertheless, these procedures are not so diffused at industrial level. This is mainly due to the
lack of knowledge about the methods for data interpretation and to the complexity of the information that these

63 analyses can provide (Chapman, 2000).

The present study is focused on the analysis of cultivable autochthonous mycobiota present in mixed liquors of the oxidation tanks of two wastewater treatment plants, which collect vegetable and chromium tannery effluents, respectively. Selective media were exploited for the isolation of fungi with biodegradation potential and/or the ability to compete with other microorganisms. Moreover, the samples were incubated at both 25 °C and 15 °C, in order to acquire information about the effect of seasonal temperature fluctuations on mycoflora development. Finally ecotoxicological aspects of the samples were assessed by means of four bioassays.

70

#### 71 2 Materials and Methods

72 2.1 Wastewaters

73

The samples (three in total) were collected from two WWTPs located in Tuscany (Italy):

74 Co – mixed liquor from the aerobic tanks of Cuoiodepur vegetal tannery WWTP (San Miniato, Pisa,
75 Italy).

F1 – effluent from the settler of the first aerobic stage of the chromium tannery WWTP of Consorzio
Aquarno SpA (Santa Croce sull'Arno, Pisa, Italy).

F2 - mixed liquor from the second biological stage of chromium tannery wastewater treatment plant of
 Consorzio Aquarno SpA (Santa Croce sull'Arno, Pisa, Italy).

80 Cuoiodepur manages a consortial WWTP, where the effluents of about 100 tanneries, operating 81 vegetable tanning process, are treated. The influent wastewaters are characterized by a high organic and nitrogen 82 load (COD > 15000 mg  $L^{-1}$ ), high salinity and by the presence of natural and synthetic tannins. The Cuoiodepur 83 WWTP treats about 5000 m<sup>3</sup> d<sup>-1</sup> of tannery wastewaters and the treatment train is composed of: pretreatments, 84 simultaneous equalization and sulphide oxidation with pure oxygen, primary settling, conventional suspended 85 activated sludge (denitrification nitrification) biological stage and coagulation flocculation as tertiary treatment. 86 The temperature of the biological reactors ranges between 19 °C and 35 °C, the hydraulic retention time (HRT) 87 is about 3 days and the solids retention time (SRT) is usually between 50 and 70 d.

Aquarno is a joint-stock company by large majority private operating in the Tuscany tannery district for
 the treatment of chromium tannery wastewaters. The Aquarno WWTP treats about 12000 m<sup>3</sup> d<sup>-1</sup> of chromium

90 tannery wastewaters. The treatment train is composed of: pretratments, aerobic activated sludge system (at low 91 SRT to remove sulphide), second conventional activated sludge system for nitrogen and carbon removal, and 92 Fenton tertiary treatment. The average temperature of the influent is in the range 25-30 °C, the HRT of 93 biological sections is about 7 days and the SRT of the second biological stage is close to 30 days.

94

#### 95 2.2 Chemical analyses

96 The pH was determined with a Hach-Lange's probe. The ammonium and metals were measured by 97 means of a nitrogen analyser (TOC-L, Shimazdu) and an inductively coupled plasma mass spectrometry (ICP-98 MS) (Perkin Elmer), respectively. The Chemical Oxygen Demand (COD) and soluble Chemical Oxygen 99 Demand (sCOD) were determined using Hach-Lange's cuvettes after filtration (at 0.45 μm). The Total 90 Suspended Solids (TSS) were measured as the dry weight (1 hour at 105 °C) of the residue of the filtered 91 sample. The Volatile Suspended Solids (VSS) were measured as the dry weight (30 min at 570 °C) of the residue 92 of the filtered sample.

103

#### 104 2.3 Ecotoxicity tests

A battery of four bioassays was performed in order to evaluate the toxicity of the samples. The target organisms were: the unicellular green alga (I) *Raphidocelis subcapitata* (Korshikov) Nygaard, Komárek, J.Kristiansen & O.M. Skulberg (UNI EN ISO 8692:2005); the dicotyledonous plants (II) *Cucumis sativus* L. and (III) *Lepidium sativum* L. (UNICHIM N. 1651, 2003); the monocotyledonous plant (IV) *minor* L. (ISO SO/WD 20079). The samples were filtered (Whatman type 1) for the execution of algal test only. In fact, filtration was needed to avoid interference with the spectrophotometric lectures performed at the end of this test.

Each dose-response curve consisted of six dilutions (in triplicate for plant test and in quadruplicate for algal test) and the control was performed with four and six repetitions for plant and algal tests, respectively.

113 Significant differences between dose-effect regression lines were analysed using T test (p<0.05 for line 114 slope, p<0.001 for translation), for all the possible pairs.

115

#### 116 2.4 Isolation and identification of autochthonous mycoflora

Aliquots of 1 mL of each sample were placed in Petri dishes (16 cm diam.) containing 30 mL of culture medium. Three different media were used: a modified Malt Extract Agar (MEAp) (agar 20 g, glucose 2 g, malt extract 2 g, peptone 0.2 g, water up to 1 L); Dichloran Rose Bengal Agar (DRBC 31.5 g, water up to 1 L); Wastewater-Agar (WA; agar 20 g, glucose 2 g, sample supernatant after 5 min at 10000 rpm up to 1 L). A set of three antibiotics was added to all media: streptomycin 0.015 g L<sup>-1</sup>, chloramphenicol 0.05 g L<sup>-1</sup>, and ampicillin 0.05 g L<sup>-1</sup>. The trial was performed with 20 replicates for each medium and sample. Ten replicates were incubated at 25 °C and ten at 15 °C, in the dark.

124 At regular intervals of time, the colony forming units (CFU) were counted and the different fungal 125 morphotypes were isolated in pure cultures. Fungi were identified conventionally, according to their 126 macroscopic and microscopic features. After determination of their genera (Domsch et al., 1980; Kiffer and 127 Morelet, 1997; von Arx 1981), they were transferred to the media recommended by the authors of selected genus 128 monographs for species identification. Molecular identification was performed by amplification and sequencing 129 of specific markers: actin for Cladosporium (Bensch et al., 2012); β-tubulin for Penicillium and Aspergillus 130 (Glass e Donaldson 1995); D1/D2 region for yeasts (Fell et al., 2000); Internal Transcribed Spacer (ITS) for 131 other genera (White et al., 1990). The resulting sequences were compared with reference sequences in online 132 databases provided by the CBS-KNAW Collection (Westerdijk Fungal Biodiversity Institute, Utrecht, The 133 Netherlands) and the NCBI National Center for Biotechnology Information (USA).

The nonparametric Mann-Whitney test ( $p \le 0.05$ ) was run to assess the significance of the quantitative (fungal load) differences between all possible permutations of the trial. Differences within factors (tannery effluent, incubation temperature, and culture medium) were evaluated by applying a Permutational Multivariate Analysis of Variance (PERMANOVA; p < 0.05) and visualised by Non-Metric Multidimensional Scaling (NMDS). The contribution of individual species (in percentage) to the diversity observed was assessed by SIMilarity PERcentage (SIMPER) analysis.

140

- 141 **3** Results and discussion
- 142 *3.1 Chemical analyses*

The results of chemical analyses are shown in Table 1. The pH of the samples ranged around neutral values, from 7.22 of F2 to 7.66 of Co. All tannery samples showed high values of COD (10000-24740 mg L<sup>-1</sup>) and sCOD (362-518 mg L<sup>-1</sup>). It is noteworthy that 50% of the sCOD was due to tannins in Co sample (Munz et al., 2009). Both in Co and F1, the TSS and VSS were very high ( $\geq$ 12.61 g L<sup>-1</sup> and  $\geq$ 9.61 g L<sup>-1</sup>, respectively), while in F2 these values decreased (7.79 g L<sup>-1</sup> and 5.77 g L<sup>-1</sup>, respectively). The ammonium was 278 mg L<sup>-1</sup> in F1 sample, whereas for Co and F2 it was 1.04 mg L<sup>-1</sup> and 9.65 mg L<sup>-1</sup>, respectively. Eventually, supernatant colour was dark brown for all the samples.

According to the literature, tannery wastewaters have very high COD, high nitrogen, and low phosphate 151 concentrations (Prigione et al., 2009; Kim et al., 2013; Lofrano et al. 2013). On the whole, the three samples, Co, 152 F2 e F1, were typical of tannery WWTPs, when compared to the data in the literature.

153

154 3.2 Sample ecotoxicity

155 In this work four ecotoxicity tests were applied in order to evaluate the effectiveness of tannery 156 wastewater secondary treatment and the evaluation of potential impact of the samples on the environment.

157 The inhibition effects of samples on *Raphidocelis subcapitata* are shown in Figure 1. The maximum 158 testable concentration was 70% for F2, 50% for Co and 40% for F1. At higher doses, the algal concentration was 159 not detectable on account of the deep colour impact on the spectrophotometric analysis.

160 The recorded data were almost linearly distributed only for F2 and Co samples (see  $R^2>0.9$  in Figure 1). 161 On the contrary, F1 points seem to be distributed according to an S-shaped curve ( $R^{2}<0.9$ ). The three tannery 162 effluents showed no significant differences among angular coefficients of dose-effect regression lines, indicating 163 a similar toxicity effect for the three samples. However, this elaboration is possible only for positive inhibition 164 values, which are the minority of the recorded data. Thus, the scarcity of data exploitable for statistical analysis 165 affected the reliability of this elaboration.

166

167

The results about the ecotoxicity test with Lepidium sativum and Cucumis sativus are shown in Figure 2.

168 The phytotoxicity tests towards terrestrial dicotyledons (Figure 2) indicated a clear linearity between 169 doses and effects only for F2 ( $R^2>0.9$ ). On the contrary, Co and F1 points are guite scattered indicating no linear 170 relation between dose and effect, in particular towards C. sativum. Moreover, most of the tested dilutions 171 resulted in negative inhibition effect (biostimulation), decreasing the number of effective data for statistical 172 elaboration. No significant difference among angular coefficients of dose-effect regression lines was found. Also 173 in this case, as for the algal test, the limited number of doses showing effects with positive value affected the 174 strength of statistical elaboration.

175 L. minor, the tested aquatic monocotyledon, was not sensitive towards the three samples with the 0% 176 inhibition of fronds number. Also the plant dry weight was not significantly affected by the exposition to the 177 samples, with an average inhibition ranging from -17% to 8%.

178 By comparing the observed data, a first consideration on the sensitivity of the tested organisms towards 179 this kind of wastewaters can be drawn. Among the three plants, L. sativum was the most sensitive, with

180 inhibition of 65-90% at 80% sample dose. Its higher sensitivity with respect to C. sativus and L. minor was 181 already signalled in a study on the landfill leachate (Tigini et al., 2014). On the contrary, L. minor was the least 182 sensitive, pointing out its inadequacy for the ecotoxicity assessment of different kinds of wastewaters (Tigini et 183 al., 2014; Tigini et al., 2016). Instead the alga showed high sensitivity to the sample, comparable to L. sativum. 184 The alga has been previously signalled as particularly sensitive towards textile effluents (Bedoui et al., 2015, 185 Novotný et al., 2006). However, a comparison with the other organisms is hard to define, on account of the 186 limits due to the analytical interference caused by the samples colour. On the other hand, this acquatic organism 187 can relieve the so-called physical toxicity, e.g. the growth inhibition due to the decrease of light penetration that 188 turns in the reduction of photosynthesis. Moreover, the unicellular alga coud be more exposed to sample toxicity 189 on account of its distinctive higher surface/volume ratio. Eventually, the presence of protozoa with dimension 190 under 11 µm, not retained by the paper filter, could have preved the alga, reducing its growth rate at the end of 191 the experiment.

192 Another consideration can be drawn on the effectiveness of the second phase of biological treatment of 193 chromium tannery wastewaters (comparing F1, sampled before the second aerobic stage treatment, and F2, 194 sampled from the mixed liquor of the second aerobic stage treatment), in terms of toxicity removal. Despite the 195 ammonium drastically decreased, the algal test did not relieve significant difference (T test) both in slope and 196 translation of their regression lines. On the contrary, both the terrestrial plant tests underlined a difference in the 197 translation of the two regression lines, with an increase in wastewater toxicity after the treatment between 35-198 38%. The reasons of this toxicity increase should be deeper investigated, since all the main chemical features 199 decreased after the treatment, but the phosphorous (which should not reasonably be responsible for this toxicity 200 increase).

201 Interestingly, no difference in chromium (F2) and vegetable (Co) tannery samples of mixed liquor was 202 detected by all tested organisms. This indicates that chromium in treated tannery wastewater is not of concern. 203 Chromium is generally present in low concentration and in trivalent form, the less toxic one (Prigione et al., 204 2009). Actually, hexavalent chromium is mostly adsorbed on the activated sludge flocks. Thus, the alga could 205 not come in contact with this pollutant, since we tested the supernatant in order to avoid spectrophotometric 206 interference. In the other tests, we analysed the entire samples, because in these cases sample turbidity did not 207 represent an analytical limitation. However, pollutants adsorbed on activated sludge flocks could be not 208 bioavailable to the test organisms. As a consequence, even in this case the toxicity due to these components 209 could be not detectable.

210 Noteworthy, at the maximum concentration, all the samples were highly toxic, even exceeding the legal 211 threshold value (i.e. 50% for the Italian law D.L. 152/06 and further modifications). This could be due to the Cl concentration, which exceeded 2900 mg  $L^{-1}$  in all the samples. As a consequence, a last comment is addressed to 212 213 the use of dewater excess sludge and treated wastewaters in agriculture. This procedure is seen as a desirable 214 option, in the light of Directive 2008/98/EC, which gives added value to wastes through their integrated 215 management, moved by the aspiration to the reduction of mineral fertilizer use (Zhang et al., 2015). The 216 hypothesized multiple benefits for agriculture and the environment are actually to be confirmed, by evaluating 217 also the toxicity of these wastes. On the base of the obtained results, the tested samples are still heavily toxic for 218 different terrestrial plants. The present results are in line with another study, which warns on the possible 219 inappropriateness of direct application of activated sludge and treated effluents (Alvarenga et al., 2007). 220 Moreover, the assessment of their impact on the human health should be evaluated.

221

#### 222 *3.3 Characterisation of cultivable autochthonous fungi*

The total fungal load of the three wastewaters was particularly high, ranging from 2630 CFU 100 mL<sup>-1</sup> (Co on WA 15 °C) to 11020 CFU 100 mL<sup>-1</sup> (F2 on DRBC 25 °C). On the whole, a total of 102 fungi were isolated from the three samples (complete list in Table A of Supplementary materials). The most abundant species and their percentage contribution to intragroup (each column) similarity are reported in Table 2.

There are few studies focusing on the characterisation of the cultivable mycobiota associated to tannery industries. Most of them are referred to the contamination of work environments, i.e. indoor/outdoor air, superficies, raw materials, etc. (Skóra et al., 2014; Gutarowska et al., 2014; Castellanos-Arevalo et al., 2016). In others, the mycoflora associated to samples of the final effluent was characterised (Prigione et al., 2009). Nevertheless, a common key aspect emerged about tanneries as high-risk work environments, due to the massive presence of fungal propagules in bioaerosol, up to 2800 CFU m<sup>-3</sup> (Skóra et al., 2014).

As far as we know, this work is one of the first reported studies focusing on a detailed analysis of the cultivable fungi present in influents and mixed liquors of a tannery WWTP.

In this study, several opportunistic pathogenic fungi, such as *Aspergillus fumigatus*, *Candida tropicalis*, *Geotrichum candidum*, *Lichtheimia corymbifera*, *Rhodotorula* sp., and *Pseudallescheria boydii/Scedosporium apiospermum* species complex were isolated. These fungi are emerging pathogens with an incredibly high ecological success in the last decades. They are often associated with polluted sites and their rapid spread is particularly alarming on account of their increasing impact on humans (Rougeron et al., 2015). Moreover, several potential toxigenic species have been identified in this study, in particular *Aspergillus flavus* (see Table
A, in Supplementary material).

Besides potential pathogens and toxicogenic species, even the presence of other fungi, characterised by high sporulating rate, can be considered of concern. Actually, they consistently contribute to the bioaerosol with 0.65-2.1 µm diameter (classified as PM 2.5), which is one of the main professional risk factors in this field. In particular, five filamentous fungi have been determined as indicator of high health risk in tannery work environments according to their prevalence, source of isolation and health implications (Sokora et al., 2014). Noteworthy, two of them, *Cladosporium cladosporioides* and *Penicilliun crustosum*, were found in this work, too.

249

250 3.3.1 Effect of sample origin

Quantitative (fungal load) differences among the three samples were significant in most cases (Figure3).

Co had a lower fungal load with respect of F2 and F1. It showed a similar fungal load to F2 only in MEA 15 °C and to F1 only in DRBC 15 °C. As regards the comparison between the two samples coming from chromium tannery, F2 and F1, they had different fungal loads (with the sole exception of WA 25 °C). However, which of them achieved the highest fungal load depended on the trial (F2>F1 in MEAp 25 °C, DRBC 25 °C, and DRBC 15°C; whereas F2<F1 in MEAp 15 °C, and WA 15 °C).

The three samples were qualitatively different, too. Among the 102 taxa, only 20 were common to all the samples. Their origin can be likely ascribable to air or working surfaces and they accidentally can be grown in WWTPs tanks. Actually, some of them are common air contaminant (e.g.. *Cladosporium* spp., *Penicillium* spp., and *Trichoderma* spp.), and can be easily sampled from outdoor environments (Skóra et al., 2014). Others are well known agents of deterioration of tanned leathers or chemicals used in this kind of industries, e.g. *Chaetomium globosum, Aspergillus* spp., *Fusarium* spp. (Orlita, 2004).

264 On the contrary, most of the isolated taxa were exclusive of a single sampling site. In particular, 30
265 were exclusive of Co, 11 were exclusive of F2, and 26 were exclusive of F1.

The differences recorded in the fungal composition of the three tanks indicate the development of three separated ecological niches. Actually, activated sludge in WWTPs tends to form a characteristic community mainly in relation to the influent composition (Evans and Seviour, 2012). This was particularly evident with the medium DRBC, on which the fungi of the three samples showed a similarity lower than 26% (Figure 4). 270 Probably, this result was due to the inhibition of fast growing fungi caused by DRBC medium that turned in the271 development of sample-specific fungi.

272

#### 273 3.3.2 Effect of medium

274 Selective media with low glucose concentration (MEAp) or fungistatic (DRBC) were adopted, in order 275 to limit the rapid growing fungi. Actually in preliminary exploring trials they were particularly abundant in all 276 the samples. Nevertheless, these fast growing fungi still caused the failure of isolation of some fungal colonies 277 with a slower growth rate (indicated as "unidentified" and with taxon name ending with sp. in Table 2 and in 278 Table A of Supplementary material). Moreover, a medium containing wastewater (WA) as principal source of 279 carbon was used in order to select fungi with a biodegradation potential toward the main organic compounds 280 present in the samples. The use of different media significantly affected both fungal load (Figure 3) and 281 biodiversity (Figure 5). Intergroup dissimilarity percentages (PERANOVA test) were in the ranges of 77-100% 282 in Co, 51-77% in F2 and 61-96% in F1.

In four out of six cases, samples inoculated on DRBC showed the highest fungal load. The two exceptions were Co 25 °C and F1 15 °C, in which fungal load detected on DRBC was not significantly higher with respect to other media. The higher fungal load retrieved on DRBC is probably due to its higher nutrient's concentration (up to 25 fold for N source and five fold for C source) with respect to the other media, in which the low nutrient concentration could represent a limiting factor for fungal growth. Besides, higher peptone concentration could activate fungal propagules of dormant species (Thanh and Nout, 2004).

In two cases of particular interest (F2 15 °C and F1 25 °C), WA showed a significantly higher fungal load with respect to MEAp. Actually WA has the same glucose concentration, but lacks of malt and peptone. Moreover, it definitively exercised a selection due to the toxicity of the wastewater used to prepare the culture medium.

Among the most frequent taxa, some highly sporulating fungi are present. Thus, the fungal load of these fungi could be overestimated, due to the culture dependent method used for the isolation and identification of fungi (Evans and Seviour, 2012). However, the massive presence of these fungi in such selective medium indicates at least a great tolerance to pollutants in WWTPs tanks.

DRBC was the richest medium in terms of taxa diversity (17-25 taxa), whereas, MEAp showed the lowest fungal biodiversity (9-21 taxa). Surprisingly, WA showed also a good biodiversity (11-21 taxa). The presence of toxic or recalcitrant substrates generally leads to the reduction of microbial biodiversity in favour of the development of few species capable of tolerating such limiting factors (Cudowski et al., 2015). However, the relative abundance changed drastically, since the dominant species on WA (*Scedosporium apiospermum* up to 62% in Co, *Pseudallescheria boydii* up to 51% in F1 and 52% in F2) were sporadically counted in other media (0-20%). These species, which are an anamorphic and teleomorphic forms, respectively, belonging to the *Pseudallescheria boydii* species complex, have been already signalled to degrade several xenobiotics (Claussen and Schmidt, 1998; Ishii et al., 2009; Santos and Linardi, 2004) and are typically present in human-impacted areas (Rougeron et al., 2015).

Even other species, present in lower percentages (8-20%), have interesting degradation capability, i.e.
 *Trematosphaeria* (Mabrouk et al., 2012), *Scopulariopsis* (Verma et al., 2017) and *Geotrichum* (Ayed et al.,
 2016).

310 The use of WA culture medium, thus, decreased the species stocastically present, enhancing the 311 development of a few species probably active in the degradation of pollutants in the WWTPs. These fungal 312 species could potentially belong to the theoretical 20% of determinative species in the community that cause the 313 80% of the WWTP effectiveness, according the 20-80% ratio among cause-effect relationship (Pareto law). The 314 knowledge of this determinative 20% species is at the base of a successful bioaugmentation strategy, since not 315 all the species retrieved in a certain environment can effectively degrade of pollutans (Herrero and Stuckey, 316 2015). However, since most of them imply a concrete risk for human health, their massive exploitation in open 317 WWPTs should be accurately evaluated.

Intuitively, the fungi isolated form the WA media consisting of chromium tannery samples, which are subsequent in the same plant (F1 and F2) showed the highest similarity (48.6%). Whereas Co was significantly different, with similarity to both F1 and F2 lower than 29% (Figure 4). The species in Co that mostly contributed to Co dissimilarity towards the other two samples were: *T. virens, S. apiospermum, T. asperellum, T. grisea* and *T. capillare*. These species could be potentially involved in tannins degradation.

On the other hand, in chromium tannery samples, the species that mostly contributed to F1/F2 dissimilarity towards Co were: *G. candidum, P. boydii, L. corymbifera A. furcatum, T. chromospermum,* P. *simplicissimum, P. dangeardii, F solani, S. brevicaulis.* These species, instead, could be rather involved in the degradation of pollutants present in chromium tannery wastewaters. None of these species have been reported for bioremediation of tannery wastewaters. However, several *Trichoderma* species have been signalled in the literature for biosorption of hexavalent chromium from tannery wastewaters (Shukla and Vankar, 2014). Moreover, other species of Penicillium and Fusarium, isolated from contaminated soil, have been used for bioremediation experiment of tannery wastewater (Sharma and Malaviya, 2016). The role of the fungi isolated in
the present study should be investigated with specific bioremediation experiments, in order to validate this
hypothesis.

333

#### 334 *3.4 Cultivable autochthonous fungi – effect of incubation temperature*

335 The effect of temperature on bacterial composition of activated sludge was recently assessed (Niu et al.,
336 2015). However, no study about the effect of temperature on fungal community in WWTPs has been performed.

337 The incubation temperature significantly affected the fungal load in five cases out of nine (CFU in 338 greater number at 15 °C for Co on DRBC and for F1 on MEAp; CFU in greater number at 25 °C for F2 on 339 MEAp and DRBC and for F1 on DRBC) (Figure 3). The temperature affected the biodiversity too, since the 340 PERANOVA test pointed out significant differences in all cases. However, the intragroup similarity was 60-81% 341 for both the temperatures. This turned in a low intergroup dissimilarity percentage 29-60%, with the exception of 342 F1 cultured on MEAp and DRBC, which achieved respectively 85% and 74% dissimilarity between the two 343 incubation temperatures. This is visible in Figure 4, in which the two incubation temperature of F1 on MEAp 344 and DRBC are clearly distinct (intergroup similarity lower than 40%). Temperature does not seem to determine 345 differences in mycoflora development as medium does, which is graphically evident in Figure 5.

Seasonal variations in bacterial composition in activated sludge treating tannery wastewater have been already observed, in particular during the summer. However, this phenomenon seems to be more related to the different influent composition and load during that period than to the high temperature (Giordano et al., 2016). Actually, starvation period can definitively modify the bacterial composition of the activated sludge (Cabezas et al., 2009). This could be true also for fungi, whose cellular structure (with a wall cell) protects them from environmental stress.

352

#### 353 4 Conclusions

This is one of the first detailed studies on the cultivable fungi present in mixed liquors in chromium and vegetable tannery WWTPs aimed at developing novel bioremediation applications. Strong differences in mycoflora indicated the development of specific fungal population in each treatment tank. This study raised some questions about the risk associated with both potential pathogens / toxigenic species and high ecotoxicity, regardless the origin or the stage of depuration. In such kind of samples, the identification of sample-specific species with lower growth rate is strongly linked to the use of fungistatic. Contrarily, incubation temperature, in 360 the range of seasonal fluctuations, does not substantially contribute to the development of different fungal 361 species.

In further studies, the use of culture-independent methods based on next-gen sequencing (e.g. 454,
SOLiD, and illumina, etc.) is needed, in order to achieve the complete identification of the fungi present in these
kinds of samples, which are considerably rich in microorganisms.

- From the applicative point of view, instead, a selective medium, consisting of agarised wastewater as principal source of carbon, allowed to select fungal strains that can be potentially exploited for bioaugmentation or biostimulation, with different suitability for vegetable or chromium tannery wastewaters. However, several of these species are potential pathogens and, thus, their application for *in situ* treatment must be evaluated with caution.
- 370

#### 371 Acknowledgements

This work was supported by University of Turin, Fondazione CRT and by Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR) with the FIRB project RBFR13V3CH\_002 with the title "*In situ* bioaugmentation to exploit the combination of fungi and bacteria for recalcitrant compounds removal".

- The Authors thanks Cuoiodepur Spa (San Miniato, Pisa, Italy) and Consorzio Aquarno SpA (Santa
  Croce sull'Arno, Pisa, Italy) which provided the samples.
- 377

#### 378 References

1. Alvarenga, P., Palma, P., Gonçalves, A.P., Fernandes, R.M., Cunha-Queda, A.C., Duarte, E., Vallini,

380 G., 2007. Evaluation of chemical and ecotoxicological characteristics of biodegradable organic residues
381 for application to agricultural land. Environ. Int. 33, 505–513.

- Anastasi, A., Spina, F., Romagnolo, A., Tigini, V., Prigione, V., Varese, G.C., 2012. Integrated fungal
   biomass and activated sludge treatment for textile wastewaters bioremediation. Bioresour.
   Technol. 123, 106-111.
- 385
  3. Awad, M.F., Kraume, M., 2011. Mycological survey of activated sludge in MBRs. Mycoses. 54, e493–
  386 e498.
- 387
  4. Ayed, L., Asses, N., Chammem, N., Hamdi, M., 2016. Improvement of green table olive processing
  388
  388 wastewater decolorization by *Geotrichum candidum*. Desalin. Water Treat. 57, 17322–17332.

- 389 5. Bedoui, A., Tigini, V., Ghedira, K., Varese, G.C., Ghedira, L.C., 2015. Evaluation of an eventual
  acotoxicity induced by textile effluents using a battery of biotests. Environ. Sci. Pollut. Res. 22, 16700–
  16708.
- Bensch, K., Braun, U., Groenewald, J.Z., Crous, P.W., 2012. The genus *Cladosporium*. Stud. Mycol.
   72, 1–401.
- 394 7. Cabezas, A., Draper, P., Etchebehere, C., 2009. Fluctuation of microbial activities after influent load
  395 variations in a full-scale SBR: recovery of the biomass after starvation. Appl. Microbiol.
  396 Biotechnol. 84, 1191-1202.
- Castellanos-Arevalo, A.P., Camarena-Pozos, D.A., Castellanos-Arevalo, D.C., Rangel-Cordova, A.A.,
   Pena-Cabriales, J.J., Arevalo-Rivas, B., de Pena, D.G., Maldonado-Vega, M., 2016.
   Microbial contamination in the indoor environment of tanneries in Leon, Mexico. Indoor Built
   Environ. 25, 524–540.
- 401
  9. Chapman, P.M., 2000. Whole effluent toxicity testing-usefulness, level of protection, and risk
  402 assessment. Environ. Toxicol. Chem. 1, 3–13.
- 403 10. Claussen, M., Schmidt, S., 1998. Biodegradation of phenol and p-cresol by the hyphomycete
   404 *Scedosporium apiospermum*. Res. Microbiol. 149, 399–406.
- 405 11. Cudowski, A., Pietryczuk, A., Hauschild, T., 2015. Aquatic fungi in relation to the physical and
  406 chemical parameters of water quality in the Augustów Canal. Fungal Ecol. 13, 193–204.
- 407 12. Djelal, H., Amrane, A., 2013. Biodegradation by bioaugmentation of dairy wastewater by fungal
  408 consortium on a bioreactor lab-scale and on a pilot-scale. J. Environ. Sci. 25, 1906–1912.
- 409 13. Domsch, K.H., Gams, W., Anderson, T.H., 1980. Compendium of soil fungi, first ed. Academic Press,
  410 New York.
- 411 14. Evans, T.N., Seviour, R.J. 2012. Estimating biodiversity of fungi in activated sludge communities using
  412 culture-independent methods. Microb. Ecol. 63, 773–786.
- 413 15. Fell, J.W., Boekhout, T., Fonseca, A., Scorzetti, G., Statzell-Tallman, A., 2000. Biodiversity and
  414 systematics of basidiomycetous yeasts as determined by large-subunit rDNA D1/D2 domain sequence
  415 analysis. Int. J. Syst. Evol. Microbiol. 50, 1351-1371.
- 416 16. Giordano, C., Boscaro, V., Munz, G., Mori, G., Vannini, C., 2016. Summer holidays as break-point in
  417 shaping a tannery sludge microbial community around a stable core microbiota. Sci. Rep. 6, 30376.

- 418 17. Glass, N.L., Donaldson, G.C., 1995. Development of prime sets designed for use with the PCR to
  419 amplify conserved genes from filamentous Ascomycetes. Appl. Environ. Microbiol. 61, 1323–1330.
- 420 18. Grabinskaloniewska, A., Slavikova, E., Furmanska, M., Slomczynski, T., 1993. Fungi in activated421 sludge biocenosis. Acta Microbiol. Pol. 42, 303–313.
- 422 19. Gutarowska, B., Skóra, J., Stępień, Ł., Twarużek, M., Błajet-Kosicka, A., Otlewska, A., Grajewski, J.,
  423 2014. Estimation of fungal contamination and mycotoxin production at workplaces in composting
  424 plants, tanneries, archives and libraries. World Mycotoxin J. 7, 345–355.
- 425 20. Herrero, M., Stuckey, D.C., 2015. Bioaugmentation and its application in wastewater treatment: A
  426 review. Chemosphere 140, 119–28.
- 427 21. Ishii, K., Furuichi, T., Tanikawa, N., Kuboshima, M., 2009. Estimation of the biodegradation rate of
  428 2,3,7,8-tetrachlorodibenzo-p-dioxin by using dioxin-degrading fungus, *Pseudallescheria boydii*. J.
  429 Hazard. Mater. 162, 328–332.
- 430 22. Kiffer, E., Morelet, M., 1997. Les deutéromycètes. Classification et clés d'identification générique, first
  431 ed. INRA, Paris.
- 432 23. Kim, I.S., Ekpeghere, K., Ha, S.Y., Kim, S.H., Kim, B.S., 2013. An eco-friendly treatment of tannery
  433 wastewater using bioaugmentation with a novel microbial consortium. J. Environ. Sci. Health Part A.
  434 48, 1732–1739.
- 435 24. Korzeniewska, E., 2011. Emission of bacteria and fungi in the air from wastewater treatment plants a
  436 review. Front. Biosci. 3, 393–407.
- 437 25. Leyva-Diaz, J., Martin-Pascual, J., Poyatos, J.M., 2017. Moving bed biofilm reactor to treat wastewater.
  438 Int. J. Environ. Sci. Technol. 14, 881–910.
- 439 26. Liu, J., Li, J., Tao, Y., Sellamuthu, B., Walsh, R., 2017. Analysis of bacterial, fungal and archaeal
  440 populations from a municipal wastewater treatment plant developing an innovative aerobic granular
  441 sludge process. World J. Microbiol. Biotechnol. 33:14.
- 442 27. Lofrano, G., Meric, S., Zengin, G.E., Orhon, D., 2013. Chemical and biological treatment technologies
  443 for leather tannery chemicals and wastewaters: A review. Sci. Total Environ. 461: 265–281.
- 444 28. Mabrouk, A.M., Kheiralla, Z.H., Hamed, E.R., Youssry, A.A., Abd El Aty, A.A., 2012.
  445 Characterization and purification of the crude *Trematosphaeria mangrovei* laccase enzyme. Egypt.
  446 Pharmaceut. J. 11, 93–98.

- 447 29. More, T.T., Yan, S., Tyagi, R.D., Surampalli, R.Y., 2010. Potential use of filamentous fungi for
  448 wastewater sludge treatment. Bioresour. Technol. 101, 7691–7700.
- 30. Munz, G., De Angelis, D., Gori, R., Mori, G., Casarci, M., Lubello, C., 2009. The role of tannins in
  conventional and membrane treatment of tannery wastewater. J Hazard. Mater. 164, 733–739.
- 451 31. Novotný, C., Dias, N., Kapanen, A., Malachová, K., Vándrovcová, M., Itävaara, M., Lima, N., 2006.
  452 Comparative use of bacterial, algal and protozoan tests to study toxicity of azo- and anthraquinone
  453 dyes. Chemosphere 63, 1436–1442.
- 454 32. Niu, L., Li, Y., Wang, P., Zhang, W., Wang, C., Wang, Q., 2015. Understanding the linkage between
  455 elevation and the activated-sludge bacterial community along a 3,600-meter elevation gradient in
  456 China. Appl. Environ. Microbiol. 81, 6567–76.
- 457 33. Orhon, D., 2015. Evolution of the activated sludge process: the first 50 years. J. Chem. Technol.
  458 Biotechnol. 90, 608–640.
- 459 34. Orlita, A., 2004. Microbial biodeterioration of leather and its control. Int. Biodeterior. Biodegrad. 53,
  460 157–163.
- 461 35. Prigione, V., Zerlottin, M., Refosco, D., Tigini, V., Anastasi, A., Varese, G.C., 2009. Chromium
  462 removal from a real tanning effluent by autochthonous and allochthonous fungi. Bioresour. Technol.
  463 100, 2770–2776.
- 36. Rougeron, A., Schuliar, G., Leto, J., Sitterle, E., Landry, D., Bougnoux, M-E., Kobi, A., Bouchara, J-P.,
  Giraud, S., 2015. Human-impacted areas of France are environmental reservoirs of the *Pseudallescheria boydii/ Scedosporium apiospermum* species complex. Environmental
  Microbiology; 17: 1039–1048.
- 37. Santos, V.L., Linardi, V.R., 2004. Biodegradation of phenol by a filamentous fungus isolated from
  industrial effluents-identification and degradation potential. Process biochem. 39, 1001–1006.
- 470 38. Sharma, S., Malaviya, P., 2016. Bioremediation of tannery wastewater by chromium resistant novel
  471 fungal consortium. Ecol. Eng. 91, 419-425.
- 472 39. Shukla, D., Vankar, P.S., 2014. Role of Trichoderma Species in Bioremediation Process: Biosorption
  473 Studies on Hexavalent Chromium, in: Gupta, V.K., Schmoll, M., HerreraEstrella, A., Upadhyay, R.S.,
- 474 Druzhinina, I., Tuohy, M.G. (Eds), Biotechnology and biology of trichoderma, Elsevier Science BV,
- 475 The Netherlands, pp: 405-412.

- 476
  40. Skóra, J., Gutarowska, B., Stępień, L., Otlewska, A., Pielech-Przybylska K., 2014. The evaluation of
  477
  477
  478
  479
  479
  479
  470
  470
  470
  471
  471
  471
  471
  472
  473
  473
  474
  474
  474
  475
  475
  475
  476
  477
  477
  477
  478
  477
  478
  479
  479
  479
  470
  470
  470
  470
  471
  471
  471
  472
  473
  473
  474
  474
  474
  475
  475
  476
  477
  477
  477
  478
  478
  478
  478
  479
  479
  479
  470
  470
  470
  470
  471
  471
  471
  471
  472
  472
  473
  474
  474
  474
  475
  474
  475
  475
  476
  477
  477
  478
  478
  478
  479
  479
  479
  470
  470
  470
  471
  471
  471
  471
  472
  472
  473
  474
  474
  474
  475
  474
  475
  475
  475
  476
  476
  476
  477
  476
  477
  477
  477
  478
  478
  478
  478
  478
  478
  478
  478
  478
  478
  478
  478
  478
  479
  479
  479
  479
  479
  470
  470
  470
  471
  471
  471
  471
  471
  471
  471
  471
  471
  471
  471
  471
  471
  471
  471
  <li
- 478 41. Svobodová, K., Petráčková, D., Kozická, B., Halada, P., Novotný, C., 2016. Mutual interactions
  479 of *Pleurotus ostreatus* with bacteria of activated sludge in solid-bed bioreactors. World J. Microbiol.
  480 Biotechnol. 32:94.
- 481 42. Thanh, N.V., Nout, M.J.R., 2004. Dormancy, activation and viability of *Rhizopus oligosporus*482 sporangiospores. Int. J. Food Microbiol. 92, 171–179.
- 483 43. Tigini, V., Franchino, M., Bona, F., Varese, G.C., 2016. Is digestate safe? A study on its ecotoxicity
  484 and environmental risk on a pig manure. Sci. Total Environ. 551, 127–132.
- 485 44. Tigini, V., Prigione, V., Varese, G.C., 2014. Mycological and ecotoxicological characterisation of
  486 landfill leachate before and after traditional treatments. Sci. Total Environ. 487, 335–341.
- 487 45. von Arx, J.A., 1981. The genera of fungi sporulating in pure culture, second ed. Port Jervis Lubrecht &
  488 Cramer Ltd, New York.
- 46. Verma, A., Singh, H., Anwar, S., Chattopadhyay, A., Tiwari, K.K., Kaur, S., Dhilon, G.S., 2017.
  Microbial keratinases: industrial enzymes with waste management potential. Crit. Rev. Biotechnol.
  37, 476–491.
- 492 47. White, T.J., Brun, T., Lee, S., Taylor, J.W., 1990. Amplification and direct sequencing of fungal
  493 ribosomal RNA genes for phylogenesis, in: Innis, M.A., Gelfand D.H., Sninsky, J.J., White, T.J. (Eds.)
- 494 PCR Protocols: A Guide to Methods and Applications, Academic Press Inc., New York, pp. 315–322.
- 495 48. Zhang, S., Gao, X., He, L., Qiu, Y., Zhu, H., Cao, Y, 2015. Novel trends for use of microbial tannases.
  496 Prep. Biochem. Biotech. 45, 221–232.



![](_page_19_Figure_1.jpeg)

![](_page_19_Figure_2.jpeg)

![](_page_19_Figure_4.jpeg)

502 Figure 2. Phytotoxicity caused by the three tannery effluents (Co, F1, and F2).

![](_page_20_Figure_0.jpeg)

505 Figure 3. Total fungal load of the three samples (Co, F1, and F2) inoculated on three different media 506 (MEAp, DRBC, and WA) and incubated at two temperatures (25 °C and 15 °C). Capital letters indicate 507 significant differences among fungal loads observed on different culture media at 25 °C incubation; small 508 letters indicate significant differences among fungal loads observed on different culture media at 15 °C 509 incubation; numbers indicate significant differences between incubation temperature; symbols indicates 510 significant differences among sample origins (Mann Whitney, P≤0.05).

![](_page_21_Figure_0.jpeg)

512 Figure 4. Graphical representation of similarity among the sample on the three culture media (MEAP,
513 DRBC and WA for the three samples (Co, F1 and F2) incubated at 25 °C.

![](_page_22_Figure_0.jpeg)

Figure 5. Graphical representation of similarity among all the trials (temperatures and culture media) for

- the three samples (Co, F1 and F2).
- Table 1. Samples chemical features.

520		Co	F1	F2
521	рН	7.66	7.33	7.22
522	Ammonium [mg L <sup>-1</sup> ]	1.04	278	9.65
523	Total phosphorus [mg L <sup>-1</sup> ]	5.92	2.43	3.46
524	Chlorides [mg L <sup>-1</sup> ]	2906	4100	4218
525	COD [mg L <sup>-1</sup> ]	14280	24740	10000
526	SCOD [mg L <sup>-1</sup> ]	362	760	518
527	TSS [g L <sup>-1</sup> ]	12.93	12.61	7.49
528 529	VSS [g L <sup>-1</sup> ]	10.78	9.61	5.77

Table 2: List of the most abundant species retrieved. For each sample (Co, F1, F2), medium (MEAp, DRBC, WA), and incubation temperature (25, 15 °C), contribution in percentage to the intragroup (each column) similarity and total number of taxa are reported.

	Sample			C	Co					F	71			F2						
	Medium	Mł	EAp	DR	BC	W	'A	MF	САр	DR	BC	W	'A	MF	ЕАр	DR	BC	W	'Α	
	Incubation temperature	25	15	25	15	25	15	25	15	25	15	25	15	25	15	25	15	25	15	
	Acremonium furcatum															27				
	Aspergillus niveus																19			
arity	Candida tropicalis var. tropicalis								35											
nil	Cladosporium sp.											12								
si)	Fusarium solani				9															
dno	Geotrichum candidum	48	41					67						63	48	24	38	34	34	
gr	Lichtheimia corymbifera									30										
ntre	Penicillium simplicissimum	21	21	14					37	28	11			16		13				
0 ir	Pseudallescheria boydii						35			20	12	36	22					47	42	
() t	Scedosporium apiospermum				22	60	40				42	12	19							
8	Scopulariopsis brevicaulis											12	13							
ion	Trematosphaeria grisea					21														
out	Trichoderma asperellum			29	33															
tril	Trichoderma capillare	17	13					12												
Con	Trichoderma chromospermum			12	8						15			17		12	18			
	Trichoderma virens			26											20					
	Undetermined												43							
	Total n° of taxa	12	18	21	19	16	20	13	21	23	17	19	21	10	9	25	17	11	20	

- 535 536
  - 36 Table A: List of the taxa retrieved. For each sample (Co, F1, F2), medium (MEAp, DRBC, WA), and
- 537 incubation temperature (25, 15 °C), the abundance of single taxa and average of total load (both expressed

538 as CFU/mL), and total number of taxa are reported.

				(	Co			F1							F2						
	Taxa	M	MEAp DRBC			W	A	MF	ЕАр	DRBC		W	/A	MF	ЕАр	DR	BC	W	'A		
		25	15	25	15	25	15	25	15	25	15	25	15	25	15	25	15	25	15		
1	Absidia glauca Hagem								0.1												
2	Acremonium berkeleyanum (P. Karst.) W. Gams						0.1														
3	Acremonium furcatum Moreau & V. Moreau ex W. Gams	0.1														33. 9					
4	Acrostalagmus luteoalbus (Link) Zare, W. Gams & Schroers											0.1									
5	Ascodesmis macrospora W. Obrist												0.1								
6	Aspergillus caespitosus Raper & Thom													0.3	0.4	0.2					
7	Aspergillus candidus Link							0.1	0.1	0.1	0.6	0.1									
8	Aspergillus					0.1															
9	Aspergillus		0.3	0.2	0.1	0.1				0.1	0.1										
10	Aspergillus flavus Link									0.2	0.2			0.2							
11	Aspergillus fumigatus Fresen.		0.1	0.7	0.1							0.1				0.2					
12	Aspergillus neoniveus Samson	1.4	0.7																		
13	Aspergillus niger Tiegh.										0.1										
14	Aspergillus niveus Blochwitz								0.1							0.2	11. 7				
15	Aspergillus ochraceus K. Wilh.			2.4	1.5			0.2	0.1	1.3	0.3	0.6	0.3		0.1	6	0.2	0.4			
16	Aspergillus ostianus Wehmer																0.1				
17	Aspergillus sojae Sakag. & K.									0.1											
18	Aspergillus terreus Thom	0.1	3.4	0.1	0.2																
19	Aspergillus versicolor (Vuill.) Tirab.					0.1															
20	Aureobasidium sp.						0.1														
21	Basifimbria sp.					0.5										0.1					
22	<i>Bjerkandera</i> <i>adusta</i> (Willd.) P. Karst.				0.1																

23	Botryotrichum									0.1									
	piluliferum																		
	Sacc. &																		
24	Marchal						0.0												
24	Byssochlamys						0.2												
	(C Ram)																		
	Samson,																		
	Houbraken &																		
	Frisvad																		
25	Byssochlamys nivea Westling															0.1			
26	Candida	1.6	0.1						14.									0.2	
	tropicalis var. tropicalis								2										
27	Chaetomium									0.1									
	bostrychodes																		
	Zopf																		
28	Chaetomium									0.4									0.2
20	<i>elatum</i> Kunze					0.1	0.2		0.2		0.7	0.2	0.5						0.1
29	glohosum					0.1	0.5		0.2		0.7	0.5	0.5						0.1
	Kunze ex Fr.																		
30	Chaetomium		0.1				0.2					4.1	3.6					0.2	4.5
	homopilatum																		
24	Omvik											0.2	0.5						
31	chaetomium											0.2	0.5						
	Booth &																		
	Shipton) X.W.																		
	Wang & Crous																		
32	Chaetomium sp.					0.4													
33	Chaetomium						0.3												
	tarraconense Stabigal																		
34	Cladosporium						0.5		04		_	_			_				_
34	cladosporioides						0.5		0.4										
	(Fresen.) G.A.																		
	de Vries																		
35	Cladosporium								1.1										
	pseudocladospo																		
36	Cladosporium				38	0.1						39					46	35	2.2
	sp.				5.0	0.1						5.9						5.0	2.2
37	Cladosporium												3.3						
	xylophilum Bens																		
	ch, Shabunin,																		
	Braun																		
38	Clonostachys			0.2															
	<i>solani</i> f.																		
	nigrovirens																		
	(J.F.H. Beyma) Schroers																		
39	Coelomvcetes								0.3				0.1						
	sp.																		
40	Cryptococcus					0.2													
	humicola																		
<u></u>	(Dasz.) Golubev												0.1						0.1
41	terricola L F H												0.1						0.1
	Beyma,																		
42	Eurotium sp.												0.2						
43	Fusarium								7.2										
	keratoplasticum																		
	D. Geiser,																		
	Short et Zhang																		
44	Fusarium solani			0.1	4.2	0.2		1.9	0.2	0.1	0.2	1.3	3.3	0.1	0.2	0.8	2.8		1.8
	(Mart.) Sacc.			<i></i>	2					5.1		1.5	5.5	5.1	<u> </u>	5.5			1.0
45	Geotrichum	24.	22.					38.	0.3	0.3	1.4		0.3	50	26.	28	53.	19.	14.
	candidum Link	2	2					3							4		1	2	6
46	Gliocladium	1.4	0.1																

	solani (Harting)																
47	Petch							0.2									
47	Guomasitx sp. Gymnascella							0.2			0.1						
-0	<i>hyalinospora</i> (Kuehn, G.F. Orr & G.R.										0.1						
	Ghosh) Currah										0.6	0.1					
49	<i>Gymnaschella</i> sp.										0.6	0.1					
50	Gymnoascus udagawae Arx					0.5											
51	Humicola fuscoatra Traaen			0.2		0.7											
52	<i>Humicola</i> grisea Traaen											0.1					
53	Isaria fumosorosea Wize									0.1							
54	Lichtheimia corymbifera (Cohn) Vuill.		0.1	0.1					32. 4		1.5	0.2		0.6		0.1	0.6
55	Mammaria echinobotryoide s Ces.											0.1					0.1
56	Monascus ruber Tiegh.			0.1	0.1	0.2	0.2				0.1	0.1	0.1				
57	Mucor circinelloides var. circinelloides		0.1														
58	<i>Mucor luteus</i> Linnem. ex Wrzosek						0.6										
59	Mucor plumbeus Bonord.	1.5	3.0		0.3												
60	Neosartorya fischeri (Wehmer) Malloch & Cain	0.1															
61	Paecilomyces carneus (Duché & R. Heim) A.H.S. Br. & G. Sm.								0.4								
62	Paecilomyces variotii Bainier								0.4					5.2			0.2
63	Penicillium atramentosum Thom		0.1	0.6				0.5							0.6		
64	Penicillium camemberti Thom												0.1				
65	Penicillium crustosum Thom		0.3														
66	Penicillium dangeardii Pitt													5.4			
67	Penicillium expansum Link					0.1											
68	Penicillium funiculosum Thom			0.1													
69	Penicillium								0.3					0.1			
	<i>mexicanum</i> Visagie, Seifert & Samson																
70	Penicillium							0,7									
71	paneum Frisvad Penicillium rugulosum				0.5								0.3	0.2	1.2		
	ruguiosum																

	Thom																		
72	Penicillium	5.1	5.4	3.8	2.5			2.3	26.	25.	4.6			3.2	3.5	16.	1.8		
	simplicissimum								4	6						1			
70	(Oudem.) Thom			0.0	0.1			0.1	0.7	0.1	0.4		0.0			0.1	0.2		
73	Penicillum sp.			0.8	0.1			0.1	0.7	0.1	0.4	0.1	0.0			0.1	0.2		
/4	Petrietta guttulata G L											0.1							
	Barron & Cain																		
75	Phoma sorghina							0.1											
	(Sacc.)																		
	Boerema,																		
	Kesteren																		
76	Phoma sp.					0.4													
77	Pseudallescheri				1.1	3.1	7.6			15.	3.9	32.	17.					32.	25
	a boydii (Shear)									4		8	5					6	
	McGinnis, A.A.																		
	Aiello																		
78	Rhodotorula sp.		1.6																
79	Sagenomella	0.7	0.3																
	diversispora																		
	(J.F.H. Beyma)																		
80	W. Gams							0.1											
81	Sarocladium							0.1										0.2	
01	strictum (W.																	0.2	
	Gams)																		
	Summerbell				10	15	0.7				20		11		<u>a</u> :	0.1	0.7	4.0	0.7
82	Sceaosporium				12	17. Q	9.5				39. 3	5.7	11. - A		2.4	0.4	0.7	4.2	0.7
	(Sacc.) Sacc. ex					ĺ ĺ					5		- T						
	Castell. &																		
	Chalm.																		
83	Scedosporium					1.0	0.2										0.1	1.6	0.1
	Gilgado																		
84	Scolecobasidio					0.1	0.7												
	<i>m</i> sp.																		
85	Scopulariopsis									0.1		4.4	5		0.1	0.4	0.5	0.7	1.3
	(Sacc.) Bainier																		
86	Sporormiella															0.1	0.1		0.1
	minima																		
	(Auersw.) S.I.																		
07	Ahmed & Cain																		0.2
07	chartarum																		0.2
	(Ehrenb.) S.																		
	Hughes			0 -															
88	Talaromyces			0.7															
	Stolk & Samson																		
89	Talaromyces sp.															0.1			
90	Talaromyces															0.7			
	wortmannii																		
	(Klocker) C.R. Beni																		
91	Thelebolus sp.						0.1												0.2
92	Trematosphaeri					4.4	4.3												0.5
	a grisea (J.E.																		
	Mackinnon,																		
	rerrada & Montemartini)																		
	S.A. Ahmed.																		
	W.W.J. van de																		
	Sande, A. Fahal																		
02	& de Hoog									0.1									
93	asperelloides									0.1									
	Samuels																		
94	Trichoderma			17.	23.					1.8	0.2					0.5	0.1		
1	asperellum			3	6														

	Samuels, Lieckf. & Nirenberg																		
95	<i>Trichoderma</i> <i>capillare</i> Samuels & C.P. Kubicek	3.9	1.7	0.1				1.3	0.2	9.0									
96	Trichoderma chromospermu m P. Chaverri & Samuels			4.0	3.6				3.0		6.8			4.3	1.8	9.8	8.7		
97	<i>Trichoderma</i> harzianum Rifai			0.7												0.2			
98	Trichoderma virens (J.H. Mill., Giddens & A.A. Foster) Arx			13. 8	4.1			0.1	1.2	0.8	1.6	0.1		4.7	6.8	0.8	1.1		
99	<i>Trichosporon</i> sp.			0.1	0.4														
10 0	Undetermined	0.4	4.1	2.6	2.6	0.3	0.6	1.4	0.2	2	10. 2	7.9	18. 7						0.1
10 1	<i>Westerdykella dispersa</i> (Clum) Cejp & Milko						0.1												
10 2	<i>Westerdykella</i> sp.																		0.1
	Total n° of taxa	12	18	21	19	17	20	13	21	23	17	19	21	10	9	25	17	11	20
	Average of total CFU/mL	40. 5	43. 7	<b>48.</b> 7	60. 9	29	26. 3	46. 7	57. 4	91. 2	70. 7	64	66. 1	63. 3	41. 7	110 .2	87. 6	62. 9	52. 7

540 Species common to all samples are underlined in blue; species exclusively present in Co are

541 underlined in red; species exclusively present in F1 are underlined in purple; species

542 exclusively present in F2 are underlined in green.

543