

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Estimation of the baseline level of aneuploidy in germ and reproductive health in domestic animals

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1507157> since 2018-03-18T16:20:06Z

Published version:

DOI:10.1007/s10577-010-9145-8

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

D. Di Bernardino, D. Nicodemo, A. Pauciullo, G. Cosenza, L. Ramunno, V. Peretti, L. Iannuzzi, J. Rubes. Estimation of the baseline level of aneuploidy in germ cells and reproductive health in domestic animals. *Chromosome Research* (2010) 18:729–729

DOI 10.1007/s10577-010-9145-8

The publisher's version is available at:

<https://link.springer.com/article/10.1007%2Fs10577-010-9145-8>

When citing, please refer to the published version.

Link to this full text:

<https://link.springer.com/article/10.1007%2Fs10577-010-9145-8>

This full text was downloaded from iris-Aperto: <https://iris.unito.it/>

Estimation of the baseline level of aneuploidy in germ cells and reproductive health in domestic animals

D. Di Berardino¹, D. Nicodemo¹, A. Pauciullo¹, G. Cosenza¹, L. Ramunno¹, V. Peretti², L. Iannuzzi³, J. Rubes⁴

¹Department of Soil, Plant, Environment and Animal Production Sciences, University of Naples “Federico II”, Portici, Italy

²Department of Animal Science and Food Inspection, University of Naples “Federico II”, Naples, Italy

³Research National Council (CNR), ISPAAM, Laboratory of Animal Cytogenetics and Gene Mapping, Naples, Italy

⁴Veterinary Research Institute, Brno, Czech Republic

The reproductive health of a given species is strictly related to the karyological integrity of the breeding animals (i.e., absence of chromosomal abnormalities) and to the precision with which the haploid chromosomes are segregated into the male (sperm) and female (oocyte) germ cells prior to fertilization. This assumption is supported by the fact that in mammals more than 70% of the embryonic mortality is attributed to aneuploidies in the germ cells which give rise to genetically unbalanced embryos (monosomic or trisomic) destined to abortion, thus reducing fertility and the reproductive/productive efficiency of the animal production industry.

Estimation of the ‘baseline’ level of aneuploidy in sperm and in in vitro matured oocytes with the corresponding first polar body in domestic animal species and breeds/genetic types is now possible by using the fluorescent in situ hybridization (FISH) technique and chromosome specific painting probes obtained via microdissection and DOP-PCR. Such a ‘baseline’ level could be used as ‘control reference’ not only for improving the in vitro production of embryos destined to the animal production industry but also, and more importantly, for monitoring future trends of the reproductive health of the domestic species/breeds engaged in zootechnical productions, especially in relation to the increasing risk of environmental challenges and hazards, such as chemical contaminants in the water, soil, air, micotoxins in the feedstuff, hormone unbalancements, nutritional and dietetical mistakes, and other factors which are known to damage the mitotic and meiotic machinery of the cells.