

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Inclusion of bilberry pomace in rabbit diets: Effects on carcass characteristics and meat quality

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1616946> since 2017-07-03T11:03:45Z

Published version:

DOI:10.1016/j.meatsci.2016.10.013

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Inclusion of bilberry pomace in rabbit diets: effects on carcass characteristics and meat quality

Sihem Dabbou^{a,b}, Francesco Gai^c, Manuela Renna^{a*}, Luca Rotolo^a, Samia Dabbou^b, Carola Lussiana^a, Attawit Kovitvadhi^{a,d}, Alberto Brugiapaglia^a, Michele De Marco^e, Ahmed Nouredine Helal^b, Ivo Zoccarato^a, Laura Gasco^{a,c}

^aDepartment of Agricultural, Forest and Food Sciences, University of Turin, largo Paolo Braccini 2, 10095 Grugliasco, Turin, Italy

^bLaboratory of Bioresources, Integrative Biology and Valorisation, Higher Institute of Biotechnology of Monastir, av. Tahar Hadded, BP 74, 5000, Monastir, Tunisia

^cInstitute of Science of Food Production, National Research Council, largo Paolo Braccini 2, 10095 Grugliasco, Turin, Italy

^dDepartment of Physiology, Faculty of Veterinary Medicine, Kasetsart University, 50 Ngam Wong Wan Road, Chatuchak, Bangkok 10900, Thailand

^eDepartment of Veterinary Sciences, University of Turin, largo Paolo Braccini 2, 10095 Grugliasco, Turin, Italy

*Corresponding author: Tel +39 011 6708582, Fax +39 011 6708563; E-mail:

manuela.renna@unito.it

E-mail addresses: sihem.dabbou@yahoo.fr; francesco.gai@ispa.cnr.it;

manuela.renna@unito.it; luca.rotolo@unito.it; samia_1509@yahoo.fr;

carola.lussiana@unito.it; attawitthai@hotmail.com; alberto.brugiapaglia@unito.it;

michele.demarco@unito.it; ahelal584@gmail.com; ivo.zoccarato@unito.it;

laura.gasco@unito.it

Abstract

A trial was carried out to evaluate bilberry pomace (BP) as an alternative source of nutrients for rabbits. One hundred and forty-four Grimaud weaned rabbits were divided into 4 groups of 36 animals each and fed *ad libitum* with a basal diet (BP0) tested against three assay diets developed by substituting 50, 100 and 150 g / kg of the BP0 diet with BP (BP5, BP10 and BP15 diets, respectively). Carcass characteristics of rabbits were not affected by treatment, with the exception of liver weight. Dietary inclusion of BP did not significantly affect the proximate composition and the oxidative status of *longissimus thoracis et lumborum* muscle. However, dietary BP significantly ameliorated, in the same muscle, the nutritional quality and health properties of fat. Consumer acceptance of cooked rabbit meat was not significantly affected by treatment. In conclusion, BP can be included in rabbit diets to improve the lipid composition of meat, without adverse effects on carcass characteristics, other physico-chemical properties, oxidative status and meat sensorial traits.

Keywords: rabbit, *Vaccinium myrtillus*, by-product, fatty acids, TBARS, sensory analysis

1. Introduction

Agricultural wastes obtained by processing fruits and vegetables are inexpensive, easy to store and then available for an acceptable length of time during the year. They may contain bioactive substances, such as antioxidants, characterized by health promoting properties and potential technological applications (Lee & Wrolstad, 2004). These wastes could be

consequently explored as useful feed sources for the production of animal derived food products with functional nutritional value.

In rabbits, some studies have been carried out to evaluate performance, meat quality or health status of animals fed different antioxidants derived from olive pomace (Dal Bosco et al., 2012), artichoke bracts (Dabbou et al., 2014), grape pomace (Eid, 2008), chestnut hydrolysable tannins (Liu et al., 2009), microalgae (Peiretti & Meineri, 2011) and green tea (Eid, Zeweil, Ahmed, Basyony, & Farok, 2011).

Bilberry (*Vaccinium myrtillus* L.) is one of the most important sources of phenolic compounds in the human diet. Phenolic compounds, being characterized by various biological activities, may confer bilberries the capability of improving human health conditions (Lee & Wrolstad, 2004; Khanal, Howard, Brownmiller, & Prior, 2009). Juice processing of bilberries generates large amounts of pomace, a by-product which still contains an assortment of beneficial phytochemicals including proanthocyanidins, anthocyanins and other flavonoids, suitable for the development of novel functional food ingredients (Vulić et al., 2011). Bilberry pomace (BP) has already been used as ingredient in extruded products which have been associated with *in vivo* health benefits in animal models, such as reduced plasma cholesterol and abdominal fat (Khanal et al., 2009; Khanal, Howard, Wilkes, Rogers, & Prior, 2012). Kim, Bartley, Rimando, and Yokoyama (2010) suggested that hepatic modulation of bile acid and cholesterol synthesis primarily contributes to the cholesterol-lowering effect of BP.

Based on the current knowledge, no studies have been performed on the use of BP either in rabbit or other livestock animals' diets. Therefore, the aim of the present investigation was to evaluate the effects of BP inclusion in rabbit diets on carcass characteristics, and on physico-chemical characteristics, fatty acid (FA) profile, oxidative stability and sensorial characteristics of rabbit meat.

2. Materials and methods

2.1. Animals and experimental design

The trial was carried out at the experimental rabbitry of the Department of Agricultural, Forest, and Food Sciences (DISAFA; University of Turin), located in Carmagnola, Turin, Italy. The experimental protocol was designed according to the guidelines of the current European and Italian laws on the care and use of experimental animals (European Directive 86 609/EEC, put into law in Italy with D.L. 116/92). The experimental protocol was approved by the Ethical Committee of the Department of Agricultural, Forest, and Food Sciences of the University of Turin (Italy). One hundred forty-four weaned crossbred (Grimaud) rabbits (35 days old) were randomly divided into 4 groups of 36 animals each; the average initial weight was 938 ± 33.4 g. The animals were housed individually in wire cages (41 cm \times 0.30 cm \times 28 cm height) and had free access to clean drinking water. The temperature and photoperiod in the rabbitry were 22 ± 2 °C and 16L:8D, respectively. Rabbits were fed *ad libitum* with a basal diet not containing BP (BP0 diet) (ingredients, g / kg fresh matter: alfalfa meal 300, wheat bran 200, barley 170, dried beet pulp 150, soybean meal 115, molasses 20, wheat straw 20, and soybean oil 5) tested against three assay diets developed by substituting 50, 100 and 150 g / kg of the BP0 diet with BP (BP5, BP10, and BP15 diets, respectively) according to Goby and Gidenne (2008). All diets also contained a vitamin-mineral premix and bicalcium phosphate (15 and 5 g / kg fresh matter, respectively). BP was included in the treated diets during the raw material mixing process. All diets were pelleted fresh, stored in dark bags and kept at environmental temperature in the feed facility to prevent auto-oxidation of the lipid sources.

2.2. Chemical analyses of feed

All **chemical** analyses were carried out on three replicates of each feed sample, according to the recommendations of the European Group on Rabbit Nutrition (2001).

2.2.1. Determination of BP phenolic compounds

BP generated during fruit juice production was obtained from a private fruit processing company ("Arc en Ciel" Soc. Agr. Coop., Cafasse, Turin, Italy). The bilberries had been harvested during July and August 2013 in wild forests with an organic certification. BP was dried in an oven at 60 °C until constant weight was reached and then finely ground.

Total phenols and *o*-diphenols were extracted from about 0.20 g of BP powder using a methanol / water solution (4:1, v:v). The solution was mixed for 30 min at 37 °C under continuous agitation and centrifuged at **3075 g** for 20 min. The methanolic fraction was recovered and stored at -20 °C until analysis. Total phenols and *o*-diphenols were determined colorimetrically at 765 nm and 500 nm respectively and expressed as mg hydroxytyrosol equivalents / g dry matter (DM), as reported by Montedoro, Servili, Baldioli, and Miniati (1992).

The same methanolic extract was used for total flavonoids determination at 510 nm (Zhishen, Mengcheng, & Jianming, 1999). The results are expressed as mg catechin equivalents / g DM. Condensed tannins were determined from 50 µl of the same methanolic extract using the vanillin method (Julkenen-Tiitto, 1985). The results are expressed as mg of catechin equivalents / g DM.

For anthocyanins determination, acidified methanol (1%) was used to prepare extracts. The absorbance was measured at 530 and 657 nm and results are expressed as mg cyanidin 3-glucoside equivalents (CyE) / g DM (Mancinelli, Huang Yang, Lindquist, Anderson, & Rabine, 1975).

All colorimetric determinations were performed using a UV–VIS spectrophotometer (UV–VIS Beckman spectrophotometer DU 650, Beckman Instruments Inc., Fullerton, CA, USA). The concentrations of BP phenolic compounds are reported in Table 1.

2.2.2. Antioxidant activity of BP

To evaluate the antioxidant activity of the BP extract, the free radical scavenging activity was determined by using the 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay according to the method described by Kontogiorgis and Hadjipavlou-Litina (2005). The loss of DPPH color caused by consumption of DPPH radical by antioxidant species present in the sample was measured using a UV–VIS spectrophotometer (UV–VIS Beckman spectrophotometer DU 650, Beckman Instruments Inc., Fullerton, CA, USA). Briefly, a solution of 20 µl of BP sample dissolved in absolute ethanol to a final volume of 1 ml was added to 1 ml of DPPH (0.1 mM, in absolute ethanol) and allowed to stand in the dark for 20 min (reaction time) before the measurement of the absorbance at 517 nm. The inhibition of DPPH radical scavenging activity was calculated as follows:

$$\% \text{ inhibition} = [(\text{absorbance of blank} - \text{absorbance of BP extracts}) / \text{absorbance of blank}] \times 100$$

where the blank is the control solution containing all reagents except the sample.

2.2.3. Proximate composition and fatty acid profile of BP and experimental diets

The BP and the experimental diets were ground with a cutting mill to pass a 1-mm screen sieve (Pulverisette 15 – Fritsch GmbH, Idar-Oberstein, Germany). They were analyzed for DM (# 930.15), ash (# 923.03), crude protein (CP, # 984.13), ether extract (EE, # 2003.05), acid detergent fiber (ADF, # 973.18) and acid detergent lignin (ADL, # 973.18) according to AOAC procedures (AOAC International, 2000; 2003). Neutral detergent fiber (NDF) was

determined according to Van Soest, Robertson, and Lewis (1991). Starch content was determined using the Ewer's polarimetric method (European Economic Community, 1972).

Gross energy (GE) was measured using an adiabatic calorimetric bomb (C7000, IKA, Staufen, Germany).

The FA composition of feedstuffs was assessed using a combined direct *trans*-esterification and solid-phase extraction method as described by Alves, Cabrita, Fonseca, and Bessa (2008). Separation, identification, and quantification of fatty acid methyl esters (FAME) were performed as reported by Renna et al. (2014). The results are expressed as g / 100 g DM and reported as g / 100 g of total detected FA.

The proximate and FA compositions of feeds are reported in Tables 2 and 3, respectively.

2.3. Slaughter procedures and muscle sampling

At 83 days of age, 12 rabbits per group (mean weight 2984 ± 138.0 g) were slaughtered in an experimental slaughterhouse without fasting. The slaughtered rabbits were bled and the skin, genitals, urinary bladder, gastrointestinal tract and distal part of the legs were removed as recommended by Blasco, Ouhayoun, and Masoero (1993). The carcass was weighed; the skin and full gastrointestinal tract weights were recorded and expressed as a percentage of slaughter weight (SW). Carcasses (with head, thoracic cage organs, liver and kidneys) were chilled at 4 °C for 24 h in a refrigerated room. The chilled carcass weight (CCW) was recorded and the dressing out percentage was calculated as the ratio between CCW and SW. Head and liver weight were expressed as a percentage of CCW. The head, thymus, trachea, esophagus, heart, lungs, liver and kidneys weights were removed from the CCW to obtain the reference carcass weight (RCW). For meat quality analyses, *Longissimus thoracis et lumborum* (LTL) muscle was removed from both left and right side. Meat quality parameters were measured on the left loin while sensory analysis was performed on the right loin.

172

173 2.4. Meat quality parameters

174 2.4.1. pH

175 Meat pH of the LTL muscle (at the level of the 7th lumbar vertebra) was measured at 24 h
176 *postmortem* (pH₂₄) in duplicate using a Crison portable pH-meter (Crison Instruments, S.A.,
177 Alella, Spain) fitted with a spear-type electrode and an automatic temperature compensation
178 probe.

179

180 2.4.2. Color

181 Meat color was measured at room temperature (20°C) on a freshly cut surface of the loin at
182 the level of the 7th lumbar vertebra using a portable colorimeter Chroma Meter CR-400
183 Konica Minolta Sensing (Minolta Sensing Inc, Osaka, Japan). Color measurements were
184 reported in terms of lightness (L*), redness (a*) and yellowness (b*) in the CIELAB color
185 space model (Commission Internationale de l'Éclairage, 1976). The values were recorded for
186 CIE standard illuminant D₆₅ and the CIE 2° standard observer. The color values were
187 obtained considering the average of three readings per sample.

188

189 2.4.3. Proximate composition

190 The proximate analyses were carried out according to the AOAC International (2000)
191 methods. Tissue samples were weighed, dried at 125°C for 5 h and reweighed to determine
192 the water content. The samples remaining from water analysis were placed into a furnace
193 oven at 525°C for 6 h for ash determination. The meat was further lyophilized and ground in a
194 blender for analyses of protein and intramuscular fat. Nitrogen was determined by Kjeldahl
195 method and CP was calculated by multiplying N × 6.25. Lipid extraction of intramuscular fat
196 was determined by Soxhlet method.

197

198 2.4.4. *Fatty acid composition*

199 The FA composition of freeze-dried LTL muscle samples was assessed as reported by Belforti
200 et al. (2015). Briefly, total lipids were extracted with dichloromethane / methanol (2:1, v:v) by
201 homogenization at room temperature. The solution was filtered in a separating funnel
202 containing 1 ml of a water solution of MgCl_2 (2%, w/v) and 20 ml of water. The organic
203 phase was separated and evaporated to dryness. Glycerides were saponified with a solution of
204 NaOH 0.5 M in methanol and then BF_3 (about 10% in methanol) were added for methylation.
205 Peaks were identified by injecting pure FAME standards as detailed by Renna et al. (2012).
206 Quantification was assessed using tridecanoic acid ($\text{C}_{13:0}$) as internal standard. The results
207 are expressed as g / 100 g of LTL muscle and reported as g / 100 g of total detected FA.

208

209 2.4.5. *Lipid oxidation*

210 Lipid oxidation was determined on meat samples (10 g) at 30 and 60 days of frozen storage,
211 by thiobarbituric acid reactive substances (TBARS) assay as described by Dabbou et al.
212 (2014). The samples were analyzed in duplicate and the absorbance was read at 532 nm with a
213 Helios spectrophotometer (Unicam Limited, Cambridge, UK). TBARS values were calculated
214 from a standard curve of 1,1,3,3-tetramethoxypropane (TMP; Sigma–Aldrich, Steinheim,
215 Germany) and expressed as mg TMP / kg of meat.

216

217 2.4.6. *Sensory evaluation*

218 A sensory panel of 68 untrained assessors, 29 males and 39 females, ranging in age from 21 to
219 60 years, was recruited among the students and staff members of the University of Turin.
220 Participants were regular consumers of rabbit meat and were already involved in surveys on
221 meat acceptability / preference tests. Affective tests were carried out in 8 distinct evaluation

222 sessions during 4 days and performed in the sensory laboratory of DISAFA under controlled
223 conditions with panelists placed in individual tasting booths. In each day, an acceptance test
224 was performed to assess the oxidative stability of meat. Muscles were kept unpackaged in a
225 dark cooler set at 4 °C for 4 days. After aging, the samples were vacuum-packed and stored at
226 -20 °C until analysis. The loins, from rabbits of the 4 groups, were thawed at 4 °C / 24 h,
227 packages were opened and the samples were simultaneously cooked without salt or spice on a
228 double plate grill, preheated at 250 °C, to a final temperature of 70 °C. Cooking temperature
229 was monitored by an iron/constantan thermocouple placed in the geometric centre of each
230 loin. After grilling, the loins were immediately cut into equal portions. The latter were labeled
231 with three-digit numbers and offered using a Williams design to balance the order of
232 presentation (MacFie, Bratchell, Greenhoff, & Vallis, 1989).

233 Immediately before the sensory tasting sessions, the panelists were asked to sign an informed
234 consent form. Each panelist received 4 warm samples corresponding to the 4 experimental
235 diets.

236 The panelists were asked to measure the degree of liking or disliking of rabbit meat by the use
237 of the 9-point hedonic scale (1 = “dislike extremely”, 2 = “dislike very much”, 3 = “dislike
238 moderately”, 4 = “dislike slightly”, 5 = “neither like nor dislike”, 6 = “like slightly”, 7 = “like
239 moderately”, 8 = “like very much”, 9 = “like extremely”; Peryam & Pilgrim, 1957).

240 Tap water was offered to the panelists to rinse their mouths between samples.

242 2.5. Statistical analyses

243 The statistical analyses were performed using the SPSS software package (version 17 for
244 Windows, SPSS Inc., Chicago, IL, USA).

245 One-way ANOVA was used to evaluate the effect of BP dietary inclusion levels on carcass
246 characteristics and meat quality traits. The assumption of equal variances was assessed by

Levene's homogeneity of variance test. If such an assumption did not hold, the Brown-Forsythe statistic was performed to test for the equality of group means instead of the F one. Pairwise multiple comparisons were performed to test the difference between each pair of means (Duncan test and Tamhane's T2 in the cases of equal variances assumed or not assumed, respectively).

The effects of dietary treatment and storage time on lipid oxidation were statistically analyzed with a mixed between-within subjects model (GLM for Repeated Measures).

Sensory data from the acceptance test were analyzed using the GLM procedure with overall liking as dependent variable, dietary treatment as fixed effect and panelist as random effect (Naes, Brockhoff, & Tomic, 2010). The scores of acceptability obtained for the 4 samples from each consumer were converted into ranked data by assigning rank order numbers to the evaluations. Ties received average rank scores. Ranking data were analyzed with the Friedman's test (Meilgaard, Civille, & Carr, 1991).

For all statistical analyses **significance** was declared at $P < 0.05$.

3. Results and discussion

3.1. Carcass characteristics and meat quality traits

Carcass characteristics are reported in Table 4. **Differences were found in liver weight that was lower in the animals fed BP containing diets; all other parameters were not affected by BP dietary inclusion.** Similar results were obtained by Heyman et al. (2014) who found a significantly reduced liver mass and liver triacylglycerol lipid accumulation in C57BL/6J mice fed a high-fat diet supplemented (20%) with different berries (lingonberry, blackcurrant, and bilberry) that implies protection against liver steatosis.

The proximate composition, pH₂₄ and color of the LTL muscle of rabbits are reported in Table 5. These meat quality traits fell within standard ranges for rabbit meat and were not

affected by treatment. Abdel-Khalek (2013) reviewed that dietary supplementation with antioxidants has no clear trend in the physical and chemical characteristics of rabbit meat.

3.2. Fatty acid profile of diets and LTL muscle

Inclusion of BP increased the total FA content and modified the FA profile of the lipids in the diets (Table 3). Results showed that BP is rich in unsaturated FA (93.38 g / 100 g total FA). The most abundant FA in BP was linoleic acid (C18:2 n6, LA; 36.33 g / 100 g total FA), followed by α -linolenic acid (C18:3 n3, ALA; 32.59 g / 100 g total FA) and oleic acid (C18:1 c9, OA; 23.50 g / 100 g total FA). An increasing percentage of MUFA and PUFA (mainly due to increases in C18:1 c9 and C18:3 n3) at the expense of SFA was observed following increasing inclusion levels of BP in the diets. The values of the Σ n6 / Σ n3 PUFA ratio were 5.13, 3.30, 2.49 and 2.14 in BP0, BP5, BP10 and BP15 diets, respectively.

In rabbits, dietary FA are directly incorporated into intramuscular lipids (Dalle Zotte et al., 2014). Expectedly, the FA profile of rabbit LTL muscle (Table 6) thus generally followed that of the dietary ingested lipids (Dal Bosco, Castellini, Bianchi, & Mugnai, 2004). Increasing PUFA and decreasing SFA proportions from BP0 to BP15 diets resulted in significantly higher (26.08 to 37.46 g / 100 g total FA) and lower (42.93 to 36.48 g / 100 g total FA) proportions of these FA groups in the muscle ($P < 0.001$), respectively. The observed trend for total MUFA was opposite, as increasing levels in the diets led to decreasing proportions in the LTL lipid fraction ($P < 0.001$). The estimated $\Delta 9$ -desaturase activity did not differ among treatments (Table 6). We may therefore exclude the hypothesis of a reduced conversion of SFA into *cis*-9 MUFA due to dietary PUFA inhibition of SCD (Papadomichelakis, Karagiannidou, Anastasopoulos, & Fegeros, 2010a). Most probably, as already reported for α -tocopherol (Dal Bosco et al., 2004), antioxidants in BP may have inhibited the peroxidation

of FA with high (PUFA) rather than low (MUFA) degree of unsaturation, with consequent proportional increasing contents of PUFA at the expense of SFA and MUFA in the muscle. The inclusion of BP in the diets induced significant modifications in the proportion of the majority of individual detected FA in the LTL muscle. The main variations regarded n3 PUFA. ALA levels in rabbit meat depend on the level of exogenous fatty acids (Liu et al., 2009; Kouba, Benatmane, Blochet, & Mourot, 2008) as rabbits effectively synthesize endogenous n3 PUFA from its precursor in the liver; the amount produced depends on the dietary $\Sigma n6 / \Sigma n3$ FA ratio (Peiretti & Meineri, 2008). Therefore, the proportional increasing content of ALA in the BP diets (Table 3) expectedly resulted in significant increasing percentages of ALA in the muscle. In particular, a double content of ALA in BP15 if compared to BP0 diets resulted in 4-times higher percentage of ALA in muscles. Rabbits are able to elongate and desaturate ALA, also due to caecal microflora and caecotrophe reingestion (Dal Bosco et al., 2004). As a consequence, the long-chain n3 PUFA (C20:5 n3 and C22:5 n3) were also significantly increased following BP inclusion in the diet ($P < 0.001$), which is consistent with the results obtained in other nutritional trials using linseed (Dal Bosco et al., 2015; Kouba et al., 2008; Dal Bosco et al., 2004) and fresh forage (Dal Bosco et al., 2014; Forrester-Anderson, McNitt, Way, & Way, 2006).

Probably due to differences in analytical and gas chromatographic conditions as well as lack or difficulties in peak identification, till now very few published trials (Papadomichelakis et al., 2010a; Papadomichelakis Karagiannidou, Anastasopoulos, & Fegeros, 2010b; Leiber et al., 2008) reported information on the odd- (OCFA) and branched-chain fatty acids (BCFA) in rabbit meat. These FA have recently received increasing attention by researchers due to their association with reduced disease risk for coronary heart disease, diabetes and cancer (Jenkins, West, & Koulman, 2015; Oku & Yanagita, 2009). The major part of these FA are produced by bacteria in the caecum and reach the duodenum and the blood via caecotrophs reingestion

321 while, most probably, endogenous synthesis takes place to a lower extent (Leiber et al., 2008).
322 Caecal bacterial response to dietary changes in lipid content and composition has been poorly
323 investigated yet. Papadomichelakis, Anastasopoulos, Karagiannidou, and Fegeros (2010c)
324 reported unchanged concentrations of OCFA and BCFA in the caecotrophs of rabbits fed
325 unsaturated lipid supplemented diets if compared to a control diet, which were reflected to
326 unchanged concentrations of OCFA and BCFA incorporated into *biceps femoris* and
327 *longissimus lumborum* muscles (Papadomichelakis et al., 2010a). In the current study, total
328 OCFA and total BCFA proportions did not differ in LTL muscle of rabbits fed BP0, BP5,
329 BP10 or BP15 diets (Table 6), suggesting lack of differences in caecotrophy activity among
330 treatments.

331 The nutritional quality of fat for human consumption is usually evaluated in terms of the
332 PUFA / SFA ratio (optimal values ≥ 0.45), the $\Sigma n6 / \Sigma n3$ FA ratio (optimal values ≤ 4), the
333 atherogenicity (AI) and thrombogenicity (TI) indexes (both as low as possible) (Lazzaroni,
334 Biagini, & Lussiana, 2009). In all treatments, the PUFA / SFA ratio fell within the
335 recommended values. Anyhow, increasing BP inclusion levels in rabbit diets led to a
336 significant progressive increase of this ratio in the LTL muscle, which was 1.7-fold higher in
337 the BP15 group if compared to BP0 ($P < 0.001$). In ordinary dietary conditions, the $\Sigma n6 / \Sigma$
338 $n3$ PUFA ratio in rabbit meat is set at around 10 (Dalle Zotte, 2002). In the current study, this
339 ratio decreased from 9.30 to 2.88 in muscles of rabbits fed the BP0 and BP15 diets,
340 respectively. BP10 and BP15 diets allowed LTL muscle having the $\Sigma n6 / \Sigma n3$ PUFA ratio
341 within optimal values for human consumption. Optimal values were also obtained in other
342 trials when ALA-rich feedstuffs were fed by rabbits (Peiretti, Gasco, Brugiapaglia, & Gai,
343 2011; Kouba et al., 2008; Peiretti & Meineri, 2008). Significant variations were also observed
344 for the AI and TI, both being lower in the LTL muscle of the rabbits fed BP diets if compared
345 to those fed the control diet. Overall the obtained results on long-chain n-3 PUFA, due to the

positive role exerted in the control of cardiovascular diseases (Endo & Arita, 2016), as well as the studied ratios and indexes show that BP inclusion in rabbit diets may be of particular significance for the related nutritional benefits associated to human consumption of rabbit meat.

3.3. Lipid oxidation

Rabbit meat, due to its high content of PUFA, is prone to lipid oxidation leading to a reduced shelf life of the product (Dalle Zotte & Szendrő, 2011). Susceptibility of rabbit meat to lipid oxidation, during refrigerated or frozen storage, can be reduced by dietary phenolic compounds supplementation (Dalle Zotte et al. 2014). In our study, the effects of the BP dietary inclusion on the oxidative stability of the frozen LTL muscle were investigated and results are shown in Table 7. In the applied experimental conditions, the dietary treatment did not affect TBARS values. Storage time instead increased significantly the oxidation of frozen muscles, with higher values recorded in the samples stored at -25 °C for 60 days.

According to the peroxidability index reported in Table 6, meat samples derived from BP treated rabbits were expected to be more susceptible to lipid peroxidation, while they showed similar TBARS levels compared to the control group. Similar findings were also reported in a study carried out in rabbits fed high dietary fat supplemented with a blackcurrant pomace extract rich in polyphenols (Jurgoński, Juśkiewicz, Zduńczyk, Matusevicius, & Kołodziejczyk, 2014). These authors observed lower concentrations of TBARS in kidneys and serum of rabbits treated with enriched extract diet if compared to control; they concluded that the suppression of lipid peroxidation was due to the blackcurrant polyphenolic compounds that may increase filtration in these organs and inactivate free radicals. Similarly, the results obtained in our study could be related to the antioxidant activity of BP polyphenolic compounds. Such antioxidant activity was confirmed by the high percentage

(65.18%) of inhibition obtained with the *in vitro* free radical DPPH assay, which is in agreement with the results obtained by Bunea et al. (2011) on wild and cultivated blueberries.

3.4. *Sensory evaluation*

The results of the affective tests are summarized in Table 8. The data correspond to a sensory analysis where panelists were asked to rate the likeness, in terms of taste and aroma, of 4 meat samples on a 9-point hedonic scale. Both median and mode values of meat samples from the rabbits fed BP5 and BP15 diets were 7, which corresponds to “like moderately” on the hedonic scale. The BP0 and BP10 meat had the same median value (6 = “like slightly”), while the mode value for BP0 meat was higher than that for BP10 meat (7 and 6, respectively). It should be noted that there were no scores in the first two liking categories and that all samples had the majority of scores located in the “liking part” of the hedonic scale (Figure 1). Looking at the distribution of frequencies in Figure 1, it is evident that meat from BP0, BP5 and BP15 treatments on one hand, and meat from BP10 treatment on the other hand had more responses in the “like moderately” and in the “like slightly” category, respectively.

The average overall liking ratings ranged from 6.1 to 6.2, which correspond to “like slightly” according to the hedonic scale. Although the meat from BP10 and BP15 treatments had different median and mode values, they received the same mean score (Table 8). The ANOVA results showed no significant differences in the average overall liking scores among treatments. The samples were considered acceptable if 50% or more of responses were in the range from 6 to 9 on the hedonic scale; all meats were accepted by panelists (votes ≥ 6 from 66% to 74%).

Although ANOVA can be correctly applied to data that slightly deviate from the Gaussian distribution (Figure 1), we decided also to analyze acceptability data using the non-parametric Friedman’s test by converting the preferences scores given by each consumer into rank order

numbers. As the calculated value of the Friedman's statistic (2.93) is lower than 7.81 (the critical value of a chi-square distribution with 3 degrees of freedom), we can confirm the ANOVA results and conclude that there were not significant differences in liking ranking among the 4 treatments.

Flavor, which comprises mainly the two sensations of taste and aroma, is an important meat quality attribute affecting consumers' meat-buying habits and preferences. The sensory evaluation indicated that the use of BP **did not affect** the flavor of cooked rabbit meats.

4. Conclusions

The obtained results show that BP is a suitable ingredient for growing rabbits' diets. Inclusion of BP up to 15% of the diet does not affect the carcass characteristics of rabbits, nor the proximate composition, lipid oxidation and consumer acceptance of rabbit meat.

Dietary inclusion of BP **improved the fatty acid profile of the *longissimus thoracis et lumborum* muscle of rabbits, largely by means of an increase of total n3 PUFA and a reduction of the Σ n6 / Σ n3 fatty acids ratio. Such modifications also determined the lowering of both the atherogenicity and thrombogenicity indexes of the meat.**

Bilberry pomace, being a rich source of **polyunsaturated fatty acids**, bioactive antioxidant compounds and natural colorants (mainly anthocyanins), possesses a good potential for the valorization of bilberry juice by-products through further uses in the feed industry.

Acknowledgements

This research was supported by a "University of Torino (ex 60%)" grant (Es. fin. 2013). Thanks are due to Dr. Gabriella Garino ("Arc en Ciel" Soc. Agr. Coop., Cafasse, Turin, Italy) for providing the bilberry pomace, Mrs. Vanda Maria Malfatto and Dr. Camilla Ragazzoni for

technical support, Mr. Dario Sola, Dr. Paolo Montersino and Mr. Mario Colombano for rabbit care.

References

Abdel-Khalek, A. M. (2013). Supplemental antioxidants in rabbit nutrition: A review. *Livestock Science*, 158, 95–105.

Alves, S. P., Cabrita, A. R. J., Fonseca, A. J. M., & Bessa, R. J. B. (2008). Improved method for fatty acid analysis in herbage based on direct transesterification followed by solid-phase extraction. *Journal of Chromatography A*, 1209, 212–219.

AOAC International (2000). *Official methods of analysis* (17th ed.). Gaithersburg (MD, USA): Association of Official Analytical Chemists.

AOAC International (2003). *Official methods of analysis* (17th ed., 2nd revision). Gaithersburg (MD, USA): Association of Official Analytical Chemists.

Belforti, M., Gai, F., Lussiana, C., Renna, M., Malfatto, V., Rotolo, L., De Marco, M., Dabbou, S., Schiavone, A., Zoccarato, I., & Gasco, L. (2015). *Tenebrio molitor* meal in rainbow trout (*Oncorhynchus mykiss*) diets: effects on animal performance, nutrient digestibility and chemical composition of fillets. *Italian Journal of Animal Science*, 14, 670–675.

Blasco, A., Ouhayoun, J., & Masoero, G. (1993). Harmonization of criteria and terminology in rabbit meat research. *World Rabbit Science*, 1, 3–10.

Bunea, A., Rugină, D. O., Pinteă, A. M., Sconța, Z., Bunea, C. I., & Socaciu, C. (2011). Comparative polyphenolic content and antioxidant activities of some wild and cultivated blueberries from Romania. *Notulae Botanicae Horti Agrobotanici*, 39, 70–76.

443 Commission Internationale de l'Éclairage (1976). Recommendations on uniform colour
 444 spaces-colour difference equations, psychometric colour terms (Supplement no. 2 to CIE
 445 publication No. 15). Commission Internationale de l'Éclairage, Paris, France.

446 Dabbou, S., Gasco, L., Gai, F., Zoccarato, I., Rotolo, L., Dabbou Fekih, S., Brugiapaglia, A.,
 447 Helal A. N., & Peiretti, P. G. (2014). Dried artichoke bracts in rabbits nutrition: effects
 448 on the carcass characteristics, meat quality and fatty-acid composition. *Animal*, 8, 1547–
 449 1553.

450 Dal Bosco, A., Castellini, C., Bianchi, L., & Mugnai, C. (2004). Effect of dietary α -linolenic
 451 acid and vitamin E on the fatty acid composition, storage stability and sensory traits of
 452 rabbit meat. *Meat Science*, 66, 407–413.

453 Dal Bosco, A., Castellini, C., Martino, M., Mattioli, S., Marconi, O., Sileoni, V., Ruggeri, S.,
 454 Tei, F., & Benincasa, P. (2015). The effect of dietary alfalfa and flax sprouts on rabbit
 455 meat antioxidant content, lipid oxidation and fatty acid composition. *Meat Science*, 106,
 456 31–37.

457 Dal Bosco, A., Mourvaki, E., Cardinali, R., Servili, M., Sebastiani, B., Ruggeri, S., Mattioli,
 458 S., Taticchi, A., Esposto, S., & Castellini, C. (2012). Effect of dietary supplementation
 459 with olive pomaces on the performance and meat quality of growing rabbits. *Meat*
 460 *Science*, 92, 783–788.

461 Dal Bosco, A., Mugnai, C., Roscini, V., Mattioli, S., Ruggeri, S., & Castellini, C. (2014).
 462 Effect of dietary alfalfa on the fatty acid composition and indexes of lipid metabolism of
 463 rabbit meat. *Meat Science*, 96, 606–609.

464 Dalle Zotte, A. (2002). Perception of rabbit meat quality and major factors influencing the
 465 rabbit carcass and meat quality. *Livestock Production Science*, 75, 11–32.

466 Dalle Zotte, A., Cullere, M., Sartori, A., Szendrő, Z., Kovács, M., Giaccone, V., & Dal Bosco,
 467 A. (2014). Dietary Spirulina (*Arthrospira platensis*) and Thyme (*Thymus vulgaris*)

468 supplementation to growing rabbits: Effects on raw and cooked meat quality, nutrient
469 true retention and oxidative stability. *Meat Science*, 98, 94–103.

470 Dalle Zotte, A., & Szendrő, Z. (2011). The role of rabbit meat as functional food. *Meat*
471 *Science*, 88, 319-331.

472 Eid, Y. (2008). Dietary grape pomace affects lipid peroxidation and antioxidative status in
473 rabbit semen. *World Rabbit Science*, 16, 155–161.

474 Eid, Y., Zeweil, H., Ahmed, M., Basyony, M., & Farok, M. (2011). Effect of plant source of
475 omega-3 fatty acids and green tea powder on the performance and meat quality of
476 growing rabbits. *Egyptian Journal of Rabbit Science*, 20, 115–134.

477 Endo, J., & Arita, M. (2016). Cardioprotective mechanism of omega-3 polyunsaturated fatty
478 acids. *Journal of Cardiology*, 67, 22–27.

479 European Economic Community (1972). Analytical determination of starch. *Official Journal*
480 *of European Communities*, L123, 7.

481 European Group on Rabbit Nutrition (2001). Technical note: attempts to harmonize chemical
482 analyses of feeds and faeces, for rabbit feed evaluation. *World Rabbit Science*, 9, 57–64.

483 Forrester-Anderson, I. T., McNitt, J., Way, R., & Way, M. (2006). Fatty acid content of
484 pasture-reared fryer rabbit meat. *Journal of Food Composition and Analysis*, 19, 715–
485 719.

486 Goby, J. P., & Gidenne, T. (2008). Nutritive value of carrot (whole plant), dried at low
487 temperature, for the growing rabbit. *Proceedings of 9th World Rabbit Congress*, Verona,
488 Italy, pp. 677–682.

489 Heyman, L., Axling, U., Blanco, N., Sterner, O., Holm, C., & Berger, K. (2014). Evaluation
490 of beneficial metabolic effects of berries in high-fat fed C57BL/6J mice. *Journal of*
491 *Nutrition and Metabolism*, Vol. 2014, Article ID 403041, 12 pages.

492 Jenkins, B, West, J. A., & Koulman, A. (2015). A review of odd-chain fatty acid metabolism
 493 and the role of pentadecanoic acid (C15:0) and heptadecanoic acid (C17:0) in health and
 494 disease. *Molecules*, 20, 2425–2444.

495 Julkunen-Tiitto, R. (1985). Phenolic constituents in the leaves of northern willows: methods
 496 for the analysis of certain phenolics. *Journal of Agricultural and Food Chemistry*, 33,
 497 213–217.

498 Jurgoński, A., Juśkiewicz, J. Zduńczyk, Z., Matusevicius, P., & Kołodziejczyk, K. (2014).
 499 Polyphenol-rich extract from blackcurrant pomace attenuates the intestinal tract and
 500 serum lipid changes induced by a high-fat diet in rabbits. *European Journal of Nutrition*,
 501 53, 1603–1613.

502 Khanal, R. C., Howard, L. R., Brownmiller, C. R., & Prior, R. L. (2009). Influence of
 503 extrusion processing on procyanidin composition and total anthocyanin contents of
 504 blueberry pomace. *Journal of Food Science*, 74, H52–H58.

505 Khanal, R. C., Howard, L. R., Wilkes, S. E., Rogers, T. J., & Prior, R. L. (2012). Effect of
 506 dietary blueberry pomace on selected metabolic factors associated with high fructose
 507 feeding in growing Sprague-Dawley rats. *Journal of Medicinal Food*, 15, 802–810.

508 Kim, H., Bartley, G. E., Rimando, A. M., & Yokoyama, W. (2010). Hepatic gene expression
 509 related to lower plasma cholesterol in hamsters fed high-fat diets supplemented with
 510 blueberry peels and peel extract. *Journal of Agricultural and Food Chemistry*, 58, 3984–
 511 3991.

512 Kontogiorgis, C., & Hadjipavlou-Litina, D. (2005). Synthesis and antiinflammatory activity
 513 of coumarin derivatives. *Journal of Medicinal Chemistry*, 48, 6400–6408.

514 Kouba, M., Benatmane, F., Blochet, J. E., & Mourot, J. (2008). Effect of a linseed diet on
 515 lipid oxidation, fatty acid composition of muscle, perirenal fat, and raw and cooked rabbit
 516 meat. *Meat Science*, 80, 829–834.

517 Lazzaroni, C., Biagini, D., & Lussiana, C. (2009). Fatty acid composition of meat and
518 perirenal fat in rabbits from two different rearing systems. *Meat Science*, 83, 135–139.

519 Lee, J., & Wrolstad, R. E. (2004). Extraction of anthocyanins and polyphenolics from
520 blueberry processing waste. *Journal of Food Science*, 69, 564–573.

521 Leiber, F., Meier, J. S., Burger, B., Wettstein, H.-R., Kreuzer, M., Hatt, J.-M., & Clauss, M.
522 (2008). Significance of coprophagy for the fatty acid profile in body tissues of rabbits fed
523 different diets. *Lipids*, 43, 853–865.

524 Liu, H. W., Gai, F., Gasco, L., Brugiapaglia, A., Lussiana, C., Guo, K. J., Tong, J. M., &
525 Zoccarato, I. (2009). Effects of chestnut tannins on carcass characteristics, meat quality,
526 lipid oxidation and fatty acid composition of rabbits. *Meat Science* 83, 678–683.

527 MacFie, H. J., Bratchell, N., Greenhoff, K., & Vallis, L. V. (1989). Designs to balance the
528 effect of order of presentation and first-order carry-over effects in hall tests. *Journal of*
529 *Sensory Studies*, 4, 129–148.

530 Mancinelli, A. L., Huang Yang, C. P., Lindquist, P., Anderson, O. R., & Rabine, I. (1975).
531 Photocontrol of anthocyanin synthesis. III. The action of streptomycin on the synthesis of
532 chlorophyll and anthocyanin. *Plant Physiology*, 55, 251–257.

533 Meilgaard, M., Civille G. V., & Carr, B. T. (1991). *Sensory evaluation techniques*. (2nd ed.).
534 Boca Raton, Florida: CRC Press.

535 Montedoro, G. F., Servili, M., Baldioli, M., & Miniati, E. (1992). Simple and hydrolysable
536 compounds in virgin olive oil. 1. Their extraction, separation and quantitative and
537 semiquantitative evaluation by HPLC. *Journal of Agricultural and Food Chemistry*, 40,
538 1571–1576.

539 Naes, T., Brockhoff, B., & Tomic, O. (2010). *Statistics for sensory and consumer science*.
540 West Sussex, United Kingdom: John Wiley and Sons Ltd.

541 Oku, H., & Yanagita, T. (2009). Branched-chain fatty acid as a functional lipid. In C. T. Hou,
542 & J.-F. Shaw (Eds.), *Biocatalysis and Agricultural Biotechnology* (pp. 259–267). CRC
543 Press.

544 Papadomichelakis, G., Karagiannidou, A., Anastasopoulos, V., & Fegeros, K. (2010a). Effect
545 of dietary soybean oil addition on the odd-numbered and branched-chain fatty acids in
546 rabbit meat. *Meat Science*, 86, 264–269.

547 Papadomichelakis, G., Karagiannidou, A., Anastasopoulos, V., & Fegeros, K. (2010b). Effect
548 of high dietary digestible fibre content on the fatty acid composition of two muscles in
549 fattening rabbits. *Livestock Science*, 129, 159–165.

550 Papadomichelakis, G., Anastasopoulos, V., Karagiannidou, A., & Fegeros, K. (2010c).
551 Effects of dietary digestible fibre and soybean oil level on the odd-numbered, branched-
552 chain and hydroxy-fatty acid composition of caecotrophs in rabbits. *Animal Feed Science*
553 *and Technology*, 158, 95–103.

554 Peiretti, P. G., Gasco, L., Brugiapaglia, A., & Gai, F. (2011). Effects of perilla (*Perilla*
555 *frutescens* L.) seeds supplementation on performance, carcass characteristics, meat
556 quality and fatty acid composition of rabbits. *Livestock Science*, 138, 118–124.

557 Peiretti, P. G., & Meineri, G. (2008). Effects on growth performance, carcass characteristics,
558 and the fat and meat fatty acid profile of rabbits fed diets with chia (*Salvia hispanica* L.)
559 seed supplements. *Meat Science*, 80, 1116–1121.

560 Peiretti, P. G., & Meineri, G. (2011). Effects of diets with increasing levels of *Spirulina*
561 *platensis* on the carcass characteristics, meat quality and fatty acid composition of
562 growing rabbits. *Livestock Science*, 140, 218–224.

563 Peryam, D. R., & Pilgrim, P. J. (1957). Hedonic scale method for measuring food preferences.
564 *Food Technology*, 11, 9–14.

Renna, M., Gasmi-Boubaker, A., Lussiana, C., Battaglini, L.M., Belfayez, K., & Fortina, R.
(2014). Fatty acid composition of the seed oils of selected *Vicia* L. taxa from Tunisia.
Italian Journal of Animal Science, 13 (2), 308–316.

Renna, M., Cornale, P., Lussiana, C., Malfatto, V., Fortina, R., Mimosi, A., & Battaglini, L.
M. (2012). Use of *Pisum sativum* (L.) as alternative protein resource in diets for dairy
sheep: effects on milk yield, gross composition and fatty acid profile. *Small Ruminant
Research*, 102, 142–150.

Van Soest, P. J., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fibre, neutral
detergent fibre and nonstarch polysaccharides in relation to animal nutrition. *Journal of
Dairy Science*, 74, 3583–3597.

Vulić, J. J., Tumbas, V. T., Savatović, S. M., Đilas, S. M., Četković, G. S., & Čanadanović-
Brunet, J. M. (2011). Polyphenolic content and antioxidant activity of the four berry
fruits pomace extracts. *Acta Periodica Technologica*, 42, 271–279.

Zhishen, J., Mengcheng, T., & Jianming, W. (1999). The determination of flavonoid contents
in mulberry and their scavenging effects on superoxide radical. *Food Chemistry*, 64, 555–
559.

590 **Table 1**

591 Mean concentration of the main phenolic compounds of bilberry pomace.

Phytochemical compounds	Content
Total phenols (mg hydroxytyrosol equivalents / g DM)	46.54
<i>o</i> -diphenols (mg hydroxytyrosol equivalents / g DM)	10.55
Total flavonoids (mg catechin equivalents / g DM)	9.45
Total anthocyanins (mg CyE / g DM)	0.14
Condensed tannins (mg catechin equivalents / g DM)	36.38

592 Abbreviations: DM = dry matter; CyE = cyanidin 3-glucoside equivalents.

593 **Table 2**

594 Proximate composition of bilberry pomace (BP) and experimental diets.

<i>Experimental diets (g / kg as fed)</i>	BP	BP0	BP5	BP10	BP15
Basal mixture ¹	-	980	930	880	830
Bilberry pomace	-	0	50	100	150
Vitamin-mineral premix ²	-	15	15	15	15
Bicalcium phosphate	-	5	5	5	5
<i>Proximate composition (g / kg DM, unless otherwise stated)</i>					
Dry matter (g / kg)	944	882	882	880	885
Ash	18	75	75	71	72
Crude protein	142	177	177	175	176
Ether extract	155	26	33	39	42
Neutral detergent fiber	626	368	368	372	391
Acid detergent fiber	433	198	208	220	233
Acid detergent lignin	258	35	46	56	68
Starch	137	199	204	207	205
Gross energy (MJ / kg DM)	22.7	17.9	18.1	18.4	18.6

595 Abbreviations: DM = dry matter.

596 ¹Containing (g / kg fresh matter): alfalfa meal 300, wheat bran 200, barley 170, dried beet
597 pulp 150, soybean meal 115, molasses 20, wheat straw 20, soybean oil 5.

598 ²Containing (per kg of diet): Vitamin A 200 U, α -tocopheryl acetate 16 mg, Niacin 72 mg,
599 Vitamin B6 16 mg, Cholin 0.48 mg, DL-methionin 600 mg, Ca 500 mg, P 920 mg, K 500 mg,
600 Na 1 g, Mg 60 mg, Mn 17 mg, Cu 0.6 mg.

601 **Table 3**

602 Fatty acid composition (g / 100 g of total FA) of bilberry pomace (BP) and experimental diets.

Fatty acid	BP	BP0	BP5	BP10	BP15
C14:0	0.06	0.39	0.32	0.33	0.24
C16:0	4.82	21.21	17.25	15.09	13.37
C16:1 <i>c</i> 9	0.09	0.19	0.16	0.14	0.14
C18:0	1.19	3.41	2.85	2.63	2.48
C18:1 <i>c</i> 9	23.50	17.28	18.78	20.02	21.50
C18:1 <i>c</i> 11	0.58	0.98	0.92	0.86	0.83
C18:2 <i>n</i> 6	36.33	45.33	44.24	42.08	40.61
C18:3 <i>n</i> 3	32.59	8.88	13.47	16.98	19.06
C20:0	0.30	0.54	0.47	0.49	0.52
C20:1 <i>c</i> 11	0.16	0.35	0.34	0.28	0.27
C20:4 <i>n</i> 6	0.02	0.17	0.14	0.12	0.09
C22:0	0.13	0.57	0.44	0.42	0.38
C24:0	0.09	0.42	0.34	0.28	0.26
Other FA ¹	0.16	0.27	0.28	0.29	0.25
SFA	6.61	26.66	21.80	19.39	17.36
MUFA	24.35	18.89	20.28	21.35	22.79
PUFA	69.03	54.45	57.92	59.27	59.85
TFA (g / 100 g DM)	15.21	2.13	2.72	3.32	3.98

603 Abbreviations: FA = fatty acids; *t* = *trans*; *c* = *cis*; SFA, saturated fatty acids; MUFA,

604 monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; TFA, total fatty acids; DM =

605 dry matter.

606 ¹Other FA (all ≤ 0.10 g / 100 g of total FA in BP and in the experimental diets): C10:0 +607 C12:0 + C16:1 *t*3 + C18:3 *n*6.

608 **Table 4**

609 Effect of dietary bilberry pomace (BP) on the carcass characteristics of rabbits (n = 12).

	BP0	BP5	BP10	BP15	SEM	P-value
SW (g)	3002	2927	2978	3035	19.675	0.256
Skin (% SW)	16.7	17.0	17.2	16.9	0.109	0.516
Full gastrointestinal tract (% SW)	13.7	13.6	13.2	13.2	0.170	0.696
CCW (g)	1878	1844	1872	1927	14.200	0.220
Dressing out (%)	62.5	63.0	62.9	63.5	0.195	0.386
Head (% CCW)	9.19	9.31	9.10	9.17	0.070	0.746
Liver (% CCW)	4.29 ^a	3.68 ^b	3.99 ^{ab}	3.61 ^b	0.080	0.003
RCW (g)	1574	1555	1580	1633	12.306	0.139
Perirenal fat (% RCW)	1.90	1.67	2.13	2.04	0.080	0.196

610 Abbreviations: SW = slaughter weight; CCW = chilled carcass weight; RCW = reference

611 carcass weight.

612 ^{a-b} Different superscripts within a row indicate significant differences (**P<0.05**).

613 **Table 5**

614 Effect of dietary bilberry pomace (BP) on the quality traits of *longissimus thoracis et*

615 *lumborum* muscle (n = 12).

	BP0	BP5	BP10	BP15	SEM	P-value
pH ₂₄	5.86	5.79	5.87	5.86	0.021	0.494
Color						
L [*]	52.13	52.67	51.60	51.49	0.491	0.827
a [*]	0.05	-0.08	-0.15	-0.01	0.082	0.862
b [*]	5.52	6.04	5.55	5.99	0.166	0.560
Proximate composition (% fresh matter)						
Water	75.11	74.94	74.68	74.86	0.070	0.158
Protein	22.50	22.53	22.97	22.70	0.070	0.060
Ether extract	0.87	0.95	0.95	1.03	0.043	0.646
Ash	1.21	1.23	1.25	1.22	0.010	0.572

616 Abbreviations: L*: lightness; a*: redness; b*: yellowness.

617 **Table 6**

618 Effect of dietary bilberry pomace (BP) on the fatty acid composition (g / 100 g of total FA) of
 619 *longissimus thoracis et lumborum* muscle of rabbits (n = 12).

	BP0	BP5	BP10	BP15	SEM	P-value
Σ SFA	42.93 ^a	39.89 ^b	38.83 ^c	36.48 ^d	0.376	<0.001
C10:0	0.13	0.11	0.10	0.11	0.006	0.202
C12:0	0.17 ^a	0.15 ^{ab}	0.13 ^b	0.13 ^b	0.005	0.010
C14:0	2.60 ^a	2.37 ^{ab}	2.33 ^b	1.98 ^c	0.052	<0.001
C15:0	1.31	1.29	1.17	1.24	0.026	0.266
C16:0	29.84 ^a	27.66 ^b	26.85 ^b	24.70 ^c	0.302	<0.001
C17:0	0.61	0.60	0.59	0.62	0.009	0.523
C18:0	6.76 ^a	6.16 ^b	6.33 ^b	6.28 ^b	0.080	0.036
C20:0	0.08 ^b	0.09 ^b	0.08 ^b	0.12 ^a	0.004	0.004
C22:0	0.18 ^a	0.11 ^b	0.09 ^c	0.07 ^c	0.007	<0.001
Σ BCFA	1.26	1.35	1.17	1.23	0.036	0.388
C15:0 <i>iso</i>	0.41	0.46	0.39	0.44	0.020	0.606
C15:0 <i>aiso</i>	0.07	0.09	0.08	0.08	0.002	0.061
C16:0 <i>iso</i>	0.36	0.40	0.34	0.39	0.017	0.630
C17:0 <i>iso</i>	0.05	0.05	0.05	0.05	0.002	0.952
C17:0 <i>aiso</i>	0.36 ^a	0.34 ^{ab}	0.31 ^{bc}	0.28 ^c	0.008	0.003
Σ MUFA	30.99 ^a	28.19 ^b	27.40 ^{bc}	25.90 ^c	0.373	<0.001
C16:1 <i>c9</i>	4.89 ^a	4.02 ^{ab}	3.58 ^{bc}	2.99 ^c	0.178	0.001
C17:1 <i>c9</i>	0.32 ^a	0.31 ^a	0.28 ^b	0.26 ^b	0.006	<0.001
C18:1 <i>t6-11</i>	0.09 ^a	0.06 ^{bc}	0.07 ^b	0.06 ^c	0.003	<0.001
C18:1 <i>t12-14</i>	0.09 ^{ab}	0.08 ^b	0.11 ^a	0.11 ^a	0.004	0.023
C18:1 <i>c9</i>	23.63 ^a	22.12 ^b	21.76 ^b	21.16 ^b	0.237	0.001
C18:1 <i>c11</i>	1.51 ^a	1.21 ^b	1.15 ^{bc}	1.04 ^c	0.031	<0.001
C18:1 <i>c12</i>	0.12 ^{ab}	0.11 ^b	0.14 ^a	0.14 ^a	0.005	0.012
C18:1 <i>c14</i> (+ <i>c13</i> + <i>c15</i>)	0.15	0.12	0.16	0.16	0.006	0.116
C18:2 ¹	0.62 ^a	0.48 ^{ab}	0.48 ^{ab}	0.38 ^b	0.028	0.023
C20:1 <i>c11</i>	0.20 ^a	0.16 ^{ab}	0.16 ^{ab}	0.14 ^b	0.006	0.037
Σ PUFA	26.08 ^c	31.92 ^b	33.77 ^b	37.46 ^a	0.670	<0.001
C18:2 <i>n6</i>	19.67 ^c	22.75 ^{ab}	22.56 ^b	23.99 ^a	0.319	<0.001
CLA <i>c9t11</i>	0.05	0.04	0.05	0.04	0.002	0.406

C18:3 n3	2.10 ^d	4.65 ^c	6.28 ^b	8.65 ^a	0.362	<0.001
C20:3 n6	0.35	0.34	0.34	0.33	0.012	0.957
C20:4 n6	2.92	3.04	3.09	3.09	0.105	0.939
C20:5 n3	0.12 ^b	0.16 ^b	0.25 ^a	0.29 ^a	0.013	<0.001
C22:5 n3	0.25 ^c	0.46 ^b	0.73 ^a	0.70 ^a	0.036	<0.001
Σ n3	2.47 ^d	5.27 ^c	7.26 ^b	9.64 ^a	0.393	<0.001
Σ n6	22.94 ^b	26.13 ^a	25.99 ^a	27.40 ^a	0.365	<0.001
Σ n6 / Σ n3	9.30 ^a	5.02 ^b	3.61 ^c	2.88 ^d	0.366	<0.001
Σ PUFA / Σ SFA	0.61 ^c	0.80 ^b	0.87 ^b	1.03 ^a	0.025	<0.001
PI ²	39.86 ^d	49.79 ^c	55.16 ^b	61.24 ^a	0.261	<0.001
AI ²	0.72 ^a	0.63 ^b	0.60 ^b	0.52 ^c	0.011	<0.001
TI ²	1.14 ^a	0.84 ^b	0.73 ^c	0.59 ^d	0.030	<0.001
Δ5- plus Δ6-desaturase ²	13.04	11.79	12.37	11.11	0.356	0.273
Δ9-desaturase ³	0.78	0.77	0.76	0.78	0.009	0.905
TFA (mg / 100 g FM)	732.38	866.78	862.61	905.40	41.268	0.494

Abbreviations: FA = fatty acids; SFA = saturated fatty acids; BCFA = branched-chain fatty

acids; MUFA = monounsaturated fatty acids; *c*, *cis*; *t*, *trans*; PUFA = polyunsaturated fatty

acids; CLA = conjugated linoleic acid; PI = peroxidability index; AI = atherogenicity index;

TI = thrombogenicity index; TFA = total fatty acids; FM = fresh matter.

^{a-d} Different superscripts within a row indicate significant differences (**P<0.05**).

¹Sum of octadecadienoic isomers *t9t12*, *c9t13*, *t8c12*, *c9t12*, *t8c13*, *t9c12*, *t11c15*.

²Calculated as reported by Dal Bosco et al. (2014):

PI = (% monoenoic × 0.025) + (% dienoic × 1) + (% trienoic × 2) + (% tetraenoic × 4) +

(% pentaenoic × 6) + (% hexaenoic × 8);

AI = (C12:0 + 4 × C14:0 + C16:0) / [(Σ MUFA + Σ n6) + Σ n3];

TI = (C14:0 + C16:0 + C18:0) / [(0.5 × Σ MUFA + 0.5 × Σ n6 + 3 × Σ n3) + (Σ n3) / Σ

n6)];

632 Estimated $\Delta 5$ -desaturase plus $\Delta 6$ -desaturase activity = $(C20:2\ n6 + C20:4\ n6 + C20:5\ n3 +$
633 $C22:5\ n3 + C22:6\ n3) / (C18:2\ n6 + C18:3\ n3 + C20:2\ n6 + C20:4\ n6 + C20:5\ n3 + C22:5$
634 $n3 + C22:6\ n3) \times 100$.

635 ³Estimated $\Delta 9$ -desaturase activity = $(C16:1\ c9 + C18:1\ c9) / (C16:0 + C18:0)$.

636 **Table 7**
637 Effect of dietary bilberry pomace (BP) on the oxidative status (TBARS, mg TMP / kg of
638 meat) of *longissimus thoracis et lumborum* muscle of rabbits during frozen storage at -25 °C
639 (n = 12).

	30 days				60 days				P-value		
	BP0	BP5	BP10	BP15	BP0	BP5	BP10	BP15	Diet	Time	Interaction
TBARS	0.30	0.30	0.33	0.28	0.34	0.46	0.35	0.35	0.151	0.013	0.269

640 **Table 8**

641 Sensory acceptance test of rabbit meat: descriptive statistics, mean score and rank sums.

	BP0	BP5	BP10	BP15	P-value
Median	6	7	6	7	-
Mode	7	7	6	7	-
Mean	6.2	6.2	6.1	6.1	0.958
Rank Sum	161	169	185	165	0.317

Figure 1

Frequency of acceptability scores for each meat sample.

