

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**New insights in phytoplasma-vector interaction: acquisition and inoculation of flavescence dorée phytoplasma by Scaphoideus titanus adults in a short window of time**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1669919> since 2018-06-22T15:50:30Z

*Published version:*

DOI:10.1111/aab.12433

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

ALMA A., LESSIO F., GONELLA E., PICCIAU L., MANDRIOLI M., TOTA F. 2018. New insights in phytoplasma-vector interaction: acquisition and inoculation of flavescence dorée phytoplasma by *Scaphoideus titanus* adults in a short window of time. *Annals of Applied Biology* 173, 55-62.

**The publisher's version is available at:**

<https://onlinelibrary.wiley.com/doi/10.1111/aab.12433>

**When citing, please refer to the published version.**

This full text was downloaded from iris-Aperto: <https://iris.unito.it/>

1  
2  
3 1 **New insights in phytoplasma-vector interaction: acquisition and**  
4  
5 2 **inoculation of Flavescence dorée phytoplasma by *Scaphoideus titanus***  
6  
7 3 **adults in a short window of time**  
8  
9  
10 4

11 5 **A. Alma<sup>1</sup>, F. Lessio<sup>1</sup>, E. Gonella<sup>1</sup>, L. Picciau<sup>1</sup>, M. Mandrioli<sup>2</sup>, F. Tota<sup>1</sup>**

12 6 <sup>1</sup>Dipartimento di Scienze Agrarie, Forestali e Alimentari (DISAFA), University of Torino,  
13 7 Grugliasco (TO), Italy

14 8 <sup>2</sup>Dipartimento di Scienze della Vita (DSV), University of Modena e Reggio Emilia, Modena, Italy  
15 9

16 10 Correspondence: A. Alma, Dipartimento di Scienze Agrarie, Forestali e Alimentari (DISAFA),  
17 11 Università degli Studi di Torino, Largo P. Braccini 2, 10095 Grugliasco (TO), Italy. E-mail  
18 12 alberto.alma@unito.it, Phone +39 011 6708534, Fax +39 011 6708535  
19 13

20 14 **Running title:** Phytoplasma acquisition and transmission by *Scaphoideus titanus* adults  
21 15

22 16 **Abstract:**

23 17 The leafhopper *Scaphoideus titanus* is able to transmit 16SrV phytoplasmas agents of grapevine's  
24 18 Flavescence dorée (FD) within 30-45 days, following an Acquisition Access Period (AAP) of a few  
25 19 days feeding on infected plants as a nymph, a Latency Period (LP) of 3-5 weeks becoming  
26 20 meanwhile an adult, and an Inoculation Access Period (IAP) of a few days on healthy plants.  
27 21 However, several aspects of FD epidemiology suggest how the whole transmission process may  
28 22 take less time, and may be start directly with adults of the insect vector. Transmission experiments  
29 23 have been set up under lab condition. Phytoplasma-free *S. titanus* adults were placed on broad bean  
30 24 (BB) plants (*Vicia faba*) infected by FD-C (16SrV-C) phytoplasmas for an AAP=7 days.  
31 25 Afterwards, they were immediately moved onto healthy BB for IAP, which were changed every 7  
32 26 days, obtaining three timings of inoculation: IAP 1, IAP 2 and IAP 3, lasting 7, 14 and 21 days from  
33 27 the end of AAP, respectively. DNA was extracted from plants and insects, and PCR were performed  
34 28 to identify FD phytoplasmas. Insects were dissected and fluorescent in situ hybridization was made  
35 29 to detect the presence of phytoplasmas in midguts and salivary glands. The rate of infection in  
36 30 insects ranged 46-68% without significant differences among IAPs. Inoculation in plants succeeded  
37 31 in all IAPs, at a rate of 16-23% (no significant differences). Phytoplasma load was significantly  
38 32 higher in IAP3 than IAP 1-2 for both plants and insects. Phytoplasmas were identified both in  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 33 midgut and salivary glands of *S. titanus* at all IAP times. The possible implications of these results  
4 34 in the epidemiology of Flavescence dorée are discussed.  
5  
6

35

7  
8 36 **Key words:** Flavescence dorée, *Scaphoideus titanus*, *Vicia faba*, acquisition by adults, latency  
9 37 access period, transmission process  
10

11 38  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## 39 Introduction

40  
41 Flavescence dorée (FD) is a serious disease of grapevine caused by 16SrV phytoplasmas  
42 (subgroups C and D) (Arnaud *et al.*, 2007) transmitted by *Scaphoideus titanus* Ball (Chuche &  
43 Thiery, 2014), and to a lesser extent by *Dictyophara europaea* (L.) (Filippin *et al.*, 2009) and  
44 *Orientus ishidae* (Matsumura) (Lessio *et al.*, 2016). However, unlike these two last ones, *S. titanus*  
45 is able to accomplish its whole life cycle only on grapevine, both *Vitis vinifera* L. and American  
46 grapevines' and/or wild rootstocks (Maixner *et al.*, 1993; Vidano, 1966; Lessio *et al.*, 2007;  
47 Chuche & Thiery, 2014). *S. titanus* is a univoltine species, and overwinters as egg, laid under the  
48 bark of 2-yr old wood, although sometimes 1-yr old wood can be used as well (Bagnoli *et al.*, 2011;  
49 Lessio & Alma, 2013). During its postembryonic development, it undergoes five nymphal instars:  
50 N1, N2, N3, N4 and N5. The time elapsing between molts depends mostly on temperature (Falzoi  
51 *et al.*, 2014). Usually, eggs start hatching in the beginning of May. Hatching dynamic depends on the  
52 winter temperatures eggs are exposed to: the colder is the winter, the shorter will be the hatching  
53 period (Chuche & Thiery, 2009).

54 Transmission of phytoplasmas by insect vectors follows a typical sequence of events: Acquisition  
55 Access Period (AAP), Latency Access Period (LAP), or Latency Period (LP), an Inoculation Access  
56 Period (IAP) (Alma *et al.*, 2015). The transmission is therefore considered as “persistent-  
57 propagative”. According to the state-of art, *S. titanus* nymphs (N3, N4 and N5) can acquire  
58 phytoplasmas by feeding on infected grapevines (both European and American), and during their  
59 LAP they become adults able to move from plant to plant and to transmit phytoplasmas to healthy  
60 plants (Chuche & Thiery, 2014; Lessio *et al.*, 2007). In *S. titanus*, firstly, the LAP was thought  
61 lasting about 28-35 days, with a minimum of 7 days for both AAP and IAP starting from N3  
62 nymphs (Schvester *et al.*, 1961; 1969). Afterwards, Caudwell *et al.* (1970) demonstrated a possible  
63 inoculation to broad bean (BB) (*Vicia faba* L.) and grapevine after 21 days of AAP + LAP and 7  
64 days of IAP (total days: 28).

65 Acquisition efficiency of phytoplasmas by *S. titanus* depends on a number of factors. The source of  
66 inoculum plays an important role: highly susceptible vine varieties are a better source than tolerant  
67 ones (Bressan *et al.*, 2005b; Galetto *et al.*, 2016); as well, a higher phytoplasma concentration  
68 increases acquisition efficiency (Galetto *et al.*, 2014; 2016), which on the other hand is lower in the  
69 case of recovered grapevines (Roggia *et al.*, 2014). Moreover, acquisition is more efficient in late  
70 summer because of a higher phytoplasma load in grapevines (Galetto *et al.*, 2014). However, in late  
71 summer there are no nymphs left, but only adults. If AAP is performed just by nymphs, this aspect  
72 would be biologically unrealistic, whereas it might be possible that adults acquire phytoplasmas too.

1  
2  
3 73 At present, the whole literature about *S. titanus* and transmission of FD phytoplasma (FDP)  
4 74 involves the nymphs for AAP, and adults for IAP, after a LAP of 28- 35 days. It is therefore  
5 75 necessary to ascertain if the whole transmission period could be completed within adults' lifespan.  
6 76 The aim of this research was to investigate if *S. titanus* adults are able to complete the transmission  
7 77 process starting from AAP and finishing with IAP. Moreover, since adults have never been used for  
8 78 AAP before, we reviewed the LAP in order to state if the whole transmission process could be  
9 79 completed within a shorter period.  
10  
11  
12  
13  
14

## 15 80 16 81 **Materials and methods**

### 17 82 18 83 Insect source

19 84  
20 85 *S. titanus* was reared under lab conditions in order to obtain healthy individuals. Grapevine canes  
21 86 containing eggs (two-year old wood, or older) were collected during winter in 2014 and 2015 in  
22 87 several Piedmontese grapevine growing areas, in particular Asti Province and Canavese district,  
23 88 where *S. titanus* was detected in great number during the previous summer, by means of yellow  
24 89 sticky traps. Canes were cut in pieces of approximately 20 cm length, placed in plastic bags,  
25 90 periodically sprinkled to avoid desiccation of eggs, and preserved into a cool chamber (+4°C)  
26 91 before use. The twigs were placed into BugDorm® insect rearing cages (47.5 x 47.5 x 138 cm)  
27 92 made of mesh and polyethylene at springtime, along with potted plants of healthy BB. Broad beans  
28 93 were used to rear *S. titanus* through its whole life cycle to avoid any phytoplasmas contamination of  
29 94 the insects before the transmission experiments, as 16SrV phytoplasmas are thought not being  
30 95 transmitted through seeds (Duduk and Bertaccini, 2009). The cages were kept either outdoors,  
31 96 under a shelter against rain and sunlight, from May 15th to September 1st, or into a greenhouse  
32 97 from March 15th to May 15th and from September 1st to November 15th, and periodically  
33 98 sprinkled. Insects were reared up to the adult stage, and then used for transmission experiments.  
34 99 Adults from such rearing were also directly collected and dissected for fluorescence in situ  
35 100 hybridization (FISH) analysis, as a negative control.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

### 48 101 49 102 Transmission experiments

50 103  
51 104 Transmission experiments were performed in a climatic chamber (T=25°C, RH=75%). Acquisition  
52 105 (AAP) of phytoplasmas by *S. titanus* adults was made on BBs previously infected with FD-C  
53 106 phytoplasmas by means of *Euscelidius variegatus* (Kirschbaum) (Caudwell *et al.*, 1972; Salar *et*

1  
2  
3 107 *al.*, 2013). Plants used for AAP were 30-40 cm high, and had 8-10 leaves; before use, phytoplasma  
4 108 infection was checked by qPCR (see below). AAP was performed by placing 1-2 infected plants  
5 109 into BugDorm® insect rearing cages (47.5 x 47.5 x 47.5 cm) along with 20-30 newly emerged  
6 110 adults of *S. titanus* for 7 days. The procedure was repeated several times until enough live infected  
7 111 adults to be used in IAPs were obtained.

8  
9  
10 112 After AAP, insects were divided into batches of 5 individuals, and each batch was placed onto a  
11 113 healthy BB seedling (height 5 cm, 2 leaves), inside a Plexiglas cylinder (h=20 cm; diameter=12 cm)  
12 114 with the top covered by a fine mesh. Every 7 days, live insects were moved onto another plant up to  
13 115 three times to perform a total of three IAPs, named IAP1-3 (Fig. 1). Cylinders were checked daily,  
14 116 and dead insects were removed and preserved into a freezer (-20°C) for molecular analyses to test  
15 117 for FDP (see later).

16  
17 118 The same set up was repeated for FISH analysis: at the end of each IAP five live leafhoppers from  
18 119 single batches (one for each of the IAPs) were collected and dissected to isolate midgut and salivary  
19 120 glands. Hence, a total of 28 batches with 140 specimens (25 batches with 125 specimens for qPCR,  
20 121 and 3 batches with 15 insects for FISH, Fig 1) started IAP 1, however the number of batches  
21 122 progressively decreased due to insect mortality or collection in IAPs 2 and 3. Inoculated plants were  
22 123 treated with an insecticide (Dichlorvos, 0.5 g/L) and kept in a climatic chamber, inside an insect-  
23 124 proof cage to avoid contamination, for three weeks from the beginning of phytoplasma inoculation  
24 125 before molecular analyses.

25  
26 126

#### 27 127 DNA extraction and quantitative Real Time PCR analysis

28 128

29 129 Total DNA extraction was performed from whole, single insects used in transmission experiments  
30 130 and from inoculated BB. Nucleic acids extraction from *S. titanus* was carried out by following a  
31 131 procedure previously described for leafhoppers (Marzachi *et al.*, 1998). Plant DNA was extracted  
32 132 from leaf tissue previously grounded with liquid nitrogen in a sterile mortar, according to the  
33 133 DNeasy Plant Mini Kit protocol instructions (Qiagen, Milan, Italy). Quantitative real-time PCR  
34 134 (qPCR) was carried out to measure the presence and concentration of phytoplasma genome units in  
35 135 insect and plant samples. A Chromo4 real-time detector (Bio-Rad, Milan, Italy) was used with  
36 136 PrecisionPlus™ -SY Mastermix (Primerdesign, Chandler's Ford, UK). Reactions targeting the 16S  
37 137 rRNA gene of group 16SrV phytoplasmas were carried out on all samples by using the fAY/rEY  
38 138 primer pair (Marccone *et al.*, 1996; Marzachi *et al.*, 2001), with the conditions described by Galetto  
39 139 *et al.* (2005). To calculate the average FDP Genome Units (GU) / sample, 16S rRNA gene copy  
40 140 numbers were divided by two, because this gene is estimated to be in double copy in the genome of

1  
2  
3 141 phytoplasmas (Schneider & Seemuller, 1994). Additionally, qPCR targeting the insect's 18S rRNA  
4 142 was performed on insect DNA gene to normalize the absolute phytoplasma density. Primers MqFw  
5 143 / MqRv were used according to Marzachi & Bosco (2005). Hence, normalized phytoplasma GU  
6 144 were calculated per pg of insect 18Sr RNA gene. On the other hand, qPCR results from BBs were  
7 145 expressed as FDp GU per 100 mg of leaf.

8  
9  
10 146 Standard curves were constructed by using dilutions of PCR-amplified 16S rRNA gene of FDP  
11 147 cloned with the pGEM T-easy Vector Cloning Kit (Promega, Milan, Italy). The detection limit was  
12 148 calculated as the highest dilution of cloned amplicons used for standard curves which was  
13 149 successfully amplified, and corresponded to  $1.0 \times 10^0$  FDP GU.

14  
15  
16  
17 150

#### 18 19 151 Fluorescent *in situ* hybridization

20  
21 152

22 153 FISH analyses were performed on *S. titanus* midguts and salivary glands. All of the experiments  
23 154 were carried out using a fluorescent probe specifically targeting group 16SrV phytoplasmas, along  
24 155 with Mollicutes-specific and eubacterial probes. Specifically, the 16SrV phytoplasma-specific  
25 156 probe ph1298, labelled with Cy5 (indodicarbocyanine, absorption/emission at 650/670nm), the  
26 157 Mollicutes probe MCP52, labelled with fluorescein isothiocyanate (FITC, absorption/ emission at  
27 158 494/520 nm), and the eubacterial probe Eub338, labelled with Texas Red (absorption and emission  
28 159 at 595 nm and 620 nm, respectively) were used as described by Gonella *et al.* (2011) and Lessio *et*  
29 160 *al.* (2016).

30  
31  
32 161 In order to perform whole mount FISH, salivary glands were dissected in a sterile saline solutions,  
33 162 then fixed for 2 min at 4°C in 4% paraformaldehyde and washed in PBS. All hybridization  
34 163 experiment steps were performed following Gonella *et al.* (2011). After hybridization, the samples  
35 164 were mounted in antifading medium and then observed in a laser scanning confocal microscope  
36 165 SP2- AOBS (Leica).

37  
38  
39  
40  
41  
42 166

#### 43 44 45 167 Statistical analysis

46  
47 168

48 169 Statistical analyses were performed by mean of SPSS Statistics 24® (IBM Corp. Released 2016,  
49 170 Armonk, NY). A generalised linear model (GLM) was run for analysing infection rates (binomial:  
50 171 positive/negative) and quantitative PCR (qPCR) data, concerning both *S. titanus* adults and plants,  
51 172 as dependent variables, whereas the inoculation access period (IAP) was the categorical variable in  
52 173 both cases, counting three levels (IAP 1, IAP 2, and IAP 3). Concerning qPCR, negative values  
53 174 were excluded from GLM analyses. A binomial distribution with a Logit link function was used for



175 infection rates, whereas a normal distribution and a logarithm link function was chosen for qPCR  
176 data. When IAP effects were significant, Helmert contrasts were performed to identify differences  
177 in infection rates and qPCR data (for both plants and insects), as follows: IAP1 vs. mean (IAP2,  
178 IAP3) and IAP2 vs. IAP3.

179

## 180 **Results**

181

### 182 FDP transmission by adult *S. titanus*

183

184 The acquisition of FDP by adult leafhoppers was performed on a total of eight experimentally  
185 infected BBs. The phytoplasma concentration in those plants was stable, ranging from  $1.46 \times 10^4$  to  
186  $1.60 \times 10^5$  FDP GU / sample, with an average density of  $6.64 \times 10^4$  FDP GU / sample.

187 Quantitative PCR analysis on insects and BBs showed that transmission of FDP did actually occur,  
188 and started from the first week after AAP (IAP1). The percentage of FDP-infected *S. titanus* adults  
189 and their corresponding BBs obtained at the end of each IAP are presented in Table 1, along with  
190 the average concentration of phytoplasma cells found in positive samples. FISH experiments  
191 confirmed the results of qPCR analyses, as at the end of all IAPs, 16SrV phytoplasmas were found  
192 not only in the midgut of leafhoppers (Fig. 2 A-C), but also in salivary glands (Fig. 2 D-F).  
193 Phytoplasma- specific hybridization was not observed in insects directly collected from the mass  
194 rearing, where the eubacterial signal was only detected (Fig. 2 G-I).

195 Considering *S. titanus* individuals, the infection rate reached 70% of the total, whereas infected BBs  
196 were up to 25%. For both insects and plants, infection rates peaked at the end of IAP2, while the  
197 lowest percentages of positive samples were recorded at the end of IAP1 as expected; however,  
198 according to binomial GLM analysis, no significant differences between IAP levels were found (*S.*  
199 *titanus*:  $\chi^2 = 5.359$ ;  $df = 2$ ;  $P = 0.07$ ; BBs:  $\chi^2 = 0.634$ ;  $df = 2$ ;  $P = 0.73$ ). Symptoms (such as  
200 yellowing and curling, especially of apical leaves) were observed on 9 BBs out of 13 which resulted  
201 FD-positive after molecular analyses (69%). Symptomatic plants were evenly distributed among  
202 IAP times (2 plants in IAP 1 and 3, 3 plants in IAP 2). On the other hand, none of the 49 FD-  
203 negative plants have shown any symptom.

204 The average concentration of phytoplasma cells ranged from  $1.30 \times 10^2$  to  $1.40 \times 10^3$  FDP GU /  
205 sample for leafhoppers and from  $5.74 \times 10^0$  to  $5.76 \times 10^2$  FDP GU / sample for inoculated plants, and  
206 in both cases IAP factor was significant (GLM, *S. titanus*:  $\chi^2 = 16.60$ ,  $df = 2$ ;  $P < 0.001$ ; BBs:  $\chi^2 =$   
207  $29.76$ ,  $df = 2$ ;  $P < 0.001$ ). Helmert tests showed significant differences between IAP levels, except  
208 from IAP 1 vs. mean (IAP 2, IAP 3) in broad beans (Table 2).

209

210 **Discussion**

211

212 The results of the present study highlighted that, under laboratory conditions, *S. titanus* is able to  
213 acquire FD-C phytoplasma from infected BB as an adult too, and not only at the nymphal stage, as  
214 previously reported (Caudwell *et al.*, 1970; Bressan *et al.*, 2005b; Galetto *et al.*, 2014). Indeed,  
215 phytoplasmas were successfully acquired after seven days of AAP, blooming to detectable loads  
216 after only seven days of LAP. Additionally, the phytoplasma load in insects increased over a period  
217 of 28 days (7 days AAP + 21 days LAP), indicating that the pathogen multiplies inside the  
218 leafhopper's body, in agreement with a persistent-propagative transmission model (Alma *et al.*,  
219 2015). However, overall phytoplasma concentrations were lower compared to those reported about  
220 other vectors, such as *E. variegatus* and *Macrosteles quadripunctulatus* (Kirschbaum), in similar  
221 experiments (Rashidi *et al.*, 2014; Bosco *et al.*, 2007). Furthermore, the mortality of *S. titanus*  
222 adults, which may be related to experimental conditions, since BB is not the favorite plant host for  
223 *S. titanus* (Chuche *et al.*, 2016), suggests limited chance for adults to reach high levels of  
224 phytoplasma load, at least in laboratory. Under natural conditions, and especially on grapevine,  
225 adults' lifespan is longer, and acquisition of FDP from infected vines may therefore be more likely.  
226 This is in agreement with previous observations: *S. titanus* adults were thought to live up to 40 days  
227 (Vidano, 1964). Yet, recent (unpublished) results show how lifespan of adults is much longer under  
228 semi-natural conditions. If these data were confirmed, adults would be able to complete their  
229 transmission cycle (AAP+LAP+IAP) acquiring and transmitting phytoplasmas to vines, especially  
230 in the last part of the season, when no insecticides are used in vineyards. This is particularly  
231 threatening because of adults incoming from wild grapevine nearby (Lessio *et al.*, 2014): a similar  
232 pattern in FD transmission implies that adults could acquire phytoplasmas directly on infected  
233 grapevines in a vineyard, even if coming from outside. Yet, another issue which needs further  
234 investigation is the influence of FDP on the fitness of *S. titanus* when AAP occurs at the adult stage.  
235 It has been demonstrated that FDP have a negative influence on lifespan and fecundity of *S. titanus*  
236 when AAP occurs at the nymphal stage (Bressan *et al.*, 2005a). If confirmed for adults too, this  
237 could explain the low survival obtained in our experimental conditions.

238 In addition, our results underlined that phytoplasma inoculation may occur after a shorter LAP than  
239 previously believed (Schvester *et al.*, 1969; Caudwell *et al.*, 1970), following an AAP performed  
240 by adults. In effect, transmission to BBs was detected as soon as seven days after the end of AAP,  
241 with no significant differences in transmission efficiency with respect to longer LAPs, in spite of  
242 the higher phytoplasma titer recorded at the last IAP. Since similar percentages of infected BBs

1  
2  
3 243 were found at the end of IAPs 1-3, it can be assumed that the presence of positive leafhoppers at the  
4 244 end of IAP1, although bearing a low titer, is not merely due to transient ingested phytoplasma cells,  
5  
6 245 but more likely to effective colonization of insects, which rapidly become infective. The  
7  
8 246 transmission of a phloem-restricted plant pathogen by adult vectors has been previously  
9  
10 247 demonstrated only for *Diaphorina citri* Kuwayama, vectoring '*Candidatus Liberibacter asiaticus*'.  
11 248 However, diverging results were obtained by different studies: while Pelz-Stelinski *et al.* (2010) and  
12 249 Wu *et al.* (2016) reported successful adult transmission, even though with lower efficiency than in  
13 250 nymphs, Inoue *et al.* (2009) found no adults transmitting the pathogen. Moreover, Pelz-Stelinski *et*  
14 251 *al.* (2010) demonstrated that a one day long IAP is enough for *D. citri* to transmit liberibacter, but  
15 252 with an efficiency lower than 10%. Our results show that *S. titanus* adults are much more efficient  
16 253 in transmitting FDP, especially with short IAP, suggesting that different insect-pathogen dynamics  
17 254 occur for liberibacters and phytoplasmas, causing divergent transmission pathways.

18 255 Temperature may have played a role concerning multiplication of FDP both in insects and in plants.  
19 256 It has been (partially) demonstrated that, under lab conditions, multiplication of FDP in *S. titanus*  
20 257 was faster at low temperatures and CO<sub>2</sub> concentrations, whereas this trend was inverted in plants  
21 258 (Galetto *et al.*, 2011). Our experiments took place at 25 °C, but under field conditions temperatures  
22 259 change along with the day and the season. It is possible therefore that the multiplication of  
23 260 phytoplasmas occur faster in the last part of summer, increasing the threat caused by long-living  
24 261 adults.

25 262 Although successful transmission was observed, the average phytoplasma concentration in infected  
26 263 BB samples was always lower if compared to densities reported about phytoplasmas in wooden  
27 264 plants (Raddadi *et al.*, 2011; Galetto *et al.*, 2016; Jawhari *et al.*, 2015), possibly because of the  
28 265 herbaceous nature of BB. Moreover, higher FDP loads were recorded in BBs experimentally  
29 266 infected by *E. variegatus* (Salar *et al.*, 2013); low phytoplasma concentration found in inoculated  
30 267 plants may be due to the limited number of infected leafhoppers hosted by each plant during our  
31 268 experiments and the reduced pathogen load in inoculating insects.

32 269 Even though it must be pointed out that our results were obtained with BB, where *S. titanus* was  
33 270 reported to acquire the phytoplasma more rapidly than from grapevine in laboratory conditions  
34 271 (Chuche & Thiery, 2014), this work suggests a potential role of *S. titanus* adults in acquiring FDP  
35 272 from infected grapevines in vineyards, and a consequent inoculation to healthy plants within  
36 273 insects' lifespan. This evidence opens to concerning scenarios for viticulture, explaining at least  
37 274 partially the epidemic development of FD which is observed in some cases, in spite of continuous  
38 275 control. Indeed, insecticide sprays are generally limited with respect to the occurrence of adults in  
39 276 the vineyard, which may be prolonged until autumn (Lessio *et al.*, 2011) due to global increase of

1  
2  
3 277 temperatures, therefore the total contribution of adults to FDP transmission can be relevant.  
4 278 Moreover, additional insecticide treatments in late summer, targeting adults, may be difficult to  
5  
6 279 apply due to food safety issues. In addition, the presence of hotbeds with overgrown grapevine's  
7  
8 280 rootstocks is a serious threat for vineyards due to incoming adults during late summer (Lessio *et al.*,  
9  
10 281 2014; Lessio *et al.*, 2015; Pavan *et al.*, 2012).

11 282 The results of this work, by showing that *S. titanus* is able to acquire and transmit FDP at the adult  
12  
13 283 stage with reduced LAP in laboratory, represent the necessary scientific background for a deeper  
14  
15 284 study of interactions involving adult leafhoppers, pathogens and host plants in different agricultural  
16  
17 285 models, including the grapevine-FD pathosystem. Further researches are needed to shed light on  
18  
19 286 such relationships, in order to improve the control of phytoplasma diseases.

20 287

### 21 288 **Acknowledgements**

22 289 The authors are grateful to Federico Bocca and Federico Gottardi for their precious help in field and  
23  
24 290 laboratory activities, and to Dario Sacco for hints and review of statistical aspects of the manuscript.

25 291  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- Alma A., Tedeschi R., Lessio F., Picciau L., Gonella E., Ferracini C. (2015) Insect vectors of plant pathogenic Mollicutes in the European-Mediterranean region. *Phytopathogenic Mollicutes*, **5**, 53-73.
- Arnaud G., Malembic-Maher S., Salar P., Bonnet P., Maixner M., Marcone C., Boudon-Padieu E., Foissac X. (2007) Multilocus sequence typing confirms the close genetic interrelatedness of three distinct flavescence doree phytoplasma strain clusters and group 16SrV phytoplasmas infecting grapevine and alder in Europe. *Applied and Environmental Microbiology*, **73**, 4001-4010.
- Bagnoli B., Angelini E., Borgo M., Ferretti L., Gargani E., Pasquini G. (2011) Valutazione del rischio di diffusione di *Scaphoideus titanus* mediante il materiale di propagazione della vite. *Rivista di viticoltura e di enologia*, **64**, 25-26.
- Bosco D., Galetto L., Leoncini P., Saracco P., Raccach B., Marzachi C. (2007) Interrelationships between "*Candidatus* Phytoplasma asteris" and its leafhopper vectors (Homoptera : Cicadellidae). *Journal of Economic Entomology*, **100**, 1504-1511.
- Bressan A., Girolami V., Boudon-Padieu E. (2005a) Reduced fitness of the leafhopper vector *Scaphoideus titanus* exposed to Flavescence doree phytoplasma. *Entomologia Experimentalis Et Applicata*, **115**, 283-290.
- Bressan A., Spiazzi S., Girolami V., Boudon-Padieu E. (2005b) Acquisition efficiency of Flavescence doree phytoplasma by *Scaphoideus titanus* Ball from infected tolerant or susceptible grapevine cultivars or experimental host plants. *Vitis*, **44**, 143-146.
- Caudwell A., Kuszala C., Bachelier J. C., Larrue J. (1970) Transmission of the golden flavescence of vines to herbaceous plants by the prolongation of the time during which *S. littoralis* can be used and the study of its survival on a large number of plant species. *Annales de Phytopathologie*, **2**, 415-428.
- Caudwell A., Kuszala C., Larrue J., Bachelier J. C. (1972) Transmission de la Flavescence dorée de la fève à la fève par des cicadelles des genres *Euscelis* et *Euscelidius*. Intervention possible de ces insectes dans l'épidémiologie du Bois noir en Bourgogne. *Annales de Phytopathologie*, **Hors série**, 181-189.
- Chuche J., Boudon-Padieu E., Thiéry D. (2016) Host preferences of the leafhopper *Scaphoideus titanus*, vector of "flavescence dorée" phytoplasma. *Phytopathogenic Mollicutes*, **6**, 38-45.
- Chuche J., Thiery D. (2009) Cold winter temperatures condition the egg-hatching dynamics of a grape disease vector. *Naturwissenschaften*, **96**, 827-834.
- Chuche J., Thiery D. (2014) Biology and ecology of the Flavescence doree vector *Scaphoideus titanus*: a review. *Agronomy for Sustainable Development*, **34**, 381-403.
- Duduk B., Bertaccini A. (2009) Phytoplasma and phytoplasma diseases: a review of recent research. *Phytopathologia Mediterranea*, **48**, 355-378.
- Falzo S., Lessio F., Spanna F., Alma A. (2014) Influence of temperature on the embryonic and post-embryonic development of *Scaphoideus titanus* (Hemiptera: Cicadellidae), vector of grapevine Flavescence doree. *International Journal of Pest Management*, **60**, 246-257.
- Filippin L., Jovic J., Cvrkovic T., Forte V., Clair D., Tosevski I., Boudon-Padieu E., Borgo M., Angelini E. (2009) Molecular characteristics of phytoplasmas associated with Flavescence doree in clematis and grapevine and preliminary results on the role of *Dictyophara europaea* as a vector. *Plant Pathology*, **58**, 826-837.
- Galetto L., Bosco D., Marzachi C. (2005) Universal and group-specific real-time PCR diagnosis of flavescence dorée (16Sr-V), bois noir (16Sr-XII) and apple proliferation (16Sr-X) phytoplasmas from field-collected plant hosts and insect vectors. *Annals of Applied Biology*, **147**, 191-201.

- 1  
2  
3 342 Galetto L., Marzachì C., Marques R., Graziano C., Bosco D. (2011) Effects of temperature and CO<sub>2</sub>  
4 343 on phytoplasma multiplication pattern in vector and plant. *Bulletin of Insectology*, **64**, 151-  
5 344 152.
- 6 345 Galetto L., Miliordos D., Pegoraro M., Sacco D., Veratti F., Marzachì C., Bosco D. (2016)  
7 346 Acquisition of Flavescence dorée phytoplasma by *Scaphoideus titanus* Ball from different  
8 347 grapevine varieties. *International Journal of Molecular Sciences*, **17**, 1563.
- 9 348 Galetto L., Miliordos D., Roggia C., Rashidi M., Sacco D., Marzachi C., Bosco D. (2014)  
10 349 Acquisition capability of the grapevine Flavescence doree by the leafhopper vector  
11 350 *Scaphoideus titanus* Ball correlates with phytoplasma titre in the source plant. *Journal of*  
12 351 *Pest Science*, **87**, 671-679.
- 13 352 Gonella E., Negri I., Marzorati M., Mandrioli M., Sacchi L., Pajoro M., Crotti E., Rizzi A.,  
14 353 Clementi E., Tedeschi R., Bandi C., Alma A., Daffonchio D. (2011) Bacterial endosymbiont  
15 354 localization in *Hyalesthes obsoletus*, the insect vector of Bois noir in *Vitis vinifera*. *Applied*  
16 355 *and Environmental Microbiology*, **77**, 1423-1435.
- 17 356 Inoue H., Ohnishi J., Ito T., Tomimura K., Miyata S., Iwanami T., Ashihara W. (2009) Enhanced  
18 357 proliferation and efficient transmission of *Candidatus Liberibacter asiaticus* by adult  
19 358 *Diaphorina citri* after acquisition feeding in the nymphal stage. *Annals of Applied Biology*,  
20 359 **155**, 29-36.
- 21 360 Jawhari M., Abrahamian P., Sater A. A., Sobh H., Tawidian P., Abou-Jawdah Y. (2015) Specific  
22 361 PCR and real-time PCR assays for detection and quantitation of 'Candidatus Phytoplasma  
23 362 phoenicium'. *Molecular and Cellular Probes*, **29**, 63-70.
- 24 363 Lessio F., Alma A. (2013) Influenza dello sviluppo del ritidoma e della termoterapia sulle uova di  
25 364 *Scaphoideus titanus* Ball. *Petria*, **23**, 157-160.
- 26 365 Lessio F., Mondino E. B., Alma A. (2011) Spatial patterns of *Scaphoideus titanus* (Hemiptera:  
27 366 Cicadellidae): a geostatistical and neural network approach. *International Journal of Pest*  
28 367 *Management*, **57**, 205-216.
- 29 368 Lessio F., Picciau L., Gonella E., Mandrioli M., Tota F., Alma A. (2016) The mosaic leafhopper  
30 369 *Orientalus ishidae*: host plants, spatial distribution, infectivity, and transmission of 16SrV  
31 370 phytoplasmas to vines. *Bulletin of Insectology*, **69**, 277-289.
- 32 371 Lessio F., Portaluri A., Paparella F., Alma A. (2015) A mathematical model of flavescence dorée  
33 372 epidemiology. *Ecological Modelling*, **312**, 41-53.
- 34 373 Lessio F., Tedeschi R., Alma A. (2007) Presence of *Scaphoideus titanus* on American grapevine in  
35 374 woodlands, and infection with "flavescence doree" phytoplasmas. *Bulletin of Insectology*,  
36 375 **60**, 373-374.
- 37 376 Lessio F., Tota F., Alma A. (2014) Tracking the dispersion of *Scaphoideus titanus* Ball (Hemiptera:  
38 377 Cicadellidae) from wild to cultivated grapevine: use of a novel mark-capture technique.  
39 378 *Bulletin of Entomological Research*, **104**, 432-443.
- 40 379 Maixner M., Pearson R. C., Boudonpadieu E., Caudwell A. (1993) *Scaphoideus titanus*, a possible  
41 380 vector of grapevine yellows in New York. *Plant Disease*, **77**, 408-413.
- 42 381 Marcone C., Ragozzino A., Schneider B., Lauer U., Smart C. D., Seemuller E. (1996) Genetic  
43 382 characterization and classification of two phytoplasmas associated with spartium witches'-  
44 383 broom disease. *Plant Disease*, **80**, 365-371.
- 45 384 Marzachi C., Bosco D. (2005) Relative quantification of chrysanthemum yellows (16Sr I)  
46 385 phytoplasma in its plant and insect host using real-time polymerase chain reaction.  
47 386 *Molecular Biotechnology*, **30**, 117-127.
- 48 387 Marzachi C., Palermo S., Boarino A., Veratti F., d'Aquilio M., Loria A., Boccardo G. (2001)  
49 388 Optimisation of a one-step PCR assay for the diagnosis of Flavescence dorée related  
50 389 phytoplasmas in field-grown grapevines and vector populations. *Vitis*, **40**, 213-217.
- 51 390 Marzachi C., Veratti F., Bosco D. (1998) Direct PCR detection of phytoplasmas in experimentally  
52 391 infected insects. *Annals of Applied Biology*, **133**, 45-54.

- 1  
2  
3 392 Pavan F., Mori N., Bigot G., Zandigiaco P. (2012) Border effect in spatial distribution of  
4 393 Flavescence dorée affected grapevines and outside source of *Scaphoideus titanus* vectors.  
5 394 *Bulletin of Insectology*, **65**, 281-290.
- 6 395 Pelz-Stelinski K. S., Brlansky R. H., Ebert T. A., Rogers M. E. (2010) Transmission Parameters for  
7 396 *Candidatus Liberibacter asiaticus* by Asian Citrus Psyllid (Hemiptera: Psyllidae). *Journal of*  
8 397 *Economic Entomology*, **103**, 1531-1541.
- 9 398 Raddadi N., Gonella E., Camerota C., Pizzinat A., Tedeschi R., Crotti E., Mandrioli M., Bianco  
10 399 P.A., Daffonchio D., Alma A. (2011) '*Candidatus Liberibacter europaeus*' sp nov that is  
11 400 associated with and transmitted by the psyllid *Cacopsylla pyri* apparently behaves as an  
12 401 endophyte rather than a pathogen. *Environmental Microbiology*, **13**, 414-426.
- 13 402 Rashidi M., D'Amelio R., Galetto L., Marzachi C., Bosco D. (2014) Interactive transmission of two  
14 403 phytoplasmas by the vector insect. *Annals of Applied Biology*, **165**, 404-413.
- 15 404 Roggia C., Caciagli P., Galetto L., Pacifico D., Veratti F., Bosco D., Marzachi C. (2014)  
16 405 Flavescence doree phytoplasma titre in field-infected Barbera and Nebbiolo grapevines.  
17 406 *Plant Pathology*, **63**, 31-41.
- 18 407 Salar P., Charenton C., Foissac X., Malembic-Maher S. (2013) Multiplication kinetics of  
19 408 Flavescence dorée phytoplasma in broad bean. Effect of phytoplasma strain and  
20 409 temperature. *European Journal of Plant Pathology*, **135**, 371-381.
- 21 410 Schneider B., Seemuller E. (1994) Presence of 2 sets of ribosomal genes in phytopathogenic  
22 411 Mollicutes. *Applied and Environmental Microbiology*, **60**, 3409-3412.
- 23 412 Schvester D., Carle P., Moutous G. (1961) Transmission de la Flavescence dorée de la vigne par  
24 413 *Scaphoideus littoralis* Ball, (Homopt. Jassidae). *Annales Epiphyties*, **14**, 175-198.
- 25 414 Schvester D., Carle P., Moutous G. (1969) Nouvelles données sur la transmission de la Flavescence  
26 415 dorée de la vigne par *Scaphoideus littoralis* Ball. *Annale de Zoologie et écologie animale*, **1**,  
27 416 445-465.
- 28 417 Vidano C. (1964) Scoperta in Italia dello *Scaphoideus littoralis* Ball, cicalina americana collegata  
29 418 alla *Flavescence dorée* della vite. *L'Italia Agricola*, **10**.
- 30 419 Vidano C. (1966) Scoperta della ecologia ampelofila del cicadellide *Scaphoideus littoralis* Ball  
31 420 nella regione neartica originaria. *Annali della Facoltà di Scienze Agrarie della Università*  
32 421 *degli Studi di Torino*, **3**.
- 33 422 Wu T. Y., Luo X. Z., Xu C. B., Wu F. N., Qureshi J. A., Cen Y. J. (2016) Feeding behavior of  
34 423 *Diaphorina citri* and its transmission of '*Candidatus Liberibacter asiaticus*' to citrus.  
35 424 *Entomologia Experimentalis Et Applicata*, **161**, 104-111.  
36 425  
37  
38  
39  
40 426  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 427 **Table 1.** Results of inoculation trials of FDP to broad beans (BB) by means of *S. titanus* adults.  
4 428 Data of qPCR analyses on leafhoppers ("*S. titanus*" columns) and inoculated plants ("BB" columns)  
5 are indicated.  
6  
7

8 430

9 431 **Table 2.** Results of Helmert contrasts between levels of the categorical variable "Inoculation  
10 Access Period (IAP)", for the dependent variable "quantitative PCR (qPCR)" concerning both *S.*  
11 432 *titanus* (ST) and broad beans (BB).  
12 433

13 434

14 435 **Figure 1.** Experimental design of FDP transmission trials. Adult *S. titanus* [25 batches with 125  
15 insects for qPCR analysis (A) and three batches with 15 insects for FISH analysis (B)] were caged  
16 436 for phytoplasma acquisition at day 0, and then from day 7 they were maintained on BB seedlings in  
17 437 groups of five. Every seven days live specimens were moved onto a new plant to perform up to  
18 438 three IAPs: 14 days (IAP1), 21 days (IAP2), and 28 days (IAP3) after the beginning of AAP,  
19 439 respectively.  
20 440  
21 441

22 442

23 443 **Figure 2.** FISH on *S. titanus* midgut and salivary glands. Exemplificative micrographs showing  
24 444 results from FISH of: midgut of a specimen at the end of IAP2 (A-C), salivary glands of an adult at  
25 445 the end of IAP3 (D-F), and midgut of a mass-reared individual (negative control) (G-I). A, D and G  
26 446 show interferential contrast micrographs of the organs. Hybridizations with Mollicutes probe (green  
27 447 signal in B) and eubacterial probe (green signal in E and H) are presented; moreover, hybridization  
28 448 with 16SrV phytoplasma probe is shown in C (cyan signal), F (red signal), and I. Bars = 75  $\mu$ m.  
29 449 The phytoplasmas were effectively detected in the organs of leafhoppers after transmission  
30 450 experiments, while they were absent in the negative control.  
31 451

32 452

33 453

34 454

35 455

36 456

37 457

38 458

39 459

40 460

41 461

42 462

43 463

44 464

45 465

46 466

47 467

48 468

49 469

50 470




Rank	<i>S. titanus</i>				BB			
	N	Positive samples	Infection rate <sup>1</sup> ± SE	Titer <sup>2</sup> ± SE	N	Positive samples	Infection rate <sup>1</sup> ± SE	Titer ± SE
IAP1	65	30	0.46 ± 0.06	1.30×10 <sup>2</sup> ± 3.99×10 <sup>1</sup>	25	4	0.16 ± 0.07	6.57×10 <sup>1</sup> ± 5.43×10 <sup>1</sup>
IAP2	38	26	0.68 ± 0.08	5.05×10 <sup>2</sup> ± 1.39×10 <sup>2</sup>	24	6	0.25 ± 0.09	5.74×10 <sup>0</sup> ± 1.33×10 <sup>0</sup>
IAP3	22	14	0.64 ± 0.10	1.40×10 <sup>3</sup> ± 6.42×10 <sup>2</sup>	13	3	0.23 ± 0.12	5.76×10 <sup>2</sup> ± 1.98×10 <sup>2</sup>

<sup>1</sup> Rate of 16SrV phytoplasma-positive individuals related to the total tested samples. <sup>2</sup> 16SrV phytoplasma GU per sample (single insect or 100 mg of plant tissue). Values below the detection limit (1.00×10<sup>0</sup> GU /sample) were considered negative (cut-off value). SE: Standard Error.

	IAP 1 vs. mean (IAP 2, IAP 3)			IAP 2 vs. IAP 3		
	$\chi^2$	df	P	$\chi^2$	df	P
ST qPCR	12.957	1	0.00	6.736	1	0.01
BB qPCR	2.078	1	0.15	29.597	1	0.00


For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Experimental design of FDP transmission trials. Adult *S. titanus* [25 batches with 125 insects for qPCR analysis (A) and three batches with 15 insects for FISH analysis (B)] were caged for phytoplasma acquisition at day 0, and then from day 7 they were maintained on BB seedlings in groups of five. Every seven days live specimens were moved onto a new plant to perform up to three IAPs: 14 days (IAP1), 21 days (IAP2), and 28 days (IAP3) after the beginning of AAP, respectively.

209x180mm (300 x 300 DPI)



FISH on *S. titanus* midgut and salivary glands. Exemplificative micrographs showing results from FISH of: midgut of a specimen at the end of IAP2 (A-C), salivary glands of an adult at the end of IAP3 (D-F), and midgut of a mass-reared individual (negative control) (G-I). A, D and G show interferential contrast micrographs of the organs. Hybridizations with Mollicutes probe (green signal in B and E) and eubacterial probe (green signal in H) are presented; moreover, hybridization with 16SrV phytoplasma probe is shown in C (cyan signal), F (red signal), and I. Bars = 75 $\mu$ m. The phytoplasmas were effectively detected in the organs of leafhoppers after transmission experiments, while they were absent in the negative control.

180x181mm (300 x 300 DPI)