



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Phospho-proteomic analysis of mantle cell lymphoma cells suggests a pro-survival role of B-cell receptor signaling.

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1656541 since 2018-07-10T13:10:04Z
Published version:
DOI:10.1007/s13402-011-0019-7
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Phospho-proteomic analysis of mantle cell lymphoma cells suggests a pro-survival role of B-cell receptor signaling

Chiara Pighi, Ting-Lei Gu, [...], and Alberto Zamò

Abstract

Background

Mantle cell lymphoma (MCL) is currently an incurable entity, and new therapeutic approaches are needed. We have applied a high-throughput phospho-proteomic technique to MCL cell lines to identify activated pathways and we have then validated our data in both cell lines and tumor tissues.

Methods

PhosphoScan analysis was performed on MCL cell lines. Results were validated by flow cytometry and western blotting. Functional validation was performed by blocking the most active pathway in MCL cell lines.

Results

PhosphoScan identified more than 300 tyrosine-phosporylated proteins, among which many protein kinases. The most abundant peptides belonged to proteins connected with B-cell receptor (BCR) signaling. Active BCR signaling was demonstrated by flow cytometry in MCL cells and by western blotting in MCL tumor tissues. Blocking BCR signaling by Syk inhibitor piceatannol induced dose/time-dependent apoptosis in MCL cell lines, as well as several modifications in the phosphorylation status of BCR pathway members and a collapse of cyclin D1 protein levels.

Conclusion

Our data support a pro-survival role of BCR signaling in MCL and suggest that this pathway might be a candidate for therapy. Our findings also suggest that Syk activation patterns might be different in MCL compared to other lymphoma subtypes.

Electronic supplementary material

The online version of this article (doi:10.1007/s13402-011-0019-7) contains supplementary material, which is available to authorized users.

Keywords: Mantle cell lymphoma, Proteomics, PhosphoScan, Syk, Piceatannol, BCR signaling

Introduction

Mantle cell lymphoma constitutes about 6% of non-Hodgkin lymphoma in the western world [1]. It shows a characteristic clinical behavior with initial response to therapy followed by relapses and death in 3–5 years. Aggressive therapeutic regimens are promising, but to-date MCL is still considered incurable, with the exception of a small group of long-surviving patients. In recent years, many high-throughput techniques have been applied to MCL in order to gain insights into its pathogenesis, and to discover suitable therapeutic targets [2–9]. These studies have clarified a role for increased proliferation and resistance to apoptosis in MCL, involving several signaling pathways.

A few studies from our group and also from others have attempted the study of MCL from a proteomic point of view [10–18]. Some of these studies were mainly focused on the diagnostic implementation of proteomic techniques [14–16], while other were more focused on the pathogenetic mechanisms [12, 18–20]. These studies have identified single up-regulated proteins in MCL compared to tonsil B-cells [12], or several proteins up-regulated in MCL cell lines compared to FL-derived cell lines followed by network analysis [21]. More recently, a proteomic study focused on the plasma membrane proteome of leukemic MCL and compared it to normal B-cells [19]; this study identified several alterations in the proteome of MCL lipid rafts that might have therapeutic potential. In a previous study we have focused our attention on the phospho-proteome of several MCL cell lines by using immobilized metal affinity chromatography (IMAC) pre-fractionation (a column-based technique that selectively enriches

all phospho-proteins by a reversible binding to metal oxydes), followed by 2D-PAGE/MS [20]. In that study we identified several activated pathways that might reflect the biology of MCL, including the alternative NF-kB pathway, the mTOR pathway and the mitochondrial pathway. In the present work we have used a complementary approach, called PhosphoScan, that involves an immunoprecipitation of peptides by using an anti-phosphotyrosine antibody followed by capillary electrophoresis and MS identification of eluted peptides [22]. Compared to IMAC, PhosphoScan should better dissect phosphorylation profiles, specifically identifying tyrosine-phosphorylated peptides. This approach has already been successfully used in the phosphoprofiling of primary [23] and metastatic [24] lung cancer, in acute leukemia samples [25], in Bcr/Abl positive cell lines [26], AML cell lines [27] and Hodgkin lymphoma cell lines [28]. Since tyrosine phosphorylation regulates the activity of many proteins, the PhosphoScan approach can provide a picture of the most active pathways in the analyzed samples. Moreover, since many kinases are themselves activated by tyrosine phosphorylation, the profile of active kinases is also provided; often these kinases are driving cell proliferation and survival, and are therefore candidate targets for therapy.

In the current work, we used this approach to recognize the most represented activated pathways in MCL cell lines. We then verified the presence of these activated molecules in MCL cell lines and tumor tissues. We subsequently explored the functional role of the most activated pathway in MCL cell lines by inhibition experiments.

Results and discussion

In the current paper we describe the results obtained from the application of PhosphoScan to MCL cells. This is to our knowledge the first use of this technique in MCL; we report a large amount of data with precise identification of phosphorylation sites for each protein, some of which novel.

PhosphoScan analysis identifies the tyrosine-activated phosphoproteome of MCL cell lines and identifies B-cell receptor signaling as the most active pathway

PhosphoScan analysis identified 421 unique tyrosine-phosphorylated peptides, corresponding to 341 proteins, ranked by overall abundance in Table 1 (for a complete list of peptides and complete MS information see Supplemental Table 1). Interestingly, several identified proteins mapped to cytogenetic loci that have been reported to be altered in MCL [29–31]. Among these, several activated protein kinases were identified (Fig. 1), that are therefore appealing as potential therapeutic targets.

Table 1 Top 02 phototo papidos identified (surfue of phototo papidos correspondentio Resignable protein is reported. For complete let and exact shorthorylation stims, sa							
Public Public	Granta-	Aks-	MAVER	Rec.	BCR	8,710	
Cel	17	38	-14	24			
0082	*0	34	14				
0043	10	35	- 10	17			
Dut.	9	19	24	3			
24	7	18	18	50			
PAD		54					
1994		15		2		×.	
Lat		10			×.		
lobela	0	22	2				
1000					*		

Table 1

Top 50 phospho-peptides identified (number of phospho peptides corresponding to the specific protein is reported. For complete list and exact phosphorylation sites, see suppl. Table 1)



Fig. 1

List of protein kinases identified by the PhosphoScan approach, ranked by overall abundance in the four MCL cell lines analyzed. Color intensity proportional to the number of peptides (see figure for color corresponding to the average number of peptides ...

The three most represented phospho-peptides are cyclin-dependent kinases Cdc2, CDK2 and CDK3. Cdc2 (CDK1) is actually part of the proliferation signature able to predict prognosis in MCL [8] and has also been validated at the immunohistochemical level [32]. *CDK2* gains have been detected in MCL in a variable proportion of cases [30, 31, 33], and have been associated with a poor prognosis [33]. *CDK3* is less studied compared to other cyclin-dependent kinases, but its locus has also been reported as subject to gains [31, 34]. Its involvement in MCL is a novel finding, although the role of *CDK3* as an oncogene has been demonstrated in other models [35, 36].

Syk and many other BCR pathway members were then found. Many kinases belonging to this pathway (e.g. Syk, Lyn, Hck, Btk, PKC-delta), were among the most represented phospho-peptides identified by the PhosphoScan approach, and this was confirmed when looking for non-random enrichment of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways [37] compared to the hypothesis of random distribution. This analysis showed that (using a cutoff value of more than five total

peptides), the most enriched pathway was hsa04662:B cell receptor signaling pathway (17.16-fold enrichment) (for complete data see Supplemental Table 2).

Many of the proteins belonging to the most enriched pathways were either present in different pathways (redundant) or are functionally connected. Manual annotation by literature search revealed that several of these, in the topmost part of the ranking, were connected to BCR signaling, even if not present in the canonical KEGG BCR pathway (Table 1 and Fig. 2).



Fig. 2

Simplified diagram showing some of the identified BCR pathway members. In red, activating members; in green, inhibiting members; for some members the precise final effect is not clear; in grey, other proteins known to be present but not identified. Full ...

The presence of active B-cell receptor signaling is validated in MCL cell lines and tissues

To further verify the functional role of BCR signaling in MCL, we analyzed the phosphorylation status of Syk, Lyn, Btk, Blnk and GSK3alfa/beta in MCL cell lines. Although not present in the top 50 phospho-peptides, Blnk was investigated because it is considered a classical downstream molecule of the BCR pathway. The presence of basal levels of phospho-Syk Y525 and Y323, as well as of phospho-Blnk (Y84) was verified by flow cytometry (Fig. 3). By this technique, we could detect no basal levels of phospho-Syk Y352. In addition basal levels of phospho-Lyn (Y396 and Y507) and also of downstream effectors phospho-Btk (S180) and phospho-GSK3alfa/beta (S9/21) were demonstrated by flow cytometry (Supplemental Figure 1). BCR pathway activation in cell lines is somehow intriguing since it is present in absence of an appropriate antigen stimulation, and is therefore probably self-sustained by tumor cells, either by side-by-side activation or by auto-activation.



Analysis of basal levels of phosphorylated Syk and Blnk residues by flow cytometry. In grey, isotypic control; in red, basal levels. On the X axis, arbitrary fluorescence units (log scale); on the Y axis, cell count

In order to verify whether we could find this activation in MCL tumors as well, we resorted to western blotting analysis of phosphorylated forms of BCR pathway members. This analysis showed that the activated forms of Syk (in 5/6 cases, 83%), Lyn (in 6/6 cases, 100%), and Blnk (in 6/6 cases, 100%) were present also in MCL tumor tissues (Fig. 4), therefore supporting the in vivo role of active BCR signaling; as far as we know, this is the first report of the presence of active (phosphorylated) BCR pathway members in MCL tissues.



Fig. 4

Western blotting analysis of MCL tissues. The presence of phospho-Syk (Y525), phospho-Lyn (Y396) and phospho-Blnk (Y84) is shown in six MCL tumor tissues. Cases 1, 2, 5 and 6 were classical MCL, while cases 3 and 4 were blastoid variants

The activation of the BCR pathway in MCL has been hypothesized in a previous paper based on cytogenetic and RNA studies [6], but to our knowledge this is the first protein-based and data-driven study that supports this hypothesis. Another proteomic study focusing only on the plasma membrane [19] showed an abnormal association of PKCbeta to the cell membrane in MCL leukemic cells, indirectly supporting an active BCR signaling.

Recent studies have shown the importance of tonic BCR signaling in DLBCL [38, 39] and B-CLL [40], with a basal activation of phospho-Syk residue Y352, while Y525 was detected only after BCR cross-linking. The presence of significant basal levels of phospho-Syk Y525 and Y323, with no detectable phospho-Syk Y352 in basal conditions in MCL cells are not concordant with what has been reported in B-CLL and DLBCL [40], and suggest a different pattern of activation of BCR signaling in MCL.

A recent report of a phase 1/2 clinical trial of fostamatinib disodium, the first clinically available oral Syk inhibitor, in patients with recurrent B-cell non-Hodgkin lymphoma, showed that only 1 in 9 MCL showed some response [41]. Several explanations might be possible for this low response rate. First, the specificity of this drug for Syk has been recently questioned [39]. Second, relapsed lymphomas might have evolved into BCR-independent clones (such as the cell line Rec-1). Third, since our data support the hypothesis that the activation pattern of Syk in MCL is different from B-CLL and DLBCL, it is possible that this phenomenon influences the response to fostamatinib.

Inhibition of Syk induces apoptosis in MCL cell lines

Since the proteins belonging to the BCR signaling pathway were shown to be active, we tested the effect of the blockade of this pathway on MCL cells. For this purpose, Syk activity was inhibited by a widely used inhibitor, piceatannol [42–45], a natural stilbene also resulting from the hepatic metabolism of resveratrol, a compound found to be pro-apoptotic for the MCL cell line Jeko-1 [11]. The Rec-1 cell line showed a marked hypersensitivity to DMSO, and it was deemed unfit for further experimental work, which was conducted on less sensitive cell lines.

Piceatannol induced apoptosis in MCL cell lines Jeko-1, UPN-1, Mino, Maver-1 and Granta-519, as evidenced by Annexin V staining, with a LD50 varying from 9.50 μ M to 28.91 μ M at 24 h and 5.51 μ M and 23.54 μ M at 48 h (Supplemental Table 3 and Fig. 5).



Fig. 5

Induction of apoptosis in MCL cell lines. Upper panel shows the percentage of live cells (Y axis) in function of the piceatannol concentration (X axis) at 24 h. Lower panel shows the same variables at 48 h of treatment. Annexin V staining ...

Interestingly, Jeko-1, which is very sensitive to piceatannol, also shows the highest number of phospho-peptides belonging to the BCR pathway (Supplemental Table 4).

Apoptosis was dose and time dependent being visible after 4 h of treatment in Jeko-1, but only after 16 h in MAVER-1 (data not shown).

Inhibition of Syk down-regulates Cyclin D1 protein levels

Cyclin D1 protein levels were massively down-regulated in treated cells, as densitometric analysis showed a drop by 85 to 90% compared to untreated cells (Supplemental Figure 2). The corresponding *CCND1* mRNA was down-regulated after piceatannol treatment, showing 22% to 71% decrease (after normalization) in five tested cell lines (Supplemental Figure 3). This decrease was however significant only in three cell lines, namely UPN-1 (54% decrease; p = 0.047), Granta-519 (71% decrease; p = 0.0078) and MAVER-1 (69% decrease; p = 0.008).

This finding is interesting, considering that the *CCND1* gene is translocated under the control of a strong enhancer such as the Igµ enhancer. A hypothetical scenario could be hypothesized in which BCR signaling and *CCND1* gene transcription are connected, possibly *via* Syk. Cyclin D1 has been shown to be a Syk target gene in breast cancer and other cells [46]. It is possible that *CCND1* gene transcription is directly repressed by Syk, in accordance to the fact that normal B cells show very low levels of this protein and at the same time nuclear localization of Syk. As previously stated, in support of this hypothesis Syk acts as a transcriptional repressor of the *CCND1* gene in breast carcinoma cells, and this activity is necessary for its tumor-suppressor function [46]. One hypothesis is that in MCL cells Syk might act *via* Stat3, a known *CCND1* gene transcription inducer, as suggested by a recent publication [47]. In support of this hypothesis, the phosphorylation of Stat3 was reduced by piceatannol in MCL cell lines (Supplemental Figure 4), while total Stat3 was reduced in Jeko-1 and UPN-1, but not significantly in Maver-1 and Granta-519. According to these data, while the phosphorylation of Stat3 was reduced in all cell lines, the total level is reduced only in sensitive ones. Several publications support a role for Stat3 in the pathogenesis of MCL [48–52].

According to our data the down-regulation of *CCND1* mRNA cannot explain the whole picture, since it was significant only in three cell lines after normalization using an unrelated transcript. An increase in Cyclin D1 degradation is another possible explanation, and might operate alongside transcription repression. In support of this complementary hypothesis is the reduction

of GSK3 phosphorylation on negative regulatory serine residues, which increases its kinase activity with subsequent phosphorylation and degradation of Cyclin D1 [53].

SYK mRNA was significantly down-regulated by piceatannol treatment in UPN-1 and MAVER-1 (59% and 75% reduction; p = 0.047 and p = 0.01 respectively). It was also reduced in Granta-519, but the values did not reach statistical significance (49% reduction; p = 0.11). *SYK* mRNA was increased after treatment in Jeko-1 and Mino (by 7% and 40% respectively) although values were not statistically significant (p = 0.8 and p = 0.32 respectively). Modifications of mRNA levels of *SYK* and *CCND1* showed a good correlation with an r = 0.88 (Supplemental Figure 3).

Piceatannol-induced apoptosis involves the activation of the intrinsic pathway, and is accompanied by the modification of several regulators of cell cycle and apoptosis

Apoptosis was triggered trough the intrinsic pathway, involving Caspase 9 cleavage, as shown by western blotting experiments. Cleaved Caspase 8 levels were actually lower in the treated samples than in the untreated ones (not shown). Several apoptosis (Bcl-2, Bcl-xL, Bax, p53, p21), cell-cycle related (p53, p21, p27) molecules were investigated by western blotting. All investigated proteins showed the same behavior in all cell lines with the exception of Bax (Supplemental Figure 5). In detail, p53 levels were increased after treatment (Jeko-1 resulted completely p53-negative due to the presence of a truncating mutation, while other cell lines with the exception of Granta-519 bear a missense mutation [54]); p21 was strongly down-regulated; p27 was cleaved with a decrease of the 27 kDa isoform and increase of the 23KDa isoform; Bcl-2 was slightly increased, Bcl-xL was decreased, while Bax was increased in Jeko-1 and decreased in other cell lines. The mechanism inducing the increase of Bax (a pro-apoptotic Bcl-2 family member) in Jeko-1 is not clear, but this phenomenon might indicate a more pronounced perturbation of mitochondrial outer membrane potential [55]. The only clear-cut genomic abnormality that might suggest a stronger Syk-dependance of Jeko-1 is the presence of the amplification of the *SYK* locus [6], although this cell line also bears a peculiar missense mutation of *TP53* (in addition to a deletion of the other allele) that causes a premature stop codon with degradation of its mRNA [5], and at variance with other cell lines shows no p53 at all; this condition might cause an abnormal regulation of *BAX*, that is a known p53 target gene [56].

Piceatannol induces complex modifications of the Syk phosphorylation pattern and alterations of downstream effectors of the B-cell receptor pathway

To verify the effects of piceatannol on the pattern of phosphorylation of Syk and other BCR pathway members, we resorted to flow cytometry using phospho-specific antibodies. These experiments showed that piceatannol induced a marked decrease in phospho-Syk tyrosine residues Y525 and Y323, while unexpectedly the levels of Y352, which were almost negative in basal conditions, increased after treatment (Fig. 6). This residue resides in the linker region of Syk, in a stretch that is essential for nuclear localization, since a mutant lacking residues 332–359 was unable to enter the nucleus, as reported by a previous article [57]. These modifications were detectable after 6 h of treatment, but not after two (data not shown). As previously stated, B-CLL and DLBCL cells show a basal activation of Syk Y352, but no information is available for comparison about how this phosphorylation might be changed by Syk inhibition.



Fig. 6

Modification of Syk phosphorylation profile following piceatannol treatment. After treatment, phosphorylation of residues Y525 and Y323 is reduced, while that of residue Y352 is increased. *Red*, untreated cells; *blue*, treated cells; *solid grey*, isotypic ...

The levels of the downstream effectors phospho-Blnk and phospho-Btk were also reduced by piceatannol treatment in Jeko-1 and to a lesser degree in MAVER-1 cells (Supplemental Figure 6).

To obtain independent information about the modifications of phospho-Syk after treatment, we used immunofluorescence microscopy. These experiments showed that in basal conditions phospho-Syk (Y525) is present in all cell lines tested, and is apparently confined to the cytoplasmic compartment of MCL cell lines (with the exception of Granta-519, that showed also a partial nuclear localization). After piceatannol treatment phospho-Syk (Y525) was down-regulated in the cytoplasm, but its presence could be demonstrated in the nucleus of MCL cell lines (Jeko-1, UPN-1, Mino, Granta-519, MAVER-1) (Fig. 7). This

finding is in accordance with the reduction of phospho-Syk (Y525) detected by antibodies used in flow cytometry experiments, which were able to enter the cytoplasm but not the nucleus in our experimental conditions.



Fig. 7

Localization of phospho-Syk following piceatannol treatment. P-Syk Y525 (shown in *green* pseudo-color) disappears from the cytoplasm and appears in the nucleus after treatment in all cell lines. Nuclei counter stained in *blue* pseudo-color (DAPI staining) ...

One possible hypothesis to explain this phenomenon is the fact that Syk has two known splice variants, of which only the longest is able to enter the nucleus [58]. Western blotting experiments in the five MCL cell lines Jeko-1, Granta-519, MAVER-1, UPN-1 and Rec-1 showed that only one band was identifiable in the expected molecular weight range. A proteolytic 36 kDa isoform of Syk was also detected (data not shown), although the significance of this isoform described in erythrocytes [59], is still not clear in our cells, because it might be either a true functional variant or an experimental artifact. The absence of the shortest splice variant of Syk was also demonstrated by isoform-specific *SYK* RT-PCR experiments (Supplemental Figure 7). This was confirmed using two amplicons of different length as previously described [58]. The nuclear exclusion phenomenon therefore is probably not related to Syk differential splicing, but more probably to functional modifications of the protein, such as phosphorylation.

Conclusions

Our study has identified a large number of phosphorylated proteins and several activated pathways in MCL cells, which deserve a more thorough investigation by functional validation experiments. Our experimental data suggest that active BCR signaling is present in MCL cell lines and tumors, and that it plays a role in the survival of MCL cells; Syk is possibly one of the key molecules in this event, and it might have a double-face role in the cytoplasmic and nuclear compartments.

Our data also suggest that piceatannol, resveratrol or their analogues could represent a therapeutic option for patients with MCL if in vitro data will be confirmed in vivo.

Methods

Cell lines MCL cell lines Jeko-1, Granta-519 and Rec-1 were purchased from DSMZ (Braunschweig, Germany). Cell line MAVER-1 was developed in our laboratory [60]. Cell line Mino was obtained from ATCC (Manassas, VA, USA). UPN-1 cells were kindly provided by Dr. Elias Campo (Barcelona, Spain). Due to over-sensitivity to DMSO-induced apoptosis, Rec-1 cells were not used for apoptosis-induction assays.

Tumor samples Tumor samples were collected from the archives of the Department of Pathology and Diagnostics of the University of Verona. Informed consent had been collected for all patients and procedures were carried out according to the ethical guidelines of the University Hospital of Verona with the approval of an ethics committee, in compliance with the Helsinky Declaration.

PhosphoScan analysis PhosphoScan analysis was performed as previously described [23] on MCL cell lines MAVER-1, Granta-519, Jeko-1, and Rec-1.The phospho-proteins identified using this analysis were categorized into KEGG Pathways using the DAVID Ease web framework [61]. For P-value and fold enrichment calculation the complete list of human genes was considered as a background. The calculation was repeated for lists obtained by selecting genes above the abundance cut-offs of 3, 4 and 5.Illustrations were produced using the R statistical software package (http://www.r-project.org).

Flow cytometry Flow cytometric analysis of several tyrosine-phosphorylated forms of Syk (Y323, Y352 and Y525) and downstream target phospho-BLNK was performed as previously described [38] using antibodies shown in Table 2. All experiments were performed in triplicate.



RNA isolation and cDNA synthesis RNA isolation was performed using the Allprep DNA/RNA/Protein Mini Kit (Qiagen GmbH, Hilden, Germany), and RNA quality was assessed by using an Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA) and RNA 6000 Nano chips (Agilent Technologies). Each cDNA was synthesized from 1 µg total RNA using random primers and the Superscript III First-strand synthesis system (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions.

RT-PCR The analysis of *SYK* isoforms was performed as previously described [58]. Reaction products were analyzed by the Agilent 2100 Bioanalyzer and the DNA1000 Chip (Agilent Technologies).

Quantitative RT-PCR Quantitative RT-PCR mRNA expression analysis was performed on ABI PRISM 7900HT Fast Real-time PCR System (Applied Biosystems, Foster City, CA) using Power SYBR green PCR Master Mix (Applied Biosystems). Oligonucleotide primers used were: CCND1-F AACTACCTGGACCGCTTCCT, and CCND1-R GGGGATGGTCTCCTTCATCT; TBP-F, GCACAGGAGCCAAGAGTGAA, and TBP-R,

TCACAGCTCCCCACCATATT.TATA box binding protein (RefSeq ID NM_003194.3) transcript level was used to normalize *SYK* expression.Calibration curves for each couple of primers were obtained by serial dilution of cDNA. Expression data were analyzed by the comparative threshold cycle (Ct) method accordingly to User Bulletin No. 2 (Applied Biosystems). All experiments were performed in triplicate. The statistical significance of the data was investigated by Student's *t*-test. All P values were two sided and considered significant when less than 0.05.

Syk inhibition experiments Piceatannol (Sigma-Aldrich Chemie GmBH, Steinheim, Germany) was resuspendend in DMSO as a 100 mM stock. Cells were diluted at $500*10^3$ /ml in RPMI 1640 (1% fetal calf serum) and let grow for 16 h. Piceatannol was then added at concentrations between 1 μ M and 80 μ M (depending on the sensitivity of cell lines). Cells were harvested at 24 and 48 h and apoptosis levels analyzed by Annexin V (Annexin V–FITC apoptosis detection kit I, BD) as previously described [11]. All experiments were performed in triplicate.

Western blotting Immunoblotting was performed by standard methodology, using the antibodies and dilutions indicated Table 2. Protein content of samples was measured using a colorimetric method (DC protein assay, Bio-Rad). All lanes were loaded with the same amount of total protein (in duplicate or triplicate), and were also visually verified by Ponceau red staining after electroblotting.

Immunofluorescence Immunofluorescence was performed using the antibodies shown in Table 2 (both primary and secondary antibodies were diluted 1:100). Briefly, cells were washed in PBS, deposited on charged slides by gravity, fixed in cold methanol (-20°C) for 20 min and then air-dried. Cells were then re-hydrated, incubated with protein blocking solution for 10', then incubated serially with primary and secondary antibodies (with three washes in PBS in-between). Slides were examined under fluorescent light using an Olympus BX61 microscope. Images were acquired using appropriate filters via a monochromatic camera cooled CCD camera (iAi, Japan) and analyzed using the Olympus Cytovision software.

Electronic supplementary material

Below is the link to the electronic supplementary material.

Supplemental Figure 1(424K, jpg)

Basal levels of phospho-Lyn (Y396 and Y507), phospho-Btk (S180) and phospho-GSK3alfa/beta (S9/21) as analyzed by flow cytometry in MCL cells. In grey, isotypic control; in red, basal levels. On the X axis, arbitrary fluorescence units (log scale); on the Y axis, cell count. (JPEG 423 kb)

Supplemental Figure 2(97K, jpg)

Western blotting analysis of Cyclin D1 protein levels in piceatannol-treated cells. (JPEG 96 kb)

Supplemental Figure 3(537K, jpg)

Modifications of *CCND1* and *SYK* mRNA after piceatannol treatment (normalized). A decrease of one $\Delta\Delta$ Ct unit corresponds to a 50% reduction of RNA levels. A correlation plot between modifications of mRNA levels is also shown. (JPEG 536 kb)

Supplemental Figure 4(571K, jpg)

Western blotting analysis of modifications of levels of phospho-Stat3 (Y705) and Stat3 after piceatannol treatment in cell lines Jeko-1, UPN-1, Maver-1 and Granta-519. (JPEG 571 kb)

Supplemental Figure 5(963K, jpg)

Western blotting analysis of apoptosis-related and cell cycle-related molecules after exposure to piceatannol by western blotting. UPN-1 was totally BCL2-negative, while Jeko-1 showed no p53 protein due to a truncating mutation. (JPEG 963 kb)

Supplemental Figure 6(355K, jpg)

Levels of phospho-Blnk and phospho-Btk after piceatannol treatment in cell lines Jeko-1 and MAVER-1 analyzed by flow cytometry. In grey, isotypic control; in red, basal levels, in blue, treated cells. On the X axis, arbitrary fluorescence units (log scale); on the Y axis, cell count. (JPEG 354 kb)

Supplemental Figure 7(348K, jpg)

Isoform-specific *SYK* RT-PCR experiments. Pseudo-gel image (upper panel) and two representative electropherograms (lower panel) of RT-PCR products are shown (images produced by microfluidic analysis). The absence of the shortest splice variant of Syk was demonstrated. (JPEG 348 kb)

Supplemental Table 1(488K, xls)

Complete list of peptides and complete MS information. (XLS 487 kb)

Supplemental Table 2(28K, xls)

List of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways enriched in MCL cell lines compared to random distribution. (XLS 28 kb)

Supplemental Table 3(6.0K, xls)

LD50 of piceatannol at 24 and 48 h for MCL cell lines. (XLS 6 kb)

Supplemental Table 4(10K, xls)

Number of identified phospho-peptides corresponding to members of the KEGG BCR pathway in the four tested MCL cell lines.. (XLS 10 kb)

Acknowledgements

Financially supported by AIRC MFAG grant to Alberto Zamò.

We would like to thank Roberto Polackiewitz from Cell Signaling Technology for helpful discussion.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Article information

Cell Oncol (Dordr). 2011 Apr; 34(2): 141–153. Published online 2011 Mar 11. doi: 10.1007/s13402-011-0019-7 PMCID: PMC3063577 PMID: 21394647

Chiara Pighi,¹ Ting-Lei Gu,² Irene Dalai,¹ Stefano Barbi,¹ Claudia Parolini,¹ Anna Bertolaso,¹ Serena Pedron,¹ Alice Parisi,¹ Jianmin Ren,² Daniela Cecconi,³ Marco Chilosi,¹ Fabio Menestrina,¹ and Alberto Zamo^[3]

¹Department of Pathology and Diagnostics, University of Verona, P. le Scuro 10, 37134 Verona, Italy
 ²Cell Signaling Technology Inc., 3 Trask Lane, Danvers, MA 01923 USA
 ³Department of Biotechnology, University of Verona, Strada Le Grazie 15, 37134 Verona, Italy
 ⁴Dipartimento di Patologia e Diagnostica, Sezione di Anatomia Patologica, Università di Verona – Policlinico G. B. Rossi, Piazzale L. A. Scuro, 10, 37134 Verona, Italy
 Verona, Italy
 Alberto Zamò, Phone: +39-045-8124835, Fax: +39-045-8027136, Email: alberto.zamo@univr.it.
 Corresponding author.

Accepted 2011 Feb 12.

Copyright © The Author(s) 2011

This article has been cited by other articles in PMC.

References

 Swerdlow SH, Campo E, Seto M, Muller-Hermelink HK. In: Mantle cell lymphoma, in: WHO classification of tumours of haematopoietic and lymphoid tissues. Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri S, Stein H, Thiele J, Vardiman J, editors. Lyon: IARC Press; 2008.

2. Hofmann WK, Vos S, Tsukasaki K, Wachsman W, Pinkus GS, Said JW, Koeffler HP. Altered apoptosis pathways in mantle cell lymphoma detected by oligonucleotide microarray. Blood. 2001;98:787–794. doi: 10.1182/blood.V98.3.787. [PubMed] [Cross Ref]

3. Martinez N, Camacho FI, Algara P, Rodriguez A, Dopazo A, Ruiz-Ballesteros E, Martin P, Martinez-Climent JA, Garcia-Conde J, Menarguez J, Solano F, Mollejo M, Piris MA. The molecular signature of mantle cell lymphoma reveals multiple signals favoring cell survival. Cancer Res. 2003;63:8226–8232. [PubMed]

4. Ortega-Paino E, Fransson J, Ek S, Borrebaeck CA. Functionally associated targets in mantle cell lymphoma as defined by DNA microarrays and RNA interference. Blood. 2008;111:1617–1624. doi: 10.1182/blood-2007-02-068791. [PubMed] [Cross Ref]

5. Pinyol M, Bea S, Pla L, Ribrag V, Bosq J, Rosenwald A, Campo E, Jares P. Inactivation of RB1 in mantle-cell lymphoma detected by nonsense-mediated mRNA decay pathway inhibition and microarray analysis. Blood. 2007;109:5422–5429. doi: 10.1182/blood-2006-11-057208. [PubMed] [Cross Ref]

6. Rinaldi A, Kwee I, Taborelli M, Largo C, Uccella S, Martin V, Poretti G, Gaidano G, Calabrese G, Martinelli G, Baldini L, Pruneri G, Capella C, Zucca E, Cotter FE, Cigudosa JC, Catapano CV, Tibiletti MG, Bertoni F. Genomic and expression profiling identifies the B-cell associated tyrosine kinase Syk as a possible therapeutic target in mantle cell lymphoma. Br. J. Haematol. 2006;132:303–316. doi: 10.1111/j.1365-2141.2005.05883.x. [PubMed] [Cross Ref]

 Rizzatti EG, Falcao RP, Panepucci RA, Proto-Siqueira R, Anselmo-Lima WT, Okamoto OK, Zago MA. Gene expression profiling of mantle cell lymphoma cells reveals aberrant expression of genes from the PI3K-AKT, WNT and TGFbeta signalling pathways. Br. J. Haematol. 2005;130:516–526. doi: 10.1111/j.1365-2141.2005.05630.x. [PubMed] [Cross Ref]

8. Rosenwald A, Wright G, Wiestner A, Chan WC, Connors JM, Campo E, Gascoyne RD, Grogan TM, Muller-Hermelink HK, Smeland EB, Chiorazzi M, Giltnane JM, Hurt EM, Zhao H, Averett L, Henrickson S, Yang L, Powell J, Wilson WH, Jaffe ES, Simon R, Klausner RD, Montserrat E, Bosch F, Greiner TC, Weisenburger DD, Sanger WG, Dave BJ, Lynch JC, Vose J, Armitage JO, Fisher RI, Miller TP, LeBlanc M, Ott G, Kvaloy S, Holte H, Delabie J, Staudt LM. The proliferation gene expression signature is a quantitative integrator of oncogenic events that predicts survival in mantle cell lymphoma. Cancer Cell. 2003;3:185–197. doi: 10.1016/S1535-6108(03)00028-X. [PubMed] [Cross Ref]

9. Thieblemont C, Nasser V, Felman P, Leroy K, Gazzo S, Callet-Bauchu E, Loriod B, Granjeaud S, Gaulard P, Haioun C, Traverse-Glehen A, Baseggio L, Bertucci F, Birnbaum D, Magrangeas F, Minvielle S, Avet-Loiseau H, Salles G, Coiffier B, Berger F, Houlgatte R. Small lymphocytic lymphoma, marginal zone B-cell lymphoma, and mantle cell lymphoma exhibit distinct gene-expression profiles allowing molecular diagnosis. Blood. 2004;103:2727–2737. doi: 10.1182/blood-2003-06-2160. [PubMed] [Cross Ref]

 Antonucci F, Chilosi M, Parolini C, Hamdan M, Astner H, Righetti PG. Two-dimensional molecular profiling of mantle cell lymphoma. Electrophoresis. 2003;24:2376–2385. doi: 10.1002/elps.200305457. [PubMed] [Cross Ref]

11. Cecconi D, Zamó A, Parisi A, Bianchi E, Parolini C, Timperio AM, Zolla L, Chilosi M. Induction of apoptosis in Jeko-1 mantle cell lymphoma cell line by resveratrol: a proteomic analysis. J. Proteome Res. 2008;7:2670–2680. doi: 10.1021/pr700712p. [PubMed] [Cross Ref]

12. Ghobrial IM, McCormick DJ, Kaufmann SH, Leontovich AA, Loegering DA, Dai NT, Krajnik KL, Stenson MJ, Melhem MF, Novak AJ, Ansell SM, Witzig TE. Proteomic analysis of mantle-cell lymphoma by protein microarray. Blood. 2005;105:3722–3730. doi: 10.1182/blood-2004-10-3999. [PMC free article] [PubMed] [Cross Ref]

13. Weinkauf M, Zimmermann Y, Hartmann E, Rosenwald A, Rieken M, Pastore A, Hutter G, Hiddemann W, Dreyling M. 2-D PAGEbased comparison of proteasome inhibitor bortezomib in sensitive and resistant mantle cell lymphoma. Electrophoresis. 2009;30:974–986. doi: 10.1002/elps.200800508. [PubMed] [Cross Ref]

14. Jansen C, Hebeda KM, Linkels M, Grefte JM, Raemaekers JM, Krieken JH, Groenen PJ. Protein profiling of B-cell lymphomas using tissue biopsies: a potential tool for small samples in pathology. Cell. Oncol. 2008;30:27–38. [PMC free article] [PubMed]

15. Jansen C, Feuth T, Raemaekers JMM, Rijntjes J, Meijer JW, Westenend PJ, Baarlen J, Krieken JHJM, Hebeda KM, Groenen PJTA. Protein profiling in pathology: analysis and evaluation of 239 frozen tissue biopsies for diagnosis of B-cell lymphomas. Proteomics: Clin. Appl. 2010;4:519–527. [PubMed]

16. Stranneheim H, Orre LM, Lehtio J, Flygare J. A comparison between protein profiles of B cell subpopulations and mantle cell lymphoma cells. Proteome Sci. 2009;7:43. doi: 10.1186/1477-5956-7-43. [PMC free article] [PubMed] [Cross Ref]

17. Miguet L, Bechade G, Fornecker L, Zink E, Felden C, Gervais C, Herbrecht R, Dorsselaer A, Mauvieux L, Sanglier-Cianferani S. Proteomic analysis of malignant B-cell derived microparticles reveals CD148 as a potentially useful antigenic biomarker for mantle cell lymphoma diagnosis. J. Proteome Res. 2009;8:3346–3354. doi: 10.1021/pr801102c. [PubMed] [Cross Ref]

 Weinkauf M, Christopeit M, Hiddemann W, Dreyling M. Proteome- and microarray-based expression analysis of lymphoma cell lines identifies a p53-centered cluster of differentially expressed proteins in mantle cell and follicular lymphoma. Electrophoresis. 2007;28:4416 –4426. doi: 10.1002/elps.200600831. [PubMed] [Cross Ref]

19. Boyd RS, Jukes-Jones R, Walewska R, Brown D, Dyer MJ, Cain K. Protein profiling of plasma membranes defines aberrant signaling pathways in mantle cell lymphoma. Mol. Cell. Proteomics. 2009;8:1501–1515. doi: 10.1074/mcp.M800515-MCP200. [PMC free article] [PubMed] [Cross Ref]

20. Cecconi D, Zamo A, Bianchi E, Parisi A, Barbi S, Milli A, Rinalducci S, Rosenwald A, Hartmann E, Zolla L, Chilosi M. Signal transduction pathways of mantle cell lymphoma: a phosphoproteome-based study. Proteomics. 2008;8:4495–4506. doi: 10.1002/pmic.200800080. [PubMed] [Cross Ref]

21. Weinkauf M, Hiddemann W, Dreyling M. Sample pooling in 2-D gel electrophoresis: a new approach to reduce nonspecific expression background. Electrophoresis. 2006;27:4555–4558. doi: 10.1002/elps.200600207. [PubMed] [Cross Ref]

22. Rush J, Moritz A, Lee KA, Guo A, Goss VL, Spek EJ, Zhang H, Zha XM, Polakiewicz RD, Comb MJ. Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. Nat. Biotechnol. 2005;23:94–101. doi: 10.1038/nbt1046. [PubMed] [Cross Ref]

23. Rikova K, Guo A, Zeng Q, Possemato A, Yu J, Haack H, Nardone J, Lee K, Reeves C, Li Y, Hu Y, Tan Z, Stokes M, Sullivan L, Mitchell J, Wetzel R, Macneill J, Ren JM, Yuan J, Bakalarski CE, Villen J, Kornhauser JM, Smith B, Li D, Zhou X, Gygi SP, Gu TL, Polakiewicz RD, Rush J, Comb MJ. Global survey of phosphotyrosine signaling identifies oncogenic kinases in lung cancer. Cell. 2007;131:1190–1203. doi: 10.1016/j.cell.2007.11.025. [PubMed] [Cross Ref]

24. J. Carretero, T. Shimamura, K. Rikova, A.L. Jackson, M.D. Wilkerson, C.L. Borgman, M.S. Buttarazzi, B.A. Sanofsky, K.L. McNamara, K.A. Brandstetter, Z.E. Walton, T.L. Gu, J.C. Silva, K. Crosby, G.I. Shapiro, S.M. Maira, H. Ji, D.H. Castrillon, C.F. Kim, C. Garcia-Echeverria, N. Bardeesy, N.E. Sharpless, N.D. Hayes, W.Y. Kim, J.A. Engelman, K.K. Wong, Integrative genomic and proteomic analyses identify targets for Lkb1-deficient metastatic lung tumors. Cancer Cell 17 547–559 [PMC free article] [PubMed]

 Walters DK, Goss VL, Stoffregen EP, Gu TL, Lee K, Nardone J, McGreevey L, Heinrich MC, Deininger MW, Polakiewicz R, Druker BJ. Phosphoproteomic analysis of AML cell lines identifies leukemic oncogenes. Leuk. Res. 2006;30:1097–1104. doi: 10.1016/j.leukres.2006.01.001. [PubMed] [Cross Ref]

26. Goss VL, Lee KA, Moritz A, Nardone J, Spek EJ, MacNeill J, Rush J, Comb MJ, Polakiewicz RD. A common phosphotyrosine signature for the Bcr-Abl kinase. Blood. 2006;107:4888–4897. doi: 10.1182/blood-2005-08-3399. [PMC free article] [PubMed] [Cross Ref]

Gu TL, Goss VL, Reeves C, Popova L, Nardone J, Macneill J, Walters DK, Wang Y, Rush J, Comb MJ, Druker BJ, Polakiewicz RD.
 Phosphotyrosine profiling identifies the KG-1 cell line as a model for the study of FGFR1 fusions in acute myeloid leukemia. Blood.
 2006;108:4202–4204. doi: 10.1182/blood-2006-06-026666. [PubMed] [Cross Ref]

28. T.L. Gu, J. Cherry, M. Tucker, J. Wu, C. Reeves, R.D. Polakiewicz, Identification of activated Tnk1 kinase in Hodgkin's lymphoma. Leukemia 24 861–865. [PubMed]

29. Bea S, Salaverria I, Armengol L, Pinyol M, Fernandez V, Hartmann EM, Jares P, Amador V, Hernandez L, Navarro A, Ott G, Rosenwald A, Estivill X, Campo E. Uniparental disomies, homozygous deletions, amplifications, and target genes in mantle cell lymphoma revealed by integrative high-resolution whole-genome profiling. Blood. 2009;113:3059–3069. doi: 10.1182/blood-2008-07-170183. [PMC free article] [PubMed] [Cross Ref]

30. Rubio-Moscardo F, Climent J, Siebert R, Piris MA, Martin-Subero JI, Nielander I, Garcia-Conde J, Dyer MJ, Terol MJ, Pinkel D, Martinez-Climent JA. Mantle-cell lymphoma genotypes identified with CGH to BAC microarrays define a leukemic subgroup of disease and predict patient outcome. Blood. 2005;105:4445–4454. doi: 10.1182/blood-2004-10-3907. [PubMed] [Cross Ref]

31. Vater I, Wagner F, Kreuz M, Berger H, Martin-Subero JI, Pott C, Martinez-Climent JA, Klapper W, Krause K, Dyer MJ, Gesk S, Harder L, Zamo A, Dreyling M, Hasenclever D, Arnold N, Siebert R. GeneChip analyses point to novel pathogenetic mechanisms in mantle cell lymphoma. Br. J. Haematol. 2009;144:317–331. doi: 10.1111/j.1365-2141.2008.07443.x. [PubMed] [Cross Ref]

32. Hui D, Reiman T, Hanson J, Linford R, Wong W, Belch A, Lai R. Immunohistochemical detection of cdc2 is useful in predicting survival in patients with mantle cell lymphoma. Mod. Pathol. 2005;18:1223–1231. doi: 10.1038/modpathol.3800409. [PubMed] [Cross Ref]

33. F. Jardin, J.M. Picquenot, F. Parmentier, P. Ruminy, M. Cornic, D. Penther, P. Bertrand, H. Lanic, O. Cassuto, C. Humbrecht, E. Lemasle, A. Wautier, C. Bastard, H. Tilly, Detection of gene copy number aberrations in mantle cell lymphoma by a single quantitative multiplex PCR assay: clinicopathological relevance and prognosis value. Br. J. Haematol. (2009) [PubMed]

34. Tagawa H, Karnan S, Suzuki R, Matsuo K, Zhang X, Ota A, Morishima Y, Nakamura S, Seto M. Genome-wide array-based CGH for mantle cell lymphoma: identification of homozygous deletions of the proapoptotic gene. BIM Oncogene. 2005;24:1348–1358. doi: 10.1038/sj.onc.1208300. [PubMed] [Cross Ref]

35. Cho YY, Tang F, Yao K, Lu C, Zhu F, Zheng D, Pugliese A, Bode AM, Dong Z. Cyclin-dependent kinase-3-mediated c-Jun phosphorylation at Ser63 and Ser73 enhances cell transformation. Cancer Res. 2009;69:272–281. doi: 10.1158/0008-5472.CAN-08-3125. [PMC free article] [PubMed] [Cross Ref]

36. Zheng D, Cho YY, Lau AT, Zhang J, Ma WY, Bode AM, Dong Z. Cyclin-dependent kinase 3-mediated activating transcription factor 1 phosphorylation enhances cell transformation. Cancer Res. 2008;68:7650–7660. doi: 10.1158/0008-5472.CAN-08-1137.
[PMC free article] [PubMed] [Cross Ref]

37. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids Res. 1999;27:29–34. doi: 10.1093/nar/27.1.29. [PMC free article] [PubMed] [Cross Ref]

38. Chen L, Monti S, Juszczynski P, Daley J, Chen W, Witzig TE, Habermann TM, Kutok JL, Shipp MA. SYK-dependent tonic B-cell receptor signaling is a rational treatment target in diffuse large B-cell lymphoma. Blood. 2008;111:2230–2237. doi: 10.1182/blood-2007-07-100115. [PMC free article] [PubMed] [Cross Ref]

39. R.E. Davis, V.N. Ngo, G. Lenz, P. Tolar, R.M. Young, P.B. Romesser, H. Kohlhammer, L. Lamy, H. Zhao, Y. Yang, W. Xu, A.L. Shaffer, G. Wright, W. Xiao, J. Powell, J.K. Jiang, C.J. Thomas, A. Rosenwald, G. Ott, H.K. Muller-Hermelink, R.D. Gascoyne, J.M. Connors, N.A. Johnson, L.M. Rimsza, E. Campo, E.S. Jaffe, W.H. Wilson, J. Delabie, E.B. Smeland, R.I. Fisher, R.M. Braziel, R.R. Tubbs, J.R. Cook, D.D. Weisenburger, W.C. Chan, S.K. Pierce, L.M. Staudt, Chronic active B-cell-receptor signalling in diffuse large B-cell lymphoma. Nature 463 88–92 [PMC free article] [PubMed]

40. Gobessi S, Laurenti L, Longo PG, Carsetti L, Berno V, Sica S, Leone G, Efremov DG. Inhibition of constitutive and BCR-induced Syk activation downregulates Mcl-1 and induces apoptosis in chronic lymphocytic leukemia B cells. Leukemia. 2009;23:686–697. doi: 10.1038/leu.2008.346. [PubMed] [Cross Ref]

41. J.W. Friedberg, J. Sharman, J. Sweetenham, P.B. Johnston, J.M. Vose, A. Lacasce, J. Schaefer-Cutillo, S. De Vos, R. Sinha, J.P. Leonard, L.D. Cripe, S.A. Gregory, M.P. Sterba, A.M. Lowe, R. Levy, M.A. Shipp, Inhibition of Syk with fostamatinib disodium has significant clinical activity in non-Hodgkin lymphoma and chronic lymphocytic leukemia. Blood **115** 2578–2585 [PMC free article] [PubMed]

42. Oliver JM, Burg DL, Wilson BS, McLaughlin JL, Geahlen RL. Inhibition of mast cell Fc epsilon R1-mediated signaling and effector function by the Syk-selective inhibitor, piceatannol. J. Biol. Chem. 1994;269:29697–29703. [PubMed]

43. Leseux L, Hamdi SM, Saati T, Capilla F, Recher C, Laurent G, Bezombes C. Syk-dependent mTOR activation in follicular lymphoma cells. Blood. 2006;108:4156–4162. doi: 10.1182/blood-2006-05-026203. [PubMed] [Cross Ref]

44. Rinaldi A, Kwee I, Taborelli M, Largo C, Uccella S, Martin V, Poretti G, Gaidano G, Calabrese G, Martinelli G, Baldini L, Pruneri G, Capella C, Zucca E, Cotter FE, Cigudosa JC, Catapano CV, Tibiletti MG, Bertoni F. Genomic and expression profiling identifies the
B-cell associated tyrosine kinase Syk as a possible therapeutic target in mantle cell lymphoma. Br. J. Haematol. 2006;132:303–316. doi: 10.1111/j.1365-2141.2005.05883.x. [PubMed] [Cross Ref]

45. Ennaciri J, Girard D. IL-4R(alpha), a new member that associates with Syk kinase: implication in IL-4-induced human neutrophil functions. J. Immunol. 2009;183:5261–5269. doi: 10.4049/jimmunol.0900109. [PubMed] [Cross Ref]

46. Wang L, Devarajan E, He J, Reddy SP, Dai JL. Transcription repressor activity of spleen tyrosine kinase mediates breast tumor suppression. Cancer Res. 2005;65:10289–10297. doi: 10.1158/0008-5472.CAN-05-2231. [PMC free article] [PubMed] [Cross Ref]

47. F.M. Uckun, S. Qazi, H. Ma, L. Tuel-Ahlgren, Z. Ozer, STAT3 is a substrate of SYK tyrosine kinase in B-lineage leukemia/lymphoma cells exposed to oxidative stress. Proc. Natl. Acad. Sci. U S A **107** 2902–2907. [PMC free article] [PubMed]

48. Baran-Marszak F, Boukhiar M, Harel S, Laguillier C, Roger C, Gressin R, Martin A, Fagard R, Varin-Blank N, Ajchenbaum-Cymbalista F, Ledoux D. Constitutive and B-cell receptor-induced activation of STAT3 are important signaling pathways targeted by bortezomib in leukemic mantle cell lymphoma. Haematologica. 2010;95:1865–1872. doi: 10.3324/haematol.2009.019745. [PMC free article] [PubMed] [Cross Ref]

49. Pham LV, Tamayo AT, Li C, Bornmann W, Priebe W, Ford RJ. Degrasyn potentiates the antitumor effects of bortezomib in mantle cell lymphoma cells in vitro and in vivo: therapeutic implications. Mol. Cancer Ther. 2010;9:2026–2036. doi: 10.1158/1535-7163.MCT-10-0238. [PMC free article] [PubMed] [Cross Ref]

50. Tauzin S, Ding H, Khatib K, Ahmad I, Burdevet D, Echten-Deckert G, Lindquist JA, Schraven B, Din NU, Borisch B, Hoessli DC. Oncogenic association of the Cbp/PAG adaptor protein with the Lyn tyrosine kinase in human B-NHL rafts. Blood. 2008;111:2310–2320. doi: 10.1182/blood-2007-05-090985. [PubMed] [Cross Ref]

51. Yared MA, Khoury JD, Medeiros LJ, Rassidakis GZ, Lai R. Activation status of the JAK/STAT3 pathway in mantle cell lymphoma. Arch. Pathol. Lab. Med. 2005;129:990–996. [PubMed]

52. Lai R, Rassidakis GZ, Medeiros LJ, Leventaki V, Keating M, McDonnell TJ. Expression of STAT3 and its phosphorylated forms in mantle cell lymphoma cell lines and tumours. J. Pathol. 2003;199:84–89. doi: 10.1002/path.1253. [PubMed] [Cross Ref]

53. Diehl JA, Cheng M, Roussel MF, Sherr CJ. Glycogen synthase kinase-3beta regulates cyclin D1 proteolysis and subcellular localization. Genes Dev. 1998;12:3499–3511. doi: 10.1101/gad.12.22.3499. [PMC free article] [PubMed] [Cross Ref]

54. Camps J, Salaverria I, Garcia MJ, Prat E, Bea S, Pole JC, Hernandez L, Rey J, Cigudosa JC, Bernues M, Caldas C, Colomer D, Miro R, Campo E. Genomic imbalances and patterns of karyotypic variability in mantle-cell lymphoma cell lines. Leuk. Res. 2006;30:923–934. doi: 10.1016/j.leukres.2005.11.013. [PubMed] [Cross Ref]

55. Tait SW, Green DR. Mitochondria and cell death: outer membrane permeabilization and beyond. Nat. Rev. Mol. Cell Biol. 2010;11:621–632. doi: 10.1038/nrm2952. [PubMed] [Cross Ref]

56. Miyashita T, Reed JC. Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. Cell. 1995;80:293–299. doi: 10.1016/0092-8674(95)90513-8. [PubMed] [Cross Ref]

57. Zhou F, Hu J, Ma H, Harrison ML, Geahlen RL. Nucleocytoplasmic trafficking of the Syk protein tyrosine kinase. Mol. Cell. Biol. 2006;26:3478–3491. doi: 10.1128/MCB.26.9.3478-3491.2006. [PMC free article] [PubMed] [Cross Ref]

58. Wang L, Duke L, Zhang PS, Arlinghaus RB, Symmans WF, Sahin A, Mendez R, Dai JL. Alternative splicing disrupts a nuclear localization signal in spleen tyrosine kinase that is required for invasion suppression in breast cancer. Cancer Res. 2003;63:4724–4730. [PubMed]

59. Bordin L, Fiore C, Bragadin M, Brunati AM, Clari G. Regulation of membrane band 3 Tyr-phosphorylation by proteolysis of p72(Syk) and possible involvement in senescence process. Acta. Biochim. Biophys. Sin. (Shanghai) 2009;41:846–851. doi: 10.1093/abbs/gmp071. [PubMed] [Cross Ref]

60. Zamo A, Ott G, Katzenberger T, Adam P, Parolini C, Scarpa A, Lestani M, Menestrina F, Chilosi M. Establishment of the MAVER-1 cell line, a model for leukemic and aggressive mantle cell lymphoma. Haematologica. 2006;91:40–47. [PubMed]

61. Dennis G, Jr, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA. DAVID: Database for Annotation, Visualization, and Integrated Discovery. Genome Biol. 2003;4:P3. doi: 10.1186/gb-2003-4-5-p3. [PubMed] [Cross Ref]