

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Induction of chromosomal aberrations and micronuclei by 2-hydroxy-4-methoxybenzophenone (oxybenzone) in human lymphocytes**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1666678> since 2018-04-19T11:55:54Z

*Published version:*

DOI:10.1080/01480545.2018.1455206

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



http://dx.doi.org/10.1080/15475277.2014.941111



**Induction of chromosomal aberrations and micronuclei by  
2-Hydroxy-4-methoxybenzophenone (Oxybenzone) in  
human lymphocytes.**

Journal:	<i>Drug and Chemical Toxicology</i>
Manuscript ID	Draft
Manuscript Type:	Original Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Santovito, Alfredo; University of Turin, Department of Life Sciences and Systems Biology Ruberto, Stefano; University of Turin, Department of Life Sciences and Systems Biology Galli, Gabriella; University of Turin, Department of Life Sciences and Systems Biology Menghi, Costanza; University of Turin, Department of Life Sciences and Systems Biology Girotti, Marilena; University of turin, Department of Life Sciences and Systems Biology Cervella, Piero; University of Turin, Department of Life Sciences and Systems Biology
Keywords:	BP-3, endocrine disruptors, genotoxicology

SCHOLARONE™  
Manuscripts

1  
2  
3  
4 **Title: Induction of chromosomal aberrations and micronuclei by 2-Hydroxy-4-**  
5 **methoxybenzophenone (Oxybenzone) in human lymphocytes.**  
6  
7  
8

9  
10  
11 **Short Title:** Effects of Oxybenzone on Human Lymphocytes  
12

13  
14  
15 Authors: Alfredo SANTOVITO<sup>a\*</sup>, Stefano RUBERTO<sup>a</sup>, Gabriella GALLI<sup>a</sup>, Costanza MENGHI<sup>a</sup>,  
16  
17 Marilena GIROTTI<sup>a</sup> and Piero CERVELLA<sup>a</sup>  
18  
19

20  
21  
22 <sup>a</sup> University of Turin, Department of Life Sciences and Systems Biology, Via Accademia Albertina  
23  
24 n. 13, 10123 Torino (Italy)  
25  
26

27  
28  
29 \*Corresponding Author:

30 Alfredo SANTOVITO

31  
32 Department of Life Sciences and Systems Biology

33 Via Accademia Albertina n. 13

34  
35 10123 – Torino (Italy)

36  
37 Tel.: +39-0116704554

38  
39 Fax: +39-0116704508  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

Benzophenone-3 (2-hydroxy-4-methoxybenzophenone; BP-3) is a UV filter used in a variety of personal care products for protection of human skin and hair from damage by ultraviolet radiation. BP-3 is suspected to exhibit endocrine disruptive properties. Indeed, it was found able to interact with the endocrine system causing alteration of its homeostasis, with consequent adverse health effects. Moreover, it is ubiquitously present in the environment, mostly in aquatic ecosystems, with consequent risks to the health of aquatic organisms and humans. In the present study, we analysed the cytogenetic effects of BP-3 on human lymphocytes using *in vitro* chromosomal aberrations and micronuclei assays. Lymphocyte cultures were exposed to five concentrations of BP-3 (0.20, 0.10, 0.05, 0.025 and 0.0125  $\mu\text{g}/\text{mL}$ ) for 24 and 48 hrs (for chromosomal aberrations and micronuclei tests, respectively). The concentration of 0.10  $\mu\text{g}/\text{mL}$  represents the ADI/TDI reference dose established by EU, whereas 0.20, 0.05, 0.025 and 0.0125  $\mu\text{g}/\text{mL}$  represent multiple and sub-multiple of this concentration value. Our results reported, for the first time in literature, the cytogenetic effects of BP-3 on cultured human lymphocytes in terms of increased micronuclei and chromosomal aberrations frequencies at all tested concentrations, including concentrations lower than those established by EU. *Vive versa*, after 48-h exposure, a significant reduction of the CBPI value in cultures treated with BP-3 was not observed, indicating that BP-3 does not seem to produce effects on the proliferation/mitotic index when its concentration is equal or less than 0.20  $\mu\text{g}/\text{mL}$ .

**Keywords:**

BP-3, endocrine disruptors, genotoxicology.

**Abbreviations**

Ab.C = Aberrant Cell

AF = Acentric Fragments

BP = Benzophenone

BP-1 = 2,4- dihydroxybenzophenone

BP-8 = 2,2'-dihydroxy-4-methoxybenzophenone

BP-2 = 2,2',4,4'-tetrahydroxybenzophenone

4-OH-BP = 4-hydroxybenzophenone

B' = chromatid breaks

B'' = chromosome breaks

bw = body weight

CAs = Chromosomal Aberrations

CBPI = Cytokinesis-Block Proliferation Index

Dic = Dicentrics

EU = European Union

EFSA = European Food Safety Authority

FCS = Foetal Calf Serum

KCl = Potassium chloride

MNs = Micronuclei

OECD = Organization for Economic Co-operation and Development

R = Rings

Re = Rearrangements

TDI = Tolerable Daily Intake

TR = tri- or tetra-radials

## Introduction

UV filters are widely used in sunscreens and personal care products, such as cosmetics and shampoos, for the protection of skin and hair from UV irradiation (Chisvert et al., 2012; Asimakopoulos et al., 2014). They are also present as chemical ingredients of insecticides, agricultural chemicals and pharmaceuticals (Careghini et al., 2015), as well as they are used to coat surfaces exposed to sunlight, including some food packaging (Vione et al., 2013).

Benzophenone (BP)-type chemicals are one of the primary components in the UV-filter family, detected at high concentrations in biological fluids of different populations worldwide distributed (Calafat et al., 2008; Wang and Kannan, 2013).

The 2-Hydroxy-4-methoxybenzophenone (BP-3), also known as oxybenzone, is a compound naturally occurs in flower pigments that can adsorb sunlight in the UVA and UVB regions (French, 1992). For this reason and for its limited phototransformation, it is one of the most commonly-used chemical components in sunscreen and cosmetic products (Careghini et al., 2015; Kim and Choi, 2014), found in 59% of sunscreens in the United States (Dewalque et al., 2014). BP-3 is also employed as photostabiliser in food packaging materials, to prevent polymer photochemical degradation, in the treatment of photodermatitis, as well as in plastic surface coatings and polymers (Vione et al., 2013).

However, from ecological point of view, the increasing use of UV filters constitutes a potential risk for the environment. Indeed, these filters are often inert in traditional wastewater treatment processes, and thus have the potential to contaminate the reclaimed water system, natural water bodies and drinking water resources (Xiao et al., 2013). Moreover, these chemicals are also directly released into surface waters through swimming, bathing, leaching of land and house coatings. As final result, BP-3 and other UV-filter components were detected in several environmental matrices such as in surface and tap waters and in sediments (Balmer et al., 2005, Kameda et al., 2011; Gago-Ferrero et al., 2011). In particular, BP3 has been detected at levels of up to some  $\mu\text{g/L}$  in raw

1  
2 wastewater, at tens to several hundreds ng/L in treated wastewater, up to a hundred ng/L in lake  
3  
4 water and at ng/g levels in solid matrices and in biota (Vione et al., 2013).

5  
6 Humans are exposed to BP-type UV filters largely through dermal absorption. Because of the  
7  
8 extensive use in personal care products, BP-3 and its metabolic derivatives, such as 2,4-  
9  
10 dihydroxybenzophenone (BP-1), 2,2',4,4'-tetrahydroxybenzophenone (BP-2), 2,2'-dihydroxy-4-  
11  
12 methoxybenzophenone (BP-8), and 4-hydroxybenzophenone (4-OH-BP), were found widely in  
13  
14 various human bodily fluids, such as urine (Calafat *et al.*, 2008; Kunisue *et al.*, 2012; Wolff *et al.*,  
15  
16 2007), breast milk (Ye et al., 2008a), blood (Ye et al., 2008b), and semen (León et al., 2010).

17  
18 In particular, these compounds have been reported in urines of a high percentage of subjects from  
19  
20 many countries, including United States (Calafat et al., 2008; Philippat et al., 2015; Wang and  
21  
22 Kannan, 2013), China (Wang and Kannan, 2013; Zhang et al., 2013), and Europe (Asimakopoulos  
23  
24 et al., 2014; Philippat et al., 2012).

25  
26 To date, the reproductive and developmental toxicity of BP-type UV filters has been revealed in  
27  
28 some animal and human studies. A few BP-type UV filters, BP-3 included, have been suspected to  
29  
30 have endocrine disrupting effects, with consequent alterations in the reproductive system and with  
31  
32 *in vitro* and *in vivo* estrogenic and anti-androgenic effects (Schlumpf et al., 2001; Suzuki et al.,  
33  
34 2005). For example, BP-3 was found able to affect the reproduction of fish (Bluthgen *et al.*, 2012;  
35  
36 Gago-Ferrero and Diaz-Cruz, 2012) and other aquatic organisms, such as the crustacean *Daphnia*  
37  
38 *magna* (Fent et al., 2010) and the chlorophyte microalgae *Scenedesmus vacuolatus* (Rodil et al.,  
39  
40 2009), as well as was found able to induce vitellogenin and impairment of reproduction in rainbow  
41  
42 trout and in Japanese medaka (Coronado et al., 2008). Moreover, after oral administration of  
43  
44 benzophenone, a significant increase of the incidence of mononuclear-cell leukaemia,  
45  
46 hepatoblastoma and hepatocellular carcinoma in male mice and histiocytic sarcoma in female mice  
47  
48 was observed (IARC, 2013).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

In humans, BP-type UV filters have been linked to various endocrine and reproductive disorders, such as reduced couple fecundity and semen quality (Buck Louis et al., 2014, 2015), birth outcomes (Wolff et al., 2008), uterine leiomyoma (Pollack et al., 2015) and endometriosis (Kunisue et al., 2012).

The Council of Europe reported maximum BP-3 levels of 0.5 mg/kg for beverages and 2 mg/kg for foods (IARC, 2013). The European Food Safety Authority estimated a dietary exposure to benzophenone in the European Union (EU) and in the USA correspondent to 23 and 11 µg per capita per day, respectively (IARC, 2013). Based on published toxicological data, the EU established a temporary tolerable daily intake (TDI) and an acceptable daily intake (ADI) for BP-3 of 0.1 mg/kg bw (Conseil of Europe, 2009).

From the genetic and cytogenetic point of view, some studies evidenced that BP-type UV-filters were able to induce mutagenic effects in *Salmonella* (Zeiger et al., 1987), sister chromatid exchanges and chromosomal aberrations (CAs) in Chinese hamster ovarian cells (French, 1992). However, no data are reported in literature about *in vitro* cytogenetic effects of BP-3 on human lymphocytes. For this reason, in the present study we assessed the possible *in vitro* clastogenic and aneugenic effects of BP-3 exposure on human lymphocyte cells, by CAs and micronuclei (MNs) assays.

The CAs assay allows the detection of cells carrying unstable aberrations (chromosome and chromatid breaks, deletions, fragments, rings, dicentrics and chromatid exchanges) that will lead to cell death during proliferation, and thus represents a useful test for the detection of potential clastogenic effects of xenobiotics. MNs represent acentric chromosomal fragments or whole chromosomes left behind during mitotic cell division and appear in the cytoplasm of interphase cells as small additional nuclei. With respect to the CAs assay, the MNs assay allows the detection of both potential clastogenic (chromosome breakage) or aneugenic (chromosome lagging due to



dysfunction of mitotic apparatus) effects of different xenobiotics. Interestingly, previous published studies provided strong evidence in support of the hypothesis that high CAs and MNs frequencies in peripheral blood lymphocytes are powerful predictors of cancer risk (Bonassi et al., 2004) and are positively associated with early events in carcinogenesis (Bonassi et al., 2011).

## Methods

### *Chemicals and Media*

The IUPAC name of Oxybenzone (CAS n. 131-57-7) is 2-Hydroxy-4-methoxybenzophenone.

The BP-3 was dissolved in Dimethyl Sulfoxide (DMSO, CAS no. 67-68-5). Gibco RPMI 1640 cell culture media supplemented with L-glutamine, foetal calf serum, phytohemagglutinin (PHA), and antibiotics were purchased from Invitrogen-Life Technologies, Milan, Italy. Cytochalasin-B, Mitomycin-C (MMC), DMSO, BP-3 and Giemsa stain solution were obtained from Sigma-Aldrich, Milan, Italy. Methanol, Acetic acid, and conventional microscope slides were purchased from Carlo Erba Reagenti, Milan, Italy. Potassium chloride (KCl) and Sørensen buffer were obtained from Merck S.p.A., Milan, Italy. Vacutainer blood collection tubes were from Terumo Europe, Rome, Italy.

### *Subjects*

Peripheral venous blood was collected from 5 healthy subjects (2 males and 3 females, mean age $\pm$ S.D., 31.50 $\pm$ 14.39), non-smoking, non-alcoholic, not under drug therapy, and with no recent history of exposure to mutagens. Informed consent was obtained from all blood donors. The study was approved by the Univeristy of Turin ethics committee and was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

### *Blood Sample Collection and Lymphocyte cultures*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Heparinized blood samples were obtained by venipuncture and collected in heparinised tubes, for genotoxicity testing. All blood samples were coded, cooled (4°C), and processed within 2 h after collection. Heparinized venous blood (0.3 mL) was cultured in 25 cm<sup>2</sup> flasks in 6 mL of RPMI-1640 medium supplemented with 20% foetal calf serum (FCS), 2% of the mitogenic agent PHA, L-glutamine (2 mM), antibiotics (100 IU/mL penicillin, and 100 µg/mL streptomycin). The cultures were incubated for 72 h at 37°C, under 5% of CO<sub>2</sub> in the air in a humidified atmosphere. After 24 h of incubation, BP-3 dissolved in DMSO (for stock solution preparation 200 µg of BP-3 were dissolved in 1 mL of DMSO) was added to the cultures to a final concentrations of 0.20 µg/mL, 0.10 µg/mL, 0.05 µg/mL, 0.025 µg/mL and 0.0125 µg/mL. The concentration at 0.10 µg/mL represents the Tolerable Daily Intake (TDI) concentration established by European Union (0.1 mg/Kg BW) for this compound (Council of Europe, 2009); 0.20 µg/mL is a multiple of the TDI, whereas 0.05, 0.025 and 0.0125 µg/mL represent sub-multiple of this value, tested in order to determine the genotoxicity threshold limit.

Three control cultures were assessed: 1) positive control, by adding only MMC (final concentration 0.1 µg/mL culture); 2) solvent control, by adding only 0.1% of DMSO; 3) negative control, culture without both BP-3 and DMSO. Only for MNs assay, after 44 h of incubation, cytochalasin-B was added to the cultures at a concentration of 6 µg/mL to block cytokinesis. After 48 h (for CAs assay) and 72 h (for MNs assay) of incubation at 37°, the cells were collected by centrifugation and treated for 10 min with a pre-warmed mild hypotonic solution (75 mM KCl). After centrifugation and removal of the supernatant, the cells were fixed with a fresh mixture of methanol/acetic acid (3:1 v/v). The treatment with the fixative was repeated three times. Finally, the supernatant was discarded and the pellet, dissolved in a minimal volume of fixative, was seeded on the slides to detect CAs and MNs by conventional staining with 5% Giemsa (pH 6.8) prepared in Sørensen buffer.

*Cytokinesis-Block Micronucleus Assay*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Microscope analysis was performed at 40X magnification on a light microscope (Dialux 20, Leica, Germany). MNs were scored in 1,000 binucleated lymphocytes with well-preserved cytoplasm per subject (total 5,000 binucleated cells per concentration), following the established criteria for MNs evaluation (Fenech et al., 2003). A total of 1,000 lymphocytes per donor per concentration were scored to evaluate the percentage of cells with 1-4 nuclei. The cytokinesis-block proliferation index (CBPI) was calculated, according to the following formula:

$[1 \times N1] + [2 \times N2] + [3 \times (N3 + N4)]/N$ , where N1–N4 represents the number of cells with 1-4 nuclei, respectively, and N is the total number of cells scored.

#### *Chromosomal Aberrations Assay*

Microscope analysis was performed at 1000X magnification on a light microscope (Dialux 20, Leica, Germany). Although the Organization for Economic Co-operation and Development (OECD) guidelines call for 100 metaphases to be scored for each dose (OECD, 2015), in order to obtain results more significant and suitable for statistical analysis, it is the normal practice in our laboratory to score 200 metaphases *per subject per dose*. Thus, in the present work, for each subject and BP-3 concentration, 200 well-spread first-division complete metaphases (for a total of 1000 metaphases for each dose) were analysed for the following categories of CAs: gaps, chromatid breaks (B'), chromosome breaks (B''), dicentrics (Dic), rings (R), tri- or tetra-radials (TR), acentric fragments (AF), rearrangements (Re) and numerical aberrations. Cells containing one of more types of CAs were scored as "aberrant cell" (Ab.C).

With regard to the opportunity to include gaps in the statistical analyses, the discussion is open. Although some authors considered gaps as the appropriate indicator of genotoxic potential of chemicals (Savage, 2004), the molecular mechanism of BP-3 to induce achromatic lesion/gaps is yet to be revealed. Thus, in our statistical analyses we decided to exclude gaps. The criteria for distinguishing chromatide breaks from gaps were the acentric piece displaced with respect to the chromosome axis and the size of the discontinuity, which exceeded the width of the chromatide. A

1  
2 dicentric with an acentric fragment was scored as one aberration.  
3  
4  
5

### 6 7 *Statistical analysis*

8  
9 Comparison of mean values of the percentage of cells with MNs, CBPI and CAs between  
10  
11 exposition levels and their controls was assessed by the non-parametric Mann-Whitney test.  
12  
13 Statistical calculations were carried out using the SPSS software package program (version 23.0,  
14  
15 Inc., Chicago, IL, USA). All *P* values were two tailed, and *P* values of 5% or less were considered  
16  
17 statistically significant for all tests carried out.  
18  
19

## 20 21 22 **Results**

### 23 24 25 26 27 *Effect of BP-3 on CAs formation*

28  
29 Table 1 shows values of CAs found in the human peripheral lymphocytes cultured in the presence  
30  
31 of different concentrations of BP-3. BP-3 was found to induce seven types of structural CAs (gaps,  
32  
33 chromatid and chromosome breaks, dicentric chromosomes, rings, tri- or tetradials, acentric  
34  
35 fragments and rearrangements). The most frequent observed aberrations were acentric fragments  
36  
37 and chromatid aberrations, respectively, whereas no numerical aberrations were found.  
38  
39

40  
41 Data obtained indicated that human lymphocytes treated *in vitro* with BP-3 significantly increased  
42  
43 the CAs and Ab.C frequencies at all tested concentrations when compared with the solvent control,  
44  
45 including the lower concentration of 0.0125 µg/mL that represents an eighth of the TDI value  
46  
47 established by EU for this substance. Moreover, a dose-effect was observed since the regression  
48  
49 analysis revealed a significant ( $P < 0.001$ ) correlation between the BP-3 concentrations and the level  
50  
51 of genomic damage (Table 3).  
52

53  
54 No significant differences were found between the DMSO solvent-control and the negative control  
55  
56 ( $P = 0.142$ ), whereas the cultures treated with the known mutagen MMC showed a significant  
57  
58  
59  
60

1  
2 increase of CAs and Ab.C with respect to all BP-3 tested concentrations ( $P < 0.05$ ), including the  
3  
4 negative and solvent control cultures ( $P = 0.008$  and  $P = 0.009$ , respectively).  
5  
6  
7

#### 8 9 *Effect of BP-3 on MNs formation*

10  
11 To verify both the aneugenic and clastogenic effects of BP-3, the MN test was assessed in parallel  
12  
13 with CAs test (Table 2). Similarly to what we already observed with the CAs assay, our results  
14  
15 indicated that BP-3 significantly increased the MNs formation at all concentrations tested, including  
16  
17 both 0.10  $\mu\text{g/mL}$ , that represents the TDI established by EU for this substance, and 0.0125  $\mu\text{g/mL}$   
18  
19 that represents an eighth of this limit value. Also in this case, a dose-effect was observed since the  
20  
21 regression analysis revealed a significant ( $P < 0.001$ ) correlation between the BP-3 concentrations  
22  
23 and the frequencies of MNs and Cells with MNs (Table 3).  
24  
25

26  
27 The DMSO solvent-control cultures did not show any difference with the negative controls ( $P =$   
28  
29 0.126), further confirming that at this low concentration DMSO has no cytogenetic effects  
30  
31 evaluable by MN test. Similarly to what we observed with CAs assay, cultures treated with the  
32  
33 mutagen MMC showed a significant increase in the MNs formation compared with the negative  
34  
35 control ( $P = 0.008$ ), solvent controls ( $P = 0.008$ ) and all tested concentrations of BP-3 ( $P = 0.009$   
36  
37 for all concentrations). Finally, after 48-h exposure, a significant reduction of the CBPI value in  
38  
39 cultures treated with BP-3 was not observed, indicating that at the tested concentrations, BP-3 does  
40  
41 not seem to produce effects on the proliferation/mitotic index, as confirmed by the regression  
42  
43 analysis (Tables 2 and 3).  
44  
45  
46  
47  
48

#### 49 **Discussion**

50  
51 BP-3 is widely used as sunscreen for protection of human skin and hair from damage by ultraviolet  
52  
53 radiation. It has been found in many cosmetic products and its production and use has been rapidly  
54  
55 increasing over the past decade. As consequence of this increased production, BP-3 was found to  
56  
57 have a wide presence in aquatic environments, affecting the water quality and, consequently, the  
58  
59  
60

1  
2 human health. For these reasons, it is considered as a personal care product of emerging  
3  
4 environmental concern and its concentration limits, in the final products for human consumption,  
5  
6 are now regulated by the United States Food and Drug Administration and the European  
7  
8 Commission.  
9

10  
11 Here, we discuss data obtained from our *in vitro* study conducted in order to establish the  
12  
13 clastogenic and aneugenic potential of BP-3 on human cultured lymphocytes. The results obtained  
14  
15 with CAs and MNs assays, evidenced a possible clastogenic effect of the BP-3 on human  
16  
17 lymphocytes, also at the concentration of 0.0125 µg/mL that represents a value eight times lower  
18  
19 with respect to the reference dose established for humans by EU (Conseil of Europe, 2009). This  
20  
21 increased cytogenetic damage observed at all tested concentrations pushes towards the adoption of  
22  
23 more safe concentration for human health with respect to the established value of 0.1 µg/mL.  
24  
25

26  
27 Indeed, increased CAs and MNs frequencies in peripheral blood lymphocytes have been positively  
28  
29 associated with increased cancer risk and early events in carcinogenesis, respectively (Bonassi et  
30  
31 al., 2004; 2011).  
32  
33

34  
35 The genotoxicity of BP-type UV-filters has previously been observed by *in vivo* and *in vitro* studies  
36  
37 using non-human organisms and cell systems (Zeiger et al., 1987; French, 1992). However, beyond  
38  
39 these old reports, no publications are present in literature about the possible *in vitro* clastogenic  
40  
41 effect of BP-3 on human lymphocytes. For this reason, we cannot compare our data with other  
42  
43 similar data.  
44  
45

46  
47 Notably, for BP-3 we observed a clastogenic effect also at the lower concentrations of 0.025 and  
48  
49 0.0125 µg/mL, whereas for the BP-A, another endocrine disruptor compound with a very similar  
50  
51 molecular weight to that of BP-3, we failed to find it (Santovito et al., 2017). This more acute  
52  
53 effect of BP-3 with respect of BP-A has been observed, by our group, also in *in vivo* experiments  
54  
55 conducted using the marine polychaete *Ophryotrocha diadema*. In this case, we found an increased  
56  
57 mortality and a lower production of eggs among polychaete treated with the same concentrations of  
58  
59  
60

1  
2 BP-3 used in the present study, compared to those treated with identical BP-A concentrations (data  
3 not showed).  
4  
5

6  
7  
8 The clastogenic properties of the BP-3 could be ascribed to its demonstrated ability to generate  
9 ROS, probably by decreasing the activities of antioxidant enzymes and increasing lipid  
10 peroxidation (Wnuk et al., 2017). Indeed, Hunson et coll. (2006) demonstred that BP-3 and other  
11 UV-filters are able to penetrate through the stratum corneum and to generate higly reactive oxygen  
12 species in the cytoplasm of the epidermic nucleated keratinocytes.  
13  
14

15  
16 Moreover, the demonstrated accessibility of the BP-3 to the labile hydrogens within the DNA  
17 (Marazzi et al., 2016) could be another possible explanation of this observed clastogenic effect, as  
18 well as of the increased levels of genomic damage observed for BP-3 with respect to BP-A. This is  
19 an important issue because increased levels of genomic damage were associated with an increased  
20 risk of cancer development (Bonassi et al., 2004; 2011), and in particular BP-3 was found able to  
21 increase metastatic potential in lung cancer cells (Phiboonchaiyanan et al., 2017).  
22  
23

24  
25 Although recent studies on BP-3 have demonstrated its cytotoxicity properties (Balázs et al., 2016)  
26 and its capacity to activate apoptosis (Wnuk et al., 2017), in the present study we did not observed a  
27 significant reduction of the CBPI, indicating that BP-3 does not seem to produce effects on the  
28 proliferation/mitotic index when its concentration is equal or less than 0.20 µg/mL (Tables 2 and 3).  
29  
30

31  
32 Finally, it should be also emphasized that the observed cytogenetic damage cannot be ascribed  
33 exclusively to the direct effect of BP-3 on lymphocytes. Indeed, *in vitro* studies showed that BP-3  
34 is converted to more hydrophilic and DNA-binding metabolites, such as BP-1, BP-2, BP-8 and 4-  
35 OH-BP (Buck Louis et al., 2014). In this scenario, the irreversible binding of BP-3 derived  
36 compounds to the DNA could cause bases loss from the DNA strand and could be responsible for  
37 some of the BP-3 observed toxic effects. It is known that many chemicals, either directly or after  
38 conversion to reactive metabolites, covalently modify nucleosides of DNA, leading to mutational  
39 changes ans, thus, playing a role in cellular toxicity or tumorigenesis induction (Atkinson and Roy,  
40 1995).  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Conclusion

The results herein reported demonstrate, for the first time in literature, the clastogenic and aneugenic effects of BP-3 on cultured human lymphocytes, by increasing CAs and MNs frequencies also at lower concentrations with respect to those established by European Union.

It should be emphasized that this observed *in vitro* genotoxic potential on human lymphocytes does not seem to manifest itself in *in vivo* experiments using animal models like drosophila and rat (Robison et al., 1994). However, although simple experimental models, like mammalian and bacterial cells, cannot accurately mimic the complex kinetics of xenobiotic compounds *in vivo*, our results suggest the need for further investigations about this compound and, eventually, the adoption of more stringent measures in order to reduce the presence of this compound in the environment and to minimize the adverse effects of the BP-3 exposure on human health.

## Acknowledgment

We are grateful to all volunteers who participated in this study.

## Conflict of Interest

The authors report no conflicts of interest.

## Funding

This research was supported by grant from the Italian Ministry of University and Scientific Research (“ex 60%”).



## References

- 1) Asimakopoulos, A.G., Thomaidis, N.S., and Kannan, K., 2014. Widespread occurrence of bisphenol A diglycidyl ethers, p-hydroxybenzoic acid esters (parabens), benzophenone type-UV filters, triclosan, and triclocarban in human urine from Athene, Grece. *Science of the Total Environmen*, 470-471, 1243-1249.
- 2) Atkinson, A and Roy. D., 1995. In vitro conversion of environmental estrogenic chemical bisphenol A to DNA binding metabolite(s). *Biochemical and Biophysical Reserch Communication*, 210(2), 424-433.
- 3) Balázs, A., et al., 2016. Hormonal activity, cytotoxicity and developmental toxicity of UV filters. *Ecotoxicology and Environmental Safety*, 131, 45-53.
- 4) Balmer, M.E., et al., 2005. Occurrence of some organic UV filters in wastewater, in surface waters, and in fish from Swiss lakes. *Environmental Science & Technology*, 39, 953-962
- 5) Blüthgen, N., Zucchi, S., and Fent, K., 2012. Effects of the UV filter benzophenone-3 (oxybenzone) at low concentrations in zebrafish (*Danio rerio*). *Toxicology and Applied Pharmacology* 263, 184-194.
- 6) Bonassi, S., et al., 2004. Chromosomal aberrations and risk of cancer in humans: an epidemiologic perspective. *Cytogenetic and Genome Research*. 104, 376-382.
- 7) Bonassi, S, El-Zein, R. and Bolognesi, C., 2011. Micronuclei frequency in peripheral blood lymphocytes and cancer risk: evidence from human studies. *Mutagenesis* 26, 93-100.
- 8) Buck Louis, G.M., et al., 2014. Urinary Concentrations of Benzophenone-Type Ultraviolet Radiation Filters and Couples' Fecundity. *American Journal of Epidemiology*, 180(12), 1168-1175
- 9) Buck Louis, G.M., et al., 2015. Urinary concentrations of benzophenone-type ultra violet light filters and semen quality. *Fertility and Sterility*, 104, 989-996.
- 10) Calafat, A.M., et al., 2008. Concentrations of the sunscreen agent benzophenone-3 in residents of the United States: National Health and Nutrition Examination Survey 2003-

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
2004. *Environmental Health and Perspectives*, 116, 893-897.
- 11) Careghini, A., et al., 2015. Bisphenol A, nonylphenols, benzophenones, and benzotriazoles in soils, groundwater, surface water, sediments, and food: a review. *Environmental Science and Pollution Research*, 22, 5711-5741.
- 12) Chisvert, A., et al., 2012. An overview of the analytical methods for the determination of organic ultraviolet filters in biological fluids and tissues. *Analytica Chimica Acta*, 752:11-29.
- 13) Concil of Europe-Report (2009). Policy Statement Concerning paper and board materials and articles intended to come into contact with foodstuffs. Version 4, pag. 21. Available on <https://rm.coe.int/CoERMPublicCommonSearchServices/DisplayDCTMContent?documentId=09000016804e4794> accessed in date 07-29-2017.
- 14) Coronado, M., et al., 2008. Estrogenic activity and reproductive effects of the UV-filter oxybenzone (2-hydroxy-4-methoxyphenyl-methanone) in fish. *Aquatic Toxicology*, 90(3), 182-187.
- 15) Dewalque, L., et al., 2014. Simultaneous determination of some phthalate metabolites, parabens and benzophenone-3 in urine by ultra high pressure liquid chromatography tandem mass spectrometry. *Journal of Chromatography B*, 949-950, 37-47.
- 16) Fenech, M., et al., 2003. HUMAN MICRONUCLEUS PROJECT (HUMN PROJECT): detailed description of the scoring criteria for the cytokinesis-block micronucleus assay using isolated human lymphocyte cultures. *Mutation Research*, 534(1-2), 65-75.
- 17) Fent, K., et al., 2010. A tentative environmental risk assessment of the UV-filters 3-(4-methylbenzylidene-camphor), 2-ethyl-hexyl-4-trimethoxycinnamate, benzophenone-3, benzophenone-4 and 3-benzylidene camphor. *Marine Environmental Research*, 69(1), S4-S6.
- 18) French, J.E., 1992. NTP Technical Report on Toxicity Studies of 2-hydroxy-4-methoxy-benzophenone. (1992) National Toxicology Report No. 21, NIH Publication No. 92, 1-52,

- 1  
2 available at [https://ntp.niehs.nih.gov/ntp/htdocs/st\\_rpts/tox021.pdf](https://ntp.niehs.nih.gov/ntp/htdocs/st_rpts/tox021.pdf). Accessed in date 07-29-  
3  
4 2017.  
5  
6  
7 19) Gago-Ferrero, P., Díaz-Cruz, M.S., and Barceló, D., 2011. Fast pressurized liquid extraction  
8 with in-cell purification and analysis by liquid chromatography-tandem mass spectrometry  
9 for the determination of UV filters and their degradation products in sediments. *Analytical*  
10 and *Bioanalytical Chemistry*, 400, 2195-2204.  
11  
12  
13  
14  
15 20) Gago-Ferrero, P., and Diaz-Cruz, M.S., 2012. An overview of UV-absorbing compounds  
16 (organic UV filters) in aquatic biota. *Analytical and Bioanalytical Chemistry*, 404, 2597-  
17 2610.  
18  
19  
20  
21  
22 21) Hanson, K.M., Gratton, E., and Bardeen, C.J., 2006. Sunscreen enhancement of UV-induced  
23 reactive oxygen species in the skin. *Free Radical Biology and Medicine* 41, 1205-1212.  
24  
25  
26  
27 22) Kameda, Y., Kimura, K., and Miyazaki, M., 2011. Occurrence and profiles of organic sun-  
28 blocking agents in surface waters and sediments in Japanese rivers and lakes. *Environmental*  
29 *Pollution*, 159(6), 1570-1576.  
30  
31  
32  
33 23) Kim, S., and Choi, K. 2014. Occurrences, toxicities, and ecological risk of benzophenone-3,  
34 a common component of organic sunscreen products: a mini-review. *Environmental*  
35 *International*, 70:143-157.  
36  
37  
38  
39 24) Kunisue, T., et al., 2012. Urinary concentrations of benzophenone-type UV filters in U.S.  
40 women and their association with endometriosis. *Environmental Science & Technology*, 46,  
41 4624-4632.  
42  
43  
44  
45  
46 25) International Agency for Research on Cancer (IARC). (2013) Some Chemicals Present in  
47 Industrial and Consumer Products, Food and Drinking Water. 101, 285-304.  
48  
49  
50  
51 26) León, Z., et al., 2010. Solid-phase extraction liquid chromatography-tandem mass  
52 spectrometry analytical method for the determination of 2-hydroxy-4-  
53 methoxybenzophenone and its metabolites in both human urine and semen. *Analytical and*  
54 *Bioanalytical Chemistry*, 398, 831-843.  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 27) Marazzi, M., et al., 2016. Hydrogen abstraction by photoexcited benzophenone: consequences for DNA photosensitization. *Physical Chemistry Chemical Physics*. 18, 7829-7836.
- 28) OECD (Organization for Economic Co-operation and Development). Guidance Documents on Revisions to OECD Genetic Toxicology Test Guidelines. [https://www.oecd.org/env/ehs/testing/Draft Guidance Document on OECD Genetic Toxicology Test Guidelines.pdf](https://www.oecd.org/env/ehs/testing/Draft%20Guidance%20Document%20on%20OECD%20Genetic%20Toxicology%20Test%20Guidelines.pdf). 2015; 31. Accessed in date 07-29-2017.
- 29) Phiboonchaiyanan, P., et al., 2017. Benzophenone-3 increases metastasis potential in lung cancer cells via epithelial to mesenchymal transition. *Cell Biology and Toxicology*, 33, 251-261.
- 30) Philippat, C., et al., 2012. Exposure to phthalates and phenols during pregnancy and offspring size at birth. *Environmental Health Perspectives*, 120, 464-470.
- 31) Philippat, C., et al., 2015. Exposure to select phthalates and phenols through use of personal care products among Californian adults and their children. *Environmental Research*, 140, 369-376.
- 32) Pollack, A.Z., et al., 2015. Bisphenol A, benzophenone-type ultraviolet filters, and phthalates in relation to uterine leiomyoma. *Environmental Research*, 137, 101-107.
- 33) Robison, S.H., et al., 1994. Assessment of the in vivo genotoxicity of 2-hydroxy 4-methoxybenzophenone. *Environmental and Molecular Mutagenesis*, 23(4):312-317.
- 34) Rodil, R., et al., 2009. Photostability and phytotoxicity of selected sunscreen agents and their degradation mixtures in water. *Analytical and Bioanalytical Chemistry*, 395(5), 1513-1524.
- 35) Santovito, A., Cannarsa, E., Schleicherova, D. and Cervella, P. (2017) Clastogenic effects of bisphenol A on human cultured lymphocytes. *Human & Experimental Toxicology*, Jan 1:960327117693069. doi: 10.1177/0960327117693069. [Epub ahead of print]
- 36) Savage, J.R., 2004. On the nature of visible chromosomal gaps and breaks. *Cytogenetics*

- 1  
2 and Genome Research, 104(1-4): 46-55.  
3  
4 37) Schlumpf, M., et al., 2001. In vitro and in vivo estrogenicity of UV screens. Environmental  
5 Health Perspectives, 109, 239-244.  
6  
7  
8 38) Suzuki, T., et al., 2005. Estrogenic and antiandrogenic activities of 17 benzophenone  
9 derivatives used as UV stabilizers and sunscreens. Toxicology and Applied Pharmacology,  
10 203, 9-17.  
11  
12  
13 39) Vione, D., et al., 2013. Phototransformation of the sunlight filter benzophenone-3 (2-  
14 hydroxy-4-methoxybenzophenone) under conditions relevant to surface waters. Science of  
15 the Total Environment, 463-464, 243-251.  
16  
17  
18 40) Wang, L., and Kannan, K. (2013). Characteristic profiles of benzophenone-3 and its  
19 derivatives in urine of children and adults from the United States and China. Environmental  
20 Science & Technology, 47, 12532–12538.  
21  
22  
23 41) Wnuk, A., et al., 2017. Apoptosis Induced by the UV Filter Benzophenone-3 in Mouse  
24 Neuronal Cells Is Mediated via Attenuation of  $E_{\alpha}/P_{\text{par}}\gamma$  and Stimulation of  $E_{\beta}/G_{\text{pr}}30$   
25 Signaling. Molecular Neurobiology, 1-22 early view. doi: 10.1007/s12035-017-0480-z.  
26  
27  
28 42) Wolff, M.S., et al., 2007. Pilot study of urinary biomarkers of phytoestrogens, phthalates,  
29 and phenols in girls. Environmental Health Perspectives, 115, 116-121.  
30  
31  
32 43) Wolff, M.S., et al., 2008. Prenatal phenol and phthalate exposures and birth outcomes.  
33 Environmental Health Perspectives, 116, 1092-1097.  
34  
35  
36 44) Xiao, M., et al., 2013. Transformation mechanism of benzophenone-4 in free chlorine  
37 promoted chlorination disinfection. Water Research, 47(16), 6223-6233.  
38  
39  
40 45) Ye, A.M. et al., 2008 a. Automated on-line column-switching HPLC–MS/MS method with  
41 peak focusing for measuring parabens, triclosan, and other environmental phenols in human  
42 milk. Analytica Chimica Acta, 622, 150-156.  
43  
44  
45 46) Ye, X., et al., 2008 b. Automated on-line column-switching HPLC–MS/MS method for  
46 measuring environmental phenols and parabens in serum. Talanta, 76, 865-871.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 47) Zhang, T., et al., 2013. Benzophenone-type UV filters in urine and blood from children,  
3  
4 adults, and pregnant women in China: partitioning between blood and urine as well as  
5  
6 maternal and fetal cord blood. *Science of the Total Environment*, 461-462, 49–55.  
7

8  
9 48) Zeiger, E., et al., 1987. Salmonella mutagenicity tests: III. Results from the testing of 255  
10  
11 chemicals. *Environmental Mutagenesis* 9(9), 1-109.  
12  
13

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only

Table 1 - Induction of chromosomal aberrations by 2-Hydroxy-4-methoxybenzophenone in human lymphocytes *in vitro*.

Test substance	Treatment		Total Scored Metaphases	CAs								Total CAs	Total CAs + Gaps	Total Ab.C	Total Ab.C + Gaps	CAs/Cell ± S.E. (%)	Ab.C/Cell ± S.E. (%)	
	Period (h)	Dose (µg/ml)		Gaps	B'	B''	DC	R	TR	AF	Re							
NC	----	----	1000	5	6	0	1	0	0	0	8	0	15	21	15	21	1.500±0.158	1.500±0.158
0.1% DMSO	24	----	1000	10	9	0	2	0	0	0	8	2	21	31	21	31	2.100±0.367	2.100±0.367
MMC	24	0.10	1000	39	37	31	12	8	18	21	17	144	183	136	175	14.400±1.336 <sup>a</sup>	13.600±1.177 <sup>a</sup>	
BP-3	24	0.20	1000	32	32	21	5	3	0	24	16	101	133	99	131	10.100±0.510 <sup>b</sup>	9.900±0.400 <sup>b</sup>	
BP-3	24	0.10	1000	22	32	11	4	4	1	26	15	93	115	93	115	9.300±0.875 <sup>b</sup>	9.300±0.875 <sup>b</sup>	
BP-3	24	0.05	1000	13	21	11	2	0	0	24	16	74	87	71	84	7.400±1.308 <sup>b</sup>	7.100±1.134 <sup>b</sup>	
BP-3	24	0.025	1000	13	20	7	2	2	0	31	11	73	85	72	85	7.300±1.446 <sup>b</sup>	7.200±1.347 <sup>b</sup>	
BP-3	24	0.0125	1000	10	11	5	0	1	0	17	1	35	45	35	45	3.500±0.548 <sup>c</sup>	3.500±0.548 <sup>c</sup>	

CAs = chromosomal aberrations; Ab.C = aberrant cells (cells with 1 ore more aberrations); NC = Negative Control; MMC = Mitomycin-C; BP-3 = 2-Hydroxy-4-methoxybenzophenone; B': chromatid break; B'': chromosome break; DC: dicentric; R: ring; TR = tri-tetradials; AF = acentric fragments; Re = rearrangements; S.E. = standard error.

<sup>a</sup>Significantly different with respect to DMSO control solvent  $P = 0.009$ ; <sup>b</sup>Significantly different with respect to DMSO control solvent  $P = 0.008$ ; <sup>c</sup>Significantly different with respect to DMSO control solvent  $P = 0.043$

Table 2 – Induction of micronuclei by 2-Hydroxy-4-methoxybenzophenone in human lymphocytes *in vitro*.

Test substance	Treatment Period (h)	Dose (µg/ml)	BNCs scored	Distribution of BNCs according to the number of MNs				MNs	Ab.C	MNs/cell ± S.E. (%)	Ab.C/cell ± S.E. (%)	CBPI ± S.E
				1	2	3	4					
NC	----	----	5000	11	0	0	0	11	11	0.220±0.000	0.220±0.000	1.732±0.025
0.1% DMSO	48	----	5000	16	0	0	0	16	16	0.320±0.000	0.320±0.000	1.811±0.062
MMC	48	0.10	5000	135	12	8	0	183	155	3.660±0.005 <sup>a</sup>	2.675±0.003	1.345±0.020
BP3	48	0.20	5000	69	4	0	0	77	73	1.540±0.001 <sup>a</sup>	1.525±0.002 <sup>a</sup>	1.684±0.017
BP3	48	0.10	5000	70	4	1	0	81	75	1.620±0.002 <sup>a</sup>	1.425±0.002 <sup>a</sup>	1.603±0.041
BP3	48	0.05	5000	44	1	0	0	46	45	0.920±0.001 <sup>a</sup>	0.850±0.001 <sup>a</sup>	1.690±0.033
BP3	48	0.025	5000	41	3	0	0	47	44	0.940±0.001 <sup>b</sup>	0.880±0.001 <sup>b</sup>	1.696±0.029
BP3	48	0.0125	5000	31	1	0	0	33	32	0.660±0.002 <sup>c</sup>	0.640±0.002 <sup>c</sup>	1.710±0.067

BNCs = Binucleated cells; MNs = micronuclei; Ab.C = Aberrant cells (cells with 1 or more MNs); NC = Negative Control;

MMC = Mitomycin-C; BP-3 = 2-Hydroxy-4-methoxybenzophenone; S.E. = Standard Error. CBPI = Cytokinesis-Block Proliferation Index.

<sup>a</sup> Significantly differ with respect to DMSO control solvent P = 0.008; <sup>b</sup> Significantly differ with respect to DMSO control solvent P = 0.011;

<sup>c</sup> Significantly differ with respect to DMSO control solvent P = 0.025



Table 3 - Multiple regression analysis evaluating the relationship between BP-3 concentrations and the level of genomic damage

<b>Biomarkers</b>	<b><math>\beta</math>-co</b>	<b><i>P</i>-value</b>	<b>95% CI</b> (Lower) – (Upper)
CAs	0.705	<0.001	(1.719) – (4.361)
Cells with CAs	0.723	<0.001	(1.751) – (4.209)
MNs	0.690	<0.001	(1.335) – (3.545)
Cells with MNs	0.724	<0.001	(1.332) – (3.188)
CBPI	0.227	0.274	(-0.041) – (-0.012)

CAs = Chromosomal Aberrations; MNs = micronuclei; CBPI = Cytokinesis-Block Proliferation Index;  $\beta$ -co =  $\beta$ -coefficient; CI = Confidence Interval