

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Everything is (still) illuminated: dual right cathodal-left anodal tDCS of PPC prevents fatigue on a visual detection task

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1677489> since 2018-10-02T19:56:30Z

Published version:

DOI:10.1016/j.brs.2018.09.017

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Accepted Manuscript

Everything is (still) illuminated: dual right cathodal-left anodal tDCS of PPC prevents fatigue on a visual detection task

Pietro Sarasso, Marco Ninghetto, Adriana Salatino, Irene Ronga, Alessandro Bongiardina, Igor Iarrobino, Marco Neppi-Modona, Raffaella Ricci



PII: S1935-861X(18)30319-X

DOI: [10.1016/j.brs.2018.09.017](https://doi.org/10.1016/j.brs.2018.09.017)

Reference: BRS 1326

To appear in: *Brain Stimulation*

Received Date: 18 September 2018

Accepted Date: 27 September 2018

Please cite this article as: Sarasso P, Ninghetto M, Salatino A, Ronga I, Bongiardina A, Iarrobino I, Neppi-Modona M, Ricci R, Everything is (still) illuminated: dual right cathodal-left anodal tDCS of PPC prevents fatigue on a visual detection task, *Brain Stimulation* (2018), doi: <https://doi.org/10.1016/j.brs.2018.09.017>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Everything is (still) illuminated:**dual right cathodal-left anodal tDCS of PPC prevents fatigue on a visual detection task**

Pietro Sarasso^{1*}, Marco Ninghetto¹, Adriana Salatino¹, Irene Ronga¹, Alessandro Bongiardina^{1§},
Igor Iarrobino^{1§}, Marco Neppi-Modona¹, Raffaella Ricci^{1*}

¹ SAMBA (SpAtial, Motor & Bodily Awareness) Research Group, Department of Psychology,
University of Turin, Italy

**Corresponding Authors:*

Raffaella Ricci

raffaella.ricci@unito.it

Pietro Sarasso

pietro.sarasso@unito.it

Dipartimento di Psicologia, Università degli Studi di Torino

Via Po, 14

10123 – Torino

Italy

§ These authors contributed equally to the study

Keywords: tDCS, visual detection, fatigue, attention

Dear Editor,

it is known that transcranial Direct Current Stimulation (tDCS) may affect attentional processing when applied to the right posterior parietal cortex (PPC) in line with current evidence on the neural bases of this cognitive function and their modulation by Transcranial Magnetic Stimulation [1]. However, research results are conflicting: some, in line with the interhemispheric rivalry account of attention [2], indicate that tDCS over PPC biases attention contra- or ipsilaterally depending upon whether (facilitatory) anodal or (inhibitory) cathodal tDCS is applied to PPC, respectively; others suggest that PPC controls attention across the entire visual field. Anodal tDCS of the right PPC boosts orienting of spatial attention leftwards [3] and enhances detection of left-sided visual targets [4], while cathodal tDCS of the right PPC induces a rightward attentional bias on line-length estimation [5]. Coherently, the application of bi-parietal, right anodal-left cathodal tDCS ameliorates detection of left-sided visual stimuli and deteriorates detection of right-sided stimuli [6]. By contrast, other findings show that cathodal stimulation of the right PPC ameliorates attentional selection of visual stimuli across the entire field [7], or that dual tDCS over PPC worsens object-motion tracking, regardless of hemifield and stimulation polarity [8].

To further investigate how modulation of PPC excitability might affect attentional performance, we employed a visual detection task that is very sensitive to attentional modulation [9], before and after bi-parietal tDCS. We asked whether dual tDCS of PPC would affect visual attention in a polarity-specific manner as predicted by the interhemispheric rivalry account of attention [2].

Forty-five right-handed healthy volunteers (24 women, mean age 23 ± 2.7) gave written informed consent to participate in the study that was approved by the ethics committee of the University of Turin. Participants seated in a dark room, 50 cm

away from a computer screen. Visual stimuli were 0.4 cm x 0.6 cm rectangles randomly presented for 50 ms (ISI: 2000 to 2500 ms) in 18 different locations distributed along the horizontal (3, 6 and 9 cm to the left or right of the central cross) and vertical (0.5 cm above or below the fixation cross) axes. There were ten possible levels of stimulus luminance, ranging from 0.8% to 1.2% of the screen maximum output. Individual subjective visual thresholds (i.e. detection rate between 40 and 60%) were determined during the training phase. During the experiment, 162 stimuli were presented prior and 162 after 13 minutes of stimulation. Stimuli were flashed at luminance threshold level (1/3), or one level below (1/3) or above (1/3) threshold. Participants pressed the keyboard spacebar, with their right index, as soon as a stimulus was detected while fixating the central cross (Fig. 1A, B). TDCS was applied at 1.5 mA by means of a *Newronika HDCstimulator* (Newronika s.r.l.) for 15 minutes. Electrodes (25 cm²), covered with conductive rubber and saline-soaked synthetic sponges, were positioned over P4 and P3 (10-20 EEG system). Participants were assigned to one of the following conditions (N=15): 1) Right Anodal-Left Cathodal tDCS (RA-LC); 2) Right Cathodal-Left Anodal tDCS (RC-LA); 3) sham stimulation (ramp-up period: 30 s).

A three-way repeated-measures ANOVA was performed on accuracy (ACC, number of detected stimuli) as dependent variable with *Time* (*pre*, *post-tDCS*) and *Side* of presentation (*left*, *right*) as within-subjects factors, and *Group* as between-subjects factor (*RC-LA*, *RA-LC*, *Sham*). The ANOVA showed significant effects of *Time* ($F_{1,42} = 15.02$, $p < 0.0001$, partial $\eta^2 = 0.263$), with higher ACC pre (94.998 ± 51.264) than post-tDCS (76.182 ± 54.78), and *Side* ($F_{1,42} = 25.57$, $p < 0.0001$; partial $\eta^2 = 0.378$), with higher ACC for right (89.376 ± 50.034) than left side (81.264 ± 57.276). Significant interactions *Time x Group* ($F_{2,42} = 3.814$, $p = 0.030$, partial $\eta^2 = 0.154$) and *Time x Side* ($F_{1,42} = 4.926$, $p = 0.032$; partial $\eta^2 = 0.105$) were also observed. Post-hoc

analyses (independent-samples t-test) showed that *Time* was significantly different only between *RC-LA* and *RA-LC* ($p=0.025$). To further analyze the effect of tDCS polarity, separate ANOVAs were performed for each group, with *Time* and *Side* as within-subjects factors (Fig. 1C). The *RA-LC* group showed significant effects of *Time* ($F_{1,14}= 12.611$, $p= 0.003$, partial $\eta^2= 0.474$), with higher detection rate pre than post-tDCS, and a significant interaction *Time* \times *Side* ($F_{1,14}= 4.709$, $p= 0.048$, partial $\eta^2= 0.252$). Post-hoc analyses revealed that pre and post-tDCS ACC differed to a lesser degree for left-sided (left-pre: 45.933 ± 11.689 , left-post: 29.467 ± 19.982) than right-sided stimuli (right-pre: 50.8 ± 9.182 , right-post: 29.267 ± 19.668 ; $t_{14}= 2.17$, $p=0.048$). The *RC-LA* group showed a significant effect of *Side* ($F_{1,14}= 5.755$, $p=0.031$, partial $\eta^2= 0.291$; i.e. better detection for right-sided than left-sided stimuli), but *Time* was not significant ($F_{1,14}=0.332$, $p=0.574$, partial $\eta^2= 0.023$). The *Sham* group manifested significant effects for *Time* ($F_{1,14}= 6.91$, $p= 0.020$, partial $\eta^2= 0.330$) and *Side* ($F_{1,14}= 21.931$, $p<0.0001$, partial $\eta^2= 0.610$). Participants showed better detection pre than post-tDCS and for right-sided than left-sided stimuli.

Figure 1 about here

Overall, we observed better detection for right-sided stimuli. This rightward bias replicates findings of our previous study [9] and has been previously described in healthy individuals as evidence of the dominance of the left hemisphere in the detection of *transient* visual stimuli [10]. Moreover, we found decreased accuracy bilaterally after *Sham* and *RA-LC* stimulation, likely due to a deterioration of sustained attention during the demanding detection task. Surprisingly, fatigue was prevented by *RC-LA* stimulation. This result, in line with Moos et al.'s finding [16], indicates that 'inhibitory'/cathodal stimulation of right PPC associated with 'facilitatory'/anodal stimulation of left PPC boosts (sustained) attention in both hemifields rather than in the right hemifield only, as expected on the basis of simple

predictions of tDCS polarity-dependent effects on interhemispheric attentional control. Future studies with large sample size, including stimulation of a non-parietal site as control condition, are needed to validate these preliminary findings and clarify, using combined neuroimaging, whether they reflect potentiation of left PPC neural activity [10] by anodal stimulation, or reduction of neural noise in right PPC by cathodal tDCS, or both. These findings may have important theoretical implications for translational research.

Acknowledgements

We would like to thank Martina Tinnirello, Chiara Rubino, and Azzurra Milano for helping with participants' recruitment and data collection.

Funding

The study was supported by a MIUR grant (RICR_RILO_17_01) to RR.

Conflict of interests

The authors report no actual or potential conflicts of interests.

REFERENCES

- [1] Salatino A, Poncini M, George MS, Ricci R. Hunting for right and left parietal hot spots using single-pulse TMS: Modulation of visuospatial perception during line bisection judgment in the healthy brain. *Front Psychol* 2014;5:1238. doi:10.3389/fpsyg.2014.01238.
- [2] Kinsbourne M. Hemi-neglect and hemisphere rivalry. *Adv Neurol* 1977; 18:41-9.
- [3] Roy LB, Sparing R, Fink GR, Hesse MD. Modulation of attention functions by anodal tDCS on right PPC. *Neuropsychologia* 2015;74:96-107. doi:10.1016/j.neuropsychologia.2015.02.028.
- [4] Filmer HL, Dux PE, Mattingley JB. Dissociable effects of anodal and cathodal tDCS reveal distinct functional roles for right parietal cortex in the detection of single and competing stimuli. *Neuropsychologia* 2015;74:120-6 doi:10.1016/j.neuropsychologia.2015.01.038.
- [5] Giglia G, Mattaliano P, Puma A, Rizzo S, Fierro B, Brighina F. Neglect-like effects induced by tDCS modulation of posterior parietal cortices in healthy subjects. *Brain Stimul* 2011, 4(4):294-9. doi:10.1016/j.brs.2011.01.003.
- [6] Sparing R, Thimm M, Hesse MD, Küst J, Karbe H, Fink GR. Bidirectional alterations of interhemispheric parietal balance by non-invasive cortical stimulation. *Brain* 2009;132(11):3011-20. doi:10.1093/brain/awp154.
- [7] Moos K, Vossel S, Weidner R, Sparing R, Fink GR. Modulation of Top-Down Control of Visual Attention by Cathodal tDCS over Right IPS. *J Neurosci* 2012; 32(46):16360-16368. doi:10.1523/JNEUROSCI.6233-11.2012.
- [8] Roe JM, Nesheim M, Mathiesen NC, Moberget T, Alnæs D, Sneve MH. The effects of tDCS upon sustained visual attention are dependent on cognitive load. *Neuropsychologia* 2016; 80:1-8. doi:10.1016/j.neuropsychologia.2015.11.005.

- [9] Ronga I, Sarasso P, Fossataro C, Salatino A, Garbarini F, Ricci R, Neppi-Modona M. Everything is illuminated: Prismatic adaptation lowers visual detection threshold in normal subjects. *J Exp Psychol Hum Percept Perform*. 2018 Jul 30. doi: 10.1037/xhp0000559
- [10] Reuter-Lorenz P, Kinsbourne M, Moscovitch M. Hemispheric control of spatial attention. *Brain Cogn* 1990;12(2):240-266. doi:10.1016/0278-2626(90)90018-J.

Figure Caption.

Figure 1. Experiment method and results. Panel A shows three examples of the visual stimuli employed in the present experiment (original proportions are not preserved in the figure); in Panel B the experiment timeline; Panel C shows participants' performance (Accuracy) in the detection task. Mean numbers of detected stimuli (out of a total of 162) and relative standard error for the three groups of participants (RC-LA= right cathodal-left anodal, RA-LC= right anodal-left cathodal, SHAM) in pre and post-tDCS conditions are reported. Stars indicate significant differences between pre and post stimulation condition for the RA-LC and the SHAM groups.





Figure 1