

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Artificial lighting triggers the presence of urban spiders and their webs on historical buildings

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1676420> since 2019-03-08T11:48:06Z

*Published version:*

DOI:10.1016/j.landurbplan.2018.09.003

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Manuscript Number: LAND-D-18-00321R1

Title: Artificial lighting triggers the presence of urban spiders and their webs on historical buildings

Article Type: Research paper

Keywords: urban ecology; digital image analysis; light pollution; Araneae; synanthropic species; aesthetic nuisance

Corresponding Author: Dr. Marco Isaia, Ph.D.

Corresponding Author's Institution: University of Torino

First Author: Stefano Mammola, Ph.D.

Order of Authors: Stefano Mammola, Ph.D.; Marco Isaia, Ph.D.; Daniele Demonte; Paolo Triolo; Marco Nervo

Abstract: Different spider species living in the urban environment spin their webs on building facades. Due to air pollution, web aggregations entrap dirt particles over time, assuming a brownish-greyish colouration and thus determining an aesthetic impact on buildings and street furniture. In Europe, the most common species causing such an aesthetic nuisance is *Brigittea civica* (Lucas) (Dictynidae). In spite of the socio-economical relevance of the problem, the ecological factors driving the proliferation of this species in the urban environment are poorly described and the effectiveness of potential cleaning activities has never been discussed in scientific literature. Over one year, we studied the environmental drivers of *B. civica* webs in the arcades of the historical down-town district of Turin (NW-Italy). We selected a number of sampling plots on arcade ceilings and we estimated the density of *B. civica* webs by means of digital image analysis. In parallel, we collected information on a number of potential explanatory variables driving the arcade colonization, namely artificial lighting at night, substrate temperature, distance from the main artificial light sources and distance from the river. Regression analysis showed that the coverage of spider webs increased significantly at plots with higher light intensity, with a major effect related to the presence of historical lampposts with incandescent lamps rather than halogen lamps. We also detected a seasonal variation in the web coverage, with significant higher values in summer. Stemming from our results, we are able to suggest good practices for the containment of this phenomenon.



---

Dr. Marco Isaia  
Department of Life Science and Systems Biology  
University of Turin  
Via Accademia Albertina, 13 - 10123 Torino  
Tel. 0116704519  
E-mail: [marco.isaia@unito.it](mailto:marco.isaia@unito.it)

May 07<sup>th</sup>, 2018

Dear Dr Xiang,

we hereby submit the revised version of manuscript LAND-D-18-00321 entitled “**Artificial lighting triggers the presence of urban spiders and their webs, causing an aesthetic damage to historical buildings**” intended for publication in *Landascape and Urban Planning*.

We thank you for handling our submission and spending time in evaluating it. In this revision, we have took into account all minor corrections suggested by referee 2 and proof-read it once more.

We confirm that the work is all original research carried out by the authors, all authors agree with the contents of the manuscript and its submission to the journal, no part of the research has been published in any form elsewhere, and the manuscript is not being considered for publication elsewhere.

Yours,

Dr Marco Isaia

On behalf of all co-authors

1 REBUTTAL LETTER FOR

2  
3 LAND-D-18-00321

4  
5 'Artificial lighting triggers the presence  
6 of urban spiders and their webs on historical buildings'

8  
9

10 Reviewer #2: Landscape and Urban Planning MS 18-0321

11  
12 This seems to be a respectable contribution suitable for publication. It is a nice combination of science  
13 addressing a real world problem. It also is a relatively clean paper. I usually find lots of mistakes in  
14 manuscripts that I review but this one had few errors. One caveat is that I have very limited statistical  
15 background so the editor needs to make sure that someone with statistical ability reviews the paper.

16  
17 **RESPONSE: Thank you for spending time to review our manuscript and for you very positive attitude  
18 toward it.**

19  
20 There are several instances of misspelling the spider family name as "Dyctinidae" which needs correcting  
21 (lines 45, 505, 506)

22  
23 **RESPONSE: Corrected.**

24  
25 line 47: cited as Hertel 1969 in the text but 1968 in the reference section

26  
27 **RESPONSE: Corrected – the right one was 1968.**

29 line 217: citation is "XXXXXXX" but is probably Hanggi which is not cited in the paper and is also 2016

30  
31 **RESPONSE: Actually, this is not a mistake. From the journal's guidelines (see highlight in colour):**

32  
33 "Landscape and Urban Planning uses a double-blind review process, and to ensure anonymity the  
34 manuscript file must not include any self-referencing, logos, headers or any other type of information  
35 or formatting that might reveal the identify or affiliation of any of the authors. Acknowledgements  
36 should not be included in the manuscript file and must be uploaded as a separate file. See Section  
37 3.9.7 below. Self-references that must be included must not be obvious in revealing any authors'  
38 identify and should refer to the authors' work only indirectly (e.g., "This work builds upon  
39 procedures developed by Smith (2010)"; NOT "I build upon my previous work (Smith, 2010)..."). **To  
40 further ensure anonymity, authors may choose to temporarily remove self-citations from the  
41 reference list and mask in-text references (e.g., "(XXX, 2009 masked for blind review)"), then restore  
42 the proper citation when the manuscript is accepted.** Although such an approach better respects the  
43 integrity of the blind review process, authors must weigh the removal of a citation against the need  
44 for reviewers to evaluate the credibility of your work."

45  
46 **On the other hand, we now reference the missing reference in the text (Hanggi).**

47  
48 line 62: change "environment" to "environments"  
49 line 152: insert a space in "Photo#2"  
50 line 175: change "generated" to "generate"  
51 line 270: change "them" to "themselves"  
52 line 287: I would change "Southern" to "southern"  
53 line 350: change "spider" to "spiders"  
54 line 352: I would change "Southern" to "southern"  
55 line 484: insert a space in "seriesmodel"

56 line 541: many other citations have a long list of authors, whereas this one just has the first author and et al.  
57 line 590s: Vetter paper should be listed before Vitousek  
58 line 620: add a space in "typeanalyses"  
59  
60 **RESPONSE: Done.**

1 **Artificial lighting triggers the presence of urban spiders and their webs on historical**  
2 **buildings**

3  
4 Stefano MAMMOLA<sup>1,2</sup>

5 stefanomammola@gmail.com

6  
7 Marco ISAIA<sup>2,\*</sup>

8 marco.isaia@unito.it

9 tel: 0116704544

10 fax: 0116704508

11

12 Daniele DEMONTE<sup>1</sup>

13 daniele.demonte@centrorestaurovenaria.it

14

15 Paolo TRIOLO<sup>1</sup>

16 paolo.triolo@centrorestaurovenaria.it

17

18 Marco NERVO<sup>1</sup>

19 marco.nervo@centrorestaurovenaria.it

20

21

22

23 1. Fondazione Centro per la Conservazione ed il Restauro dei Beni Culturali "La Venaria  
24 Reale". Venaria Reale, Italy

25

26 2. Department of Life Sciences and Systems Biology, University of Turin. Torino, Italy

27

28 \* corresponding author

29

30

31

## \*Highlights (for review)

1. Spider web aggregations affect the aesthetic value of historical building
2. We studied the ecological factors driving the abundance of spider webs on historical arcade ceilings
3. Web density was estimated using an innovative photographic-based methodology
4. By attracting prey, incandescent artificial lighting is the major factor driving of the increase of spider webs
5. Changing the lighting system type may help in reducing this problem

1 **Artificial lighting triggers the presence of urban spiders and their webs on historical**  
2 **buildings**

3  
4  
5 **Abstract**

6  
7 Different spider species living in the urban environment spin their webs on building  
8 facades. Due to air pollution, web aggregations entrap dirt particles over time, assuming a  
9 brownish-greyish colouration and thus determining an aesthetic impact on buildings and  
10 street furniture. In Europe, the most common species causing such an aesthetic nuisance  
11 is *Brigittea civica* (Lucas) (Dictynidae). In spite of the socio-economical relevance of the  
12 problem, the ecological factors driving the proliferation of this species in the urban  
13 environment are poorly described and the effectiveness of potential cleaning activities has  
14 never been discussed in scientific literature. Over one year, we studied the environmental  
15 drivers of *B. civica* webs in the arcades of the historical down-town district of Turin (NW-  
16 Italy). We selected a number of sampling plots on arcade ceilings and we estimated the  
17 density of *B. civica* webs by means of digital image analysis. In parallel, we collected  
18 information on a number of potential explanatory variables driving the arcade colonization,  
19 namely artificial lighting at night, substrate temperature, distance from the main artificial  
20 light sources and distance from the river. Regression analysis showed that the coverage of  
21 spider webs increased significantly at plots with higher light intensity, with a major effect  
22 related to the presence of historical lampposts with incandescent lamps rather than  
23 halogen lamps. We also detected a seasonal variation in the web coverage, with  
24 significant higher values in summer. Stemming from our results, we are able to suggest  
25 good practices for the containment of this phenomenon.

26  
27 **Keywords:** urban ecology; digital image analysis; light pollution; Araneae; synanthropic  
28 species, aesthetic nuisance



## 29 1. INTRODUCTION

30

31 Environmental modifications driven by urbanization have a significant effect on biodiversity  
32 (Güneralp & Seto, 2013; Seto, Gueneralp, & Hutyra, 2012; Vitousek, 1997), driving large  
33 changes in species abundances and distributions within the original biological communities  
34 (McKinney, 2008). Whilst urbanization is considered to be a major determinant of  
35 biodiversity loss (Grimm et al., 2008; Newbold et al., 2015), a number of organisms are  
36 able to coexist alongside us in urban environments (e.g., Aronson et al., 2014; Bertone et  
37 al., 2016; McKinney, 2002). Owing to their high ecological plasticity (Turnbull, 1973),  
38 several species of spiders are able to dwell in cities, representing an important component  
39 of the urban wildlife (McIntyre, 2000; Shochat, Stefanov, Whitehouse, & Faeth, 2004;  
40 Taucare-Ríos, Brescovit, & Canals, 2013). With the exception of some species of medical  
41 importance (e.g. Isbister et al., 2005; Sams et al., 2001; Vetter & Isbister, 2008), urban  
42 spiders usually have little socio-economic impacts and often remain unnoticed. A  
43 remarkable exception is found in those spider that due to their webs may cause aesthetic  
44 alterations to buildings facades — see discussion in Nentwig (2015).

45 In Europe, one of the most noticeable species causing aesthetic nuisance to  
46 buildings is *Brigittea civica* (Lucas) (Araneae: Dictynidae) (Figure 1A) (Samu, Jozsa, &  
47 Csànyi, 2004). This is a small cribellate spider (body length 2.3–3.5 mm; Nentwig, Blick,  
48 Gloor, Hänggi, & Kropf, 2018) of South European origin (Hertel, 1968), which spins a  
49 circular, tangled cribellate cobweb on flat surfaces (Billaudelle, 1957; Krumpálová, 2001).  
50 Although being relatively small in size (ca. 5 cm in diameter), cobwebs of *B. civica* may  
51 occur at very high density on wall facades and can persist for long periods of time (Figure  
52 1B–D). The fact that multiple individuals are able to coexist and spin their cobwebs at a  
53 very close distance to one another can be explained in light of the peculiar behaviour of *B.*  
54 *civica*, with different individuals being able to share prey without fighting each other  
55 (Billaudelle, 1957). Due to air pollution, these large web aggregations entrap dust and dirt

56 particles over time, assuming a brownish-greyish coloration and thus significantly reducing  
57 the aesthetic value of buildings (Havlová & Hula, 2010; Kostanjšek & Celestina, 2008;  
58 Nedvěd et al., 2011; Novotný, Hula, & Niedobová, 2017; Samu et al., 2004; Figure 1D, E).

59 The factors determining this phenomenon are as yet poorly described, and thus no  
60 good practices have been put forward to address this problem and to maximize the  
61 effectiveness of potential cleaning activities. To the best of our knowledge, the only study  
62 referring to habitat selection by *B. civica* was conducted by Samu et al. (2004) in urban  
63 environments in Hungary. The authors demonstrated quantitatively how web density is  
64 significantly higher in facades with a southern exposure and sheltered to external  
65 weathering (especially rain), whereas they found no clear pattern in the selection of  
66 different surface-types.

67 Because of its artistic heritage from one side and of its predominantly  
68 Mediterranean climate suitable for *B. civica* on the other, Italy is potentially among the  
69 most affected countries by this issue. In several Italian heritage cities, webs of this spider  
70 are found on churches, arcades, palaces and other historical buildings exploited for  
71 touristic purposes, resulting in possible economic impacts connected to the cleaning  
72 activities aimed at removing the webs.

73 We conducted a one-year field study to identify the environmental factors driving the  
74 proliferation of *B. civica* webs in the arcades of the historical down-town district of Turin  
75 (NW-Italy) (Figure 1E). Observations made by the authors during a preliminary site  
76 inspection, and similar recorded observations published by Samu et al. (2004), lead to an  
77 initial prediction that i) the density of webs is significantly higher in the vicinity of artificial  
78 lighting systems and in other areas with high illuminance. Moreover, given that webs of *B.*  
79 *civica* may last for long time on the surfaces, we further hypothesize that ii) webs are  
80 present through the year, but there should be variation in their density connected with the  
81 phenology of the species. In particular, increases in the density of webs may be expected

82 right after the breeding period for this species, approximately from April to June  
83 (Kostanjšek & Celestina, 2008; Nentwig et al., 2018; Wiehle, 1953).

## 84 2. MATERIAL & METHODS

85

### 86 2.1 Study area

87 This study was conducted in the municipality of Turin (Torino), NW-Italy (45° 04' N, 7° 42'  
88 E). The city has a long history, testified by the traditional orthogonal plant of the ancient  
89 Roman camps ("*castrum*"), and the remarkable Baroque architecture, which was  
90 developed under the Duchy of Savoy (1416–1860). The down-town district of Turin has 18  
91 kilometres of historical Baroque arcades ("*portici*"), mostly interconnected with each other,  
92 making it one of the largest pedestrian areas in Europe. Arched ceilings represent suitable  
93 habitats to *Brigittea civica* spiders, being naturally sheltered from rains, direct solar  
94 irradiation, and air currents (Billaudelle, 1957; Samu et al., 2004). As a result, most  
95 arcades ceilings are heavily colonized by *B. civica* cobwebs (Figure 1B–D). In order to  
96 improve the aesthetic value of the historical district, the competent authorities promoted  
97 the cleaning of the arcades in 2006, when the city hosted the XX Olympic Winter Games.

98 We conducted our study in the heart of the historical down-town district, on the  
99 arcades connecting Piazza Vittorio Veneto to Piazza Castello all along Via Po, for  
100 approximately 2 kilometres. In this area, arcades are predominantly Serlian-type arches,  
101 ranging from 3 to 5 m in height (Figure 1E). The majority of the arcades have vaulted  
102 ceilings, with some exceptions (coffered ceilings) along a small stretch of Via Po, which we  
103 did not consider in this analysis. At night, arcades are illuminated by historical lampposts  
104 with incandescent lights and/or halogen lamps, which are installed at different heights.

105

### 106 2.2 Data collection

107 We conducted three monitoring sessions over one year (Winter: 10 Dec 2016; Spring: 30  
108 Apr 2017; Summer: 7 Jul 2017), at night. In order to provide a homogeneous coverage of  
109 the entire study area, we selected seventy-two arcades as basic sample units,

110 approximately equidistant. All sampling plots were georeferenced. Within each basic  
111 sample unit, at each survey we randomly identified one sampling plot of 1.0 x 0.7 m on a  
112 flat area of the ceiling (i.e. avoiding angles and cracks and crevices in the plaster). Using  
113 this study design, we were able to control for substrate type (all arcade ceilings are  
114 characterized by the same plaster type) and general exposure to external weather (all  
115 arcades are protected by rains, direct solar irradiation, and air currents).

116 We conducted the monitoring by means of a photography-based methodology  
117 (Figure 2), by taking two photos of each sampling plot in order to estimate the density of  
118 *Brigittea civica* webs (photo #1) and the intensity of artificial lighting illuminating the wall  
119 surface of the plot (photo #2). Full details on the calculation of these variables are given in  
120 the section "*Photographic analysis*". At each survey, we further measured the substrate  
121 temperature with an infrared thermometer at the centre of the sampling plot, and we  
122 calculated the linear distance from 1) the closest historical lamppost (incandescent lighting  
123 system) and/or 2) the closest halogen light (Figure 2). The distance of each sampling plot  
124 from the Po River, which is flowing close to the study area (see Figure 3), was further  
125 calculated in a GIS environment (ArcView 3.3 ESRI). This latter variable was introduced to  
126 account for the potential influence of local microclimatic factors related to the proximity of  
127 water on web abundance and insect prey availability — e.g. as observed in other web  
128 weaver spiders (Akamatsu, Toda, & Okino 2004, Gillespie, 1987; Kleinteich, 2010).

## 129 **2.3 Photographic analysis**

130 We carried out the photographic analysis in National Instruments<sup>TM</sup> LabVIEW environment  
131 (Elliott, Vijayakumar, Zink, & Hansen, 2007). In order to obtain a value representing the  
132 web coverage of each plot, we acquired a photo of the plot (photo #1) in raw mode with a  
133 Nikon D810 camera equipped with a Nikon sb910 flash; the image was then converted to  
134 16 bit greyscale. Initially we conducted a set of exploratory tests, applying different pixel  
135 thresholds for converting the 16 bit greyscale image into a black and white (B/W) image.  
136 The aim of the B/W conversion was to recolour the pixels depicting webs of *Brigittea civica*  
137 in black and the remaining pixels in white, thus allowing an estimation of the web coverage  
138 within the plot. After the preliminary trial, we set the B/W conversion threshold at 49000  
139 (where 0 is black and 65535 is white), which proved to be the best trade-off value to  
140 separate the webs from other features of the surface within the image — i.e. dust particles  
141 on the substrate or darker areas due to the natural ruggedness of the ceiling plaster. We  
142 used this fixed threshold for converting all images into B/W, and we summed up the  
143 number of black pixels via an automated function.

144         Despite the evaluation of the light intensity that reaches a surface is normally  
145 performed directly using a photoradiometer, in our case this resulted highly impractical due  
146 to the height of the arcades and the number of plots. Starting from the assumption that the  
147 intensity of the light reflected by a surface is correlated with the intensity of the lighting  
148 source, we used an evaluation method based on the acquisition of the reflected light with a  
149 camera. By choosing surfaces having more or less the same colour and the same surface  
150 morphology, and by setting up a fixed exposure time and no flash, it is possible to use the  
151 acquired image as an indirect measure of the light intensity of the plot. Consequently,  
152 images with low values of R, G and B channels correspond to low values of light intensity  
153 reaching the surface, while high values of R, G and B channels correspond to high values  
154 of light intensity. For this calculation, we acquired a second image of the plot (photo #2)

155 focusing on a web-free surface (Figure 2). Photo #2 was taken in raw mode with a Nikon  
156 D3X camera without flash and using a fixed exposure. We converted the digital image to a  
157 16 bit RGB, and we derived the histogram representing the distribution of the overall pixel  
158 values (range 0 – 65535). We used the modal value — i.e., the most recurrent value in the  
159 image — as an indirect measure of the artificial light intensity that illuminated the  
160 photographed surface. We repeated the same operation for each of the RGB channels in  
161 order to explore possible relationships between the web coverage and different colours of  
162 light.

163

## 164 **2.4 Regression analysis**

165 We studied the relation between environmental factors and *Brigittea civica* web coverage  
166 by performing a regression-type analysis (Zuur & Ieno, 2016) in R (R core team, 2017).  
167 We expressed the dependent variable as the counts of black pixels (i.e. pixel covered with  
168 webs) within photo #1 — hereinafter “web coverage” (WEB). We selected the following  
169 covariates (explanatory variables) as potential drivers of the web coverage in the sampled  
170 plots: distance from the river (Dst; continuous variable), substrate temperature ( $T^{\circ}$ ;  
171 continuous variable), sampling session (Sampling; categorical variable of three levels),  
172 artificial light intensity (ILL; continuous variable calculated from photo #2), Red, Green and  
173 Blue light components (R, G, B, respectively; continuous variables calculated from photo  
174 #2), distance from the nearest historical lamppost (Dst\_lamp; continuous variable) and  
175 distance from halogen light sources (Dst\_halo; continuous variable). In order to account for  
176 potential density-dependent effects in driving the web coverage (i.e. spatial relationships  
177 among webs), we further included a variable reflecting the intercrossed distance of each  
178 plot from the others (InterDst; continuous variables). To generate this variable, we  
179 calculated the distance matrix of each sampling plot using the spatial coordinates of the  
180 plots, and derived the mean distance of each plot from the others. We tested for spatial

181 autocorrelation in web coverage *via* Moran's I test in the *ape* R package (Paradis, Claude,  
182 & Strimmer, 2004), using the Gittleman and Kot (1990) method.

183 We initially explored the dataset using the standard protocol described by Zuur,  
184 leno, and Elphick (2010). We constructed Cleveland' dotplots to assess the presence of  
185 outliers within the dataset. We investigated multi-collinearity among explanatory variables  
186 by means of scatterplots, Pearson correlation tests ( $r$ ), by setting the threshold for  
187 collinearity at  $r > |0.7|$  (Zuur, leno, Walker, Savaliev, & Smith, 2009). Boxplots were also  
188 constructed to graphically assess collinearity between continuous and categorical  
189 variables.

190 To model the response of the web coverage to the explanatory variables, we initially  
191 fitted a Poisson generalised linear model (GLM), including all non-collinear covariates of  
192 interest. In order to test for the potential effect of the different light source illuminating the  
193 plot, we allowed for interactions between the artificial light intensity and the distance from  
194 the different light sources ( $\text{Dst\_lamp} * \text{ILL}$ ;  $\text{Dst\_halo} * \text{ILL}$ ). The Poisson GLM was highly  
195 over-dispersed [dispersion statistic (DS)=600.95], and thus a negative binomial GLM was  
196 considered (Zuur et al., 2010). We fitted the negative binomial GLM in the *MASS* R  
197 package (Venables & Ripley, 2002). Over-dispersion in the negative binomial GLM was  
198 minimal (DS=1.62), so we chose this error distribution in all subsequent analysis.

199 Once we fitted the initial negative binomial GLM including all the covariates and  
200 interactions of interest, we performed model selection in order to select which variables  
201 should be included in the final model (Johnson & Omland, 2004). We used a stepwise  
202 backward elimination procedure, whereby we progressively excluded variables and  
203 interactions from the model according to the corrected Akaike information criterion for finite  
204 sample size (AICc values; Burnham & Anderson, 2002; Hurvich & Tsai, 1989). We  
205 reiterated the procedure until we obtained a Minimum Adequate Model (MAM) including



206 only significant variables. We conducted model selection using the *MuMIn* R package  
207 (Bartoń, 2017).

208 Model validation was conducted on the MAM (Zuur et al., 2009). In particular, we  
209 tested it for over-dispersion, we constructed standard validation plots using model  
210 residuals and we investigated the existence of possible non-linear responses of our  
211 covariates by means of generalized additive models (GAMs). GAMs were fitted with the  
212 *gam* R package (Hastie, 2013), using the same model structure identified during model  
213 selection.

214

## 215 **2.5 GIS analysis**

216 For each sampling plot, we predicted the value of web coverage using the most  
217 appropriate model structure supported by the observations derived from the model  
218 selection. In order to provide a graphical representation of the investigated phenomenon in  
219 the study area, we interpolated these predicted values in a GIS environment using the  
220 methodology detailed in Mammola and Isaia (2016). For this analysis, we drew the vector  
221 layer of the arcades on the raster topographical map of the study area, and we  
222 interpolated the projected values for each sampling plot relative to each sampling session.  
223 For the interpolation, we used an Inverse Distance Weighted function (IDW) using a  
224 sample of 12 plots (power 2) to estimate cell values and obtain the renderings of the model  
225 prediction.

### 226 3. RESULTS

227

228 Following the initial data exploration (Zuur et al., 2010), we dropped R, B and G light  
229 components from the GLM analysis, being collinear with artificial light intensity (all Pearson  
230  $r > 0.9$ ). Moreover, we dropped the distance from the river (Dst), being collinear with the  
231 inter-plot crossed distance (InterDst) and we excluded substrate temperature ( $T^\circ$ ) from the  
232 analysis, being collinear with the sampling session. We further applied a log-  
233 transformation to artificial light intensity to achieve homogenization of its distribution (Zuur  
234 et al., 2009), and removed one outlier from the dataset.

235 According to model selection (Table 1), the MAM had the following structure:  $WEB \sim$   
236  $\log(ILL) \times Dst\_lamp + Sampling$ . Specifically, we found a positive and significant  
237 interaction between the artificial light intensity and the distance from the historical  
238 lamppost, whereby higher web coverage was predicted at higher values of light intensity  
239 and in the vicinity of historical lamppost. The effect of this interaction can be visualized in  
240 Figure 4. We also detected a pattern of seasonal variation in the density of webs on the  
241 arcades, with significantly higher predicted values in summer with respect to spring  
242 (reference category). Coverage in winter was not significantly different from the reference  
243 category (Figure 4). Estimated regression parameters and  $p$  values are reported in Table  
244 2, and a graphical representation of the model prediction is shown in Figure 3.

245 There was a significant spatial association in the web coverage among plots  
246 (Moran's I test, observed =  $-0.022$ ; expected =  $-0.006$ ; sd =  $0.005$   $p < 0.05$ ). However, this  
247 effect was not recovered in the regression analysis, given that the variable InterDST was  
248 not significant and thus dropped from the model during the backward elimination model  
249 selection procedure. Spatial association of webs can be visualized graphically in Figure 3  
250 (note the conditional size of the dots representing the plots).

#### 251 4. DISCUSSION

252

253 The two strongest predictors of the web coverage of *Brigittea civica* in the historical  
254 arcades of Turin were the intensity of artificial light and the distance from the nearest  
255 historical (incandescent) lamppost. Moreover, we found a variation in the web coverage  
256 with respect to the sampling session (Figure 3). It is well demonstrated that artificial  
257 illuminance plays an important ecological role in urban settings (e.g., Gaston & Bennie,  
258 2014; Gaston, Bennie, Davies, & Hopkins 2013; Sanders & Gaston, 2018) and other  
259 studies have reported about the significant association between different urban species  
260 and artificial lighting (e.g. Frank, 2009; Heiling, 1999; but see Voss, Main, & Dadour,  
261 2007). As far as *Brigittea civica* is concerned, Samu et al. (2004) reported that (p. 355):  
262 "[...] casual observation [...] showed that [B. civica] webs were aggregated around artificial  
263 lights." Our work provides statistical support to this observation, as we demonstrated that  
264 the web coverage of the plots was significantly higher in the plots where artificial light was  
265 more intense.

266 It was observed that spiders thriving in urban habitats may benefit from increased  
267 trophic resources in cities (Lowe, Wilder, & Hochuli, 2016; Trubl, Gburek, Miles, &  
268 Johnson, 2012; Voss et al., 2007). Based upon this premise, the relationship between web  
269 coverage and artificial light that we observed, could be explained as a function of the  
270 higher prey availability found in the nearby of light sources (Heiling, 1999). The attraction  
271 of aerial and terrestrial arthropods to artificial lighting is indeed a well-documented  
272 phenomenon (Davies, Bennie, & Gaston, 2012; Eisenbeis, 2006; Frank, 2006; Shimoda &  
273 Honda, 2013). It has been shown that *B. civica* is able to feed upon a wide range of small  
274 flying insects — including dipterans, but also flying ants, small lepidopterans and aphids.  
275 Also, if previously starved of food, these spiders are able to feed upon prey nearly three

276 times bigger than themselves (Billaudelle, 1957). This wide range of potential prey is  
277 expected to be available in the areas surrounding street lights (Davies et al., 2012).

278 Our data also demonstrate that the type of light source is important in explaining the  
279 coverage of webs. In particular, spider webs were more abundant in the vicinity of  
280 incandescent historical lampposts (Figure 3) rather than halogen lamps. We assume that  
281 the light emitted by lampposts has its greater effects on attracting spiders due to its higher  
282 potential of attracting insect prey. It is well-documented that nocturnal insects are able to  
283 see ultraviolet radiation, being often attracted to light sources that emit large amounts of  
284 UV radiation (Shimoda & Honda, 2013). At the same time, different light spectra have  
285 different attraction potential to nocturnal arthropods (e.g. Longcore et al., 2015), with  
286 incandescent lights often attracting most nocturnal insects (Justice & Justice, 2016). This  
287 would convincingly explain the interaction we observed between light intensity and the  
288 distance from the incandescent light sources.

289 One could argue that the incandescent artificial lighting system may affect the  
290 distribution of the spiders because of the general influence on thermal conditions, rather  
291 than because of its light spectrum. In fact, there should be higher ambient temperature in  
292 the vicinity of lights, offering a more favourable habitat for a southern European species  
293 such as *B. civica*. However, we rejected this alternative explanation given that the  
294 temperature of the substrate was only limitedly anti-correlated with the distance from the  
295 incandescent artificial lighting system ( $r = -0.26$ ).

296 There is evidence indicating that spiders may be able to recognise the quality of  
297 foraging patches and change web site on the basis of prey availability (e.g., Enders, 1977;  
298 Gillespie & Caraco, 1987; Harwood, Sunderland, & Symondson, 2001; but see Vollrath,  
299 1985). Conversely, it may also be that habitat patches with higher light intensity, and thus  
300 higher prey availability, are not actively selected by the spiders, but simply support larger  
301 colonies of spiders, due to their more favourable condition. It stands to reason that the

302 individual fitness of a spider should be higher in a prey-rich rather than in prey-poor habitat  
303 patch. However, further studies are needed to determine whether the optimal patches are  
304 actively selected by the spiders, or if the higher web coverage found in the vicinity of  
305 incandescent lampposts is the actual results of a higher persistence of the local  
306 population.

307         It is likely that the effect of artificial lighting was particularly clear-cut in our case  
308 because, through the design of the study, we were able to exclude other confounding  
309 factors. It has been shown that *B. civica* avoids areas exposed to rains, winds and direct  
310 solar irradiation (Billaudelle, 1957; Samu et al., 2004). By using our study design (all plot in  
311 sheltered arcades), we were able to exclude these factors from the analysis. Moreover, we  
312 deliberately avoided plots with significant cracks and crevices in the plaster, which are  
313 preferentially selected as supporting framework for the construction of webs and may thus  
314 influence the fine distribution of this species.

315

#### 316 **4.1 Management implications**

317 It has been argued that there is not an easy solution to the contamination of urban wall  
318 surfaces by spider webs of *Brigittea civica*, mainly because this species does not show a  
319 selective preference for a particular wall material or painting (Samu et al., 2004). As far as  
320 we are aware, the only method so far implemented to deal with this issue is the  
321 mechanical removal of the spider webs from the wall surfaces. Whilst the mechanical  
322 removal is certainly effective, such methodology is rather costly, time consuming and  
323 problematic, especially in the case of high buildings. Secondly, the mechanical removal  
324 only represents a temporary remedy to the problem, needing to be reiterated over time.

325         There are three general findings relevant to the management of building surfaces  
326 arising from this work. First, the timing of the mechanical cleaning can be important. Whilst  
327 this may seem self-evident, we suggest that it is important to consider carefully the

328 phenology of the species when planning the cleaning activities. We documented a higher  
329 prevalence of webs during the warm seasons (Figure 4). In particular, a higher web  
330 coverage was observed in summer, at the end of the breeding season of *B. civica*  
331 (Kostanjšek & Celestina, 2008; Nentwig et al., 2018; Wiehle, 1953). This observation  
332 suggests that, in order to maximise the effect of the mechanical removal of the webs, one  
333 should perform the cleaning after the summer peak of density.

334         Second, we showed that the incandescent artificial lighting systems illuminating  
335 most of the arcades in our study area is the main trigger of the increase of the web  
336 coverage (Figure 4). Thus, a renovation of the artificial lighting illuminating in the down-  
337 town districts of heritage cities towards halogen or light emitting diode (LED) lights may  
338 provide a near-permanent solution to this problem, or it may at least contribute to mitigate  
339 the contamination. According to our results, arcades and building facades should become  
340 a less attractive habitats for spiders, thereby reducing the aesthetic nuisance caused by  
341 the webs.

342         Third, casual observations during this study revealed that cracks and crevices in the  
343 plaster are preferentially used as supporting framework for the construction of webs.  
344 Maintaining an intact plaster, at least in the touristic areas, will likely help in reducing the  
345 intensity of the phenomenon.

346         Ultimately, it can be argued that heavy traffic exacerbates the phenomenon, given  
347 that webs became more visible due to air pollution — see argumentation in Samu et al.  
348 (2004). Pedestrian areas have been introduced in most of the historical down-town  
349 districts of cities all around the world, with the aim of increasing commercial and touristic  
350 activities, meanwhile reducing pollution to preserve historical sites (Pagliaria & Biggiero,  
351 2013). Accordingly, an increase of pedestrian areas associated to a reduction of the local  
352 air pollution in down-town districts will, in turn, limit the aesthetic impact associated with *B.*  
353 *civica* webs.

354

## 355 **4.2 Significance statement**

356 One may argue that the results of this study might not be applied to spiders more widely  
357 and that *Brigittea civica* might represent only a very specific case. Still, it is worth noting  
358 that, despite being of southern European origin, this species has been spreading  
359 northward during the last few years. Currently, *B. civica* reaches central Siberia and central  
360 Asia (Buchar & Růžička, 2002; Zamani & Mozaffarian, 2017). Recent evidence indicate  
361 that the species is much more common than was previously known; for instance, in  
362 Central Europe it is likely that its presence has been underestimated (data from Czech  
363 Republic; Novotný et al., 2017; see also discussions in Hänggi & Straub, 2016). Moreover,  
364 the spider was recently found in North America (World Spider Catalog, 2018) and South  
365 Africa (Foord, 2014), which poses additional concerns in light of the potential economic  
366 importance of the potential aesthetic damage caused by this species.

367 Moreover, this study exemplifies a methodological approach that is efficient and  
368 inexpensive, and thus that can be easily reproduced in other cases. More studies similar to  
369 this one would be useful when considering other species which are known to have  
370 potentially negative aesthetic impacts, or that may even cause potential structural  
371 damages. For instance, the photography-based methodology herein described can be  
372 easily used to estimate the density of other organisms which may occur at high densities  
373 — even forming biofilms — on wall surfaces, especially lichens, mosses, and fungi (e.g. de  
374 los Ríos et al., 2009 Gaylarde & Morton, 1999; Lisci, Monte, & Pacini, 2003).

375

376

377

378

379

380

381 **LITERATURE CITED**

382

383 Akamatsu, F., Toda, H., & Okino, T. (2004). Food source of riparian spiders analyzed by  
384 using stable isotope ratios. *Ecological Research*, 19(6), 655-662. doi - 10.1111/j.1440-  
385 1703.2004.00680

386

387 Aronson, M. F. J., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A.,  
388 Warren, P. S., Williams, N. S. G., Cilliers, S., Clarkson, B., Dobbs, C., Dolan, R., Hedblom,  
389 M., Klotz, S., Kooijmans, J. K., Kühn, I., MacGregor-Fors, I., McDonnell, M., Mörtberg, U.,  
390 Pyšek, P., Siebert, S., Sushinsky, J., Werner, P., & Winter, M. (2014). A global analysis of  
391 the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers.  
392 *Proceedings of the Royal Society B Biological Science*, 281, 20133330. doi -  
393 10.1098/rspb.2013.3330

394

395 Bartoń, K. (2017). *MuMIn: Multi-Model Inference*. R package version 1.40.0.  
396 <https://CRAN.R-project.org/package=MuMIn>

397

398 Bertone, M. A., Leong, M., Bayless, K. M., Malow, T. L. F., Dunn, R. R., & Trautwein, M. D.  
399 (2016). Arthropods of the great indoors: characterizing diversity inside urban and suburban  
400 homes. *PeerJ*, 4, e1582. Doi - 10.7717/peerj.1582

401

402 Billaudelle, H. (1957). Zur biologie der mauerspinne *Dictyna civica* (H. LUC.) (Dictynidae:  
403 Araneida). *Zeitschrift für Angewandte Entomologie*, 41, 475-512.

404

405 Buchar, J., & Kubcová, I. (2002). Pavoučí tvář Prahy (The Spidery Face of Prague). *Živa*,  
406 50, 217-218.

407



408 Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multi model inference:*  
409 *apactical information-theoretic approach*. New York - Springer-Verlag.  
410

411 Davies, T. W., Bennie, J., & Gaston, K. J. (2012). Street lighting changes the composition  
412 of invertebrate communities. *Biology Letters*, 8(5), 764-767. doi - 10.1098/rsbl.2012.0216  
413

414 de los Ríos, A., Cámara, B., del Cura, M. Á. G., Rico, V. J., Galván, V., & Ascaso, C.  
415 (2009). Deteriorating effects of lichen and microbial colonization of carbonate building  
416 rocks in the Romanesque churches of Segovia (Spain). *Science of the Total Environment*,  
417 407(3), 1123-1134. doi - 10.1016/j.scitotenv.2008.09.042  
418

419 Eisenbeis, G. (2006). Artificial night lighting and insects: attraction of insects to street  
420 lamps in a rural setting in Germany. In Rich, C., & Longcore, T. *Ecological consequences*  
421 *of artificial night lighting* (pp. 281-304). Washington, DC - Island Press.  
422

423 Elliott, C., Vijayakumar, V., Zink, W., & Hansen, R. (2007). National instruments LabVIEW:  
424 a programming environment for laboratory automation and measurement. *JALA: Journal of*  
425 *the Association for Laboratory Automation*, 12(1), 17-24. doi - 10.1016/j.jala.2006.07.012  
426

427 Enders, F. (1977). Web-site selection by orb-web spiders, particularly *Argiope aurantia*  
428 Lucas. *Animal Behaviour*, 25, 694-712. doi - 10.1002/dev.420020212  
429

430 Foord, S. (2014). *Recent records for three exotic spider species in South Africa*. Cape  
431 *Town Invasives News*. Retrieved 17 November 2016 from  
432 [http://www.capetowninvasives.org.za/news/2014/05/recent-records-three-exotic-](http://www.capetowninvasives.org.za/news/2014/05/recent-records-three-exotic-spider-species-south-africa)  
433 [spider-species-south-africa](http://www.capetowninvasives.org.za/news/2014/05/recent-records-three-exotic-spider-species-south-africa)

434

435 Frank, K. D. (2006). Effects of artificial night lighting on moths. In Rich, C., & Longcore, T.  
436 Ecological consequences of artificial night lighting (pp. 305-344). Washington, DC - Island  
437 Press.

438

439 Frank, K. D. (2009). Exploitation of artificial light at night by a jumping spider. *Peckhamia*,  
440 78(1), 1-3. doi - 10.1007/978-3-540-48348-9\_1

441

442 Gaston, K. J., & Bennie, J. (2014). Demographic effects of artificial nighttime lighting on  
443 animal populations. *Environmental Reviews*, 22(4), 323-330. doi - 10.1139/er-2014-0005

444

445 Gaston, K. J., Bennie, J., Davies, T. W., & Hopkins, J. (2013). The ecological impacts of  
446 nighttime light pollution: a mechanistic appraisal. *Biological Reviews*, 88(4), 912-927. doi -  
447 10.1111/brv.12036

448

449 Gaylarde, C. C., & Morton, L. G. (1999). Deteriogenic biofilms on buildings and their  
450 control: a review. *Biofouling*, 14(1), 59-74. doi - 10.1080/08927019909378397

451

452 Gillespie, R. G. (1987). The mechanism of habitat selection in the long-jawed orb-weaving  
453 spider *Tetragnatha elongata* (Araneae, Tetragnathidae). *Journal of Arachnology*, 15(1), 81-  
454 90.

455

456 Gillespie, R. G., & Caraco, T. (1987). Risk-sensitive foraging strategies of two spider  
457 populations. *Ecology*, 68, 887-899. doi - 10.2307/1938360

458

459 Gittleman, J. L., & Kot, M. (1990). Adaptation: statistics and a null model for estimating  
460 phylogenetic effects. *Systematics Zoology*, 39, 227-241. doi - 10.2307/2992183  
461

462 Grimm, N. B., Faeth, S. H., Golubiewski, N. E., Redman, C. L., Wu, J., Bai, X., & Briggs,  
463 J. M. (2008). Global change and the ecology of cities. *Science*, 319(5864), 756-760. doi -  
464 10.1126/science.1150195  
465

466 Güneralp, B., & Seto, K. C. (2013). Futures of global urban expansion: uncertainties and  
467 implications for biodiversity conservation. *Environmental Research Letters*, 8, 014025. doi  
468 - 10.1088/1748-9326/8/1/014025  
469

470 Hänggi, A., & Straub, S. (2016). Storage buildings and greenhouses as stepping stones for  
471 non-native, potentially invasive spiders (Araneae)—a baseline study in Basel, Switzerland.  
472 *Arachnologische Mitteilungen*, 51, 1-8. doi - 10.5431/aramit5101  
473

474 Harwood, J. D., Sunderland, K. D., & Symondson, W. O. C. (2001). Living where the food  
475 is: web location by linyphiid spiders in relation to prey availability in winter wheat. *Journal*  
476 *of Applied Ecology*, 38(1), 88–99. doi - 10.1046/j.1365-2664.2001.00572  
477

478 Hastie, T. (2013). *gam: Generalized Additive Models*. R package version 1.09. [http://](http://CRAN.R-project.org/package=gam)  
479 [CRAN.R-project.org/package=gam](http://CRAN.R-project.org/package=gam).  
480

481 Havlová, V., & Hula, V. (2010). The *Dictyna civica*, a cribellate web building spider, causes  
482 dirty facades. *Živa*, 2010(4), 172–173.  
483

484 Heiling, A. M. (1999). Why do nocturnal orb-web spiders (Araneidae) search for light?  
485 *Behavioral Ecology and Sociobiology*, 46(1), 43-49. doi - 10.1007/s002650050590  
486

487 Hertel, R. (1968). Über das Auftreten der südeuropäischen Spinne *Dictyna civica* (H. Luc.)  
488 in Dresden (Dictynidae, Araneida). *Abhandlungen des Naturkundemuseums Görlitz*, 44,  
489 89-94.  
490

491 Hurvich, C. M., & Tsai, C. L. (1989). Regression and time series model selection in small  
492 samples. *Biometrika*, 76, 297-307. doi - 10.1093/biomet/76.2.297  
493

494 Isbister, G. K., Gray M. R., Balit C. R., Raven R. J., Stokes B. J., Porges K., Tankel A. S.,  
495 Turner E., White J., & Fisher M. M. (2005). Funnel-web spider bite: a systematic review of  
496 recorded clinical cases. *The Medical Journal of Australia*, 182(8), 407-411.  
497

498 Johnson, J. B., & Omland, K. S. (2004). Model selection in ecology and evolution. *Trends*  
499 *in Ecology and Evolution*, 19(2), 101-108. doi - 10.1016/j.tree.2003.10.013.  
500

501 Justice, M. J., & Justice, T. C. (2016). Attraction of insects to incandescent, compact  
502 fluorescent, halogen, and led lamps in a light trap: implications for light pollution and urban  
503 ecologies. *Entomological News*, 125(5), 315-326. doi - 10.3157/021.125.0502  
504

505 Kleinteich, A. (2010). *Life history of the bridge spider, Larinioides sclopetarius (Clerck,*  
506 *1757)*. PhD thesis, University of Hamburg, Hamburg, Germany. Retrieved from: [https://d-](https://d-nb.info/999993666/34)  
507 [nb.info/999993666/34](https://d-nb.info/999993666/34)  
508

509 Kostanjšek, R., & Celestina, A. (2008). New records on synanthropic spider species  
510 (Arachnida: Araneae) in Slovenia. *Natura Sloveniae*, 10(1), 51–55.  
511

512 Krumpálová, Z. (2001). The synantropic spider *Dictyna civica* (Lucas, 1850) (Araneae,  
513 Dytinidae) in Slovakia. *Sborník přírodovědného klubu v Uherském Hradišti*, 6, 82-85.  
514

515 Lisci, M., Monte, M., & Pacini, E. (2003). Lichens and higher plants on stone: a review.  
516 *International Biodeterioration & Biodegradation*, 51(1), 1-17. doi - 10.1016/S0964-  
517 8305(02)00071-9  
518

519 Longcore, T., Aldern, H. L., Eggers, J. F., Flores, S., Franco, L., Hirshfield-Yamanishi, E.,  
520 Petrinec, L. N., Wilson, Y. A., & Barroso, A. M. (2015). Tuning the white light spectrum of  
521 light emitting diode lamps to reduce attraction of nocturnal arthropods. *Philosophical*  
522 *Transactions of the Royal Society B: Biological Sciences*, 370(1667), 20140125. doi -  
523 10.1098/rstb.2014.0125  
524

525 Lowe, E. C., Wilder, S. M., & Hochuli, D. F. (2016). Persistence and survival of the spider  
526 *Nephila plumipes*. *Urban Ecosystems*, 19(2), 705-720. doi - 10.1007/s11252-015-0518-9  
527

528 Mammola, S., & Isaia, M. (2016). The ecological niche of a special- ized subterranean  
529 spider. *Invertebrate Biology*, 135, 20-30. doi - 10.1111/ivb.12113  
530

531 McIntyre, N. E. (2000). Ecology of urban arthropods: a review and a call to action. *Annals*  
532 *of the Entomological Society of America*, 93(4), 825-835. doi - 10.1603/0013-  
533 8746(2000)093  
534

535 McKinney, M. L. (2002). Urbanization, biodiversity, and conservation. *Bioscience*, 52, 883-  
536 890. doi - 10.1641/0006-3568(2002)052  
537

538 McKinney, M. (2008). Effects of urbanization on species richness: a review of plants and  
539 animals. *Urban Ecosystems*, 11, 161-76. doi - 10.1007/s11252-007-0045-4  
540

541 Nedvěd, O., Pekár, S., Bezděčka, P., Líznarová, E., Řezáč, M., Schmitt, M., Sentenská, L.  
542 (2011). Ecology of Arachnida alien to Europe. *Biological Control*, 56(4), 539-550. doi -  
543 10.1007/s10526-011-9385-3  
544

545 Nentwig, W. (2015) Introduction, establishment rate, pathways and impact of spiders alien  
546 to Europe. *Biological Invasions*, 17(9), 2757-2778. doi - 10.1007/s10530-015-0912-5  
547

548 Nentwig, W., Blick, T., Gloor, D., Hänggi, A., & Kropf, C. (2018). Spiders of Europe (Version  
549 01.2018). Retrieved April 30, 2018 from [www.araneae.unibe.ch](http://www.araneae.unibe.ch). doi - 10.24436/1  
550

551 Newbold, T., et al. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*,  
552 520, 45-50. doi - 10.1038/nature14324  
553

554 Novotný, B., Hula, V., & Niedobová, J. (2017). Insufficiency in distributional faunistic data  
555 in synanthropic spiders: a case study of the occurrence of *Brigittea civica* (Araneae,  
556 Dictynidae) in South Moravia, Czech Republic. *Acta Universitatis Agriculturae et*  
557 *Silviculturae Mendelianae Brunensis*, 65(3), 899-905. doi - 10.11118/actaun201765030899  
558

559 Pagliara, F., & Biggiero, L. (2013). Limited traffic zones and local economy development:  
560 what's the Impact?. *International Journal of Scientific Research*, 2(12), 129-131.

561

562 Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and  
563 evolution in R language. *Bioinformatics*, 20, 289-290.

564

565 R Core Team (2017). *R: A language and environment for statistical computing*. R  
566 Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.

567

568

569 Sams, H. H., Hearsh, S. B., Long, L. L., Wilson, D. C., Sanders, D. H., & King, L. E. (2001).  
570 Nineteen documented cases of *Loxosceles reclusa* envenomation. *Journal of the*  
571 *American Academy of Dermatology*, 44(4), 603-608. doi - 10.1067/mjd.2001.112380

572

573 Samu, F., Jozsa, Z., & Csányi, E. (2004). Spider web contamination of house facades:  
574 habitat selection of spiders on urban wall surfaces. In Samu, F., & Szinetár, C. *European*  
575 *Arachnology 2002* (pp. 351-356). Budapest - Plant Protection Institute and Berzsenyi  
576 College.

577

578 Sanders, D., & Gaston, K. J. (2018). How ecological communities respond to artificial light  
579 at night. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology*.  
580 doi - 10.1002/jez.2157

581

582 Shochat, E., Stefanov, W. L., Whitehouse, M. E. A., & Faeth, S. H. (2004). Urbanization  
583 and spider diversity: influences of human modification of habitat structure and productivity.  
584 *Ecological Applications*, 14(1), 268-280. doi - 10.1890/02-5341

585

586 Shimoda, M., & Honda, K. I. (2013). Insect reactions to light and its applications to pest  
587 management. *Applied Entomology and Zoology*, 48(4), 413–421. doi - 10.1007/s13355-  
588 013-0219  
589

590 Seto, K. C., Gueneralp, B., & Hutyra, L. R. (2012). Global forecasts of urban expansion to  
591 2030 and direct impacts on biodiversity and carbon pools. *Proceedings of the National*  
592 *Academy of Sciences of the United States of America*, 109(40), 16083-16088. doi -  
593 10.1073/pnas.1211658109  
594

595 Taucare-Ríos, A., Brescovit, A., & Canals, M. (2013). Synanthropic spiders (Arachnida:  
596 Araneae) from Chile. *Revista Ibérica de Aracnología*, 23, 49–53.  
597 doi - 10.13140/RG.2.1.3082.7124  
598

599 Trubl, P., Gburek, T., Miles, L., & Johnson, J. (2012). Black widow spiders in an urban  
600 desert: population variation in an arthropod pest across metropolitan Phoenix, AZ. *Urban*  
601 *Ecosystems*, 15(3), 599–609. doi - 10.1007/s11252-011-0223-2  
602

603 Turnbull, A. L. (1973). Ecology of the true spiders (Araneomorphae). *Annual Review of*  
604 *Entomology*, 18(1), 305-348. doi - 10.1146/annurev.en.18.010173.001513  
605

606 Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S. Fourth Edition.*  
607 Springer - New York.  
608

609 Vetter, R. S., & Isbister, G. K. (2008). Medical aspects of spider bites. *Annual Review of*  
610 *Entomology*, 53, 409-429. doi - 10.1146/annurev.ento.53.103106.093503  
611



612 Vitousek, P. M. (1997) Human domination of Earth's ecosystems. *Science*, 277, 494-499.  
613 doi - 10.1126/science.277.5325.494  
614  
615 Vollrath, F. (1985). Web spider's dilemma: a risky move or site dependent growth.  
616 *Oecologia*, 68(1): 69–72. doi - 10.1007/BF00379476  
617  
618 Voss, S. C., Main, B. Y., & Dadour, I. R. (2007). Habitat preferences of the urban wall  
619 spider *Oecobius navus* (Araneae, Oecobiidae). *Austral Entomology*, 46(4), 261–268. doi -  
620 10.1111/j.1440-6055.2007.00616  
621  
622 Wiehle, H. (1953) Spinnentiere oder Arachnoidea (Araneae), IX. Orthognatha-Cribellata-  
623 Haplogynae, Entelegynae (Pholcidae, Zodariidae, Oxyopidae, Mimetidae, Nesticidae). *Die*  
624 *Tierwelt Deutschlands*, 42, 1–150.  
625  
626 World Spider Catalog (2018). *World Spider Catalog*. Natural History Museum Bern.  
627 Retrieved April 30, 2018 from <http://wsc.nmbe.ch> (version 18.5). doi - 10.24436/2  
628  
629 Zamani, A., & Mozaffarian, F. (2017). Further spider (Arachnida: Araneae) material  
630 deposited in the Agricultural Zoology Museum of Iran (AZMI), Iranian Research Institute of  
631 Plant Protection. *Arachnologische Mitteilungen*, 54, 8-20. doi - 10.5431/aramit5403  
632  
633 Zuur, A. F., & Ieno, E. N. (2016). A protocol for conducting and presenting results of  
634 regression type analyses. *Methods in Ecology and Evolution*, 7, 636-645. doi -  
635 10.1111/2041-210X.12577  
636

637 Zuur, A. F., Ieno, E. N., & Elphick, S. C. (2010). A protocol for data exploration to avoid  
638 common statistical problems. *Methods in Ecology and Evolution*, 1, 3-14. doi -  
639 10.1111/j.2041-210X.2009.00001

640

641 Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., Smith, G. M. (2009) *Mixed effect*  
642 *models and extensions in ecology with R*. Berlin - Springer.

643

644

645

646

647

648 **TABLES**

649 **Table 1.** Model selection according to the corrected Akaike criterion for finite sample size (AICc; Burnham &  
 650 Anderson, 2002; Hurvich & Tsai, 1989). Models are ordered from the most to the least appropriate.  
 651

Model	df	AICc	$\Delta$ AICc	wi
y~log(ILL)*Dst_halo + log(ILL)*Dst_lamp + Sampling + InterDst	10	2362.37	3.41	0.08
y~log(ILL)*Dst_halo +Dst_lamp + Sampling + InterDst	9	2360.35	1.40	0.21
y~log(ILL)*Dst_halo + Dst_lamp + Sampling	8	2359.72	0.77	0.28
y~log(ILL)*Dst_halo + Sampling	7	2358.95	0.00	0.42

652  
 653 Df = degrees of freedom; AICc = Corrected Akaike information criterion for finite sample size;  $\Delta$ AICc = (AICc  
 654 of themodel)—(AICc of the best model); wi = Akaike weight (*sensu* Burnham & Anderson, 2002). See text for  
 655 abbreviations of the explanatory variables.

656

657

658 **Table 2.** Estimated regression parameters and p-values according to GLM analysis. See text for  
 659 abbreviations of the explanatory variables.

Variable	Estimated $\beta$	Standard Error	z-value	p-value
Intercept	-7.3130	2.675	-	-
log(ILL)	1.3224	0.254	5.201	<0.001
Dst_lamp	1.3912	0.548	2.537	0.011
log(ILL) x Dst_lamp	-0.1528	0.052	-2.291	0.003
Sampling (level: December)	-0.1816	0.137	-1.324	0.185
Sampling (level: July)	1.3761	0.138	9.979	<0.001

660

661

662

663

664

665

666

667 **FIGURE CAPTIONS**

668

669 **Figure 1.** **A)** *Brigittea civica* (Lucas) (Araneae: Dictynidae) (photo credits:  
670 Dr. Hans-Juergen Thorns). **B)** Sampling plot on an arcade ceiling with a reduced density of cobwebs of *B.*  
671 *civica*. **C)** Sampling plot on an arcade ceiling with a significant contamination of cobwebs of *B. civica*. **D)**  
672 Sampling plot on an arcade ceiling entirely covered by cobwebs of *B. civica*. **E)** The historical arcades of  
673 Turin in the area close to Palazzo Carignano. Arrows point at area covered by webs of *B. civica* (Photo  
674 credits: Nicola Paccagnella — [www.nicola.photos](http://www.nicola.photos)).

675

676 **Figure 2.** Schematic summary of the monitoring protocol.

677

678 **Figure 3.** Maps of the studied arcades showing interpolated surfaces of the predicted coverage of webs in  
679 winter (**a**), spring (**b**) and summer (**c**) according to GLM results. Size of each sampling plot is proportional to  
680 the observed web coverage.

681

682 **Figure 4.** Predicted relationship between the coverage of webs of *Brigittea civica* and the intensity of artificial  
683 light (log-transformed) in interaction with the distance from the incandescent historical lamppost across the  
684 three sampling sessions. To generate the predictions, two arbitrary values of distance from the incandescent  
685 lamppost were used, namely zero (filled line) and six (dashed line) meters. Shaded grey surfaces are 95%  
686 confidence intervals.

687






Figure 2













Figure 3



**Figure 4**





# Graphical Abstract



Density of webs

Intensity of artificial lighting



Distance from the incandescent lamp post

1 **ACKNOWLEDGMENTS**

2

3 This study is part of a wider research project on urban decay funded by Compagnia di San  
4 Paolo. Special thanks go to Dr. Sara L. Goodacre for proofreading our English, to  
5 Francesca Cochis for fieldwork assistance and to Prof. Wolfgang Nentwig for pre-revising  
6 the manuscript. We thank Dr. Hans-Juergen Thorns and Nicola Paccagnella for sharing  
7 with us photographs of *Brigittea civica* and its habitat (Figure 1).

8