

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Study of Lactococcus lactis during advanced ripening stages of model cheeses characterized by GC-MS

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1679257> since 2018-10-29T14:25:30Z

Published version:

DOI:10.1016/j.fm.2018.03.012

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1
2
3 This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is
4 posted here by agreement between Elsevier and the University of Turin. Changes
5 resulting from the publishing process - such as editing, corrections, structural
6 formatting, and other quality control mechanisms - may not be reflected in this version
7 of the text. The definitive version of the text was subsequently published in:

8
9 [Food Microbiology, 2018, 74, 132-142, 10.1016/j.fm.2018.03.012](#)

10
11 You may download, copy and otherwise use the AAM for non-commercial purposes provided that your
12 license is limited by the following restrictions:

13
14 (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.

15
16 (2) The integrity of the work and identification of the author, copyright owner, and publisher must be
17 preserved in any copy.

18
19 (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND
20 license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>),
21 [10.1016/j.fm.2018.03.012](#)

22
23 The publisher's version is available at:

24 <https://www.sciencedirect.com/science/article/pii/S0740002017301144>

25 When citing, please refer to the published version.

26
27 Link to this full text:

28 <https://iris.unito.it/preview-item/353380?queryId=mysubmissions&>

29
30
31 This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

32

33 **Study of *Lactococcus lactis* during advanced ripening stages of model cheeses**
34 **characterized by GC-MS.**

35 Marianna Ruggirello*, Manuela Giordano, Marta Bertolino, Ilario Ferrocino, Luca Cocolin, Paola

36 Dolci

37 Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Largo

38 Paolo Braccini 2, 10095 Grugliasco, Italy

39

40

41 Corresponding author: paola.dolci@unito.it

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61 **ABSTRACT**

62 *Lactococcus lactis*, extensively used as dairy starter, has recently been detected in cheese, as
63 metabolically active cells, in advanced ripening stages, by culture-independent methods. In this
64 study, we assessed the viability of *L. lactis* subsp. *lactis* in model cheeses, up to 180 days of
65 ripening. In addition, we studied the expression of *metC* and *als* genes involved in the production
66 of aroma compounds by Gas Chromatography-Mass Spectrometry (GC-MS). Three *L. lactis*
67 subsp. *lactis* commercial starters were inoculated in pasteurized milk and model cheeses were
68 manufactured and ripened for six months. Samples were analyzed at manufacturing and ripening
69 steps, in terms of viability of *L. lactis* by both traditional plating and direct analysis of RNA by
70 reverse transcription quantitative PCR (RT-qPCR) and in terms of aroma profile by GC-MS.
71 Relatively to RT-qPCR analysis, *L. lactis* was found viable throughout the whole process of
72 cheesemaking and aging, with final average loads of 3-4 Log CFU/g at 180 days. On the contrary,
73 the microorganism was not detected, in ripened samples, by traditional plating on M17 medium,
74 suggesting its entering in a viable but not cultivable (VBNC) state. The aroma profiles of the
75 cheeses highlighted the presence of volatile compounds related to cheese flavor as acetoin,
76 diacetyl, 2,3-butanediol and dimethyl disulfide, whose presence was partially correlated to *metC*
77 and *als* genes expression. These results add new insights on the capability of *L. lactis* to persist
78 during late cheese ripening and suggest a potential contribution of the microorganism to cheese
79 flavor formation.

80

81 **Keywords:** *Lactococcus lactis*; ripening; volatile aroma compounds; RT-qPCR.

82

83 Chemical compounds studied in this article:

84 Lactose (PubChem CID: 84571); Glucose (PubChem CID: 53782692); Galactose (PubChem CID:
85 439357); Lactic acid (PubChem CID: 61503); Citric acid (PubChem CID: 311); Butyric acid
86 (PubChem CID: 264); Acetic acid (PubChem CID: 176); Propionic acid (PubChem CID: 1032);
87 Pyruvic acid (PubChem CID: 1060); 2-Butanone (PubChem CID: 6569); 2-Heptanone (PubChem
88 CID: 8051); 2-Nonanone (PubChem CID: 13187); 2-Propanone, 1-hydroxy- (PubChem CID: 8299);

89 Acetoin (PubChem CID: 179); Acetone (PubChem CID: 180); Diacetyl (PubChem CID: 650); 1-
90 Butanol (PubChem CID: 263); 1-Hexanol (PubChem CID: 8103); 1-Hexanol, 2-ethyl- (PubChem
91 CID: 7720); 1-Octanol (PubChem CID: 957); 1-Octen 3-ol (PubChem CID: 18827); 2,3-Butanediol
92 (PubChem CID: 262); 2-Buten-1-ol, 3-methyl- (PubChem CID: 11173); 3-Buten-1-ol, 3-methyl-
93 (PubChem CID: 12988); Diethylene glycol (PubChem CID: 8117); Ethanol (PubChem CID: 702);
94 Ethanol, 2-(2-ethoxyethoxy)- (PubChem CID: 8146); Ethanol, 2-butoxy- (PubChem CID: 8133);
95 Isobutanol (PubChem CID: 6560); Isopentyl alcohol (PubChem CID: 31260); alpha.-Pinene
96 (PubChem CID: 6654); delta 3-carene (PubChem CID: 26049); Limonene (PubChem CID: 22311);
97 Menthol (PubChem CID: 1254); Disulfide, dimethyl (PubChem CID: 12232); Dimethyl sulfone
98 (PubChem CID: 6213); 2,3-Butanediol diacetate (PubChem CID: 66193); Butyl acetate (PubChem
99 CID: 31272); Ethyl acetate (PubChem CID: 8857); Ethyl decanoate (PubChem CID: 8048); Ethyl
100 hexanoate (PubChem CID: 31265); Ethyl octanoate (PubChem CID: 7799); Methyl 2-hydroxy 2-
101 methyl- propionate (PubChem CID: 75024); Hexanal (PubChem CID: 6184); Nonanal (PubChem
102 CID: 31289); Butanoic acid (PubChem CID: 264); Decanoic acid (PubChem CID: 2969);
103 Heptanoic acid (PubChem CID: 8094); Hexanoic acid (PubChem CID: 8892); Isobutyric acid
104 (PubChem CID: 6590); Isovaleric acid (PubChem CID: 10430); Nonanoic acid (PubChem CID:
105 8158); Octanoic Acid (PubChem CID: 379); Pentanoic acid (PubChem CID: 7991); Propanoic acid
106 (PubChem CID: 1032); Benzaldehyde (PubChem CID: 240); Benzenecarboxylic acid (PubChem
107 CID: 243); Benzyl Alcohol (PubChem CID: 244); beta-Phenoxy ethyl alcohol (PubChem CID:
108 31226); Phenol (PubChem CID: 996); Phenyl ethyl Alcohol (PubChem CID: 6054); delta-
109 Decalactone (PubChem CID: 12810); delta-Octanolactone (PubChem CID: 12777); Undecane
110 (PubChem CID: 14257).

111

112

113

114

115

116

117 1. INTRODUCTION

118 Flavour development in dairy fermentations, most notably cheeses, results from a series of (bio)
119 chemical processes in which starter and secondary cultures provide enzymes. *Lactococcus lactis*
120 is the predominant microorganism in starter cultures used for cheese making and its enzymes,
121 which act during cheese manufacture and early ripening, affect the organoleptic properties of the
122 products in terms of texture, flavour and aroma development (Kunji et al., 1993; Ercan et al., 2015;
123 García-cayuela et al., 2012; Rijnen et al., 2003). As the result of its technological ability, *L. lactis*
124 has been the subject of numerous studies to determine the main relevant characteristics useful for
125 the selection of the best strains for dairy applications (Taïbi et al., 2011; Van Hylckama Vlieg et al.,
126 2006). Although *L. lactis* is a key microorganism in cheese making and early ripening (when it is
127 found with high count values by traditional plating), in late ripening the microorganism is outcome
128 by secondary microbiota, mainly lactobacilli, and its presence has been highlighted by some
129 authors (Desfossés-Foucault et al., 2013; Dolci et al., 2014; Flórez and Mayo, 2006; Masoud et al.,
130 2011) exclusively by culture-independent methods. Thus, it was hypothesized that *L. lactis* was
131 present, in ripened cheeses, in a viable but not cultivable (VBNC) state and trials to resuscitate it
132 were performed in previous studies (Ruggirello et al., 2016). In the present study, we investigated
133 the persistence of *L. lactis* subsp. *lactis* in model cheeses in order to further confirm the evidences
134 obtained in our previous researches (Ruggirello et al., 2016, 2014). In addition, we studied the
135 expression, in *L. lactis*, of genes involved in the production of aroma compounds to evaluate if any
136 possible correlation could be hypothesized between their expression and volatile aroma
137 compounds detected in cheese samples. Precisely, we chose *metC* gene encoding cystathionine
138 β -lyase (C β L) and *als* gene coding for α -acetolactate synthase (α -ALS) (Platteeuw et al., 1995;
139 Toonen et al., 1994). C β L is responsible to catalyse the deamination of methionine to methanethiol
140 (Smit et al., 2005; Weimer, 2007) toward the production of sulphur aroma compounds (dimethyl
141 sulphide, DMDS and dimethyl trisulphide, DMTS) while α -ALS represents the first enzyme of the
142 diacetyl/acetoin biosynthetic route (Hugenholtz and Starrenburg, 1992; Platteeuw et al., 1995;
143 Snoep et al., 1992; Verhue and Tjan, 1991). Finally, we studied the aroma profiles of model
144 cheeses by Gas Chromatography-Mass Spectrometry (GC-MS) and we evaluated if any possible

145 correlation could be hypothesized between *als* and *metC* gene expression in *L. lactis* and volatile
146 aroma compounds detected in cheese samples in order to define a potential technological role of
147 the microorganism also in late ripening.

148

149 **2. MATERIALS AND METHODS**

150 **2.1 Cheese making and sampling.**

151 Commercial starter cultures, named in this paper K, M and B, were purchased as lyophilized cells
152 of *L. lactis* subsp. *lactis* and used as inoculum for cheese-making of Toma-like miniature cheeses
153 according to the protocol described by Bertolino et al. (2008) with some modifications as reported
154 by Ruggirello et al. (2016). Overall, six productions (2 for each starter) were carried out in aseptic
155 conditions. Milk, curd and cheese samples were collected, for each production, in duplicate, as
156 follow: milk before inoculum of the starter (BLANK), milk after inoculum of the starter (MILK), curd
157 after draining (DRY), cheese immediately after salting (SALT) and cheese at 15, 30, 60, 90, 120,
158 150 and 180 days of ripening. Pasteurized milk used for cheese making was checked, before
159 starter inoculum, for the absence of *L. lactis* cells by qPCR as reported by Ruggirello et al., (2014).
160 Values of pH were measured in milk samples immediately after inoculum, curd after salting and
161 cheese at the end of ripening.

162

163 **2.2 Microbial analyses and *L. lactis* detection by qPCR.**

164 Ten millilitres (milk) or grams (curd and cheese) of each sample were homogenised in 90 mL of
165 sterile Ringer solution (Oxoid, Milan, Italy) in a Stomacher (Interscience, Rockland, MA, USA) for 5
166 min. The resulting suspension was serially diluted and included in M17 agar (Biolife, Milan, Italy)
167 supplemented with lactose (5 g/L). In addition, the 10^{-1} dilution was plated for bulk formation as
168 described by Ercolini et al. (2001). The plates were incubated, aerobically, at 30 °C for 48h and
169 colony-forming units (CFU) were subsequently counted. Bulk formations were covered with 1 mL of
170 Ringer, scraped off with a sterile L-shape spreader and harvested with a pipette in tubes, which
171 were stored at - 20 °C before DNA extraction. DNA extracted according to the method reported by
172 Cocolin et al. (2001) was then submitted to *L. lactis* selective qPCR analysis with the protocol

173 optimized and described by Ruggirello et al. (2014), in order to investigate the presence of *L. lactis*
174 cultivable cells.

175

176 **2.3 Analysis of *tuf*, *metC* and *als* gene expression by RT-qPCR.**

177 **2.3.1 RNA extraction.** Ten millilitres of milk and ten grams of curd and cheese samples were flash
178 frozen in liquid nitrogen to inhibit further microbial biochemical activities (in particular DNA
179 transcription) and, then, stored at -80 °C until RNA extraction. Just before extraction, samples were
180 slowly thawed at 4 °C. RNA extraction was performed by using TRIzol® reagent (FMB, Trevose,
181 USA) according to the manufacturer's instructions with some modifications. Briefly, once thawed,
182 each matrix was resuspended in 25 mL of cold TRIzol® reagent and homogenised for 5 min at 260
183 rpm with a Stomacher® 400 Circulator (Seward Ltd, Worthing, UK). Then, 1 mL of homogenised
184 slurries was moved in a 2 mL tube containing 0.3 g of 1 mm diameter glass beads and three
185 homogenization steps of 50, 45 and 30 sec were performed at high speed, using a Tissue Lyser LT
186 (Qiagen, Germany). Subsequently, samples were centrifuged at 12,000 x g for 10 min at 4 °C and
187 the supernatants collected and transferred in a new tube after removing fat top layer by a sterile
188 loop. Two hundred millilitres of chloroform were added to the supernatants and the tubes shaken,
189 vigorously, by hand, for 15 sec and incubated for 2–3 min at room temperature. After centrifugation
190 at 12,000 x g for 15 min at 4 °C, the aqueous phase was moved in a new tube and added of 0.5
191 mL 100% isopropanol for RNA precipitation. In order to remove contaminants, the solution was
192 purified using RNeasy columns (QIAGEN, Hilden, Germany), following the manufacturer's
193 recommendations with an additional final lithium chloride (LiCl) precipitation. Briefly, LiCl (Life
194 technologies, Dublin, Ireland) was added to the final volume of 30 µl of the sample with a final
195 concentration of 3 M and all the samples were chilled 1 h at -20 °C. After centrifugation at 16,000 x
196 g for 20 min at 4 °C, supernatant was discarded and RNA pellet resuspended in sterile water. The
197 addition of LiCl allows to remove inhibitors of RNA translation and cDNA synthesis from RNA
198 preparations (Cathala et al., 1983). Remaining DNA was digested using TURBO-DNase (Ambion,
199 Milan, Italy) according to the manufacturer's protocol. After treatment, 15 mM EDTA was used to
200 deactivate DNase enzyme and 1U RiboGuard RNase Inhibitor (Epicentre, Madison, WI, USA) was

201 added to the samples subsequently incubated at 75 °C for 10 min and, finally, stored at -80 °C.
202 RNA extractions were performed in duplicate for each dairy sample. The amount and integrity of
203 total RNA extracted were analysed by using Experion™ Automated Electrophoresis System with
204 Experion HighSens chips (Bio-Rad, Milan, Italy). The absence of residual DNA was verified by
205 qPCR using the protocol described by Ruggirello et al. (2014).

206 **2.3.2 *Tuf* gene analysis.** In order to study the expression, in *L. lactis*, of the housekeeping gene
207 *tuf*, encoding elongation factor Tu, RNA samples were submitted to retrotranscription by M-MLV
208 Reverse Transcriptase kit (Promega, Milan, Italy). Briefly, RT reactions were performed as follows:
209 0.5 µL of RNA were mixed with 1 µL of TUF2r (100 mM) (Ruggirello et al., 2014) in a reaction
210 volume of 10 µL by addition of ultrapure water. The mixtures were maintained at 75 °C for 5 min for
211 RNA denaturation and immediately placed on ice for 10 min. Five microliters of M-MLV RT Buffer
212 (1X), 5 µL of dNTPs (10mM each) and 1 µL of M-MLV Reverse Transcriptase (8 U/µL) were added
213 to the mixtures in a final volume of 25 µL by addition of ultrapure water. RT reactions were carried
214 out at 42 °C for 1 h in a Biorad DNA Engine thermal cycler (Bio-Rad) and the cDNA submitted to
215 qPCR by using primers TUF2f and TUF2r, according to the protocol optimized by Ruggirello et al.
216 (2014).

217

218 **2.3.3 *metC* and *als* gene analysis**

219 Milk, curd and cheese samples were then investigated for the expression, in *L. lactis*, of *metC* and
220 *als* genes involved in aroma production pathways. The *metC* and *als* genes expression was
221 compared with the expression of *tuf* gene which was used as reference gene.

222 **2.3.3.1 Primer efficiency.** *L. lactis* selective primers were used (Table 1) and, before performing in
223 RT-qPCR, their efficiency (E) was determined on DNA extracted from *L. lactis* starter B used as
224 template. The E values were calculated for each gene according to Pfaffl (2001). Samples of DNA
225 were diluted (100, 50, 25, 5, 1 and 0.25 ng/ml) and used to construct the standard curves.
226 Amplifications were performed in a Biorad DNA Engine thermal cycler (Bio-Rad), in 10 µl reaction
227 mixture containing 1 µl of DNA, 5 µl of SsoAdvanced™ SYBR® Green Supermix (Bio-Rad,
228 Hercules, CA, USA), 400 nM and 50 nM of Tuf2f and Tuf2r primers or 900 nM of metCr-metCf and

229 alsr-alsf primer pair. A negative control with no DNA was also included in each run, as well as a
230 positive control with genomic *L. lactis* DNA. Thermal cycle consisted of a 20 sec denaturation step
231 at 95 °C, followed by 40 two-step cycles of 5 sec at 95 °C and 20 sec at 54.2 °C. In addition,
232 melting curve analysis was performed to check the presence of a single amplification product for
233 each primer pair. Dilutions were replicated three times and used singly.

234 **2.3.3.2 RT-qPCR.** Retrotranscription was performed by M-MLV Reverse Transcriptase kit
235 (Promega, Milan, Italy) as reported above (2.3.2) by using primers metCr and alsr for *metC* and *als*
236 gene analysis, respectively. Complementary DNA (cDNA) of MILK, SALT and 15, 30, 60, 90, 150,
237 180 ripening day samples was then amplified by qPCR in a Biorad DNA Engine thermal cycler
238 (Bio-Rad) using 96-well plates and following the protocol reported above (2.3.3.1). In addition,
239 melting curve analysis was performed to check the presence of a single amplification product for
240 each primer pair.

241

242 **2.4 Analysis of volatile compounds.**

243 Volatile organic compounds (VOCs) were studied in curd and cheese samples. Two grams of each
244 grated and homogenised sample were placed in a 20 mL vial, added of 10 µl of internal standard
245 solution (2-octanol) at 10 ppm, fitted with a PTFE silicone septum (Supelco, Bellefonte, PA, USA)
246 through which the headspace solid phase micro extraction (HS-SPME) syringe needle, fitted with a
247 Stable Flex 1 cm–50/30 µm divinylbenzene-carboxen-polydimethylsiloxane (DVB-CAR-PDMS)
248 fibre (Supelco, Bellefonte, PA, USA), was introduced. The vials were placed in a heat/stir plate at
249 40 °C for 20 min for the extraction phase. After exposure to the headspace (HS), the fibre was
250 retracted and transferred to the GC/qMS injector operated in the splitless mode at 260 °C for 2
251 min. GC/qMS analysis was performed with a Shimadzu GC-2010 gas chromatograph equipped
252 with a Shimadzu QP-2010 Plus quadrupole mass spectrometer (Shimadzu Corporation, Kyoto,
253 Japan) and a DB-WAXETR capillary column (30 m × 0.25 mm, 0.25 µm film thickness, J&W
254 Scientific Inc., Folsom, CA, USA). The temperature program started at 40 °C for 5 min, increased
255 at a rate of 10 °C min⁻¹ to 80 °C, then at rate of 5 °C min⁻¹ to 240 °C. The carrier gas (He) flow-rate
256 was 1 mL min⁻¹. The injection port temperature was 260 °C; the ion source and the interface

257 temperature were 240 °C. The detection was carried out by electron impact mass spectrometry in
258 total ion current (TIC) mode, using an ionization energy of 70 eV. The mass acquisition range was
259 m/z 33–300. Peak identification of each volatile metabolite was performed by comparison of the
260 retention time and mass spectra of eluting compounds to those of the pure standard. Compounds
261 were semi quantified by calculating the area under the ion quantifier peak of each compound
262 (Yarlagadda et al., 2014). All analyses were performed in duplicate for each samples.

263

264 **2.5 Organic acid and sugar profiles.**

265 The organic acids (citric, pyruvic, lactic, acetic, butyric and propionic) and sugars (lactose, glucose
266 and galactose) were determined for all dairy samples by high performance liquid chromatography
267 (HPLC), according to the method described by Marchiani et al. (2016). Briefly, samples (5 g) were
268 added to 25 mL of 0.013 NH₂SO₄ (mobile phase) and mixed for 10 min with a Stomacher® 400
269 Circulator (Seward Ltd, Worthing, UK). The slurries were subsequently centrifuged at 5,000 x g for
270 30 min at 10 °C, and the supernatant was filtered through 0.45 µm polypropylene membrane filter
271 (VWR, Milan, Italy). The HPLC system (Thermo Finnigan Spectra System, San Jose, USA) was
272 equipped with an isocratic pump (P4000), a multiple autosampler (AS3000) fitted with a 20 µL
273 loop, a UV detector (UV100) set at 210 and a refractive index detector RI-150. The analyses were
274 performed isocratically at 0.8 mL min⁻¹ at 65 °C, with a 300 × 7.8 mm i.e. cation exchange column
275 (Aminex HPX-87H) equipped with a Cation H⁺ Microguard cartridge (Bio-Rad Laboratories,
276 Hercules, USA). Three replicates were analysed for each sample. Data processing was carried out
277 using Chrom Quest™ chromatography data system (ThermoQuest Corporation, San Jose, USA).
278 Analytical grade reagents were used as standards (Sigma-Aldrich, St. Louis, USA). All analyses
279 were performed in duplicate for each samples.

280

281 **2.6 Statistical analysis.**

282 Statistical analysis was performed using the Microsoft Excel 2010, SPSS v23.0 (SPSS, Inc.,
283 Chicago, Ill., USA) R package (www.r-project.org) and STATISTICA (StatSoft Inc., Tulsa, USA)
284 programs.

285 Threshold cycle (Ct) values from RT-qPCR were exported to Excel for relative quantification. In
286 order to use parametric tests for gene expression analysis (ANOVA), off-scale data were replaced
287 with fictive Ct values (Kubista and Sindelka, 2007). Fictive Ct values are set to the highest Ct
288 observed for a truly positive sample, which is assumed the level of detection (LOD), plus one. In
289 the current study, the cut off was set at 36 cycles. As reported by some authors (Kubista and
290 Sindelka, 2007; Mataragas et al., 2015), this corresponds to assigning a concentration that is half
291 of the LOD to the off-scale samples. This is no more erroneous than assigning a zero
292 concentration to these samples, because there is no evidence that they are blank. It is only known
293 that the amount of the target in these samples is lower than the detection limit.

294 Data were converted to relative expression and log-values (fold change) (Kubista and Sindelka,
295 2007; Livak and Schmittgen, 2001). In order to measure gene expression as a function of time and
296 the three *L. lactis* starters, relative quantification was carried out. Thus, target gene expression in
297 curd and cheese samples was compared with the expression detected in MILK samples (after
298 inoculum of the starters) by one-way ANOVA. Tuckey's test for means comparison was carried out
299 at a P value of < 0.05. Mixed repeated measures ANOVA was performed to investigate whether
300 *metC* and *als* gene expression was dependent on different ripening times (factor A) and/or different
301 starters (factor B).

302 VOCs production was visualized by heatmap designed by R package and Ward's criterion was
303 used for clustering. Principal component analysis (PCA) was performed specifically on DMDS,
304 acetoin, diacetyl and 2,3-bunanediol compounds.

305 Mixed repeated measures ANOVA and Tuckey's test for means comparison were carried out for
306 HPLC values, in order to explore differences related to cheeses manufactured by different starters
307 and sampling time. For all the analyses significant results were associated to P < 0.05 or P < 0.001
308 or P < 0.001.

309

310 **3. Results**

311 **3.1 *L. lactis* detection throughout manufacturing and ripening**

312 **3.1.1 Culture-dependent approach.** By plating on M17 agar, the three starters were found, after
313 inoculum in MILK, with loads of about 7-8 Log CFU/mL and they reached maximum peaks of about
314 8-9 Log CFU/mL in curd after drying (DRY) and salting step (SALT); then, at 15 and 30 ripening
315 days, they showed a downward trend, reaching the lowest counts of 3-4 Log CFU/g at 90 ripening
316 days (B and M) and of 5 Log CFU/g at 60 ripening days (K). After that, the starters did not grow
317 anymore on M17 (Table 2). The results of bulk formation analysis confirmed microbial count data
318 (Table 2), since no *L. lactis* cultivable cells were detected by qPCR in those samples where no
319 colonies grew on M17. A decrease of pH from 6.5 ± 0.2 (MILK) to 5.1 ± 0.3 (SALT) and to 4.2 ± 0.2
320 (180) was observed.

321 **3.1.2 Culture-independent approach.** The presence of metabolically active cells of *L. lactis*
322 starters was investigated by RT-qPCR directly on milk, curd and cheese samples. Their presence
323 was quantified by interpolating the Ct values obtained in the standard curve equation optimised
324 and described in a previous work (Ruggirello et al., 2014). The results are presented in Table 2
325 and compared with the data obtained from culture-dependent approach. Starters were found in
326 MILK with minimal loads (about 1-2 Log CFU/mL), probably due to the fact that, after the inoculum
327 of the starter, *L. lactis* cells had not yet metabolically reactivated (sampling time was 5 minutes
328 subsequent the inoculum). Then, the three starters reached values of about 7-8 Log CFU/g during
329 manufacturing and their performance, throughout ripening, seemed not to be affected from the
330 initial low loads in milk. They reached values of about 6 Log CFU/g at 120 days of ripening and at
331 150 days, starter B was still found at 6 Log CFU/g while K and M loads dropped to 4 Log CFU/g
332 (Table 2). Finally, they were all present after 180 ripening days with 3-4 Log CFU/g.

333

334 **3.2 Relative expression of *L. lactis metC* and *als* genes in miniature cheeses.**

335 The expression of *metC* and *als* genes in *L. lactis* starters was studied by RT-qPCR throughout
336 manufacturing and ripening of miniature Toma-like cheeses. Efficiency and R^2 values are shown in
337 Table 1. No primer-dimers were generated during the 40 cycles.

338 Figure 1 shows the changes of the two gene expression from manufacturing to 180 ripening days,
339 compared with their expression in the samples taken as control (MILK). Because no C_t values were

340 registered for the *als* gene in cheeses made with the starter M, these samples were excluded from
341 the graph.

342 One-way ANOVA highlighted a significant over expression of *metC* gene in cheese sample SALT
343 produced with K starter and in samples at 150 and 180 ripening days produced with starter B. No
344 significant differences were found for *als* gene expression along ripening time.

345 Mixed repeated measures ANOVA underlined a significant influence on *metC* expression due to
346 ripening time and starter strain ($F= 7.46$, $P < 0.05$), as well as a significant interaction between the
347 two variables (factor A and B) ($F= 20.694$, $P < 0.05$), while *als* gene expression was not statistically
348 affected neither by factor A nor factor B. The *metC* gene was over-expressed by starters K and B
349 after salting time ($F= 25.349$, $P < 0,05$) whereas it was down regulated in starter M. K was over
350 expressed at 15 and 30 days ($F= 27.311$ and $F= 11.99$ respectively, $P < 0,05$) unlike B and M that
351 were down regulated. At 60 days all the starters up regulated the gene. Down regulation started
352 from 90 days in K and from 150 days in M. The *metC* gene expression increased in starter B from
353 60 to 180 days of ripening with a significance difference compared to starter M at 150 days ($F=$
354 $78,97$, $P<0,05$) and starters M and K at 180 days ($F= 133.73$, $P < 0,05$).

355 The *als* gene was down regulated during both manufacturing and ripening by B and K starters
356 (values for K not detected at 60, 90 and 150 days), while it was over expressed in 180-old cheeses
357 made with B ($F= 65.21$, $P < 0.05$). As reported above, no C_1 values were detected in starter M

358

359 **3.3 Volatile compounds analysis.**

360 HS-SPME and GC-MS allowed the identification of 56 different VOCs: 11 acids, 7 esters, 7
361 ketones, 2 aldehydes, 14 alcohols, 2 lactones, 4 monoterpenes, 2 sulphur compounds, 6 aromatic
362 compounds and 1 undecane compound not ascribable to these chemical classes. Mixed repeated
363 measures ANOVA was carried out to investigate whether VOCs production in cheeses was
364 different depending on the different ripening time (factor A) and/or on the different *L. lactis* starter
365 strain (factor B). The results showed that 82.1% ($F=32.69$, $P< 0.05$) and 53.5% ($F=18.21$, $P< 0.05$)
366 of the VOCs were significantly influenced by factor A and factor B, respectively. Moreover, a
367 significant interaction was highlighted between the two factors ($F= 8.63$, $P < 0.05$) for 57.1% of the

368 VOCs, showing that ripening time had a different effect on VOCs production, depending on *L. lactis*
369 starter strain used in miniature cheeses. Ketones were the chemical class mainly represented
370 during manufacturing (48.2% w/w of VOCs) and ripening (53.6% w/w of VOCs). Alcohols were
371 found at percentage of 36.4% and 24.9% and acids at 9.8% w/w and 20.2% w/w in manufacturing
372 and ripening, respectively. Regarding VOCs production, an interesting separation between milk,
373 manufacturing and ripening time was showed by heatmap in figure 2. Early ripened samples were
374 characterized by high concentration of ketones as 1-hydroxy- 2-propanone, acetoin, acetone and
375 diacetyl; among alcohols, 2-ethyl-1-hexanol, 2,3-butanediol, 3- methyl-2-buten-1-ol and ethanol
376 were the most abundant followed by monoterpenes (limonene and menthol), 2,3-butanediol
377 diacetate ester and hexanal aldehyde. During cheese ripening, an increment of acids was
378 observed. The highest amounts were found for acetic, butanoic and hexanoic acids with higher
379 levels of butanoic acid in cheeses produced with starter B more than the ones obtained with K and
380 M ($F= 36.75$, $P < 0.05$). It was interesting to note (figure 2) that starter K moved earlier to the
381 cluster on the left (after just 30 days) compared to B and M due to its high production of acetate,
382 ethanol, acetoin, diacetyl and 2,3-butanediol compounds.

383 Aroma compounds derived from cysteine and methionine catabolism (DMDS) and citrate (acetoin,
384 diacetyl and 2,3-butanediol) pathways were specifically analysed by PCA analysis (Figure 3).
385 Results underlined a certain degree of separation among the three *L. lactis* starters. DMDS
386 originated from cysteine and methionine catabolism, represented 2.3% of total VOCs during
387 manufacturing and ripening. The compound was significantly affected by both factor A ($F= 30.16$,
388 $P < 0.05$) and B ($F=130.73$, $P < 0.05$), being produced with higher concentrations by starters K and
389 B than in cheese produced with starter M. Also acetoin was significantly affected by both factors
390 ($F= 27.69$ and $F= 24.01$ respectively for starter and time factors, $P < 0.05$). 2-Butanone varied
391 significantly among the starters being more produced by K ($F= 18.08$, $P < 0.05$) whereas 2,3-
392 butanediol and diacetyl amounts varied only along ripening time ($F= 3.56$ and $F= 6.86$ respectively,
393 $P < =0.05$) without significant differences among the starter. Among ketones, acetoin, which forms
394 from diacetyl together with 2-butanone and 2, 3-butanediol, was the most abundant compound
395 detected in ripened samples. Its highest amount was found in samples at 15 days and decreased,

396 during ripening, in cheeses produced with starters B and M while remained high in cheeses made
397 with starter K ($P < 0.05$). Diacetyl was found starting from MILK samples after inoculum and
398 increased in ripened cheeses. Lastly, 2,3-butanediol, was detected in manufacturing (DRY
399 samples) and showed constant values until 180 days of ripening.

400

401 **3.4 Organic acid and sugar profiles.** Organic acid and sugar contents are reported in Table 3.
402 Mixed repeated measures ANOVA was carried out to investigate whether organic acids and sugars
403 production in cheeses was different depending on the different ripening time (factor A) and/or on
404 the different *L. lactis* starter strain (factor B). Results showed that lactose was completely
405 converted into lactate by starters metabolism or lost into whey after salting step by starters K and
406 M while after 15 days by B ($F= 3179.22$, $P < 0.001$). Glucose was not detected neither in milk nor in
407 cheese samples, whereas galactose was present at low concentrations from manufacturing (DRY)
408 until the end of ripening and affected more significantly by factor time ($F=183.47$, $P < 0,05$) than
409 factor starter ($F=15.39$, $P < 0,05$). Lactic acid was the most abundant organic acid in all the ripened
410 samples analysed. Its production, significantly affected by factor time, increased until to
411 represent 30%, 25% and 25% of the total organic acid content in 180 days ripened cheeses made
412 with starter B, M and K, respectively. Citric acid, already detected in milk samples, was significantly
413 consumed throughout manufacturing and ripening ($F= 651.18$, $P < 0.01$), due to its conversion into
414 acetate, acetoin and diacetyl. Acetic acid, not significantly affected neither by factor A nor factor B,
415 was present at low concentrations and reached the maximum values of 0.62, 1.65 and 1.70 mg/g
416 in cheeses at 180 days, manufactured with starters B, K and M, respectively. Highest
417 concentrations of pyruvic acid were detected during manufacturing (DRY samples); then, a light
418 decrease was observed until 180 days of ripening when cheeses made with starter B showed
419 higher concentration than K and M ($F= 133.33$, $P < 0.001$). Propionic acid concentration, affected
420 by factor time ($F= 9103.23$, $P < 0.01$), increased throughout manufacture and early ripening without
421 significant differences between starters. Then, its trend slightly decreased and at 180 days of
422 ripening, it represented 3.6% of total organic acids. Butyric acid, affected by both factors, time and

423 starter, formed during manufacturing and increased significantly during late ripening stages (F=
424 7925.47, P< 0.001) reaching higher value for starter B at 180 days (F= 252, P< 0.001).

425

426 **4. Discussion**

427 Microbiological analysis showed that *L. lactis* was present with the highest loads of 8-9 Log CFU/g
428 during manufacturing and early ripening and it started to decrease, markedly, at 30 days of
429 ripening until not being anymore detected in late stages. After salting, at 15 ripening days, an initial
430 slight decrement was observed. It is known that salt addition promotes starter permeabilisation and
431 cell autolysis during early stages of ripening (Rulikowska et al., 2013; Wilkinson et al., 1994), which
432 allow intracellular enzymes to access to the substrates within the cheese matrix (Guinee, 2004;
433 Sheehan et al., 2005). Several authors (McCarthy et al., 2015; Møller et al., 2007; Ramírez-Nuñez
434 and Ruth Romero-Medran, 2011; Rulikowska et al., 2013; Yanachkina et al., 2016) studied the
435 effect, on *L. lactis* cell autolysis, of changes in cheese parameter as salt concentration, pH
436 gradient, fat content, ripening time, and they suggested their significant impact on *L. lactis* growth
437 dynamics. They also underlined that the impact is strictly strain dependent since there is a
438 tolerance degree by *L. lactis* in response to cheese changes. Moreover, Taibi et al. (2011)
439 hypothesized that the reduction of viable counts, following osmotic stress, could be attributed to the
440 entering of the cells in a catabolically and/or anabolically inactive state, which made them unable to
441 multiply, more than to a process of autolysis. According with these findings, we observed, in a
442 previous work (Ruggirello et al., 2016), a strain dependent attitude of *L. lactis* to enter in a VBNC
443 state as well as to resuscitate on an enrichment medium.

444 In agreement with these observations and considering microbial count results, starters B and M
445 behaved more similarly than K, dropping at 60 days from 6-7 to 3 Log CFU/g, whereas K
446 decreased slowly to 5 Log CFU/g and disappeared in 90 day cheeses. This could be due to the
447 different capability of *L. lactis* starters to response to environmental variations which occur during
448 ripening process (Dugat-Bony et al., 2015). Particularly, the high NaCl content may transform a
449 part of the lactococcal population into VBNC cells or intact dead cells, which are only detected by
450 culture-independent methods as PCR-TTGE (Casalta et al., 2009). Nevertheless, some authors

451 (Broadbent et al., 2013; Muehlenkamp-Ulate and Warthesen, 1999) found *L. lactis* starters, by
452 plating technique, until six months of ripening in Cheddar cheese, with loads of 3-6 Log CFU/g.
453 Their explanation was that microbial population dynamics are altered by perturbations in cheese
454 microenvironment (e.g., acid and salt in moisture (S/M) contents, redox, and water activity).
455 A RT-qPCR protocol, selective for *L. lactis*, was applied to the total microbial RNA extracted
456 directly from milk, curd and cheese matrices, overcoming the issues related to the cultivation step.
457 No statistical differences were observed between three starters by RT-qPCR. Results showed the
458 presence of *L. lactis* in all miniature cheeses, up to the sixth month of ripening confirming the
459 presence of viable cells of this microorganism during the whole ripening process. No statistical
460 differences were observed between the three starters by RT-qPCR. Data obtained by direct *L.*
461 *lactis* RNA analysis were compared with data observed by traditional plating and a similar trend
462 was observed in samples taken at manufacturing and early ripening stages. On the contrary,
463 starting from cheeses at 60 days of ripening, counts detected by direct RNA analysis were
464 markedly higher than values obtained on M17 and, finally, *L. lactis* was not more found by
465 traditional plating. This discrepancy highlighted, one more time, a probably entry of *L. lactis* in a
466 VBNC state as hypothesized in our previous work (Ruggirello et al., 2016) and by other authors
467 (Desfossés-Foucault et al., 2013; Dolci et al., 2014, 2010; Flórez and Mayo, 2006; Masoud et al.,
468 2011; Rantsiou et al., 2008). These evidences support the thesis that cultural media cannot be
469 used to enumerate or detect VBNC cells (Colwel, 2009) and underline the usefulness of direct
470 RNA analysis in assessing the viability of microorganisms, which could retain the ability of
471 synthesizing RNA, although losing the capability to form colonies on synthetic medium. However,
472 purity and integrity of RNA are critical elements for the overall success of RNA-based analyses
473 (Fleige and Pfaffl, 2006) and getting good quality nucleic acids can be tough and laborious when
474 the matrix is complex as cheese. A polyphasic approach, combining culture-dependent and -
475 independent methods, should minimize the biases in both methods, showing a more accurate view
476 of the structure of microbial communities (Ampe et al., 1999; Delbès et al., 2007; Ercolini D,
477 Moschetti G, Blaiotta G, 2001; Flórez and Mayo, 2006).

478 Our data were in agreement with results reported by other authors. Desfossés-Foucault et al.
479 (2014, 2013) found, by RT-qPCR, that *L. lactis* retained transcriptional activity for six month of
480 Cheddar ripening while populations rapidly declined to totally disappear, in early ripening stages,
481 on M17 medium; Dolci et al. (2014) found *L. lactis*, by RT-PCR-DGGE and pyrosequencing, in
482 Fontina PDO cheese, up to three months of ripening. Similarly, other authors detected the
483 microorganism, by culture-independent methods, in the following ripened products: Danish
484 cheeses (Masoud et al., 2011), Castelmagno PDO (Dolci et al., 2010), Feta PDO (Rantsiou et al.,
485 2008), Toma PDO, Raschera PDO, Asiago PDO and Pecorino sardo PDO (Ruggirello et al., 2014)
486 and Celenzana (Casalta et al., 2009).

487 Our results partially differed from data found in our previous study (Ruggirello et al., 2016) where
488 the same *L. lactis* starters were used in model cheeses with the only difference that no strict
489 aseptic conditions were used during manufacturing and, probably, for this reason, higher microbial
490 counts were found on M17 during the ripening due to the growth of non- starter lactic acid bacteria
491 (NSLAB). Furthermore, in the present research, the three starters entered in VBNC state earlier
492 compared to the performance showed in the previous study (Ruggirello et al., 2016). This fact
493 could be related to physiological stresses they underwent during the two similar but probably not
494 identical experimental manufacturing conditions. These results support, once again, the hypothesis
495 that *L. lactis* entering in VBNC state is not only strain dependent, but is also affected by the
496 different environmental parameters and changes. Stuart et al. (1999) demonstrated the different
497 ability of *L. lactis* strains to exhaust lactose, which delayed their entry in VBNC state. Sugar
498 starvation has been recognized as a key stress in lactococci, although also others stress factors,
499 as pH or temperature, can contribute to VBNC state, during fermentation process (Ganesan et al.,
500 2007; Weimer, 2011).

501 In order to evaluate if *L. lactis*, in addition to being alive during ripening time, was also
502 metabolically active in terms of technological gene expression, *metC* and *als* genes were chosen
503 to be studied because involved in important aroma production pathways. Particularly, *metC* gene
504 encodes C β L (Alting et al., 1995) that can convert cystathionine to homocysteine but is also able to
505 catalyse an α,γ elimination. Starting from methionine as substrate, it catalyses the production of

506 volatile sulphur compounds (DMDS and DMTS) which are important flavours in cheese
507 (McSweeney and Sousa, 2000; Rutten et al., 2000). DMDS is associated with sensory attribute of
508 garlic, cheesy or sulphur aroma (Rijnen et al., 2003) while DMTS is related to aroma of cooked
509 cabbage, broccoli or cauliflower (McSweeney and Sousa, 2000). Fernández et al. (2011) found
510 that inactivation of C β L does not affect the formation of VOCs and that other genes, homologous to
511 cystathionine lyase and aminotransferases, are present in the genome of *L. lactis*, and they could
512 also contributed to VOCs production (Bolotin et al., 2001). *Als* gene encodes α -ALS (Platteeuw et
513 al., 1995; Toonen et al., 1994) involved in the biosynthetic pathway of diacetyl/acetoin and 2,3-
514 butanediol, which are the principal flavour compounds produced from citrate metabolism (Hassan
515 et al., 2013). Diacetyl is an important aroma compound in a number of cheese varieties, including
516 Dutch-type cheeses, Quarg and Cottage cheese (McSweeney and Sousa, 2000). The *metC* and
517 *als* genes expression results were showed in terms of fold change during manufacturing and
518 ripening time respect to time 0, corresponding to milk samples after inoculum of the starter (MILK).
519 The *metC* gene was either down or over expressed during manufacturing and ripening time in
520 respect to the inoculated milk samples. These dynamics reflect the several perturbations which
521 happen during cheese making and ripening processes. The three starters behaved differently. In
522 particular starter K showed an up regulation after salting step and starter B during late ripening.
523 Differently, *als* gene was down regulated in all samples with the exception of cheeses at 180 days
524 of ripening and produced with starter B, where the gene was over expressed. Then, the presence
525 of aroma compounds related with *metC* and *als* gene pathways were evaluated. Notably, DMDS
526 was detected, even if in small amounts, in all cheeses, throughout manufacturing and ripening with
527 higher amounts in starter B and K than in starter M. On the contrary, acetoin and diacetyl ketones
528 and 2,3-butanediol alcohol were among the most abundant molecules found during the whole
529 process. Usually, acetoin is produced in much higher quantities than diacetyl whereas production
530 of 2,3- butanediol has never been studied in detail (Hassan et al., 2013). Also, despite its
531 importance, the exact reactions which result in the formation of diacetyl remain unclear. In addition,
532 aroma production varied according to the starters used for cheesemaking. Thus, higher amounts of
533 acetoin were detected in cheeses produced with starter K, while diacetyl and 2,3-butanediol were

534 more abundant in cheeses produced with starter K and M. Finally, we evaluated if any correlation
535 could be hypothesized between gene expression and aroma analysis. While *metC* gene
536 expression could be related to the production of DMDS, on the contrary, the presence of acetoin,
537 diacetyl and 2,3-butanediol in cheeses could not be correlated with down regulation of *als* gene
538 during manufacturing and ripening time. It could be hypothesized that the early *als* gene
539 expression, after starter inoculum in milk, led to release of α -acetolactate, which was then
540 converted to aroma compounds during ripening time. Surprisingly, a marked up expression of *als*
541 gene in cheeses produced at 180 days with starter B was detected. These results highlight a
542 transcriptional activity, even if minimal, in *L. lactis* also during late ripening, supposedly related to
543 the production of flavour compounds, according with observations reported by Desfossés-Foucault
544 et al. (2014). Moreover, our results confirmed, especially for *metC*, a starter's effect. This
545 underlines the importance to emphasize, as already known, starter characteristics in strain
546 selection.

547 Volatile compound analysis by GC-MS showed an interesting separation, in terms of aroma
548 production, between milk, early production and ripening time. Higher amounts of ketones and
549 alcohols were detected during manufacturing and in early ripened cheeses while high contents of
550 acids were observed during late ripening. Ketones are formed by enzymatic oxidation of free fatty
551 acids to β -ketoacids and their consequent decarboxylation to ketones. They are very important
552 compounds for dairy products because they have very particular odours and low perception
553 thresholds (McSweeney, 2004; McSweeney and Sousa, 2000). Their presence during
554 manufacturing could corroborate higher *als* gene expression in MILK than during ripening time.
555 Finally, the trend of the different compounds detected and quantified by HPLC was comparable
556 with the one usually found in commercial cheeses, confirming the experimental reliability of these
557 model cheeses.

558

559 **5. Conclusion**

560 The results reported in the present study add new insights on the capability of *L. lactis* to persist in
561 cheese, during late ripening and up to six months, as metabolically active cells in a hypothetical

562 VBNC state, suggesting a possible technological contribution of the microorganism also during
563 ripening. The expression of *metC* and *als* genes in *L. lactis*, partially associated with the presence
564 of volatile aroma compounds in ripened cheeses, could support our thesis. In addition, the *L. lactis*
565 starters used in this study for the production of model cheeses behaved differently, in part,
566 highlighting, once again, the importance of a selection at strain level in new starter formulation.
567 Data obtained from this study could be considered a first preliminary contribute to the development
568 of new potential tools for starter selection. Nevertheless, additional efforts will be needed to better
569 interpret *L. lactis* VBNC state in cheese and to investigate its potential contribution to flavour
570 formation in late ripening.

571

572 **Author Contributions**

573 Conceived and designed the experiments: LC, PD, MR. Performed the experiments: MR, MG, MB.
574 Analysed the data: MR, IF, MB. Contributed reagents/materials/analysis tools: LC, PD. Wrote the
575 paper: MR.

576

577 **References**

- 578 Alting, A.C., Engels, W.J.M., Van Schalkwijk, S., Exterkate, F.A., 1995. Purification and
579 characterization of cystathionine α -lyase from *Lactococcus lactis* subsp. *cremoris* B78 and its
580 possible role in flavor development in cheese. *Appl. Environ. Microbiol.* 61, 4037–4042.
- 581 Ampe, F., Ben Omar, N., Moizan, C., Wacher, C., Guyot, J.P., 1999. Polyphasic study of the
582 spatial distribution of microorganisms in Mexican pozol, a fermented maize dough,
583 demonstrates the need for cultivation-independent methods to investigate traditional
584 fermentations. *Appl. Environ. Microbiol.* 65, 5464–5473.
- 585 Bertolino, M., Zeppa, G., Gerbi, V., McSweeney, P.L.H., 2008. Study of proteolysis in miniature
586 toma piemontese cheese made using wild bacteria. *Ital. J. Food Sci.* 20, 57–73.
- 587 Bolotin, A., Wincker, P., Mauger, S., Jaillon, O., Malarme, K., Weissenbach, J., Ehrlich, S.D.,
588 Sorokin, A., 2001. The Complete Genome Sequence of the Lactic Acid Bacterium. *Genome*
589 *Res.* 731–753.

590 Broadbent, J.R., Brighton, C., McMahon, D.J., Farkye, N.Y., Johnson, M.E., Steele, J.L.L., J, D.,
591 Farkye, Y., 2013. Microbiology of Cheddar cheese made with different fat contents using a
592 *Lactococcus lactis* single-strain starter. J. Dairy Sci. 96, 4212–4222.

593 Casalta, E., Sorba, J.-M., Aigle, M., Ogier, J.-C., 2009. Diversity and dynamics of the microbial
594 community during the manufacture of Calenzana, an artisanal Corsican cheese. Int. J. Food
595 Microbiol. 133, 243–51.

596 Cathala, G., Savouret, J.-F., Mendez, B., West, B.L., Karin, M., Martial, J.A., Baxter, J.D., 1983. A
597 Method for Isolation of Intact, Translationally Active Ribonucleic Acid. DNA 2, 329–335.

598 Cocolin, L., Manzano, M., Cantoni, C., Comi, G., 2001. Denaturing gradient gel electrophoresis
599 analysis of the 16S rRNA gene V1 region to monitor dynamic changes in the bacterial
600 population during fermentation of Italian sausages. Appl. Environ. Microbiol. 67, 5113–21.

601 Colwel, R.R., 2009. Viable but Not Cultivable Bacteria, in: Uncultivated Microorganisms. Volume
602 10 of the Series Microbiology Monographs. pp. 121–129.

603 Delbès, C., Ali-Mandjee, L., Montel, M.C., 2007. Monitoring bacterial communities in raw milk and
604 cheese by culture-dependent and -independent 16S rRNA gene-based analyses. Appl.
605 Environ. Microbiol. 73, 1882–1891.

606 Desfossés-Foucault, É., Lapointe, G., Roy, D., 2013. Dynamics and rRNA transcriptional activity of
607 lactococci and lactobacilli during Cheddar cheese ripening. Int. J. Food Microbiol. 166, 117–
608 124.

609 Desfossés-Foucault, É., LaPointe, G., Roy, D., 2014. Transcription profiling of interactions between
610 *Lactococcus lactis* subsp. *cremoris* SK11 and *Lactobacillus paracasei* ATCC 334 during
611 Cheddar cheese simulation. Int. J. Food Microbiol. 178, 76–86.

612 Dolci, P., Alessandria, V., Rantsiou, K., Bertolino, M., Cocolin, L., 2010. Microbial diversity,
613 dynamics and activity throughout manufacturing and ripening of Castelmagno PDO cheese.
614 Int. J. Food Microbiol. 143, 71–5.

615 Dolci, P., De Filippis, F., La Storia, A., Ercolini, D., Cocolin, L., 2014. rRNA-based monitoring of the
616 microbiota involved in Fontina PDO cheese production in relation to different stages of cow
617 lactation. Int. J. Food Microbiol. 185, 127–35.

618 Dugat-Bony, E., Straub, C., Teissandier, A., Onésime, D., Loux, V., Monnet, C., Irlinger, F.,
619 Landaud, S., Leclercq-Perlat, M.-N., Bento, P., Fraud, S., Gibrat, J.-F., Aubert, J., Fer, F.,
620 Guédon, E., Pons, N., Kennedy, S., Beckerich, J.-M., Swennen, D., Bonnarne, P., 2015.
621 Overview of a Surface-Ripened Cheese Community Functioning by Meta-Omics Analyses.
622 PLoS One 10, e0124360.

623 Edmund R.S. Kunji, Mierau, I., Hagfing, A., Poolman, B., Konings, W.N., 1993. The proteolytic
624 system of lactic acid bacteria. *Microbiologia* 9, 90–106.

625 Ercan, O., Wels, M., Smid, E.J., Kleerebezem, M., 2015. Genome-Wide Transcriptional Responses
626 to Carbon Starvation in Nongrowing *Lactococcus lactis*. *Appl. Environ. Microbiol.* 81, 2554–
627 2561.

628 Ercolini D, Moschetti G, Blaiotta G, C.S., 2001. The potential of a polyphasic PCR-dGGE approach
629 in evaluating microbial diversity of natural whey cultures for water-buffalo Mozzarella cheese
630 production: bias of culture-dependent and culture-independent analyses. *Syst. Appl.*
631 *Microbiol.* 24, 610–7.

632 Fernández, E., Alegría, Á., Delgado, S., Martín, M.C., Mayo, B., 2011. Comparative phenotypic
633 and molecular genetic profiling of wild *Lactococcus lactis* subsp. *lactis* strains of the *L. lactis*
634 subsp. *lactis* and *L. lactis* subsp. *cremoris* genotypes, isolated from starter-free cheeses made
635 of raw milk. *Appl. Environ. Microbiol.* 77, 5324–5335.

636 Fleige, S., Pfaffl, M.W., 2006. RNA integrity and the effect on the real-time qRT-PCR performance.
637 *Mol. Aspects Med.* 27, 126–139.

638 Flórez, A.B., Mayo, B., 2006. Microbial diversity and succession during the manufacture and
639 ripening of traditional, Spanish, blue-veined Cabrales cheese, as determined by PCR-DGGE.
640 *Int. J. Food Microbiol.* 110, 165–71.

641 Ganesan, B., Stuart, M.R., Weimer, B.C., 2007. Carbohydrate starvation causes a metabolically
642 active but nonculturable state in *Lactococcus lactis*. *Appl. Environ. Microbiol.* 73, 2498–512.

643 García-cayuela, T., Cadiñanos, L.P.G. De, Peláez, C., Requena, T., 2012. Expression in
644 *Lactococcus lactis* of functional genes related to amino acid catabolism and cheese aroma
645 formation is influenced by branched chain amino acids Glutamic acid Methanethiol Carboxylic

646 acid Alcohol. Int. J. Food Microbiol. 159, 207–213.

647 Guinee, T.P., 2004. Salting and the role of salt in cheese. Int. J. Dairy Technol. 57, 99–109.

648 Hassan, F., El-Gawad, M.A., Enab, A., 2013. Flavour Compounds in Cheese (Review). Res.
649 Precis. Instrum. Mach. 2, 15–29.

650 Hugenholtz, J., Starrenburg, M.J.C., 1992. Diacetyl production by different strains of *Lactococcus*
651 *lactis* subsp. *lactis* var. *diacetylactis* and *Leuconostoc* spp. Appl. Microbiol. Biotechnol. 38,
652 17–22.

653 Kubista, M., Sindelka, R., 2007. The Prime Technique. G.I.T. Lab. J. 9–10, 33–35.

654 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
655 quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods 25, 402–408.

656 Marchiani, R., Bertolino, M., Ghirardello, D., McSweeney, P.L.H., Zeppa, G., 2016.
657 Physicochemical and nutritional qualities of grape pomace powder-fortified semi-hard
658 cheeses. J. Food Sci. Technol. 53, 1585–1596.

659 Masoud, W., Takamiya, M., Vogensen, F.K., Lillevang, S., Al-Soud, W.A., Sørensen, S.J.,
660 Jakobsen, M., 2011. Characterization of bacterial populations in Danish raw milk cheeses
661 made with different starter cultures by denaturing gradient gel electrophoresis and
662 pyrosequencing. Int. Dairy J. 21, 142–148.

663 Mataragas, M., Rovetto, F., Bellio, A., Alessandria, V., Rantsiou, K., Decastelli, L., Cocolin, L.,
664 2015. Differential gene expression profiling of *Listeria monocytogenes* in Cacciatore and
665 Felino salami to reveal potential stress resistance biomarkers. Food Microbiol. 46, 408–417.

666 McCarthy, C.M., Wilkinson, M.G., Kelly, P.M., Guinee, T.P., 2015. Effect of salt and fat reduction
667 on the composition, lactose metabolism, water activity and microbiology of Cheddar cheese.
668 Dairy Sci. Technol. 95, 587–611.

669 McSweeney, P.L.H., 2004. Biochemistry of cheese ripening. Int. J. Dairy Technol. 57, 127–144.

670 McSweeney, P.L.H., Sousa, M.J., 2000. Biochemical pathways for the production of flavour
671 compounds in cheeses during ripening: A review. Lait 80, 293–324.

672 Møller, K.K., Rattray, F.P., Bredie, W.L.P., Høier, E., Ardö, Y., 2007. Physicochemical and sensory
673 characterization of Cheddar cheese with variable NaCl levels and equal moisture content. J.

674 Dairy Sci. 96, 1953–1971.

675 Muehlenkamp-Ulate, M.R., Warthesen, J.J., 1999. Evaluation of several non starter lactobacilli for
676 their influence on Cheddar cheese slurry proteolysis. J. Dairy Sci. 82, 1370–1378.

677 Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR.
678 Nucleic Acids Res. 29, e45.

679 Platteeuw, C., Hugenholtz, J., Starrenburg, M., Alen-boerrigter, I.V.A.N., Vos, W.M.D.E., 1995.
680 Metabolic Engineering of *Lactococcus lactis*: Influence of the Overproduction of α -
681 Acetolactate Synthase in Strains Deficient in Lactate Dehydrogenase as a Function of Culture
682 Conditions. Appl. Environ. Microbiol. 61, 3967–3971.

683 Ramírez-Nuñez, J., Ruth Romero-Medran, G.V.N.-M.N.G.-M., 2011. Effect of pH and salt gradient
684 on the autolysis of *Lactococcus lactis* strains. Brazilian J. Microbiol. 42, 1495–1499.

685 Rantsiou, K., Urso, R., Dolci, P., Comi, G., Cocolin, L., 2008. Microflora of Feta cheese from four
686 Greek manufacturers. Int. J. Food Microbiol. 126, 36–42.

687 Rijnen, L., Yvon, M., van Kranenburg, R., Courtin, P., Verheul, A., Chambellon, E., Smit, G.,
688 Kranenburg, R. Van, Courtin, P., Verheul, A., Chambellon, E., Smit, G., 2003. Lactococcal
689 aminotransferases AraT and BcaT are key enzymes for the formation of aroma compounds
690 from amino acids in cheese. Int. Dairy J. 13, 805–812.

691 Ruggirello, M., Cocolin, L., Dolci, P., 2016. Fate of *Lactococcus lactis* starter cultures during late
692 ripening in cheese models. Food Microbiol. 59, 112–118.

693 Ruggirello, M., Dolci, P., Cocolin, L., 2014. Detection and Viability of *Lactococcus lactis* throughout
694 Cheese Ripening. PLoS One 9, e114280.

695 Rulikowska, A., Kilcawley, K.N., Doolan, I.A., Alonso-gomez, M., Nongonierma, A.B., Hannon, J.A.,
696 Wilkinson, M.G., 2013. The impact of reduced sodium chloride content on Cheddar cheese
697 quality. Int. Dairy J. 28, 45–55.

698 Rutten, G.E.R.A.M., Marugg, J.D., Alting, A.C., Kuipers, O.P., 2000. Molecular and Functional
699 Analyses of the *metC* Gene of *Lactococcus lactis*, Encoding Cystathionine β -Lyase. Appl.
700 Environ. Microbiol. 66, 42–48.

701 Sheehan, A., O'loughlin, C., O'Cuinn, G., Fitzgerald, R.J., Wilkinson, M.G., 2005. Cheddar cheese

702 cooking temperature induces differential lactococcal cell permeabilization and autolytic
703 responses as detected by flow cytometry: implications for intracellular enzyme accessibility. J.
704 Appl. Microbiol. 99, 1007–18.

705 Smit, G., Smit, B.A., Engels, W.J.M., 2005. Flavour formation by lactic acid bacteria and
706 biochemical flavour profiling of cheese products. FEMS Microbiol. Rev. 29, 591–610.

707 Snoep, J.L., Teixeira de Mattos, M.J., Starrenburg, M.J., Hugenholtz, J., 1992. Isolation,
708 characterization, and physiological role of the pyruvate dehydrogenase complex and alpha-
709 acetolactate synthase of *Lactococcus lactis* subsp. *lactis* bv. *diacetylactis*. J. Bacteriol. 174,
710 4838–41.

711 Taïbi, A., Dabour, N., Lamoureux, M., Roy, D., LaPointe, G., 2011. Comparative transcriptome
712 analysis of *Lactococcus lactis* subsp. *cremoris* strains under conditions simulating Cheddar
713 cheese manufacture. Int. J. Food Microbiol. 146, 263–75.

714 Toonen, M.Y., Verhue, W.M., Verrips, C.T., 1994. Identification and Characterization of the ox-
715 Acetolactate Synthase Gene from *Lactococcus lactis* subsp. *lactis* biovar *diacetylactis*. Appl.
716 Environ. Microbiol. 60, 1390–1394.

717 Van Hylckama Vlieg, J.E., Rademaker, J.L., Bachmann, H., Molenaar, D., Kelly, W.J., Siezen, R.J.,
718 2006. Natural diversity and adaptive responses of *Lactococcus lactis*. Curr. Opin. Biotechnol.
719 17, 183–190.

720 Verhue, W.M., Tjan, F.S.B., 1991. Study of the citrate metabolism of *Lactococcus lactis* subsp.
721 *lactis* biovar *diacetylactis* by means of ¹³C nuclear magnetic resonance. Appl. Environ.
722 Microbiol. 57, 3371–3377.

723 Weimer, B.C., 2007. Improving the flavour of cheese. CRC Press/Woodhead, Boca Raton.

724 Weimer, B.C., 2011. Responses of lactic acid bacteria to starvation., in: Tsakalidou, E.,
725 Papadimitriou, K. (Eds.), Stress Responses of Lactic Acid Bacteria. Springer US, pp. 129–
726 144.

727 Wilkinson, M.G., Guinee, T.P., Fox, P.F., 1994. Factors which may influence the determination of
728 autolysis of starter bacteria during cheddar cheese ripening. Int. Dairy J. 4, 141–160.

729 Yanachkina, P., McCarthy, C., Guinee, T., Wilkinson, M., 2016. Effect of varying the salt and fat

730 content in Cheddar cheese on aspects of the performance of a commercial starter culture
731 preparation during ripening. *Int. J. Food Microbiol.* 224, 7–15.

732 Yarlagadda, A.B., Wilkinson, M.G., Ryan, S.P., Doolan, I.A., Sullivan, M.G.O., 2014. Utilisation of a
733 cell-free extract of lactic acid bacteria entrapped in yeast to enhance flavour development in
734 Cheddar cheese. *Int. J. Dairy Technol.* 67, 21–30.

735