



UNIVERSITÀ DEGLI STUDI DI TORINO

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Parity, breastfeeding and risk of coronary heart disease: A pan-European case-cohort study

This is a pre print version of the following article:
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1624631since 2020-02-28T15:32:28Z
Published version:
DOI:10.1177/2047487316658571
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 2

Parity, breastfeeding and risk of coronary heart disease: a pan-European case-cohort study

2

Sanne A.E. Peters,^{1,2} Yvonne T. van der Schouw,² Angela M. Wood,^{3,4} Michael J. Sweeting, ^{3,4} Karel G.M. Moons,² Elisabete 3 Weiderpass,^{5,6,7,8} Larraitz Arriola,^{*9} Vassiliki Benetou,^{*10,11}, Heiner Boeing,^{*12} Salma T. Butt,^{*13} Françoise Clavel-Chapelon,^{*14} 4 Isabel Drake,^{*15} Diana Gavrila,^{*16,17} Timothy J Key,^{*18} Eleni Klinaki,^{*11} Vittorio Krogh,^{*19} Tilman Kühn,^{*20} Camille Lassale,^{*21} 5 Giovanna Masala,^{*22} Giuseppe Matullo,^{*23,24} Melissa Merritt,^{*21} Elena Molina-Portillo,^{*19,25} Conchi Moreno-Iribas,^{*26,27} Therese 6 H. Nøst,^{*5} Anja Olsen,^{*28} N. Charlotte Onland-Moret,^{*2} Kim Overvad,^{*29,30} Salvatore Panico,^{*31} M Luisa Redondo,^{*32} Anne 7 Tjønneland,^{*28} Antonia Trichopoulou,^{*10,11} Rosario Tumino,^{*33} Renée Turzanski-Fortner,^{*20} Ioanna Tzoulaki,^{*21} Maria 8 Wennberg,^{*34} Patrik Wennberg,^{*34} Simon G. Thompson,^{3,4} Emanuele Di Angelantonio,^{3,4} Elio Riboli,²¹ Nicholas J. Wareham,³⁵ 9 John Danesh, ^{3,4} Adam S. Butterworth ^{3,4,36} 10

11 *Denotes authors are listed alphabetically

- 12 1. The George Institute for Global Health, University of Oxford, Oxford, UK
- 13 2. Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands
- 14 3. Cardiovascular Epidemiology Unit, Department of Public Health & Primary Care, University of Cambridge, UK
- The National Institute for Health Research Blood and Transplant Unit (NIHR BTRU) in Donor Health and Genomics at the University of Cambridge, UK
- Department of Community Medicine, Faculty of Health Sciences, University of Tromsø, The Arctic University of
 Norway, Tromsø, Norway
- 19 6. Department of Research, Cancer Registry of Norway, Oslo, Norway
- 20 7. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
- 21 8. Genetic Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland
- 22 9. Public Health Division of Gipuzkoa, Instituto Bio-Donostia, Basque Government, CIBERESP, Spain
- WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health,
 Department of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Athens, Greece
- 25 11. Hellenic Health Foundation, Athens, Greece
- 26 12. Department of Epidemiology, German Institute of Human Nutrition (DIfE), Potsdam-Rehbrücke, Germany
- 27 13. Department of Surgery, Clinical Sciences, Lund University, Skåne University Hospital, Malmö, Sweden
- 14. INSERM, Centre for Research in Epidemiology and Population Health (CESP), U1018, Nutrition, Hormones, and
 Women's Health Team, Institut Gustave Roussy, Villejuif, France
- 30 15. Department of Clinical Science, Lund University, Malmö, Sweden
- 31 16. Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia, Spain
- 32 17. CIBER Epidemiología y Salud Pública (CIBERESP), Spain
- 33 18. Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford
- 34 19. Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy
- 35 20. German Cancer Research Center (DKFZ), Division of Cancer Epidemiology, Heidelberg, Germany
- 36 21. Department of Epidemiology and Biostatistics, Imperial College London, London, UK
- 37 22. Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute ISPO, Florence, Italy
- 38 23. Human Genetics Foundation, Turin, Italy
- 39 24. Department of Medical Sciences, University of Turin, Italy
- 40 25. Escuela Andaluza de Salud Pública. Instituto de Investigación Biosanitaria ibs.GRANADA. Hospitales Universitarios de
 41 Granada/Universidad de Granada, Granada, Spain
- 42 26. Public Health Institute of Navarra, Pamplona, Spain
- 43 27. Red de Investigación en Servicios de Salud en Enfermedades Crónicas, Madrid, Spain
- 44 28. Diet, Genes and Environment, Danish Cancer Society Research Center, Copenhagen, Denmark

- 45 29. Department of Public Health, Section for Epidemiology, Aarhus University, Aarhus, Denmark
- 46 30. Department of Cardiology, Aalborg University Hospital, Aalborg, Denmark
- 47 31. Dipartimento di Medicina Clinica e Chirurgia, Federico II University, Naples, Italy
- 48 32. Public Health Directorate, Asturias, Spain
- 49 33. Cancer Registry and Histopathology Unit, Civic- M.P.Arezzo Hospital, ASP Ragusa , Italy
- 50 34. Nutritional Research, Umeå University, Umeå, Sweden
- 51 35. Medical Research Council Epidemiology Unit, University of Cambridge, Cambridge, UK
- 52 36. Wellcome Trust Sanger Institute, Genome Campus, Hinxton, UK
- 53 Address for correspondence
- 54 Dr. Sanne A.E. Peters
- 55 The George Institute for Global Health, University of Oxford
- 56 34 Broad Street, Oxford OX1 3BD, United Kingdom
- 57 T: +44 1865 617 200 | F: +44 1865 617 202
- 58 E: <u>sanne.peters@georgeinstitute.ox.ac.uk</u>

59 Abstract

- 60 Aim: There is uncertainty about the direction and magnitude the associations between parity, breastfeeding and the risk of
- coronary heart disease (CHD). We examined the separate and combined associations of parity and breastfeeding practices
 with the incidence of CHD later in life among women in a large pan-European cohort study.
- 63 *Methods:* Data were used from EPIC-CVD, a case-cohort study nested within the EPIC prospective study of 520,000
- 64 participants from 10 countries. Information on reproductive history was available for 14,917 women, including 5,138 incident
- 65 cases of CHD. Using Prentice-weighted Cox regression stratified by country, we calculated hazard ratios (HRs) and 95%
- 66 confidence intervals (CIs) for CHD, after adjustment for age, study centre, and several socioeconomic and biological risk
- 67 factors.
- 68
- 69 *Results:* Compared with nulliparous women, the adjusted HR was 1.19 (95% CI: 1.03-1.38) among parous women; HRs were
- higher among women with more children (e.g., adjusted HR: 1.04, 1.27-3.27, for women with ≥5 children). Compared with
- 71 women who did not breastfeed, the adjusted HR was 0.70 (0.50-0.97) among women who breastfed. For childbearing women
- 72 who never breastfed, the adjusted HR was 1.27 (0.72-2.24) compared with nulliparous women, whereas for childbearing
- 73 women who breastfed the adjusted HR was 0.92 (0.62-1.37).
- 74
- 75 *Conclusion:* Having more children was associated with a higher risk of CHD later in life, whereas breastfeeding was associated
- 76 with a lower CHD risk. Women who both had children and breastfed did not have a higher risk of CHD.

77 Introduction

Pregnancy is associated with profound changes in the maternal metabolic system, including weight gain, accumulation of 78 abdominal fat, increased insulin resistance, and higher circulating lipid levels.^{1, 2} While these metabolic changes of pregnancy 79 80 support the growth of the foetus and prepare the mother's body for breastfeeding in the short-term, they may also have a 81 prolonged effect on maternal risk of cardiovascular diseases (CVD). However, previous studies have reported conflicting associations on parity (i.e., the number of live children to whom a woman has given birth) and risk of CVD later in life.³⁻¹⁰ 82 Conversely, since the metabolic changes in pregnancy appear to reverse more quickly and more completely with 83 breastfeeding, it has been proposed that breastfeeding could reduce maternal risk of cardiometabolic diseases.¹¹ Studies 84 85 have reported that, compared with women who have never breastfed, women who have breastfed have favourable cardiometabolic profiles,^{12, 13} and exhibit a lower burden of subclinical cardiovascular disease.¹⁴ There may be a lower risk of 86 developing metabolic syndrome,¹⁵ hypertension,^{16, 17} and type 2 diabetes¹⁸⁻²¹ among women who have breastfed for longer 87 88 cumulative durations. However, it is uncertain whether there is an association between breastfeeding and incident CVD outcomes.^{10, 22-24} Also, while lifetime duration of breastfeeding will be affected by the number of children, it is unknown 89 whether extended duration of breastfeeding for one child is associated with the same extent of inverse association with CHD 90 91 risk as multiple periods of shorter breastfeeding across several pregnancies. Moreover, whether breastfeeding could 92 compensate for the potential effect of parity on CHD risk has not been examined. 93 The EPIC (European Prospective Investigation into Cancer and Nutrition)-CVD study provides an opportunity to 94 address these outstanding issues and to evaluate the separate and combined associations of parity and breastfeeding on the

risk of incident CHD in a large sample of women from diverse European countries. 95

96

97 Methods

98 EPIC-CVD is a large, prospective, case-cohort study nested within the European Prospective Investigation into Cancer and Nutrition (EPIC) study.^{25, 26} Briefly, the EPIC study involves 366,521 women and 153,457 men, mostly aged 35–70 years, 99 100 recruited by 23 centres in 10 European countries (Denmark, France, Germany, Greece, Italy, the Netherlands, Norway, Spain, 101 Sweden, and the United Kingdom) between 1991 and 1999. Participants completed questionnaires on their diet, lifestyle, and 102 medical history, and data were centralized at the International Agency for Research on Cancer (IARC) in Lyon, France. A 103 representative random subcohort of 18,249 participants (62% women), stratified by centre, was selected for the EPIC-CVD 104 project.²⁷ After exclusion of 609 participants with a prior history of myocardial infarction or stroke at baseline, 17,640 105 subcohort members remained. This study complied with the Declaration of Helsinki; ethical review boards of IARC and all 106 local institutions where participants had been recruited gave approval for the study, and all participants gave written 107 informed consent. 108

- 109
- 110

111 Definition and ascertainment of CHD events

- 112 First-time CHD events, whether non-fatal or fatal, as defined by codes 410-414 of the International Classification of Diseases
- 113 Ninth Edition (ICD-9), and codes I20-I25 of the Tenth Edition (ICD-10) were the primary study endpoint. Individual centres
- used different methods to ascertain first-time non-fatal CHD events, including self-report and linkage with morbidity or
- hospital registries. Non-fatal CHD events were further validated by additional review of medical records and/or linkage with
- registries.²⁶ Fatal CHD events were generally ascertained through mortality registries. End of follow-up for CHD events varied
- 117 between centres and ranged between 2003 and 2010.
- 118

119 Study population and measurement

- 120 Of the 16,504 women in EPIC-CVD who did not have a known history of CHD or stroke at baseline and 14,917 women
- 121 provided data on reproductive history. Two EPIC centres (Bilthoven, the Netherlands, and Umea, Sweden) did not assess
- 122 parity and breastfeeding history and thus did not contribute to the analyses. Reproductive history, socioeconomic and
- 123 lifestyle factors, and medical history were assessed once using a self-administrated questionnaire at study baseline. Trained
- health professionals measured blood pressure, weight, height, and waist circumference during a visit to each study centre,
- except in the France and Oxford centres where anthropometry was self-reported. Blood pressure measurements were not
- 126 available for the Norway, Asturias, and Navarra centres. High blood pressure was defined as self-reported hypertension at
- 127 baseline, systolic blood pressure>140 or diastolic blood pressure >90, or self-reported use of hypertension medication. Body
- mass index was calculated as weight divided by the square of height in meters. HDL cholesterol and total cholesterol levels,
- 129 measured in baseline serum samples using a Roche MODULAR ANALYTICS EVO analyser, were available in all centres except
- 130 Norway.
- 131

132 Statistical analyses

Baseline characteristics for the subcohort by parity were presented as means (standard deviation) or medians (interquartile 133 134 range) for continuous variables and as percentages for categorical variables. Cox proportional hazards models, modified for the case-cohort design using the Prentice method²⁸, were used to estimate hazard ratios and 95% confidence intervals for 135 136 first-time CHD by parity and breastfeeding history. Participants contributed only the first CHD outcome (whether non-fatal or 137 fatal) experienced during follow-up, so fatal events that followed non-fatal events were not included. Given the multilevel 138 structure of the data, models were first fitted separately within each country before pooling the country-specific estimates by 139 multivariate random-effects meta-analysis using inverse variance weights. Age was used as the underlying time variable, with 140 entry time defined as the participant's age at recruitment, and exit time as the age of first-time CHD, loss-to-follow-up or censoring at the end of the follow-up, whichever came first. The l^2 statistic was used to quantify the percentage of total 141 142 variability between countries due to between-country heterogeneity. Parity, defined as the number of live births, was 143 categorized as nulliparous (reference), 1 child, 2 children, 3 children, 4 children, or 5 or more children. Breastfeeding history 144 was examined among parous women, comparing women who ever breastfed to women who had never breastfed, and 145 categorized into groups of lifetime duration of breastfeeding (never [reference], >0-<3 months, 3-<6 months, 6-<12 months,

146 12-<23 months, and 23 months or more), and into groups of mean duration of breastfeeding per live-born child (never

147 [reference], >0-<1 months, 1-<3 months, 3-<6 months, and 6 months or more). Group-specific 95% confidence intervals were

148 estimated only from the variances that correspond to the amount of information underlying each group (including the

reference group).²⁹ To assess the combined effect of parity and breastfeeding on CHD risk, we also examined the association

between parity, history of breastfeeding and risk of first-time CHD in models including nulliparous women as the reference

151 group.

152 Models were adjusted for age at study entry and centre (Model I), and then additionally for level of attained 153 education, smoking status, and parity (for breastfeeding history only) (Model II), followed by further adjustment for other 154 confounders and potential mediators (history of high blood pressure, HDL cholesterol, total cholesterol, history of diabetes 155 mellitus, and BMI) (Model III). Model III was used as our primary, most conservative, analyses model. To account for the 156 impact of missing covariate data on our results, we restricted our main analyses to individuals with complete data for all 157 models. Secondary analyses allowed the set of individuals to vary between models and used all individuals with non-missing 158 values for the covariates separately for each model. To investigate whether age or number of live births modified the 159 association between parity or breastfeeding with CHD risk, we calculated the HRs for CHD in women <55 years versus \geq 55 years of age, and in women with 1 or 2 children versus those with 3 children or more. Statistical interaction was evaluated by 160 adding a cross-product term to the country-specific regression models and pooling these using random-effects meta-analysis. 161 162 In a sensitivity analysis, we excluded women younger than 45 in whom reproductive history may not yet be complete. All

163 statistical analyses were performed using STATA, version 12.0 (Stata, College Station, TX).

164

165 Results

The baseline characteristics of the 9,985 women in the subcohort with information on reproductive history are shown in
Table 1. The mean (SD) age at entry was 52.7 (9.1) years, 88% of these women were parous, of these 87% had ever breastfed.
Women who were parous had a lower level of attained education, had higher levels of BMI, and were less likely to smoke

than those who were nulliparous. During a median follow-up of 11.1 years (interquartile range 8.0-13.4), 5,138 of 14,890

170 women developed CHD, of whom 206 were also in the subcohort. Supplementary Table 1 shows the descriptive statistics for

- 171 main characteristics by centre.
- 172

173 Parity and risk of coronary heart disease

174 Of 12,481 women with complete data, 3,336 developed CHD during follow-up. The HR (95% CI) for CHD in parous versus

nulliparous women was 1.25 (1.07, 1.46) in the age- and country stratified model and attenuated to 1.19 (1.03, 1.38)

176 following adjustment for potential confounders and mediators (Table 2). The I² statistic for the multiple-adjusted analyses

177 was 29% (0%-67%), indicating that there was moderate heterogeneity between countries. Compared to nulliparous women,

the multiple-adjusted HRs for CHD were 1.16 (0.89, 1.51), 1.14 (1.02, 1.28), 1.16 (0.96, 1.41), 1.40 (1.18, 1.66), and 2.03 (1.27,

179 3.26) for women with 1, 2, 3, 4, or 5 or more children, respectively (Table 2 and Figure 1). Results were similar in analyses

- 180 including the largest set of women with complete data available, irrespective of incomplete data in subsequent models, or
- 181 when restricting the analyses to women aged 45 years or older at study entry (Supplementary Tables 2 and 3).
- 182

183 Breastfeeding history and risk of coronary heart disease

- Data on breastfeeding history were available on 12,779 parous women and 4348 CHD events; 8,074 women (2,404 CHD
 events) had complete data on all variables. Parous women who had ever breastfed had a multiple-adjusted HR for CHD of
- 186 0.70 (95% CI: 0.50, 0.97) compared with parous women who never breastfed (Table 2). There was substantial heterogeneity
- 187 between countries; the I² statistic was 82% (65%-91%). Compared with parous women who had never breastfed, women with
- a lifetime duration of breastfeeding of >0-<3 months, 3-<6 months, 6-<12 months, 12-<23 months, or 23 months or more had
- a multiple-adjusted HR for CHD of 0.71 (0.59, 0.86), 0.67 (0.57, 0.79), 0.68 (0.55, 0.84), 0.61 (0.52, 0.71), and 0.59 (0.44, 0.79),
- respectively (Figure 2 and Table 2). Similar results were obtained in the analyses on the mean duration of breastfeeding per
- 191 child and CHD risk (Figure 3 and Table 2). Analyses restricted to women 45 years or older or including women with missing
- data on some covariables did not change the findings materially (Supplementary Tables 2 and 3).
- 193
- 194 Combination of parity and breastfeeding and risk of coronary heart disease
- 195 Analyses of the combination of parity and breastfeeding indicated that parous women who had ever breastfed were at a
- similar risk of CHD compared to nulliparous women (HR: 0.92 [0.62, 1.37]; Table 3). There was no significant evidence for a
- 197 higher risk of CHD in parous women who had never breastfed (HR: 1.27 [0.72, 2.24]), although power to detect weak effects
- 198
- 199
- 200 Subgroup analyses

was limited.

Analyses in subgroups of women younger than 55 years versus those 55 years or above at study entry provided no evidence for a different effect of parity by age, and neither did the association between number of children and risk of CHD differ between the age groups (Table 4). A history of breastfeeding was associated with similar reductions in risk of CHD in younger as in older women, and in women with 1 or 2 children as in those with 3 or more children. The associations between lifetime duration of breastfeeding and duration of breastfeeding per child and CHD risk also did not differ between age groups or by parity (Table 4).

207

208 Discussion

- 209 In this case-cohort analysis, nested within the 10-country EPIC prospective cohort study, we examined the separate and
- 210 combined associations of parity and breastfeeding on the risk of incident CHD. Our main findings are threefold.
- 211
- 212 First, we report that parous women are at higher risk of CHD as compared to nulliparous women; with the highest risk seen
- among women who had the most offspring. These findings, therefore, add to the accumulating evidence that repeated

pregnancies could result in an accumulation of cardiometabolic changes, including elevated pro-atherogenic lipid levels,
 accumulation of abdominal fat, endothelial dysfunction, and increased systemic inflammation,^{1, 30} that may have permanent
 effects on the cardiovascular system, leading to a higher risk of CHD later in life.³⁻⁷

217

Second, in agreement with previous observations that breastfeeding is associated with lower risk of the metabolic 218 syndrome,¹⁵ hypertension,^{16, 17} and diabetes,¹⁸⁻²¹ our study supports the existence of inverse associations between 219 breastfeeding and CHD risk. Previous studies have reported that prolonged periods of breastfeeding could have beneficial 220 effects on maternal cardiovascular risk factors, including on lipids, blood pressure, insulin and glucose homeostasis, and body 221 mass index.^{12, 31} Our results extend previous work by supporting the possibility of stronger inverse associations between 222 breastfeeding and CHD risk among women who had longer cumulative lifelong duration of breastfeeding. However, the dose-223 224 response curve between breastfeeding and CHD risk requires further investigation, since it may be context-dependent. For 225 example, a previous study of more than 20,000 Norwegian women reported that breastfeeding was associated with a lower 226 risk of CVD mortality, potentially in a U-shaped fashion, but only among parous women younger than 65 years at study baseline.²² Results from the Nurses' Health Study suggested that the association between lifetime duration of breastfeeding 227 and risk of incident CHD was characterized by a threshold effect; only women with lifetime duration of two years or more had 228 a significantly lower risk of incident CHD compared to women who had never breastfed.²³ Conversely, while a history of 229 breastfeeding was associated with a slightly lower risk of fatal CHD in a cohort of 267,400 women from Shanghai, increasing 230 duration of breastfeeding did not strengthen the association.¹⁰ 231

232

Third, our study constitutes one of the few available analyses of the combined associations of parity and breastfeeding on the incidence of CHD. We found that women who both had children and breastfed did not have a higher risk of CHD, which is consistent with the notion that the physiological changes in pregnancy could reverse more quickly and more completely with breastfeeding, which in turn may confer cardiovascular protection later in life.¹¹

237

Our observations are consistent with the conclusions of previous studies that attempted to disentangle biological processes 238 related to pregnancy from lifestyle factors related to childrearing by comparing results from men and women within the same 239 cohort. For example, a study among men from prospective cohorts in the US found no relationship between the number of 240 children and the paternal risk of CHD,³² whereas there was a positive association between having 6 or more pregnancies and 241 CHD risk among women from the same cohort.⁵ A cross-sectional analysis of men and women from the British Women's Heart 242 243 and Health Study and the British Regional Heart Study reported that having more offspring was associated with higher body mass index for both male and female parents, and with more adverse lipid profiles and diabetes in women only.⁶ However, as 244 245 there was a positive association between parity and CHD among women (but not men) in these cohorts, the authors 246 concluded that the biological effects of pregnancy in women persist into later life.

247

248 The strengths and potential limitations of our study merit consideration. Our analysis maximized power and efficiency by 249 conducting a case-cohort analysis of incident CHD in the large prospective EPIC cohort, thereby focusing measurement of 250 lipids and other biochemical risk factors on the most relevant subset of the cohort. The validity of our findings was enhanced 251 by our ability to adjust for a range of relevant covariates, and by the robustness of our results to a variety of sensitivity and 252 subgroup analyses. The generalisability of our findings was enhanced by the inclusion of women from 10 diverse European 253 countries. However, we cannot discount the possibility that the associations observed in this study were, at least partly, due 254 to unmeasured or residual confounding. For example, confounding is a major concern in studies of breastfeeding and health outcomes, because mothers who breastfeed are more likely to engage in other health-promoting behaviours.³³⁻³⁵ 255 256 Nevertheless, we found that adjustment in our study for several relevant factors did not materially affect the relationships we 257 observed. Our study had insufficient data to account for CHD risk factors before or during pregnancy that determine 258 breastfeeding initiation and duration as well as future CHD risk, leaving our results potential liable to "reverse causality". For 259 example, women with pre-existing CHD risk factors such as obesity, type 1 diabetes, preeclampsia, or polycystic ovary 260 syndrome, might be less likely to initiate breastfeeding and or could breastfeed for shorter durations than women without 261 these CHD risk factors. Because our study involved self-reported information on parity and breastfeeding, information that 262 was sometimes recalled and recorded decades after childbirth and weaning (with the added limitation that duration of 263 breastfeeding was recorded in EPIC only for a woman's first three children and final child), the true strength of any 264 associations we observed could have been underestimated. Data on the number of children in men was not available, so we 265 were no able to dissect whether the association between parity and CHD was due to biological effects of childbearing or factors related to childrearing. Finally, we noted between-country heterogeneity in some of our results, a finding which 266 267 requires further investigation. 268

In conclusion, this analysis of women from 10 European countries found that having more children was associated with a
 higher risk of CHD later in life, whereas breastfeeding was associated with a lower CHD risk. Women who both had children
 and breastfed did not have a higher risk of CHD, suggesting the need for studies to determine whether breastfeeding can
 compensate for the CHD risk associated with greater parity.

273

274 Funding

- 275 EPIC-CVD has been supported by the European Union Framework 7 (HEALTH-F2-2012-279233), the European Research
- 276 Council (268834), the UK Medical Research Council (G0800270 and MR/L003120/1), the British Heart Foundation (SP/09/002
- and RG/08/014 and RG13/13/30194), and the UK National Institute of Health Research. EPIC Asturias was also supported by
- the Regional Government of Asturias. EPIC-Greece is also supported by the Hellenic Health Foundation. EPIC-Oxford was also
- supported by the UK Medical Research Council (MR/M012190/1) and Cancer Research UK (570/A16491). EPIC-Ragusa was
- also supported by the Sicilian Government, AIRE ONLUS Ragusa, and AVIS Ragusa. EPIC-Turin was supported also by the
- 281 Compagnia di San Paolo and the Human Genetics Foundation-Torino (HuGeF).
- 282

283 Acknowledgements

- 284 We thank all EPIC participants and staff for their contribution to the study. We thank staff from the EPIC-CVD and EPIC-
- 285 InterAct Coordinating Centres for carrying out sample preparation and data-handling work, particularly Sarah Spackman
- 286 (EPIC-CVD Data Manager).
- 287
- 288 Conflicts of interest
- 289 None

290 References

- 1. Lain KY, Catalano PM. Metabolic changes in pregnancy. *Clinical obstetrics and gynecology* 2007; **50**(4): 938-48.
- 292 2. Sanghavi M, Rutherford JD. Cardiovascular physiology of pregnancy. *Circulation* 2014; **130**(12): 1003-8.

293 3. Parikh NI, Cnattingius S, Dickman PW, Mittleman MA, Ludvigsson JF, Ingelsson E. Parity and risk of later-life maternal 294 cardiovascular disease. *American heart journal* 2010; **159**(2): 215-21.e6.

Jaffe DH, Eisenbach Z, Manor O. The effect of parity on cause-specific mortality among married men and women.
 Maternal and child health journal 2011; **15**(3): 376-85.

297 5. Ness RB, Harris T, Cobb J, et al. Number of pregnancies and the subsequent risk of cardiovascular disease. *The New* 298 *England journal of medicine* 1993; **328**(21): 1528-33.

299 6. Lawlor DA, Emberson JR, Ebrahim S, et al. Is the association between parity and coronary heart disease due to 300 biological effects of pregnancy or adverse lifestyle risk factors associated with child-rearing? Findings from the British

301 Women's Heart and Health Study and the British Regional Heart Study. *Circulation* 2003; **107**(9): 1260-4.

Koski-Rahikkala H, Pouta A, Pietilainen K, Hartikainen AL. Does parity affect mortality among parous women? *Journal of epidemiology and community health* 2006; **60**(11): 968-73.

- Steenland K, Lally C, Thun M. Parity and coronary heart disease among women in the American Cancer Society CPS II
 population. *Epidemiology (Cambridge, Mass)* 1996; **7**(6): 641-3.
- 306 9. Colditz GA, Willett WC, Stampfer MJ, Rosner B, Speizer FE, Hennekens CH. A prospective study of age at menarche,
- parity, age at first birth, and coronary heart disease in women. *American journal of epidemiology* 1987; **126**(5): 861-70.
- Gallagher LG, Davis LB, Ray RM, et al. Reproductive history and mortality from cardiovascular disease among women
 textile workers in Shanghai, China. *International journal of epidemiology* 2011; **40**(6): 1510-8.
- Stuebe AM, Rich-Edwards JW. The reset hypothesis: lactation and maternal metabolism. *American journal of perinatology* 2009; 26(1): 81-8.
- Gunderson EP, Lewis CE, Wei GS, Whitmer RA, Quesenberry CP, Sidney S. Lactation and changes in maternal
 metabolic risk factors. *Obstetrics and gynecology* 2007; **109**(3): 729-38.
- Schwarz EB, Ray RM, Stuebe AM, et al. Duration of lactation and risk factors for maternal cardiovascular disease.
 Obstetrics and gynecology 2009; **113**(5): 974-82.
- McClure CK, Catov JM, Ness RB, Schwarz EB. Lactation and maternal subclinical cardiovascular disease among
 premenopausal women. *American journal of obstetrics and gynecology* 2012; **207**(1): 46.e1-8.
- 318 15. Gunderson EP, Jacobs DR, Jr., Chiang V, et al. Duration of lactation and incidence of the metabolic syndrome in 319 women of reproductive age according to gestational diabetes mellitus status: a 20-Year prospective study in CARDIA
- 320 (Coronary Artery Risk Development in Young Adults). *Diabetes* 2010; **59**(2): 495-504.
- 16. Lee SY, Kim MT, Jee SH, Yang HP. Does long-term lactation protect premenopausal women against hypertension risk?
 A Korean women's cohort study. *Preventive medicine* 2005; **41**(2): 433-8.
- 323 17. Stuebe AM, Schwarz EB, Grewen K, et al. Duration of lactation and incidence of maternal hypertension: a longitudinal
 324 cohort study. *American journal of epidemiology* 2011; **174**(10): 1147-58.
- 18. Jager S, Jacobs S, Kroger J, et al. Breast-feeding and maternal risk of type 2 diabetes: a prospective study and metaanalysis. *Diabetologia* 2014; **57**(7): 1355-65.
- Stuebe AM, Rich-Edwards JW, Willett WC, Manson JE, Michels KB. Duration of lactation and incidence of type 2
 diabetes. *JAMA : the journal of the American Medical Association* 2005; **294**(20): 2601-10.
- Villegas R, Gao YT, Yang G, et al. Duration of breast-feeding and the incidence of type 2 diabetes mellitus in the
 Shanghai Women's Health Study. *Diabetologia* 2008; **51**(2): 258-66.
- Aune D, Norat T, Romundstad P, Vatten LJ. Breastfeeding and the maternal risk of type 2 diabetes: a systematic
 review and dose-response meta-analysis of cohort studies. *Nutrition, metabolism, and cardiovascular diseases : NMCD* 2014;
- **24**(2): 107-15.
- 22. Natland Fagerhaug T, Forsmo S, Jacobsen GW, Midthjell K, Andersen LF, Ivar Lund Nilsen T. A prospective population-
- based cohort study of lactation and cardiovascular disease mortality: the HUNT study. *BMC public health* 2013; **13**: 1070.
- 336 23. Stuebe AM, Michels KB, Willett WC, Manson JE, Rexrode K, Rich-Edwards JW. Duration of lactation and incidence of 337 myocardial infarction in middle to late adulthood. *American journal of obstetrics and gynecology* 2009; **200**(2): 138 e1-8.
- myocardial infarction in middle to late adulthood. *American journal of obstetrics and gynecology* 2009; **200**(2): 138 e1
 Merritt MA, Riboli E, Murphy N, et al. Reproductive factors and risk of mortality in the European Prospective
- 339 Investigation into Cancer and Nutrition; a cohort study. *BMC medicine* 2015; **13**: 252.

- 25. Riboli E, Hunt KJ, Slimani N, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study
- 341 populations and data collection. *Public health nutrition* 2002; **5**(6b): 1113-24.
- 342 26. Danesh J, Saracci R, Berglund G, et al. EPIC-Heart: the cardiovascular component of a prospective study of nutritional,
- lifestyle and biological factors in 520,000 middle-aged participants from 10 European countries. *European journal of epidemiology* 2007; **22**(2): 129-41.
- 27. Langenberg C, Sharp S, Forouhi NG, et al. Design and cohort description of the InterAct Project: an examination of the
- interaction of genetic and lifestyle factors on the incidence of type 2 diabetes in the EPIC Study. *Diabetologia* 2011; 54(9):
 2272-82.
- Prentice RL. A case-cohort design for epidemiologic cohort studies and disease prevention trials. *Biometrika* 1986; **73**(1): 1-11.
- 29. Easton DF, Peto J, Babiker AG. Floating absolute risk: an alternative to relative risk in survival and case-control analysis avoiding an arbitrary reference group. *Statistics in medicine* 1991; **10**(7): 1025-35.
- 352 30. Stewart FM, Freeman DJ, Ramsay JE, Greer IA, Caslake M, Ferrell WR. Longitudinal assessment of maternal
- endothelial function and markers of inflammation and placental function throughout pregnancy in lean and obese mothers.
 The Journal of clinical endocrinology and metabolism 2007; **92**(3): 969-75.
- 355 31. Natland ST, Nilsen TI, Midthjell K, Andersen LF, Forsmo S. Lactation and cardiovascular risk factors in mothers in a
- population-based study: the HUNT-study. *International breastfeeding journal* 2012; **7**(1): 8.
- 357 32. Ness RB, Cobb J, Harris T, D'Agostino RB. Does number of children increase the rate of coronary heart disease in men?
 358 *Epidemiology (Cambridge, Mass)* 1995; 6(4): 442-5.
- 359 33. Beck LF, Morrow B, Lipscomb LE, et al. Prevalence of selected maternal behaviors and experiences, Pregnancy Risk
- Assessment Monitoring System (PRAMS), 1999. *Morbidity and mortality weekly report Surveillance summaries (Washington, DC : 2002*) 2002; **51**(2): 1-27.
- 362 34. Pesa JA, Shelton MM. Health-enhancing behaviors correlated with breastfeeding among a national sample of
- 363 mothers. Public health nursing (Boston, Mass) 1999; 16(2): 120-4.
- 364 35. Yngve A, Sjostrom M. Breastfeeding determinants and a suggested framework for action in Europe. *Public health*
- 365 *nutrition* 2001; **4**(2b): 729-39.

Figure legend

Figure 1: Adjusted hazard ratios (with group-specific 95% confidence intervals) for incident coronary heart disease associated with number of live born children.

The highest category of live born children (5 or more) is plotted at 5.5.

Figure 2: Adjusted hazard ratios (with group-specific 95% confidence intervals) for incident coronary heart disease associated with lifetime duration of breastfeeding in parous women.

Categories are never, 0-<3 months, 3-<6 months, 6-<12 months, 12-<23 months, and 23 months or more. On the x-axis, results are placed on the mean lifetime duration of breastfeeding within category.

Figure 3: Adjusted hazard ratios (with group-specific 95% confidence intervals) for incident coronary heart disease associated with mean duration of breastfeeding per live born child.

Categories are never, 0-<1 months, 1-<3 months, 3-<6 months, and 6 months or more. On the x-axis, results are placed on the mean duration of breastfeeding per child within category.

Page 13 of 23

	Overall subcohort	Nulliparous	Parous
N (% of overall subcohort)	9,985	1,220 (12.2)	8,765 (87.8)
Age at study entry, years	52.7 (9.1)	52.7 (10.1)	52.7 (9.0)
Education level, %			
None	11.3	7.0	11.9
Primary	34.3	23.9	35.8
Secondary	15.4	18.2	15.0
Tertiary	39.0	50.9	37.4
Current smoker, %	21.7	25.4	21.1
History of diabetes, %	2.6	2.3	2.7
BMI, kg/m ²	26.1 (4.7)	24.9 (4.3)	26.2 (4.7)
Systolic blood pressure, mmHg	130.8 (20.0)	129.6 (20.4)	131.0 (19.9)
Diastolic blood pressure, mmHg	80.3 (10.6)	79.2 (10.4)	80.5 (10.6)
History of high blood pressure, %	34.3	30.1	34.9
Total cholesterol, mmol/L	6.0 (1.1)	6.0 (1.2)	6.0 (1.1)
HDL cholesterol, mmol/L	1.6 (0.4)	1.7 (0.4)	1.6 (0.4)
Postmenopausal, %	52.7	53.9	52.5
Age at menopause [†]	48.5 (4.9)	48.1 (5.3)	48.6 (4.9)
Number of children*, %			
1 child	-	-	14.8
2 children	-	-	47.5
≥3 children	-	-	37.7
History of breastfeeding*, %	-	-	87.2
Lifetime duration of breastfeeding*, months	-	-	6.4 (3.0, 13.0)
Duration of breastfeeding per child*, months	-	-	3.0 (1.5, 6.0)

Table 1: Baseline characteristics of women in the EPIC-CVD subcohort by parity

Values are mean (standard deviation) or median (25th and 75th percentile) for continuous variables. + postmenopausal women only, *parous women only.

	Model I	Model II	Model III
Parity			
N	12,481	12,481	12,481
Parous (yes vs. no)	1.25 (1.07, 1.46)	1.23 (1.05, 1.43)	1.19 (1.03, 1.38)
I ² for heterogeneity (95% CI)	40% (0%, 72%)	40% (0%, 72%)	29% (0%, 67%)
Number of children			
None	1.00 (0.84, 1.19)	1.00 (0.83, 1.20)	1.00 (0.85, 1.18)
1 child	1.18 (0.97, 1.43)	1.13 (0.93, 1.38)	1.16 (0.89, 1.51)
2 children	1.20 (1.10, 1.32)	1.16 (1.12, 1.20)	1.14 (1.02, 1.28)
3 children	1.23 (1.07, 1.41)	1.21 (1.04, 1.41)	1.16 (0.96, 1.41)
4 children	1.51 (1.27, 1.79)	1.49 (1.23, 1.79)	1.40 (1.18, 1.66)
5 or more children	2.28 (1.51, 3.45)	2.05 (1.31, 3.22)	2.03 (1.27, 3.26)
History of breastfeeding			
Ν	8,074	8,074	8,074
Ever breastfed (yes vs. no)	0.68 (0.59, 0.80)	0.61 (0.50, 0.74)	0.70 (0.50, 0.97)
I ² for heterogeneity	54% (0%, 80%)	62% (12%, 83%)	82% (65%, 91%)
Lifetime duration of breastfeeding			
Never breastfed	1.00 (0.83, 1.21)	1.00 (0.83, 1.23)	1.00 (0.73, 1.37)
>0 to <3 months	0.67 (0.59, 0.77)	0.66 (0.57, 0.77)	0.71 (0.59, 0.86)
≥3 to <6 months	0.70 (0.58, 0.85)	0.69 (0.59, 0.81)	0.67 (0.57, 0.79)
≥6 to <12 months	0.72 (0.61, 0.84)	0.67 (0.57, 0.79)	0.68 (0.55, 0.84)
≥12 to <23 months	0.63 (0.57, 0.70)	0.60 (0.52, 0.70)	0.61 (0.52, 0.71)
≥23 months	0.63 (0.48, 0.83)	0.56 (0.43, 0.73)	0.59 (0.44, 0.79)
Duration of breastfeeding per child			
Never breastfed	1.00 (0.88, 1.14)	1.00 (0.85, 1.18)	1.00 (0.74, 1.35)
>0 to <1 months	0.82 (0.71, 0.95)	0.75 (0.64, 0.89)	0.82 (0.68, 0.98)
≥1 to <3 months	0.73 (0.65, 0.81)	0.71 (0.64, 0.79)	0.70 (0.65, 0.76)
≥3 to <6 months	0.69 (0.57, 0.84)	0.66 (0.57, 0.78)	0.69 (0.60, 0.80)
≥6 months	0.64 (0.53, 0.78)	0.66 (0.54, 0.81)	0.66 (0.57, 0.77)

Table 2: Hazard ratios (95% confidence intervals) for incident coronary heart disease associated with parity in all women and history of breastfeeding in parous women only

Hazard ratios are from Prentice-weighted Cox proportional hazards models stratified by country and with age as timescale. Analyses of parity are conducted in all women. Analyses of breastfeeding are conducted in parous women only.

Model I: Adjusted for age at study entry and centre; Model II: model I + level of attained education, smoking status, and number of live born children (for breastfeeding only); Model III: model II + high blood pressure, HDL cholesterol, total cholesterol, history of diabetes mellitus, and BMI.

Table 3: Hazard ratios (95% confidence intervals) for incident coronary heart disease associated with the combined effects of parity and a history of breastfeeding

	Nulliparous	Parous – never breastfed	Parous – ever breastfed
Model I	1.00 [reference]	1.63 (1.26, 2.11)	1.13 (0.95, 1.35)
Model II	1.00 [reference]	1.50 (1.07, 2.11)	0.91 (0.63, 1.31)
Model III	1.00 [reference]	1.27 (0.72, 2.24)	0.92 (0.62, 1.37)

Hazard ratios are from Prentice-weighted Cox proportional hazards models stratified by country and with age as timescale. Model I: Adjusted for age at study entry and centre; Model II: model I + level of attained education, smoking status, and number of live born children; Model III: model II + high blood pressure, HDL cholesterol, total cholesterol, history of diabetes mellitus, and BMI. Table 4: Hazard ratios (95% confidence intervals) for incident coronary heart disease by parity and history of breastfeeding in subgroups

	Age			Parity		
	<55 years	55 years or older	P for	1-2 children	3 or more	P for
			interaction		children	interaction
Parous (yes vs. no)	1.17 (0.91, 1.52)	1.19 (0.89, 1.60)	0.94	-	-	-
Number of children						
1 child	1.37 (0.79, 2.38)	1.07 (0.63, 1.80)	0.76	-	-	-
2 children	1.12 (0.64, 1.96)	1.67 (0.79, 3.53)		-	-	
3 children	1.15 (0.75, 1.77)	1.36 (0.94, 1.96)		-	-	
4 children	1.13 (0.81, 1.57)	1.53 (0.45, 5.19)		-	-	
5 or more children	1.30 (0.91, 1.87)	1.84 (1.20, 2.82)		-	-	
History of breastfeeding (yes vs.						
no)	0.68 (0.50, 0.93)	0.72 (0.50, 1.04)	0.70	0.66 (0.45, 0.95)	0.67 (0.41, 1.08)	0.91
Lifetime duration of						
breastfeeding						
>0 to <3 months	0.74 (0.49, 1.11)	0.67 (0.41, 1.10)	0.31	0.67 (0.45, 0.99)	0.87 (0.53, 1.41)	0.30
≥3 to <6 months	0.51 (0.35, 0.74)	0.68 (0.40, 1.15)		0.63 (0.42, 0.94)	0.79 (0.51, 1.22)	
≥6 to <12 months	0.75 (0.46, 1.22)	0.61 (0.38, 1.01)		0.64 (0.36, 1.14)	0.72 (0.45, 1.17)	
≥12 to <23 months	0.54 (0.34, 0.86)	0.60 (0.34, 1.05)		0.61 (0.37, 1.01)	0.65 (0.37, 1.14)	
≥23 months	0.55 (0.29, 1.04)	0.55 (0.37, 0.81)		1.09 (0.48, 2.48)	0.62 (0.43, 0.89)	
Duration of breastfeeding per						
child						
>0 to <1 months	0.79 (0.54, 1.14)	0.71 (0.44, 1.15)	0.68	0.65 (0.42, 1.01)	0.96 (0.66, 1.40)	0.46
≥1 to <3 months	0.59 (0.42, 0.82)	0.66 (0.43, 1.03)		0.67 (0.46, 0.97)	0.70 (0.44, 1.10)	
≥3 to <6 months	0.62 (0.40, 0.98)	0.63 (0.40, 0.99)		0.63 (0.40, 0.99)	0.70 (0.41, 1.17)	
≥6 months	0.64 (0.44, 0.93)	0.59 (0.36, 0.97)		0.67 (0.43, 1.03)	0.63 (0.39, 1.01)	

Hazard ratios are from Prentice-weighted Cox proportional hazards models stratified by country and with age as timescale. Analyses of parity are conducted in all women; nulliparous women are the reference category. Analyses of breastfeeding are conducted in parous women only; parous women with no history of breastfeeding are the reference category.

Models are adjusted for age at study entry, centre, level of attained education, smoking status, number of live born children (for breastfeeding analyses only), high blood pressure, HDL cholesterol, total cholesterol, history of diabetes mellitus, and BMI.













				-		
Supplamantan	1 Tabla 1	Dacalina	charactorictic	ss for mair	wariahlas h	VEDIC contro
Supplementary	гарет.	Daseille	Characteristic	LS IOF III all	i variabies b	V EPIC CEIILIE

	1							
					Subcohort characteristics			
Country	EPIC centre	Total, N	Subcohort, N	CHD, N	Age at study entry, years	Current smoker, %	Parous, %	Ever breastfed*, %
Denmark	Aarhus	475	312	170	56.2 (4.4)	29.9	87.2	94.1
	Copenhagen	1,048	654	408	56.6 (4.3)	37.7	87.9	92.4
France	France	593	557	38	56.4 (6.5)	9.0	90.5	77.2
Germany	Heidelberg	554	489	68	49.3 (8.4)	20.3	81.8	81.8
	Potsdam	785	709	78	48.7 (9.2)	16.8	90.7	83.7
Greece	Greece	882	766	121	52.6 (12.2)	20.4	90.6	88.3
Italy	Florence	493	420	77	51.6 (7.6)	26.0	85.2	88.2
	Varese	462	293	173	51.3 (8.2)	21.2	90.4	81.9
	Ragusa	216	178	39	45.9 (7.9)	23.0	85.4	89.5
	Turin	271	237	36	51.1 (7.7)	20.3	83.5	72.7
	Naples	313	223	93	50.1 (7.7)	35.4	91.0	91.6
Norway	Norway	105	60	45	48.7 (4.6)	29.3	93.3	96.4
Spain	Asturias	568	480	96	47.6 (8.1)	19.0	90.6	84.7
	Granada	526	436	100	49.4 (8.4)	14.5	93.1	90.4
	Murcia	585	536	56	47.9 (8.5)	18.1	89.4	89.5
	Navarra	450	410	41	48.6 (7.9)	23.2	85.6	93.1
	San Sebastian	466	414	59	48.2 (7.9)	18.1	89.6	86.0
Sweden	Malmo	1,765	1,128	679	57.1 (8.0)	28.9	86.9	95.4
The Netherlands	Utrecht	1,761	914	891	57.7 (6.0)	22.0	85.7	82.6
United Kingdom	Cambridge	1,457	523	970	59.0 (9.4)	14.4	87.8	83.3
	Oxford	1,142	246	900	49.1 (11.6)	6.5	73.6	90.2
Overall		14,917	9,985	5,138	52.7 (9.1)	21.7	87.8	87.2

* Amongst parous women

	Model I	Model II	Model III
Parity			
N	14,890	14,319	12,481
Parous (yes vs. no)	1.19 (1.00, 1.40)	1.18 (0.99, 1.40)	1.19 (1.03, 1.38)
I ² for heterogeneity (95% CI)	60% (21%, 80%)	59% (18%, 80%)	29% (0%, 67%)
Number of children			
None			
1 child	1.23 (1.06, 1.42)	1.14 (0.99, 1.31)	1.16 (0.89, 1.51)
2 children	1.11 (1.00, 1.23)	1.11 (1.09, 1.13)	1.14 (1.02, 1.28)
3 children	1.14 (0.98, 1.32)	1.11 (0.98, 1.25)	1.16 (0.96, 1.41)
4 children	1.47 (1.24, 1.75)	1.45 (1.24, 1.70)	1.40 (1.18, 1.66)
5 or more children	1.92 (1.34, 2.76)	2.05 (1.30, 3.24)	2.03 (1.27, 3.26)
History of breastfeeding			
Ν	12,779	9,466	8,074
Ever breastfed (yes vs. no)	0.68 (0.59, 0.80)	0.61 (0.50, 0.74)	0.70 (0.50, 0.97)
I ² for heterogeneity	55% (8%, 78%)	61% (16%, 82%)	82% (65%, 91%)
Lifetime duration of breastfeeding			
Never breastfed	1.00 (0.86, 1.16)	1.00 (0.83, 1.21)	1.00 (0.73, 1.37)
>0 to <3 months	0.71 (0.59, 0.84)	0.64 (0.53, 0.77)	0.71 (0.59, 0.86)
≥3 to <6 months	0.67 (0.56, 0.81)	0.62 (0.52, 0.74)	0.67 (0.57, 0.79)
≥6 to <12 months	0.66 (0.59, 0.75)	0.62 (0.55, 0.69)	0.68 (0.55, 0.84)
≥12 to <23 months	0.62 (0.58, 0.67)	0.54 (0.49, 0.59)	0.61 (0.52, 0.71)
≥23 months	0.56 (0.46, 0.70)	0.51 (0.41, 0.65)	0.59 (0.44, 0.79)
Mean duration of breastfeeding per child			
Never breastfed	1.00 (0.93, 1.08)	1.00 (0.84, 1.19)	1.00 (0.74, 1.35)
>0 to <1 months	0.79 (0.68, 0.93)	0.68 (0.58, 0.81)	0.82 (0.68, 0.98)
≥1 to <3 months	0.69 (0.64, 0.75)	0.64 (0.57, 0.72)	0.70 (0.65, 0.76)
≥3 to <6 months	0.59 (0.49, 0.71)	0.59 (0.52, 0.67)	0.69 (0.60, 0.80)
≥6 months	0.56 (0.47, 0.67)	0.58 (0.50, 0.67)	0.66 (0.57, 0.77)

Supplementary Table 2: Hazard ratios (95% confidence intervals) for incident coronary heart disease associated with parity and history of breastfeeding in women with complete data on all covariates

27 individuals with no follow-up were excluded from the analyses. Hazard ratios are from Prentice-weighted Cox proportional hazards models stratified by country and with age at study entry as timescale. Analyses of parity are conducted in all women. Analyses of breastfeeding are conducted in parous women only.

Model I: Adjusted for age at study entry and centre; Model II: model I + level of attained education, smoking status, and number of live born children (for breastfeeding analyses only); Model III: model II + high blood pressure, HDL cholesterol, total cholesterol, history of diabetes mellitus, and BMI.

Supplementary Table 3: Hazard ratios (95% confidence intervals) for incident coronary heart disease associated with parity and history of breastfeeding in women aged 45 years or older at study entry

	Model I	Model II	Model III
Parity			
N	12,608	12,049	10,351
Parous (yes vs. no)	1.19 (1.00, 1.42)	1.19 (1.01, 1.41)	1.19 (1.01, 1.39)
I ² for heterogeneity (95% CI)	62% (24%, 81%)	55% (8% <i>,</i> 78%)	38% (0%, 72%)
Number of children			
None	1.00 (0.82, 1.22)	1.00 (0.85, 1.18)	1.00 (0.85, 1.18)
1 child	1.26 (1.11, 1.44)	1.18 (1.01, 1.37)	1.19 (0.93, 1.53)
2 children	1.11 (1.04, 1.18)	1.13 (1.11, 1.15)	1.13 (1.02, 1.26)
3 children	1.15 (0.98, 1.36)	1.14 (1.01, 1.28)	1.15 (0.95, 1.40)
4 children	1.50 (1.21, 1.86)	1.51 (1.28, 1.79)	1.40 (1.17, 1.67)
5 or more children	1.91 (1.34, 2.71)	2.02 (1.29, 3.15)	2.03 (1.27, 3.27)
History of breastfeeding			
Ν	10,832	8,307	6,998
Ever breastfed (yes vs. no)	0.69 (0.59, 0.81)	0.61 (0.50, 0.76)	0.71 (0.51, 1.00)
I ² for heterogeneity	58% (15%, 79%)	66% (27%, 84%)	83% (66%, 91%)
Lifetime duration of breastfeeding			
Never breastfed	1.00 (0.89, 1.13)	1.00 (0.81, 1.24)	1.00 (0.72, 1.38)
>0 to <3 months	0.70 (0.60, 0.82)	0.64 (0.51, 0.80)	0.71 (0.57, 0.87)
≥3 to <6 months	0.68 (0.60, 0.79)	0.61 (0.50, 0.75)	0.69 (0.58, 0.82)
≥6 to <12 months	0.70 (0.58, 0.84)	0.63 (0.56, 0.70)	0.70 (0.57, 0.86)
≥12 to <23 months	0.65 (0.57, 0.74)	0.55 (0.50, 0.61)	0.64 (0.54, 0.75)
≥23 months	0.59 (0.46, 0.76)	0.50 (0.38, 0.65)	0.61 (0.45, 0.82)
Mean duration of breastfeeding per child			
Never breastfed	1.00 (0.90, 1.11)	1.00 (0.83, 1.21)	1.00 (0.73, 1.37)
>0 to <1 months	0.79 (0.67, 0.94)	0.67 (0.56, 0.81)	0.82 (0.67, 1.01)
≥1 to <3 months	0.71 (0.67, 0.75)	0.66 (0.59, 0.73)	0.73 (0.68, 0.79)
≥3 to <6 months	0.61 (0.50, 0.74)	0.61 (0.53, 0.69)	0.71 (0.62, 0.83)
≥6 months	0.59 (0.49, 0.72)	0.59 (0.51, 0.69)	0.70 (0.60, 0.81)

Hazard ratios are from Prentice-weighted Cox proportional hazards models stratified by country and with age at study entry as timescale. Analyses of parity are conducted in all women. Analyses of breastfeeding are conducted in parous women only.

Model I: Adjusted for age at study entry and centre; Model II: model I + level of attained education, smoking status, and number of live born children (for breastfeeding only); Model III: model II + high blood pressure, HDL cholesterol, total cholesterol, history of diabetes mellitus, and BMI.