

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Lung ultrasound in the emergency setting: Accuracy cannot exclude expertise: Response

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1619480> since 2018-11-15T11:57:23Z

Published version:

DOI:10.1378/chest.15-1225

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Lung Ultrasound in the Emergency Setting: Response

Emanuele Pivetta MD^{abcd} Alberto Goffi MD^{ef} Enrico Lupia MD,
PhD^{dg} Maria Tizzani MD^d Giulio Porrino MD^d Enrico Ferreri MD^d Giovanni Volpicelli MD,
FCCP^h Paolo Balzaretto MDⁱ Alessandra Banderali MD^j Antonello Iacobucci MD^k Stefania Locatelli MD^d Giovanna Casoli MD^l Michael B. Stone MD^c Milena M. Maule PhD^a Ileana Baldi PhD^{am} Franco Merletti MD^a Gian Alfonso Cibinel MD^b for the SIMEU Group for Lung Ultrasound in the Emergency Department in Piedmont

^aCancer Epidemiology Unit, Department of Medical Sciences, CeRMS and University of Turin, Turin, Italy

^bDepartment of Emergency Medicine, "E. Agnelli" General Hospital, Pinerolo

^cDepartment of Emergency Medicine, Brigham and Women's Hospital, Boston, MA

^dDepartment of Emergency Medicine A.O.U. Città della Salute e della Scienza di Torino and University of Turin, Turin, Italy

^eInterdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, ON, Canada

^fToronto Western Hospital, University Health Network

^gDepartment of Medical Sciences, University of Turin

^hDepartment of Emergency Medicine, "San Luigi Gonzaga" University Hospital, Orbassano, Turin, Italy

ⁱDepartment of Emergency Medicine, "Ordine Mauriziano" Hospital, Turin, Italy

^jDepartment of Emergency Medicine, "Cardinal Massaia" Hospital, Asti, Italy

^kDepartment of Emergency Medicine, "Santa Croce e Carle" Hospital, Cuneo, Italy

^lDepartment of Emergency Medicine, "Martini" Hospital, Turin, Italy

^mDepartment of Cardiac, Thoracic, and Vascular Sciences, University of Padova

To the Editor:

We appreciate the comments of Drs Cipriani and Ghittoni, which give us the opportunity to further clarify the message of our study.¹ We believe our study demonstrates that an integrated approach combining lung ultrasonography (LUS) and the standard clinical assessment significantly improves the accuracy of the diagnostic process for acute decompensated heart failure (ADHF). We do not believe that the study outcome can be summarized as "chest radiography (CXR) showed moderate accuracy in the identification of ADHF." The poor diagnostic performance of CXR for identifying ADHF is not a new finding.^{2,3} Although very specific, detection of pulmonary venous congestion, interstitial edema, or alveolar edema on CXR has unacceptably low sensitivity.³ Current guidelines, indeed, caution physicians from using CXR in the diagnosis of ADHF, instead highlighting the value of CXR in the identification of alternative pulmonary causes of a given patient's dyspnea.⁴

In their letter, Drs Cipriani and Ghittoni suggest that the "low value of the standard workup could, in selected cases, possibly be related to poor professional accuracy of operator in charge of performing the workup, interpreting the results, or both." This statement is unsubstantiated. In our study, the sensitivity and specificity of the standard evaluation were as high as 85.3% and 90%, and LUS implementation was able to further increase them to 97% and 97.4%.¹ Although our study included ED physicians with varying levels of expertise, we believe this is part of real-world practice and should have equally affected both clinical and LUS-implemented approaches. In addition, LUS is highly reproducible and easy to learn as compared with CXR interpretation, even when performed by physicians with minimal training.^{1,5} Finally, we remind Drs Cipriani and Ghittoni that our study was conducted in seven EDs from academic, university-affiliated, and community¹ hospitals, providing a level of external validity not previously available in similar studies.

Drs Cipriani and Ghittoni also question the methodology used to compare the performance of CXR and LUS. In our study, the CXR reports completed by staff radiologists were compared with the original LUS findings obtained by ED physicians. No new a posteriori revision of LUS loops was

performed; we simply categorized the presence of diffuse interstitial syndrome as ADHF on the basis of case report forms completed at the time of the examination.

In conclusion, we believe that our study, rather than encouraging indiscriminate LUS use, suggests a sound and prudent use of LUS integrated in the diagnostic evaluation of patients presenting to the ED with acute dyspnea. LUS implementation should not raise unjustified alarms but rather be evaluated, as with all novel medical technologies, using a scientific and unbiased approach.

References

1. E Pivetta, A Goffi, E Lupia, *et al.*, for the SIMEU Group for Lung Ultrasound in the Emergency Department in Piedmont **Lung ultrasound-implemented diagnosis of acute decompensated heart failure in ED: a SIMEU multicenter study.** *Chest*, 148 (1) (2015), pp. 202-210
2. SP Collins, CJ Lindsell, AB Storrow, WT Abraham, ADHERE Scientific Advisory Committee, Investigators and Study Group **Prevalence of negative chest radiography results in the emergency department patient with decompensated heart failure.** *Ann Emerg Med*, 47 (1) (2006), pp. 13-18
3. CS Wang, JM FitzGerald, M Schulzer, E Mak, NT Ayas **Does this dyspneic patient in the emergency department have congestive heart failure?** *JAMA*, 294 (15) (2005), pp. 1944-1956
4. JJ McMurray, S Adamopoulos, SD Anker, *et al.*, ESC Committee for Practice Guidelines **ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC.** *Eur Heart J*, 33 (14) (2012), pp. 1787-1847
5. JL Martindale, VE Noble, A Liteplo **Diagnosing pulmonary edema: lung ultrasound versus chest radiography.** *Eur J Emerg Med*, 20 (5) (2013), pp. 356-360