



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The complexity of the embeddability relation between torsion-free abelian groups of uncountable size

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.het/2318/1684419 since 2018-12-11112:57:452
Published version:
DOI:10.1017/jsl.2018.9
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from convright
protection by the applicable law.

(Article begins on next page)

THE COMPLEXITY OF THE EMBEDDABILITY RELATION BETWEEN TORSION-FREE ABELIAN GROUPS OF UNCOUNTABLE SIZE

FILIPPO CALDERONI

Abstract. We prove that for every uncountable cardinal κ such that $\kappa^{-\kappa} = \kappa$, the quasi-order of embeddability on the κ -space of κ -sized graphs Borel reduces to the embeddability on the κ -space of κ -sized torsion-free abelian groups. Then we use the same techniques to prove that the former Borel reduces to the embeddability relation on the κ -space of κ -sized *R*-modules, for every S-cotorsion-free ring *R* of cardinality less than the continuum. As a consequence we get that all the previous are complete Σ_1^1 quasi-orders.

§1. Introduction. A subset of a topological space is κ -*Borel* if it is in the smallest κ -algebra containing the open sets. Given two spaces X, Y, a function $f: X \to Y$ is κ -*Borel* (measurable) if the preimage through f of every open subset of Y is κ -Borel. Two spaces X, Y are said κ -*Borel isomorphic* if there is a κ -Borel bijection $X \to Y$ whose inverse is κ -Borel too. When $\kappa = \aleph_1$ these notions coincide with the ones of Borel sets, Borel functions, and Borel isomorphism (see the classical reference [15]). More background details and examples will be given in the next section.

Let κ be an infinite cardinal. A topological space *X* is a κ -space if it admits a basis of size $\leq \kappa$. We denote by ${}^{\kappa}\kappa$ the *generalized Baire space*; i.e., the set of functions from κ to itself endowed with the topology generated by the sets of those functions extending a fixed function from a bounded subset of κ to κ . We assume the hypothesis $\kappa^{<\kappa} = \kappa$, which implies that ${}^{\kappa}\kappa$ is a κ -space. A κ -space is *standard Borel* if it is κ^+ -Borel isomorphic to a κ^+ -Borel subset of ${}^{\kappa}\kappa$. If *X* is a standard Borel κ -space, we say that $A \subseteq X$ is (κ)-analytic (or Σ_1^1) if it is a continuous image of a closed subset of ${}^{\kappa}\kappa$. The set of κ -analytic subsets of *X* is usually denoted by $\Sigma_1^1(X)$. We are interested in *quasi-orders* (i.e., reflexive and transitive binary relations) defined over standard Borel κ -spaces. A quasi-order *Q* on *X* is *analytic*, or Σ_1^1 , if and only if $Q \in \Sigma_1^1(X \times X)$. Let *P*, *Q* be Σ_1^1 quasi-orders on the standard Borel κ -spaces *X* and *Y*, respectively. We say that *P Borel reduces* to *Q* if there is a κ^+ -Borel function $f: X \to Y$ such that $\forall x_0x_1 \in X(x_0 P x_1 \Leftrightarrow f(x_0) Q f(x_1)$).

THEOREM 1.1 (Essentially Williams [21]). For every infinite cardinal κ such that $\kappa^{<\kappa} = \kappa$, the embeddability quasi-order on the κ -space of κ -sized graphs Borel reduces to the embeddability on the κ -space of κ -sized groups.

The above result was proved in [21, Theorem 5.1] for $\kappa = \omega$, and the same proof works for κ uncountable as well. In view of a result of Louveau and Rosendal (see [16, Theorem 3.1]), Theorem 1.1 in case $\kappa = \omega$ yields that the embeddability relation on

© 2018, Association for Symbolic Logic 0022-4812/18/8302-0016 DOI:10.1017/jsl.2018.9

Received April 6, 2017.

²⁰¹⁰ Mathematics Subject Classification. 03E15, 20K20, 20K40.

Key words and phrases. Borel reducibility, torsion-free abelian groups, generalized descriptive set theory.

countable groups is a complete Σ_1^1 quasi-order (i.e., a maximum among all Σ_1^1 quasiorders up to Borel reducibility). To prove Theorem 1.1, Williams maps every countable graph to a group generated by the vertices of the graph and satisfying some small cancellation hypothesis. Such groups are not abelian and have many torsion elements that are used to encode the edge relation of the corresponding graphs. So one may wonder whether there exists another Borel reduction from embeddability between countable graphs to embeddability on countable torsion-free abelian groups.

At about the same time as Theorem 1.1 was proved, Przeździecki showed in [18] that the category of graphs almost-fully embeds into the category of abelian groups. I.e., there exists a functor $G: \mathscr{G}raphs \to \mathscr{A}b$ such that for every two graphs T, V there is a natural isomorphism

$$\mathbb{Z}[\operatorname{Hom}(T, V)] \cong \operatorname{Hom}(GT, GV),$$

where $\mathbb{Z}[B]$ is defined as the free abelian group with basis *B*. A closer look into the construction of the functor reveals that it takes values in the subcategory of torsion-free abelian groups. Unfortunately for us, the restriction of *G* to the standard Borel space of countable graphs is not a Borel reduction in the classical sense because countable graphs are sent to groups of size the continuum.

In this article we work in the framework of generalized descriptive set theory. By tweaking the construction of [18] we show the following.

THEOREM 1.2. For every uncountable κ such that $\kappa^{\kappa} = \kappa$, there is a Borel reduction from the embeddability quasi-order on the κ -space of graphs of size κ to embeddability on the κ -space of κ -sized torsion-free abelian groups.

So, in view of the results of [17] and [3], the embeddability relation between κ -sized torsion-free abelian groups is a complete Σ_1^1 quasi-order.

In Section 2 we introduce the main definitions of Borel reducibility in the framework of generalized descriptive set theory. In Section 3 we recall some theorems on the existence of *R*-modules with prescribed endomorphism ring. Such theorems will be used to define the reductions we present in the ensuing sections. Section 4 is dedicated to the proof of Theorem 1.2: we define a Borel reduction from the embeddability relation on κ -sized graphs to embeddability on κ -sized torsion-free abelian groups. In Section 5 we follow the main ideas of [10], and we exploit the techniques used in Section 4 to prove an analogue of Theorem 1.2 for the embeddability relation on *R*-modules, for every S-cotorsion-free ring *R* of cardinality less than the continuum.

THEOREM 1.3. Let *R* be a commutative S-cotorsion-free ring of cardinality less than the continuum. For every uncountable κ such that $\kappa^{<\kappa} = \kappa$, there is a Borel reduction from the embeddability quasi-order on the κ -space of graphs of size κ to embeddability on the κ -space of κ -sized *R*-modules.

In Section 6 we address the problem of determining the Borel complexity of $\cong_{\mathsf{TFA}}^{\kappa}$ the isomorphism on κ -sized torsion-free abelian groups. We point out that a result of [14] implies the following.

THEOREM 1.4. Assume that V = L and κ is inaccessible. Then the isomorphism relation on κ -sized torsion-free abelian groups is a complete Σ_1^1 equivalence relation.

§2. Preliminaries. We consider the *generalized Baire space* $\kappa := \{x \mid x: \kappa \to \kappa\}$ for an uncountable cardinal κ . Unless otherwise specified, κ is endowed with the *bounded topology* τ_b , i.e., the topology generated by the basic open sets

$$N_s = \{ x \in {}^{\kappa} \kappa \mid x \supseteq s \},\$$

where $s \in {}^{<\kappa}\kappa$. Notice that since κ is uncountable, the topology τ_b is strictly finer than the product topology. We assume the hypothesis

$$\kappa^{<\kappa} = \kappa, \tag{1}$$

consequently we get that ${}^{\kappa}\kappa$ is a κ -space, as the basis $\{N_s \mid s \in {}^{<\kappa}\kappa\}$ has size κ . The *generalized Cantor space* ${}^{\kappa}2 := \{x \in {}^{\kappa}\kappa \mid x : \kappa \to 2\}$ is a closed subset of ${}^{\kappa}\kappa$ and therefore it is standard Borel with the relative topology. We recall the following proposition which gives some characterizations of κ -analytic sets.

PROPOSITION 2.1. Let X be a standard Borel κ -space and $A \subseteq X$ nonempty. Then, the following are equivalent:

- (i) A is κ -analytic;
- (ii) *A* is a continuous image of some κ^+ -Borel $B \subseteq {}^{\kappa}\kappa$;
- (iii) *A* is a κ^+ -Borel image of some κ^+ -Borel $B \subseteq {}^{\kappa}\kappa$;
- (iv) A is the projection $p(F) = \{x \in X \mid \exists y \in {}^{\kappa}\kappa((x, y) \in F)\}$ of some closed subset $F \subseteq X \times {}^{\kappa}\kappa$.

A proof of Proposition 2.1 is given in [17, Section 3]. It is specially worth to note that in view of 2.1 we are allowed to use a generalization of the celebrated Tarski–Kuratowski algorithm (see [15, Appendix C]). That is, a set $A \subseteq {}^{\kappa}\kappa$ is κ -analytic if it is defined by an expression involving only κ -analytic sets, atomic connectives, $\exists \alpha, \forall \alpha$ (where α varies over a set of cardinality $\leq \kappa$), and existential quantification over a standard Borel κ -space.

In the remainder of this article we fix some uncountable κ and study standard Borel κ -spaces with the assumption $\kappa^{<\kappa} = \kappa$. For ease of exposition we simply say Borel and analytic instead of saying, respectively, κ^+ -Borel and κ -analytic, whenever κ is clear from the context.

2.1. Spaces of κ -sized structures. In this subsection we recall briefly how to define the standard Borel κ -spaces of uncountable structures of size κ . While in [1, 7, 17] the authors are concerned only with countable languages, we extend the basic definitions to uncountable ones. Our approach is motivated by the aim to develop a unique framework to treat algebraic objects with the most diverse features. For example, following our approach it is possible to define the standard Borel κ -space of κ -sized *R*-modules, for any fixed ring *R* with $|R| < \kappa$. Such spaces will be taken into account in Section 5.

DEFINITION 2.2. If A is a set of size κ , then any bijection $f: \kappa \to A$ induces a bijection from ${}^{\kappa}\kappa$ to ${}^{A}\kappa$, so that the *bounded topology* can be copied on ${}^{A}\kappa$. A basis for such topologies is given by

$$\{N_s^A \mid \exists \alpha < \kappa \ (f''\alpha = \operatorname{dom} s)\},\$$

where $N_s^A = \{x \in {}^A \kappa \mid s \subseteq x\}.$

We briefly recall some useful applications of Definition 2.2.

(a) If G is a group of cardinality κ then, we define the κ -space of subgroups of G by identifying each subgroup of G with its characteristic function and setting

$$\operatorname{Sub}_G = \{ H \in {}^G2 \mid 1_G \in H \land \forall x, y \in G \ (x, y \in H \to xy^{-1} \in H) \},\$$

which is a closed subset of $^{G}2$ and therefore is standard Borel.

(b) Fix a language consisting of finitary relation symbols L = {R_i | i ∈ I}, |I| < κ, and let n_i be the arity of R_i. We denote by X^κ_L the κ-space of L-structures with domain κ. Every A ∈ X^κ_L is a pair (κ, {R^A_i | i ∈ I}) where each R^A_i is an n_i-ary relation on κ, so it can be identified with an element of Π_{i∈I} (^{n_iκ)}2 in the obvious way. It follows that X^κ_L can be endowed with the product of the bounded topologies on its factors (^{n_iκ)}2.

For an infinite cardinal κ , we consider the infinitary logic $L_{\kappa^+\kappa}$. In such logic formulas are defined inductively with the usual formation rules for terms, atomic formulas, negations, disjunctions and conjunctions of size $\leq \kappa$, and quantifications over less than κ many variables.

DEFINITION 2.3. Given an infinite cardinal κ and an $L_{\kappa^+\kappa}$ -sentence φ , we define *the* κ -space of κ -sized models of φ by

$$X_{\omega}^{\kappa} \coloneqq \{\mathcal{A} \in X_{L}^{\kappa} \mid \mathcal{A} \models \varphi\}.$$

The following theorem is a generalization of a classical result by López-Escobar for spaces of uncountable structures.

THEOREM 2.4 ($\kappa^{<\kappa} = \kappa$). A set $B \subseteq X_L^{\kappa}$ is Borel and closed under isomorphism if and only if there is an $L_{\kappa^+\kappa}$ -sentence φ such that $B = X_{\omega}^{\kappa}$.

To see a proof of Theorem 2.4 we refer the reader to [7, Theorem 24] or [1, Theorem 8.7]. A straightforward consequence of it is that the space defined in Definition 2.3 is standard Borel.

We conclude this subsection with a list of those spaces of models of $L_{\kappa^+\kappa}$ -sentences that we will use in the ensuing sections. We denote by $X_{\mathsf{GFAPHS}}^{\kappa}$ the *space of* κ -*sized graphs*. By graph we mean an undirected graph whose edge relation is irreflexive. We denote by $X_{\mathsf{TFA}}^{\kappa}$ the *space of* κ -*sized torsion-free abelian groups*. For any fixed ring Rsuch that $|R| < \kappa$, we denote by $X_{R-\mathsf{MOD}}^{\kappa}$, the *space of* κ -*sized* R-*modules*. Here observe that every $r \in R$ is regarded as a unary functional symbol, interpreted as the left scalar multiplication by r. The axioms of R-modules are the following:

- φ_{AB} , i.e., the first order formula defining abelian groups,
- $\forall x \forall y (r(x + y) = rx + ry),$
- $\forall x ((r+q)x = rx + qx),$
- $\forall x (r(qx) = (rq)x),$
- 1x = x.

Since *r* and *q* vary in *R*, which has size $< \kappa$, the formula defining the class of *R*-modules is a formula in the logic $L_{\kappa^+\kappa}$.

2.2. Borel reducibility. Let *X*, *Y* be standard Borel κ -spaces and *L* a fixed language such that $|L| < \kappa$. Given $\mathcal{A}, \mathcal{B} \in X_L^{\kappa}$, we say that \mathcal{A} is embeddable into \mathcal{B} , in symbols $\mathcal{A} \sqsubseteq_L^{\kappa} \mathcal{B}$, if there is $x \in {}^{\kappa}\kappa$ which realizes an isomorphism between \mathcal{A} and $\mathcal{B} \upharpoonright \operatorname{Im} x$. As pointed out in [1, Section 7.2.2] one can show directly that \sqsubseteq_L^{κ} is the projection on $X_L^{\kappa} \times X_L^{\kappa}$ of a closed subset of $X_L^{\kappa} \times X_L^{\kappa} \times \kappa$, therefore the quasi-order \sqsubseteq_L^{κ} of embeddability between κ -sized *L*-structures is Σ_1^1 . We denote by \sqsubseteq_Z^{κ} the quasi-order of embeddability on the κ -space X_Z^{κ} , where $Z \in \{\text{GRAPHS}, \text{TFA}, R\text{-MOD}\}$.

Let P, Q be Σ_1^1 quasi-orders on the standard Borel κ -spaces X and Y, respectively. Recall that P Borel reduces to Q, in symbols $P \leq_B Q$, if and only if there is a κ^+ -Borel function $f: X \to Y$ such that $\forall x_0 x_1 \in X (x_0 P x_1 \Leftrightarrow f(x_0) Q f(x_1))$. Moreover, Q is

Downloaded from https://www.cambridge.org/core. Universitá di Torino, on 28 Nov 2018 at 09:53:56, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/jsl.2018.9

706

a *complete* Σ_1^1 *quasi-order* if for every Σ_1^1 quasi-order P on a standard Borel κ -space, $P \leq_B Q$. Similarly, we say that E is a *complete* Σ_1^1 *equivalence relation* if $F \leq_B E$, for every Σ_1^1 equivalence relation F. Any quasi-order Q on X induces canonically an equivalence relation on X, which is denoted by E_Q , and defined by setting $x E_Q y$ if and only if x Q y and y Q x for all $x, y \in X$. It can be easily verified that if Q is a complete Σ_1^1 quasi-order then E_Q is a complete Σ_1^1 equivalence relation. The following theorem is a generalization of [16, Theorem 3.1] to the embeddability relation on uncountable graphs.

THEOREM 2.5 (Mildenberger–Motto Ros [3]). If κ is uncountable such that $\kappa^{<\kappa} = \kappa$, then the relation of embeddability $\sqsubseteq_{\mathsf{GRAPHS}}^{\kappa}$ on the κ -space of κ -sized graphs is a complete Σ_1^1 quasi-order.

A first version of Theorem 2.5 was obtained by Motto Ros in [17, Corollary 9.5] provided that κ is weakly compact.

§3. Existence of algebras with prescribed endomorphism ring. Some crucial results that will be used in the next sections are about the existence of algebras with prescribed endomorphism ring.

3.1. *S***-completions.** We recall the basic definitions and some simple facts on *S*-completions following the treatise of [11, Chapter 1].

NOTATION 3.1. In the remainder of this section R is a commutative ring with unit 1 and S is a subset of $R \setminus \{0\}$ containing 1 and closed under multiplication.

DEFINITION 3.2. We say that *R* is *S*-reduced if $\bigcap_{s \in S} sR = 0$, and *R* is *S*-torsion-free if for all $s \in S$ and $r \in R$, sr = 0 implies r = 0. Furthermore, we say that *R* is an *S*-ring provided that *R* is both *S*-reduced and *S*-torsion-free.

In most of the applications *S* is assumed to be countable. In such case, we denote *S* by S as in [11, Chapter 1]. Examples of S-rings include every noetherian domain *R* with $S = \{a^n \mid n \in \omega\}$, for any $a \in R$ such that aR is a proper principal ideal of *R* (see [11, Corollary 1.3]).

DEFINITION 3.3. Let *R* be an *S*-ring and *M* be an *R*-module. We say that *M* is *S*-reduced if $\bigcap_{s \in S} sM = 0$, and *M* is *S*-torsion-free if for every $s \in S$ and $m \in M$, sm = 0 implies m = 0.

We denote by \widehat{M} the *S*-completion of *M*, which is defined as follows. Given $s, q \in S$, we write $q \leq s$ if there is $t \in S$ such that s = qt. Then we set

$$M := \lim_{s \in S} M/sM$$

the inverse limit of inverse system of *R*-modules $(\{M/sM\}_{s\in S}, \{\pi_a^s\}_{q\leq s})$, where

$$\pi_a^s: M/sM \to M/qM, \qquad m+sM \mapsto m+qM.$$

Any *R*-module can be given the natural *linear S-topology*, i.e., the one generated by $\{sM \mid s \in S\}$ as a basis of neighborhoods of 0. A *Cauchy net* in *M* is a sequence $(m_s \mid s \in S)$ taking values in *M*, and such that $m_q - m_{qs} \in qM$, for all $q, s \in S$. We say that the Cauchy net $(m_s \mid s \in S)$ has *limit* $m \in M$ if and only if $m - m_s \in sM$, for every $s \in S$. Finally, we say that *M* is *S-complete* if it is complete with respect to the S-topology, that is, every Cauchy net in M has a unique limit. If M is S-reduced and S-torsion-free, then \widehat{M} is S-reduced, S-torsion-free and S complete (see [11, Lemma 1.6]).

The canonical map

 $\eta_M \colon M \to \widehat{M}, \qquad m \mapsto (m + sM \mid s \in S),$

which is always a homomorphism of *R*-modules, is injective if and only if *M* is *S*-reduced; and it is a ring homomorphism, whenever R = M. Moreover, if *M* is *S*-complete, then η_M is an isomorphism.

For any *R*-module *M*, its *S*-completion \widehat{M} carries a natural a \widehat{R} -module structure. I.e., given $\overline{r} = (r_s + sR \mid s \in S) \in \widehat{R}$ and $\overline{m} = (m_s + sM \mid s \in S) \in \widehat{M}$, we define the scalar multiplication by

$$\bar{r}\bar{m} := (r_s m_s + sM \mid s \in S).$$

3.2. Existence of abelian groups with prescribed endomorphism ring. If *G* is a \mathbb{Z} -module (i.e., an abelian group) and $\mathbb{S} = \mathbb{N} \setminus \{0\}$, then the \mathbb{S} -topology on *G* is usually called \mathbb{Z} -adic topology and the \mathbb{S} -completion of *G* is called the \mathbb{Z} -adic completion. We refer to [8, Theorem 39.5] and [9, Section 2.7] as comprehensive sources on \mathbb{Z} -adic completions.

The next theorem was pointed out by Przeździecki in [18]. It states a slightly different result of a classical theorem by Corner [5, Theorem A], whose proof can be adapted to show the following.

THEOREM 3.4 (Przeździecki [18, Theorem 2.3]). Let A be a ring of cardinality at most 2^{\aleph_0} such that its additive group is free. Then, there is a torsion-free abelian group $M \subseteq \widehat{A}$ such that

- (i) $A \subseteq M$ as (left) A-modules,
- (ii) End $M \cong A$,

(iii) |A| = |M|.

A few comments on Theorem 3.4 may be of some help. We stress the fact that M is torsion-free. By construction M inherits the natural (left) A-module structure from \widehat{A} , which is the one defined by setting

$$a * m = \eta_A(a)m,\tag{2}$$

for every $a \in A$ and $m \in M$. Moreover, condition 3.4 of Theorem 3.4 is proved by showing that for every $h \in \text{End } M$, there exists $a \in A$ such that h(m) = a * m for all $m \in M$.

REMARK 3.5. The reader familiar with the bibliography may find our notation nonstandard. In module theory people usually consider endomorphisms of R-modules as acting on the opposite side from the scalars (e.g., see [11, Chapter 1]). Nevertheless, we prefer to follow the notation of [18] and to have endomorphisms of M acting on the left.

3.3. Existence of *R*-modules with prescribed endomorphism ring. An *R*-module *M* is \mathbb{S} -cotorsion-free if it is \mathbb{S} -reduced and Hom $(\widehat{R}, M) = 0$. This definition extends naturally to any *R*-algebra *A* by saying that *A* is \mathbb{S} -cotorsion-free if *A* has this property as an *R*-module. It is shown in [11, Corollary 1.26] that whenever *M* is an *R*-module of size $< 2^{\aleph_0}$, then *M* is \mathbb{S} -cotorsion-free if and only if it is \mathbb{S} -torsion-free and \mathbb{S} -reduced.

708

We recall one more theorem of existence of *R*-algebras with prescribed endomorphism ring, that is a result of the same kind of Theorem 3.4.

THEOREM 3.6 (Göbel–Przeździecki [10, Corollary 4.5]). Let *R* be an S-cotorsion-free ring such that $|R| < 2^{\aleph_0}$. If *A* is an *R*-algebra of cardinality at most the continuum with a free additive structure over *R*, then there exists an *R*-module *M* such that

- (i) $A \subseteq M \subseteq \widehat{A}$ as (left) A-modules,
- (ii) $\operatorname{End}_R M \cong A$,
- (iii) |A| = |M|.

§4. The embeddability relation between torsion-free abelian groups. In this section we focus on the embeddability relation between κ -sized torsion-free abelian groups and we prove Theorem 1.2. To this purpose we adapt the embedding from the category of graphs into the category of abelian groups defined in [18]. For the sake of exposition we avoid the notion of colimit commonly used in category theory, instead we use the classical notion of direct limit for direct systems of abelian groups which gives more insights on the possibility to define the reduction in a κ^+ -Borel way.

Let Γ be a skeleton of the category of countable graphs; i.e., a full subcategory of the category of countable graphs with exactly one object for every isomorphism class. Without loss of generality, assume that every object in Γ is a graph over a subset of ω . Since we work under the assumption (1), there is a κ -sized universal graph, i.e., a graph of cardinality κ , which contains all graphs of cardinality κ as induced subgraphs. We denote by W_{κ} the κ -sized universal graph on κ and by $[W_{\kappa}]^{\kappa}$ the subspace of induced subgraphs of W_{κ} of cardinality κ . We identify $[W_{\kappa}]^{\kappa}$ with the κ^+ -Borel subset of subsets of W_{κ} of cardinality κ . Therefore we can consider $[W_{\kappa}]^{\kappa}$ as the standard Borel κ -space of graphs of cardinality κ .

For every graph *T* and every infinite cardinal λ , we denote by $[T]^{<\lambda}$ the set of induced subgraphs of *T* of cardinality $< \lambda$. Next, for every $S \in [W_{\kappa}]^{<\omega_1}$, we fix an isomorphism $\theta_S : S \to \sigma(S)$, where $\sigma(S)$ denotes the unique graph in Γ which is isomorphic to *S*.

Now define

$$A \coloneqq \mathbb{Z}[\operatorname{Arw}(\Gamma) \cup \{1\} \cup \mathcal{P}_{fin}(\omega)].$$
(3)

That is, the free abelian group generated by the arrows in Γ , a distinguished element 1, and the finite subsets of ω . We endow *A* with a ring structure by multiplying the elements of the basis as follows, and then extending the multiplication to the whole *A* by linearity. For every $a, b \in \operatorname{Arw}(\Gamma) \cup \mathcal{P}_{fin}(\omega)$ let

$$ab = \begin{cases} a \circ b & \text{if} \\ a \circ b & \text{if} \\ a''b & \text{if} \\ 0 & \text{otherwise.} \end{cases} \begin{pmatrix} a, b \in \operatorname{Arw}(\Gamma) \\ a \text{ and } b \text{ are composable,} \\ b \subseteq \operatorname{dom} a \\ a \upharpoonright b \text{ is an isomorphism,} \end{cases}$$
(4)

$$a1 = 1a = a. \tag{5}$$

REMARK 4.1. The definition of A in (3) differs from the one of [18, Section 3] for including $\mathcal{P}_{fin}(\omega)$ in the generating set. These elements will play the crucial role of embeddability detectors in Lemma 4.8.

Now observe that the ring A has cardinality 2^{\aleph_0} and its additive group is free. So let M be a group having endomorphism ring isomorphic to A as in Theorem 3.4. Notice that the elements of A act on M on the left as in (2).

DEFINITION 4.2. For every $C \in \Gamma$, let

 $G_C \coloneqq id_C * M.$

Notice that G_C is a subgroup of M, for all $C \in \Gamma$. Moreover, if $C, D \in \Gamma$ and $\gamma: C \to D$, then $\gamma \in A$ and thus induces a group homomorphism $G\gamma$ from G_C to G_D by left-multiplication

$$G\gamma: G_C \to G_D \qquad id_C * m \mapsto \gamma * (id_C * m).$$
 (6)

We make sure that such map is well defined as $\gamma * (id_C * m) = id_D * (\gamma * (id_C * m))$, which is clearly an element of G_D .

Now fix any $T \in [W_{\kappa}]^{\kappa}$. For every $S, S' \in [T]^{<\omega_1}$ such that $S \subseteq S'$, the inclusion map $i_{S'}^S: S \to S'$ induces a map $\gamma_{S'}^S$ from $\sigma(S)$ to $\sigma(S')$, the one that makes the diagram below commute.



The map $\gamma_{S'}^{S}$ is in Γ and thus it induces functorially a group homomorphism $G\gamma_{S'}^{S}$, as described in (6). For all $S, S' \in [T]^{<\omega_1}$ such that $S \subseteq S'$, let $\tau_{S'}^{S} = G\gamma_{S'}^{S}$. We claim that $(\{G_{\sigma(S)}\}, \{\tau_{S'}^{S}\}_{S \subseteq S'})_{S,S' \in [T]^{<\omega_1}}$ is a direct system of torsion-free abelian groups indices by the poset $[T]^{<\omega_1}$, which is ordered by inclusion.

DEFINITION 4.3. For every $T \in [W_{\kappa}]^{\kappa}$, let¹

$$GT := \lim_{S \in [T]^{<\omega_1}} G_{\sigma(S)}.$$
(7)

For the sake of definiteness, every element of the direct limit in (7) is regarded as the equivalence class [(m, S)] of an element of the disjoint union $\bigsqcup_{S \in [T]^{\leq \omega_1}} G_{\sigma(S)}$ factored out by the equivalence relation \sim_T , which is defined by setting $(m, S) \sim_T (m', S')$ provided that there is $S'' \supseteq S, S'$ such that $\tau_{S''}^S(m) = \tau_{S''}^{S'}(m')$. Such characterization for (7) holds because the poset of indexes is directed (see [19, Corollary 5.31]).

Notice that for every $T \in [W_{\kappa}]^{\kappa}$, the group GT is abelian by definition, and it is torsion-free as torsion-freeness is preserved by taking subgroups and colimits. Moreover, we claim that GT has cardinality κ . It is clear that |GT| is bounded by $|\bigsqcup_{\kappa} M| = \kappa$. Furthermore, we observe that each GT has at least κ distinct elements. To see this, consider id_{C_0} , where C_0 stands for the unique graph with one vertex and no edges in Γ . For sake of definiteness, suppose that C_0 is the graph with no edges whose unique vertex is 0. For every $\alpha < \kappa$, let $\{\alpha\}$ denote the subgraph of T with the only vertex α . It is

¹As the referee kindly pointed out the notation *GT* instead of G(T) or G_T is nonstandard in descriptive set theory. Nevertheless we prefer to stick to the common practice in category theory to denote functors by juxtaposition.

clear that $id_{C_0} \in G_{\sigma(\{\alpha\})}$. Moreover, for any distinct $\alpha, \beta \in \kappa$, we have that $(id_{C_0}, \{\alpha\})$ and $(id_{C_0}, \{\beta\})$ represent two distinct elements of G_T . For, if $S \supseteq \{\alpha\}, \{\beta\}$, then one has

$$\begin{aligned} \tau_{S}^{\{\alpha\}}(id_{C_{0}}) &= \gamma_{S}^{\{\alpha\}}id_{C_{0}} = (0 \mapsto \theta_{S}(\alpha)), \\ \tau_{A}^{\{\beta\}}(id_{C_{0}}) &= \gamma_{S}^{\{\beta\}}id_{C_{0}} = (0 \mapsto \theta_{S}(\beta)), \end{aligned}$$

which are not equal as θ_S is bijective.

The next lemma basically states that G can be defined in a κ^+ -Borel way.

LEMMA 4.4. There is a κ^+ -Borel map

$$[W_{\kappa}]^{\kappa} \to X_{\mathsf{TFA}}^{\kappa} \qquad T \mapsto \mathcal{G}T$$

such that, for every $T \in [W_{\kappa}]^{\kappa}$, the group GT is isomorphic to GT.

PROOF. Let \leq be a well-ordering of $B = \bigsqcup_{S \in [W_k]^{<\omega_1}} G_{\sigma(S)}$. First consider the map

$$f: [W_{\kappa}]^{\kappa} \to 2^{B}, \qquad T \mapsto \bigsqcup_{S \in [T]^{<\omega_{1}}} G_{\sigma(S)}.$$

To see that f is κ^+ -Borel consider the subbasis of 2^B given by the sets $\{x: B \to 2 \mid x((m, S)) = 1\}$ and $\{x: B \to 2 \mid x((m, S)) = 0\}$, for every $(m, S) \in B$. For any fixed $(m_0, S_0) \in B$, one has

$$f^{-1}(\{x: B \to 2 \mid x((m_0, S_0)) = 1\}) = \{T \in [W_{\kappa}]^{\kappa} \mid S \subseteq T\},\$$

which is κ^+ -Borel.

Then let $g: \text{Im } f \to 2^B$ be the map defined by mapping f(T) to the subset of f(T) which is obtained by deleting all of the (m, S) that are \sim_T -equivalent (i.e., equivalent in the relation used to define the direct limit indexed by $[T]^{<\omega_1}$) to some point appearing before in the well-ordering \leq . One has

$$g(f(T))((m,S)) = 1 \iff S \subseteq T \land \forall (m',S') \prec (m,S)((m',S') \not\sim_T (m,S)),$$

where $(m', S') \neq_T (m, S)$ is a shorthand for

$$\nexists S'' \supseteq S, S''(\tau_{S''}^S(m) = \tau_{S''}^{S'}(m')).$$

Then, for every *T*, we define a group *GT* with underlying set κ and operation \star_T by setting $\alpha \star_T \beta = \gamma$ if and only if the product of the α -th element and the β -th element in g(f(T)) according to \leq is \sim_T -equivalent to the γ -th element in g(f(T)). Notice that there is a unique element in g(f(T)) which is \sim_T -equivalent to such product, thus the map $T \mapsto \mathcal{G}T$ is well defined and is κ^+ -Borel.

Next lemma is derived essentially as in [18, Lemma 3.6].

LEMMA 4.5. If $T, V \in X_{\mathsf{GRAPHS}}^{\kappa}$ and $T \sqsubseteq_{\mathsf{GRAPHS}}^{\kappa} V$, then $GT \sqsubseteq_{\mathsf{TFA}}^{\kappa} GV$.

PROOF. We first claim that if $C, D \in \Gamma$ and $\gamma: C \to D$ is an embedding then $G\gamma: G_C \to G_D$ is one-to-one. Notice that by 3.4 of Theorem 3.4 and the definition of G_C one obtains

$$\mathbb{Z}[\Gamma_C \cup \mathcal{P}_{fin}(C)] \subseteq G_C \subseteq \widetilde{\mathbb{Z}[\Gamma_C \cup \mathcal{P}_{fin}(C)]}.$$

Acting by left-multiplication, γ induces the injective map

$$\langle \gamma \rangle \colon \mathbb{Z}[\Gamma_C \cup \mathcal{P}_{fin}(C)] \to \mathbb{Z}[\Gamma_D \cup \mathcal{P}_{fin}(D)], \quad a \mapsto \gamma a,$$

which in turn induces the injective map on the Z-adic completions

$$\widehat{\langle \gamma \rangle} \colon \widehat{\mathbb{Z}[\Gamma_C \cup \mathcal{P}_{fin}(C)]} \to \widehat{\mathbb{Z}[\Gamma_D \cup \mathcal{P}_{fin}(D)]}, \qquad \bar{a} \mapsto \gamma * \bar{a}.$$
(8)

Comparing (8) with (6) it follows that $G\gamma$ is indeed the restriction of $\langle \overline{\gamma} \rangle$ on G_C , which implies that $G\gamma$ is injective because so is $\langle \overline{\gamma} \rangle$.

Now let $\varphi: T \to V$ be a graph embedding. Then there exists a group homomorphism

$$G\varphi: GT \to GV, \qquad [(g,S)] \mapsto [(G\gamma^{S}_{\varphi''S}(g), \varphi''S)],$$

where $\varphi''S$ is the point-wise image of *S* through φ and $\gamma_{\varphi''S}^S : \sigma(S) \to \sigma(\varphi''S)$ is the map induced by $\varphi \upharpoonright S$, which is clearly a graph embedding. We are left to prove that $G\varphi$ is one-to-one. So fix any $[(g,S)], [(g',S')] \in GT$ such that $[(g,S)] \neq [(g',S')]$. By directedness of $[T]^{<\omega_1}$ we can assume that S = S' without any loss of generality. One has

$$G\varphi([(g, S)]) = [(G\gamma^{S}_{\varphi''S}(g), \varphi''S)],$$

$$G\varphi([(g', S)]) = [(G\gamma^{S}_{\varphi''S}(g'), \varphi''S)],$$

н

which are different elements of GV because $G\gamma^{S}_{\omega''S}$ is injective.

Now we are left to prove that $GT \sqsubseteq_{\mathsf{TFA}}^{\kappa} GV$ implies that $T \sqsubseteq_{\mathsf{GRAPHS}}^{\kappa} V$. Given any linear combination $\sum k_i \varphi_i$, $k_i \in \mathbb{Z}$ and $\varphi_i \in \operatorname{Hom}(T, V)$, one can define a group homomorphism $\Psi(\sum k_i \varphi_i) \colon GT \to GV$ as follows. For any φ_i and $S \in [T]^{<\omega_1}$, let δ_i^S be the function such that the diagram commutes



Since δ_i^S is an arrow in Γ , it induces a group homomorphism

$$G\delta_i^S : G_{\sigma(S)} \to G_{\sigma(\varphi_i'S)} \qquad m \mapsto \delta_i^S * m.$$

as observed in (6). Then we define

$$\Psi(\sum k_i\varphi_i)\colon GT\to GV \qquad [(m,S)]\mapsto \sum k_i[(G\delta_i^S(m),\varphi_i''S)].$$

THEOREM 4.6 (Przeździecki [18, Theorem 3.14]). There is a natural isomorphism

 $\Psi \colon \mathbb{Z}[\operatorname{Hom}(T, V)] \xrightarrow{\cong} \operatorname{Hom}(GT, GV).$

Remark 4.7. Theorem 4.6 states that G is an almost-full embedding, according to the terminology of [10, 18]. It can be proved arguing as in [18, Section 3].

Now we come to the point where our modification from (4) becomes crucial. Since *A* contains the finite subsets of ω , we use them and the property of almost-fullness of *G* to detect an embedding among $\varphi_0, \ldots, \varphi_n$ when $\Psi(\sum_{i \le n} k_i \varphi_i)$ is one-to-one.

LEMMA 4.8. For every two graphs T and V in $X_{\text{GRAPHS}}^{\kappa}$, if $GT \sqsubseteq_{\text{GROUPS}}^{\kappa} GV$ holds then $T \sqsubseteq_{\text{GRAPHS}}^{\kappa} V$.

712

PROOF. Let T, V be as in the hypothesis and $h: GT \to GV$ a group embedding. By Theorem 4.6 we have

$$h = \Psi(\sum_{i \in I} k_i \varphi_i),$$

for some linear combination of graph homomorphisms $\varphi_i \in \text{Hom}(T, V)$. We claim that there must be some $i \in I$ such that φ_i is a graph embedding from T into V. Suppose that it is not true, aiming for a contradiction. Since $[T]^{<\omega_1}$ is directed, there is some finite $S \in [T]^{<\omega}$ such that, for every $i \in I$, the restriction map $\varphi_i \upharpoonright S$ is not one-to-one or does not preserve nonedges. Call d the vertex set of $\sigma(S)$. Such d is a finite subset of ω and is an element of $G_{\sigma(S)}$ because $d = id_{\sigma(S)} * d$. Now consider [(d, S)], the element of GTrepresented by $d \in G_{\sigma(S)}$. Then [(d, S)] is a nontrivial element and

$$\begin{split} h([(d,S)]) &= \sum k_i [(G\delta_i^S(d),\varphi_i''S)] = \\ &= \sum k_i [(\delta_i^S * d,\varphi_i''S)] = 0 \end{split}$$

because if $\varphi_i \upharpoonright S$ is not an embedding then neither is the induced map δ_i^S . This contradicts the fact that *h* is one-to-one.

Summing up the results of this section we can prove the main theorem.

PROOF OF THEOREM 1.2. In view of the Lemma 4.4, we can assume that *G* is κ^+ -Borel. By Lemma 4.5, *G* is a homomorphism from $\sqsubseteq_{\mathsf{GRAPHS}}^{\kappa}$ to $\sqsubseteq_{\mathsf{TFA}}^{\kappa}$, and Lemma 4.8 yields that *G* is a reduction.

COROLLARY 4.9. For every uncountable κ such that $\kappa^{\kappa} = \kappa$, the embeddability relation $\sqsubseteq_{\mathsf{TFA}}^{\kappa}$ on the κ -space of κ -sized torsion-free abelian groups is a complete Σ_1^1 quasi-order.

PROOF. Combining Theorem 1.2 with Theorem 2.5, it follows that $\sqsubseteq_{\mathsf{TFA}}^{\kappa}$ is a complete Σ_{1}^{1} quasi-order provided that $\kappa^{<\kappa} = \kappa$ holds.

It is worth mentioning that the analogue of Corollary 4.9, where $\kappa = \omega$, has been proved recently in [4].

§5. The embeddability relation between *R*-modules. In this section we use the second Corner's type theorem stated in section 3 to prove that the quasi-order of embeddability between *R*-modules, for any S-ring *R* of cardinality less than the continuum, is a complete Σ_1^1 quasi-order.

PROOF OF THEOREM 1.3. Let Γ , W_{κ} , and σ be as in Section 4. Define

$$A := R[\operatorname{Arw}(\Gamma) \cup \{1\} \cup \mathcal{P}_{fin}(\omega)].$$
(9)

That is, the free *R*-module generated by the arrows in Γ , a distinguished element 1 and the finite subsets of ω . The *R*-module *A* can be endowed a ring structure by defining a multiplication on the element of its basis as in (4). Such multiplication is compatible with the *R*-module structure, therefore we can regard *A* as an *R*-algebra. Notice that *A* has cardinality the continuum so we apply Theorem 3.6 which yields the existence of an *R*-module $M \cong \text{End}_R A$. We continue defining *G* similarly to how we did in Section 4. That is, for every $C \in \Gamma$,

$$G_C \coloneqq id_C * M$$
,

and for every $T \in [W_{\kappa}]^{\kappa}$, let

$$GT := \lim_{S \in [T]^{<\omega_1}} G_{\sigma(S)}.$$

Since |M| = |A|, for every $T \in [W_{\kappa}]^{\kappa}$, GT has size κ . Then one can argue as in Lemmas 4.4, 4.5, and 4.8 to prove that *G* is a Borel reduction from $\sqsubseteq_{\mathsf{GRAPHS}}^{\kappa}$ to $\sqsubseteq_{R-\mathsf{MOD}}^{\kappa}$. In this case the almost-fullness for *G* was proved essentially in [10, Theorem 3.16] with the same argument used in [18].

COROLLARY 5.1. For every uncountable cardinal κ such that $\kappa^{<\kappa} = \kappa$ and every ring R as in the statement of Theorem 1.3, the embeddability relation $\sqsubseteq_{R-\text{MOD}}^{\kappa}$ on the κ -space of κ -sized R-module is a complete Σ_1^1 quasi order.

§6. The isomorphism problem. At the end of his article Przeździecki posed the question if every two isomorphic groups in the target of the functor have isomorphic inverse images (see [18, Section 8]). We still do not know whether after our modification the answer is positive, i.e., whether the map G defined in Definition 4.3 is a reduction for the isomorphisms. Then we ask a more general question in terms of Borel reducibility.

QUESTION 6.1. Is there any Borel reduction from isomorphism $\cong_{\mathsf{GRAPHS}}^{\kappa}$ on κ -sized graphs to isomorphism $\cong_{\mathsf{TFA}}^{\kappa}$ on κ -sized torsion-free abelian groups?

Question 6.1 is still open even in the case $\kappa = \omega$, where a positive answer will yield that the relation \cong_{TFA} of isomorphism on countable torsion-free abelian groups is a maximum up to Borel reducibility among all the equivalence relations induced by a Borel action of S_{∞} on a standard Borel space. A remarkable result in such direction is one by Hjorth, who proved in [12] that \cong_{TFA} is not Borel. In fact, this was extended by Downey–Montalbán [6], who showed that \cong_{TFA} is complete Σ_1^1 as a set of pairs. We ought to mention that there are several results in classical Borel reducibility concerning the isomorphism relation on torsion-free abelian groups with finite rank. For every $n < \omega$, denote by \cong_{TFA_n} the isomorphism on countable torsion-free abelian group of rank n. An old result by Baer [2] establishes that \cong_{TFA_1} is essentially E_0 (i.e., it is Borel bireducible with E_0). Moreover Thomas proved in [20] that for every $n \ge 1$, $(\cong_{\mathsf{TFA}_n}) <_B$ $(\cong_{\mathsf{TFA}_{n+1}})$.

Now we go back to Question 6.1. Some results of [7] and [14] use certain model theoretic properties of complete theories to obtain information about the isomorphism relation between the models of those. First let us mention that in [13] the authors give the following example, among many others, of a complete Σ_1^1 equivalence relation in the constructible universe.

DEFINITION 6.2. Let E_{ω}^{κ} the equivalence relation defined on ${}^{\kappa}\kappa$ by

 $x E_{\omega}^{\kappa} y \iff \{\alpha < \kappa \mid x(\alpha) = y(\alpha)\}$ contains an ω -club.

In [13, Theorem 7], under the assumption V = L, the equivalence relation E_{ω}^{κ} is shown to be complete Σ_1^1 for every inaccessible cardinal κ . Then, in [14, Definition 5.4] the authors defined the *orthogonal chain property* (OCP) for stable theories and proved the following result.

THEOREM 6.3 (Hyttinen–Moreno [14, Corollary 5.10]). Assume that κ is inaccessible. For every stable theory T with OCP, the equivalence relation E_{ω}^{κ} reduces continuously to \cong_{T}^{κ} .

As it was kindly pointed out by Hyttinen to the author of this article, the theory of \mathbb{Z}_p (i.e., the group of *p*-adic integers) has OCP and is stable. Thus one obtains

Theorem 1.4 as a corollary of Theorem 6.3. This gives an affirmative partial answer to Question 6.1.

§7. Acknowledgments. The author is very grateful to Adam J. Przeździecki for explaining some crucial parts of [18] and for some invaluable discussions which eventually lead to the proof of the main results of this article. The author would like to thank Tapani Hyttinen for some useful comments that contributed to the last section. Moreover, the author thanks Andrew Brooke-Taylor, Raphaël Carroy, Simon Thomas, and Matteo Viale for comments and interesting discussions. The results presented in this article will be part of the PhD thesis of the author and carried out under the supervision of Luca Motto Ros. This work was supported by the "National Group for the Algebraic and Geometric Structures and their Applications" (GNSAGA–INDAM).

REFERENCES

[1] A. ANDRETTA and L. M. Ros, *Classifying uncountable structures up to bi-embeddability*, preprint, 2016, arXiv:1609.09292v1.

[2] R. BAER, Abelian groups without elements of finite order. Duke Mathematical Journal, vol. 3 (1937), no. 1, pp. 68–122.

[3] F. CALDERONI, H. MILDENBERGER, and L. M. Ros, Uncountable structures are not classifiable up to bi-embeddability, submitted.

[4] F. CALDERONI and S. THOMAS, *The bi-embeddability relation for countable abelian groups*. *Transactions of the American Mathematical Society*, to appear.

[5] A. L. S. CORNER, Every countable reduced torsion-free ring is an endomorphism ring. Proceedings of the London Mathematical Society. Third Series, vol. 13 (1963), pp. 687–710.

[6] R. DOWNEY and A. MONTALBÁN, *The isomorphism problem for torsion-free abelian groups is analytic complete*. Journal of Algebra, vol. 320 (2008), no. 6, pp. 2291–2300.

[7] S.-D. FRIEDMAN, T. HYTTINEN, and V. KULIKOV, *Generalized descriptive set theory and classification theory*. *Memoirs of the American Mathematical Society*, vol. 230 (2014), no. 1081, pp. vi+80.

[8] L. FUCHS, *Infinite Abelian Groups. Vol. I*, Pure and Applied Mathematics, vol. 36, Academic Press, New York, London, 1970.

[9] — , Abelian Groups, Springer Monographs in Mathematics, Springer, Cham, 2015.

[10] R. GÖBEL and A. J. PRZEŹDZIECKI, An axiomatic construction of an almost full embedding of the category of graphs into the category of *R*-objects. Journal of Pure and Applied Algebra, vol. 218 (2014), no. 2, pp. 208–217.

[11] R. Göbel and J. TRLIFAJ, *Approximations and Endomorphism Algebras of Modules. Volume 2*, extended ed., de Gruyter Expositions in Mathematics, vol. 41, Walter de Gruyter GmbH & Co. KG, Berlin, 2012, Predictions.

[12] G. HJORTH, *The isomorphism relation on countable torsion free abelian groups*. *Fundamenta Mathematicae*, vol. 175 (2002), no. 3, pp. 241–257.

[13] T. HYTTINEN and V. KULIKOV, On Σ_1^1 -complete equivalence relations on the generalized Baire space. *Mathematical Logic Quarterly*, vol. 61 (2015), no. 1–2, pp. 66–81.

[14] T. HYTTINEN and M. MORENO, On the reducibility of isomorphism relations. Mathematical Logic Quarterly, vol. 63 (2017), no. 3–4, pp. 175–192.

[15] A. S. KECHRIS, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.

[16] A. LOUVEAU and C. ROSENDAL, Complete analytic equivalence relations. Transactions of the American Mathematical Society, vol. 357 (2005), no. 12, pp. 4839–4866.

[17] L. M. Ros, *The descriptive set-theoretical complexity of the embeddability relation on models of large size*. *Annals of Pure and Applied Logic*, vol. 164 (2013), no. 12, pp. 1454–1492.

[18] A. J. PRZEŹDZIECKI, An almost full embedding of the category of graphs into the category of abelian groups. Advances in Mathematics, vol. 257 (2014), pp. 527–545.

[19] J. J. ROTMAN, An Introduction to Homological Algebra, second ed., Universitext, Springer, New York, 2009.

Downloaded from https://www.cambridge.org/core. Universitá di Torino, on 28 Nov 2018 at 09:53:56, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/jsl.2018.9

FILIPPO CALDERONI

[20] S. THOMAS, The classification problem for torsion-free abelian groups of finite rank. Journal of the American Mathematical Society, vol. 16 (2003), no. 1, pp. 233–258.

[21] J. WILLIAMS, Universal countable Borel quasi-orders, this Journal, vol. 79 (2014), no. 3, pp. 928–954.

DIPARTIMENTO DI MATEMATICA «GIUSEPPE PEANO» UNIVERSITÀ DI TORINO VIA CARLO ALBERTO 10, 10123 TORINO, ITALY *E-mail:* filippo.calderoni@unito.it