

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

On the automorphism group of a closed G2-structure

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1676269> since 2019-04-03T09:35:52Z

Published version:

DOI:10.1093/qmath/hay045

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

ON THE AUTOMORPHISM GROUP OF A CLOSED G_2 -STRUCTURE

FABIO PODESTÀ AND ALBERTO RAFFERO

ABSTRACT. We study the automorphism group of a compact 7-manifold M endowed with a closed non-parallel G_2 -structure, showing that its identity component is abelian with dimension bounded by $\min\{6, b_2(M)\}$. This implies the non-existence of compact homogeneous manifolds endowed with an invariant closed non-parallel G_2 -structure. We also discuss some relevant examples.

1. INTRODUCTION

A seven-dimensional smooth manifold M admits a G_2 -structure if the structure group of its frame bundle can be reduced to the exceptional Lie group $G_2 \subset \mathrm{SO}(7)$. Such a reduction is characterized by the existence of a global 3-form $\varphi \in \Omega^3(M)$ satisfying a suitable non-degeneracy condition and giving rise to a Riemannian metric g_φ and to a volume form dV_φ on M via the identity

$$g_\varphi(X, Y) dV_\varphi = \frac{1}{6} \iota_X \varphi \wedge \iota_Y \varphi \wedge \varphi,$$

for all $X, Y \in \mathfrak{X}(M)$ (see e.g. [2, 13]).

By [10], the intrinsic torsion of a G_2 -structure φ can be identified with the covariant derivative $\nabla^{g_\varphi} \varphi$, and it vanishes identically if and only if both $d\varphi = 0$ and $d *_\varphi \varphi = 0$, $*_\varphi$ being the Hodge operator defined by g_φ and dV_φ . On a compact manifold, this last fact is equivalent to $\Delta_\varphi \varphi = 0$, where $\Delta_\varphi = d^*d + dd^*$ is the Hodge Laplacian of g_φ . A G_2 -structure φ satisfying any of these conditions is said to be *parallel* and its associated Riemannian metric g_φ has holonomy contained in G_2 . Consequently, g_φ is Ricci-flat and the automorphism group $\mathrm{Aut}(M, \varphi) := \{f \in \mathrm{Diff}(M) \mid f^* \varphi = \varphi\}$ of (M, φ) is finite when M is compact and $\mathrm{Hol}(g_\varphi) = G_2$.

Parallel G_2 -structures play a central role in the construction of compact manifolds with holonomy G_2 , and various known methods to achieve this result involve *closed* G_2 -structures, i.e., those whose defining 3-form φ satisfies $d\varphi = 0$ (see e.g. [2, 3, 15, 19]).

A G_2 -structure whose defining 3-form φ satisfies the equation $d *_\varphi \varphi = 0$ is called *co-closed*. On every compact 7-manifold admitting G_2 -structures there exists a co-closed one (cf. [6]), while general results on the existence of closed G_2 -structures are not known.

Due to the recent developments on the G_2 -Laplacian flow and related open problems [11, 14, 16, 17, 19, 20, 21], it is of foremost interest to provide examples of compact manifolds admitting closed G_2 -structures. Most of the known examples consist of simply connected Lie groups endowed with a left-invariant closed G_2 -form φ [5, 8, 9, 12, 17]. Compact

2010 *Mathematics Subject Classification.* 53C10, 57S15.

Key words and phrases. closed G_2 -structure, automorphism.

The authors were supported by GNSAGA of INdAM.

locally homogeneous examples can be obtained considering the quotient of such groups by a co-compact discrete subgroup, whenever this exists. Further non-homogeneous closed G_2 -structures on the 7-torus can be constructed starting from the symplectic half-flat $SU(3)$ -structure on \mathbb{T}^6 described in [7, Ex. 5.1] (see Example 2.4 for details).

Up to now, the existence of compact homogeneous 7-manifolds admitting an invariant closed non-parallel G_2 -structure was not known (cf. [17, Question 3.1] and [18, 26]). Moreover, among the G_2 -manifolds acted on by a cohomogeneity one simple group of automorphisms studied in [4] no compact examples admitting a closed G_2 -structure occur.

In this short note, we investigate the properties of the automorphism group $\text{Aut}(M, \varphi)$ of a compact 7-manifold M endowed with a closed non-parallel G_2 -structure φ . Our main results are contained in Theorem 2.1, where we show that the identity component $\text{Aut}^0(M, \varphi)$ is necessarily abelian with dimension bounded by $\min\{6, b_2(M)\}$. In particular, this answers negatively [17, Question 3.1] and explains why compact examples cannot occur in [4]. Moreover, we also prove some interesting properties of the automorphism group action, and we describe some relevant examples.

These results shed some light on the structure of compact 7-manifolds admitting closed G_2 -structures and can be of some help in the construction of new examples.

2. THE AUTOMORPHISM GROUP

Let M be a seven-dimensional manifold endowed with a closed G_2 -structure φ , and consider its automorphism group

$$\text{Aut}(M, \varphi) := \{f \in \text{Diff}(M) \mid f^*\varphi = \varphi\}.$$

Notice that $\text{Aut}(M, \varphi)$ is a closed Lie subgroup of the isometry group $\text{Iso}(M, g_\varphi)$ of g_φ , and that the Lie algebra of its identity component $\mathfrak{g} := \text{Aut}^0(M, \varphi)$ is

$$\mathfrak{g} = \{X \in \mathfrak{X}(M) \mid \mathcal{L}_X \varphi = 0\}.$$

In particular, every $X \in \mathfrak{g}$ is a Killing vector field for the metric g_φ (cf. [19, Lemma 9.3]).

When M is compact, the Lie group $\text{Aut}(M, \varphi) \subset \text{Iso}(M, g_\varphi)$ is also compact, and we can show the following.

Theorem 2.1. *Let M be a compact seven-dimensional manifold endowed with a closed non-parallel G_2 -structure φ . Then, there exists an injective map*

$$F : \mathfrak{g} \rightarrow \mathcal{H}^2(M), \quad X \mapsto \iota_X \varphi,$$

where $\mathcal{H}^2(M)$ is the space of Δ_φ -harmonic 2-forms. As a consequence, the following properties hold:

- 1) $\dim(\mathfrak{g}) \leq b_2(M)$;
- 2) \mathfrak{g} is abelian with $\dim(\mathfrak{g}) \leq 6$;
- 3) for every $p \in M$, the isotropy subalgebra \mathfrak{g}_p has dimension $\dim(\mathfrak{g}_p) \leq 2$, with equality only when $\dim(\mathfrak{g}) = 2, 3$;
- 4) the G -action is free when $\dim(\mathfrak{g}) \geq 5$. Moreover, when $\dim(\mathfrak{g}) = 6$ the manifold M is diffeomorphic to \mathbb{T}^7 .

Proof. Let $X \in \mathfrak{g}$. Then, $0 = \mathcal{L}_X \varphi = d(\iota_X \varphi)$, as φ is closed. We claim that $\iota_X \varphi$ is co-closed (see also [19, Lemma 9.3]). Indeed, for every $p \in M$ the 2-form $\iota_X \varphi|_p$ belongs to the unique seven-dimensional G_2 -irreducible submodule $\Lambda_7^2(T_p^* M) \subset \Lambda^2(T_p^* M)$, and therefore (see e.g. [1, p. 541]) we have

$$\iota_X \varphi \wedge \varphi = 2 *_{\varphi} (\iota_X \varphi),$$

from which it follows that

$$0 = d(\iota_X \varphi \wedge \varphi) = 2 d *_{\varphi} (\iota_X \varphi).$$

Consequently, the 2-form $\iota_X \varphi$ is Δ_{φ} -harmonic and F is the restriction of the injective map $Z \mapsto \iota_Z \varphi$ to \mathfrak{g} . From this 1) follows.

As for 2), we begin observing that $\mathcal{L}_Y(\iota_X \varphi) = 0$ for all $X, Y \in \mathfrak{g}$, since every Killing field on a compact manifold preserves every harmonic form. Hence, we have

$$0 = \mathcal{L}_Y(\iota_X \varphi) = \iota_{[Y, X]} \varphi + \iota_X(\mathcal{L}_Y \varphi) = \iota_{[Y, X]} \varphi.$$

This proves that \mathfrak{g} is abelian, the map $Z \mapsto \iota_Z \varphi$ being injective. Now, G is compact abelian and it acts effectively on the compact manifold M . Therefore, the principal isotropy is trivial and $\dim(\mathfrak{g}) \leq 7$. When $\dim(\mathfrak{g}) = 7$, M can be identified with the 7-torus \mathbb{T}^7 endowed with a left-invariant metric, which is automatically flat. Hence, if φ is closed non-parallel, then $\dim(\mathfrak{g}) \leq 6$.

In order to prove 3), we fix a point p of M and we observe that the image of the isotropy representation $\rho : G_p \rightarrow O(7)$ is conjugate into G_2 . Since G_2 has rank two and G_p is abelian, the dimension of \mathfrak{g}_p is at most two. If $\dim(\mathfrak{g}_p) = 2$, then the image of ρ is conjugate to a maximal torus of G_2 and its fixed point set in $T_p M$ is one-dimensional. As $T_p(G \cdot p) \subseteq (T_p M)^{G_p}$, the dimension of the orbit $G \cdot p$ is at most one, which implies that $\dim(\mathfrak{g})$ is either two or three.

The first assertion in 4) is equivalent to proving that G_p is trivial for every $p \in M$ whenever $\dim(\mathfrak{g}) \geq 5$. In this case, $\dim(\mathfrak{g}_p) \leq 1$ by 3) and therefore the dimension of the orbit $G \cdot p$ is at least four. Then, for every $h \in G_p$, the element $\rho(h) \in G_2$ has a fixed point set containing $T_p(G \cdot p)$, hence with dimension at least four. On the other hand, a non-trivial element in G_2 is easily seen to have a fixed point set in \mathbb{R}^7 of dimension at most three. Indeed, every $u \in G_2$ is conjugate to an element of a maximal torus of G_2 contained in the maximal rank subgroup $SU(3) \subset G_2$, i.e., it can be supposed to be of the form

$$\text{diag}(z, w, \bar{z} \cdot \bar{w}) \in SU(3),$$

for some $z, w \in \mathbb{C}$ of unit norm. Thus, u fixes at least the real line $V \subset \mathbb{R}^7$ that is fixed by $SU(3)$. Moreover, if u is non-trivial, its fixed point set in the $SU(3)$ -module V^{\perp} has complex dimension at most one. This shows that $G_p = \{1_G\}$. The last assertion follows from [23]. \square

The following corollary answers negatively a question posed by Lauret in [17].

Corollary 2.2. *There are no compact homogeneous 7-manifolds endowed with an invariant closed non-parallel G_2 -structure.*

Proof. The assertion follows immediately from point 2) of Theorem 2.1. \square

In contrast to the last result, it is possible to exhibit non-compact homogeneous examples. Consider for instance a six-dimensional non-compact homogeneous space H/K endowed with an invariant symplectic half-flat $SU(3)$ -structure, namely an $SU(3)$ -structure (ω, ψ) such that $d\omega = 0$ and $d\psi = 0$ (see [25] for the classification of such spaces when H is semisimple and for more information on symplectic half-flat structures). If (ω, ψ) is not torsion-free, i.e., if $d(J\psi) \neq 0$, then the non-compact homogeneous space $(H \times \mathbb{S}^1)/K$ admits an invariant closed non-parallel G_2 -structure defined by the 3-form

$$\varphi := \omega \wedge ds + \psi,$$

where ds denotes the global 1-form on \mathbb{S}^1 .

Remark 2.3. In [4], the authors investigated G_2 -manifolds acted on by a cohomogeneity one simple group of automorphisms. Theorem 2.1 explains why compact examples in the case of closed non-parallel G_2 -structures do not occur.

The next example shows that G can be non-trivial, that the upper bound on its dimension given in 2) can be attained, and that 4) is only a sufficient condition.

Example 2.4. In [7], the authors constructed a symplectic half-flat $SU(3)$ -structure (ω, ψ) on the 6-torus \mathbb{T}^6 as follows. Let (x^1, \dots, x^6) be the standard coordinates on \mathbb{R}^6 , and let $a(x^1)$, $b(x^2)$ and $c(x^3)$ be three smooth functions on \mathbb{R}^6 such that

$$\lambda_1 := b(x^2) - c(x^3), \quad \lambda_2 := c(x^3) - a(x^1), \quad \lambda_3 := a(x^1) - b(x^2),$$

are \mathbb{Z}^6 -periodic and non-constant. Then, the following pair of \mathbb{Z}^6 -invariant differential forms on \mathbb{R}^6 induces an $SU(3)$ -structure on $\mathbb{T}^6 = \mathbb{R}^6/\mathbb{Z}^6$:

$$\begin{aligned} \omega &= dx^{14} + dx^{25} + dx^{36}, \\ \psi &= -e^{\lambda_3} dx^{126} + e^{\lambda_2} dx^{135} - e^{\lambda_1} dx^{234} + dx^{456}, \end{aligned}$$

where $dx^{ijk\dots}$ is a shorthand for the wedge product $dx^i \wedge dx^j \wedge dx^k \wedge \dots$. It is immediate to check that both ω and ψ are closed and that $d(J\psi) \neq 0$ whenever at least one of the functions $a(x^1)$, $b(x^2)$, $c(x^3)$ is not identically zero. Thus, the pair (ω, ψ) defines a symplectic half-flat $SU(3)$ -structure on the 6-torus. The automorphism group of $(\mathbb{T}^6, \omega, \psi)$ is \mathbb{T}^3 when $a(x^1)b(x^2)c(x^3) \neq 0$, while it becomes \mathbb{T}^4 (\mathbb{T}^5) when one (two) of them vanishes identically.

Now, we can consider the closed G_2 -structure on $\mathbb{T}^7 = \mathbb{T}^6 \times \mathbb{S}^1$ defined by the 3-form $\varphi = \omega \wedge ds + \psi$. Depending on the vanishing of none, one or two of the functions $a(x^1)$, $b(x^2)$, $c(x^3)$, φ is a closed non-parallel G_2 -structure and the automorphism group of (\mathbb{T}^7, φ) is \mathbb{T}^4 , \mathbb{T}^5 or \mathbb{T}^6 , respectively.

Finally, we observe that there exist examples where the upper bound on the dimension of \mathfrak{g} given in 1) is more restrictive than the upper bound given in 2).

Example 2.5. In [5], the authors obtained the classification of seven-dimensional nilpotent Lie algebras admitting closed G_2 -structures. An inspection of all possible cases shows that the Lie algebras whose second Betti number is lower than seven are those appearing in Table 1.

Let \mathfrak{n} be one of the Lie algebras in Table 1, and consider a closed non-parallel G_2 -structure φ on it. Then, left multiplication extends φ to a left-invariant G_2 -structure of the same type on the simply connected nilpotent Lie group N corresponding to \mathfrak{n} . Moreover, as the

nilpotent Lie algebra \mathfrak{n}	$b_2(\mathfrak{n})$
$(0, 0, e^{12}, e^{13}, e^{23}, e^{15} + e^{24}, e^{16} + e^{34})$	3
$(0, 0, e^{12}, e^{13}, e^{23}, e^{15} + e^{24}, e^{16} + e^{34} + e^{25})$	3
$(0, 0, e^{12}, 0, e^{13} + e^{24}, e^{14}, e^{46} + e^{34} + e^{15} + e^{23})$	5
$(0, 0, e^{12}, 0, e^{13}, e^{24} + e^{23}, e^{25} + e^{34} + e^{15} + e^{16} - 3e^{26})$	6

TABLE 1.

structure constants of \mathfrak{n} are integers, there exists a co-compact discrete subgroup $\Gamma \subset N$ giving rise to a compact nilmanifold $\Gamma \backslash N$ [22]. The left-invariant 3-form φ on N passes to the quotient defining an invariant closed non-parallel G_2 -structure on $\Gamma \backslash N$. By Nomizu Theorem [24], the de Rham cohomology group $H_{\text{dR}}^k(\Gamma \backslash N)$ is isomorphic to the cohomology group $H^k(\mathfrak{n}^*)$ of the Chevalley-Eilenberg complex of \mathfrak{n} . Hence, $b_2(\Gamma \backslash N) = b_2(\mathfrak{n})$.

Acknowledgements. The authors would like to thank Anna Fino for her comments.

REFERENCES

- [1] R. L. Bryant. Metrics with exceptional holonomy. *Ann. of Math.*, **126**(3), 525–576, 1987.
- [2] R. L. Bryant. Some remarks on G_2 -structures. In *Proceedings of Gökova Geometry-Topology Conference 2005*, pages 75–109. Gökova Geometry/Topology Conference (GGT), Gökova, 2006.
- [3] R. L. Bryant and F. Xu. Laplacian flow for closed G_2 -structures: Short time behavior. arXiv:1101.2004.
- [4] R. Cleyton and A. Swann. Cohomogeneity-one G_2 -structures. *J. Geom. Phys.*, **44**(2-3), 202–220, 2002.
- [5] D. Conti and M. Fernández. Nilmanifolds with a calibrated G_2 -structure. *Differential Geom. Appl.*, **29**(4), 493–506, 2011.
- [6] D. Crowley and J. Nordström. New invariants of G_2 -structures. *Geom. Topol.*, **19**(5), 2949–2992, 2015.
- [7] P. de Bartolomeis and A. Tomassini. On the Maslov index of Lagrangian submanifolds of generalized Calabi-Yau manifolds. *Internat. J. Math.*, **17**(8), 921–947, 2006.
- [8] M. Fernández. An example of a compact calibrated manifold associated with the exceptional Lie group G_2 . *J. Differential Geom.*, **26**(2), 367–370, 1987.
- [9] M. Fernández. A family of compact solvable G_2 -calibrated manifolds. *Tohoku Math. J. (2)*, **39**(2), 287–289, 1987.
- [10] M. Fernández and A. Gray. Riemannian manifolds with structure group G_2 . *Ann. Mat. Pura Appl. (4)*, **132**, 19–45, 1982.
- [11] J. Fine and C. Yao. Hypersymplectic 4-manifolds, the G_2 -Laplacian flow and extension assuming bounded scalar curvature. arXiv:1704.07620.
- [12] A. Fino and A. Raffero. Closed G_2 -structures on non-solvable Lie groups. arXiv:1712.09664.
- [13] N. Hitchin. Stable forms and special metrics. In *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, v. 288 of *Contemp. Math.*, pp. 70–89. Amer. Math. Soc., 2001.
- [14] H. Huang, Y. Wang and C. Yao. Cohomogeneity-one G_2 -Laplacian flow on the 7-torus. *J. London Math. Soc.*, 2018. doi:10.1112/jlms.12137
- [15] D. D. Joyce. Compact Riemannian 7-manifolds with holonomy G_2 . I, II. *J. Differential Geom.*, **43**(2), 291–328, 329–375, 1996.
- [16] J. Lauret. Laplacian flow of homogeneous G_2 -structures and its solitons. *Proc. Lond. Math. Soc. (3)*, **114**(3), 527–560, 2017.

- [17] J. Lauret. Laplacian solitons: questions and homogeneous examples. *Differential Geom. Appl.*, **54**(B), 345–360, 2017.
- [18] H. V. Lê and M. Munir. Classification of compact homogeneous spaces with invariant G_2 -structures. *Adv. Geom.*, **12**(2), 302–328, 2012.
- [19] J. D. Lotay and Y. Wei. Laplacian flow for closed G_2 structures: Shi-type estimates, uniqueness and compactness. *Geom. Funct. Anal.*, **27**(1), 165–233, 2017.
- [20] J. D. Lotay and Y. Wei. Stability of torsion-free G_2 structures along the Laplacian flow. arXiv:1504.07771. To appear in *J. Differential Geom.*
- [21] J. D. Lotay and Y. Wei. Laplacian flow for closed G_2 structures: real analyticity. arXiv:1601.04258. To appear in *Comm. Anal. Geom.*
- [22] A. I. Malčev. On a class of homogeneous spaces. *Amer. Math. Soc. Translation*, **1951**(39), 33 pp., 1951.
- [23] P. S. Mostert. On a compact Lie group action on a manifold. *Ann. of Math.*, **65**(2), 447–455, 1957; Errata, ibid. **66**(2), 589, 1957.
- [24] K. Nomizu. On the cohomology of compact homogeneous spaces of nilpotent Lie groups. *Ann. of Math. (2)*, **59**, 531–538, 1954.
- [25] F. Podestà, A. Raffero. Homogeneous symplectic half-flat 6-manifolds. To appear in *Ann. Global Anal. Geom.* doi: 10.1007/s10455-018-9615-3.
- [26] F. Reidegeld. Spaces admitting homogeneous G_2 -structures. *Differential Geom. Appl.*, **28**(3), 301–312, 2010.

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DEGLI STUDI DI FIRENZE,
 VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY
E-mail address: `podesta@math.unifi.it`, `alberto.raffero@unifi.it`