

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

On the automorphism group of a symplectic half-flat 6-manifold

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1684879> since 2019-01-24T14:41:32Z

Published version:

DOI:10.1515/forum-2018-0137

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

ON THE AUTOMORPHISM GROUP OF A SYMPLECTIC HALF-FLAT 6-MANIFOLD

FABIO PODESTÀ AND ALBERTO RAFFERO

ABSTRACT. We prove that the automorphism group of a compact 6-manifold M endowed with a symplectic half-flat $SU(3)$ -structure has abelian Lie algebra with dimension bounded by $\min\{5, b_1(M)\}$. Moreover, we study the properties of the automorphism group action and we discuss relevant examples. In particular, we provide new complete examples on $T\mathbb{S}^3$ which are invariant under a cohomogeneity one action of $SO(4)$.

1. INTRODUCTION

An $SU(3)$ -structure on a six-dimensional smooth manifold M is the data of an almost Hermitian structure (g, J) with fundamental 2-form $\omega := g(J\cdot, \cdot)$ and a complex volume form $\Psi = \psi + i\widehat{\psi} \in \Omega^{3,0}(M)$ such that

$$(1.1) \quad \psi \wedge \widehat{\psi} = \frac{2}{3} \omega^3.$$

By [11], the whole data (g, J, Ψ) is completely determined by the real 2-form ω and the real 3-form ψ , provided that they satisfy suitable conditions (see §3 for more details).

An $SU(3)$ -structure (ω, ψ) is said to be *symplectic half-flat* if both ω and ψ are closed. In this case, the intrinsic torsion can be identified with a unique real $(1, 1)$ -form σ which is primitive with respect to ω , i.e., $\sigma \wedge \omega^2 = 0$, and fulfills $d\widehat{\psi} = \sigma \wedge \omega$ (see e.g. [4]). This $SU(3)$ -structure is *half-flat* according to [4, Def. 4.1], namely $d(\omega^2) = 0$ and $d\psi = 0$, and the corresponding almost complex structure J is integrable if and only if σ vanishes identically. When this happens, (M, ω, ψ) is a *Calabi-Yau 3-fold*. Otherwise, the symplectic half-flat structure is said to be *strict*.

In recent years, symplectic half-flat structures turned out to be of interest in supersymmetric string theory. For instance, in [10] the authors proved that supersymmetric flux vacua with constant intermediate $SU(2)$ -structure [2] are related to the existence of special classes of half-flat structures on the internal 6-manifold. In particular, they showed that solutions of Type IIA SUSY equations always admit a symplectic half-flat structure. In [12], the definition of symplectic half-flat structures, which are called supersymmetric of Type IIA, is generalized in higher dimensions, and it is proved that semi-flat supersymmetric structures of Type IIA correspond to semi-flat supersymmetric structures of Type IIB via the SYZ and Fourier-Mukai transformations.

2010 *Mathematics Subject Classification.* 53C10, 57S15.

Key words and phrases. $SU(3)$ -structure, automorphism group, cohomogeneity one action.

The authors were supported by GNSAGA of INdAM.

In mathematical literature, symplectic half-flat structures were first introduced and studied in [6] and then in [8], while explicit examples were exhibited in [5, 7, 9, 16, 20]. Most of them consist of simply connected solvable Lie groups endowed with a left-invariant symplectic half-flat structure. Moreover, in [9] it was proved that every six-dimensional compact solvmanifold with an invariant symplectic half-flat structure also admits a solution of Type IIA SUSY equations.

Let M be a 6-manifold endowed with a strict symplectic half-flat structure (ω, ψ) . In the present paper, we are interested in studying the properties of the automorphism group $\text{Aut}(M, \omega, \psi) := \{f \in \text{Diff}(M) \mid f^*\omega = \omega, f^*\psi = \psi\}$, aiming at understanding how to construct non-trivial examples with high degree of symmetry.

In [16], we proved the non-existence of compact homogeneous examples and we classified all non-compact cases which are homogeneous under the action of a semisimple Lie group of automorphisms. Here, in Theorem 2.1 we show that the Lie algebra of $\text{Aut}(M, \omega, \psi)$ is abelian with dimension bounded by $\min\{5, b_1(M)\}$ whenever M is compact. This allows to obtain a direct proof of the aforementioned non-existence result. In the same theorem, we also provide useful information on geometric properties of the $\text{Aut}^0(M, \omega, \psi)$ -action on the manifold, proving in particular that the automorphism group acts by cohomogeneity one only when M is diffeomorphic to a torus. Some relevant examples are then discussed in order to show that the automorphism group can be non-trivial and that the upper bound on its dimension can be actually attained.

As our previous result on non-compact homogeneous spaces suggests, the non-compact ambient might provide a natural setting where looking for new examples. In section 3, we obtain new complete examples of symplectic half-flat structures on the tangent bundle $T\mathbb{S}^3$ which are invariant under the natural cohomogeneity one action of $\text{SO}(4)$. These include also the well-known Calabi-Yau example constructed by Stenzel [19].

2. THE AUTOMORPHISM GROUP

Let M be a six-dimensional manifold endowed with an $\text{SU}(3)$ -structure (ω, ψ) . The *automorphism group* of (M, ω, ψ) consists of the diffeomorphisms of M preserving the $\text{SU}(3)$ -structure, namely

$$\text{Aut}(M, \omega, \psi) := \{f \in \text{Diff}(M) \mid f^*\omega = \omega, f^*\psi = \psi\}.$$

Clearly, $\text{Aut}(M, \omega, \psi)$ is a closed Lie subgroup of the isometry group $\text{Iso}(M, g)$, as every automorphism preserves the Riemannian metric g induced by the pair (ω, ψ) . The Lie algebra of the identity component $\mathfrak{G} := \text{Aut}^0(M, \omega, \psi)$ is

$$\mathfrak{g} = \{X \in \mathfrak{X}(M) \mid \mathcal{L}_X\omega = 0, \mathcal{L}_X\psi = 0\},$$

and every $X \in \mathfrak{g}$ is a Killing vector field for the metric g . Moreover, the Lie group $\text{Aut}(M, \omega, \psi) \subset \text{Iso}(M, g)$ is compact whenever M is compact.

If (M, ω, ψ) is a Calabi-Yau 3-fold, i.e., if ω , ψ and $\widehat{\psi}$ are all closed, then the Riemannian metric g is Ricci-flat and $\text{Hol}(g) \subseteq \text{SU}(3)$. When M is compact and the holonomy group is precisely $\text{SU}(3)$, it follows from Bochner's Theorem that $\text{Aut}(M, \omega, \psi)$ is finite.

We now focus on *strict symplectic half-flat* structures, namely $SU(3)$ -structures (ω, ψ) such that

$$d\omega = 0, \quad d\psi = 0, \quad d\widehat{\psi} = \sigma \wedge \omega,$$

with $\sigma \in [\Omega_0^{1,1}(M)] := \{\kappa \in \Omega^2(M) \mid J\kappa = \kappa, \kappa \wedge \omega^2 = 0\}$ not identically vanishing. Notice that the condition on σ is equivalent to requiring that the almost complex structure J induced by (ω, ψ) is non-integrable (cf. e.g. [4]). In this case, we can show the following result.

Theorem 2.1. *Let M be a compact six-dimensional manifold endowed with a strict symplectic half-flat structure (ω, ψ) . Then, there exists an injective map*

$$\mathcal{F} : \mathfrak{g} \rightarrow \mathcal{H}^1(M), \quad X \mapsto \iota_X \omega,$$

where $\mathcal{H}^1(M)$ is the space of Δ_g -harmonic 1-forms. Consequently, the following properties hold:

- 1) $\dim(\mathfrak{g}) \leq b_1(M)$;
- 2) \mathfrak{g} is abelian with $\dim(\mathfrak{g}) \leq 5$;
- 3) for every $p \in M$, the isotropy subalgebra \mathfrak{g}_p has dimension $\dim(\mathfrak{g}_p) \leq 2$. If $\dim(\mathfrak{g}_p) = 2$ for some p , then $G_p = G$;
- 4) the G -action is free when $\dim(\mathfrak{g}) \geq 4$. In particular, when $\dim(\mathfrak{g}) = 5$ the manifold M is diffeomorphic to \mathbb{T}^6 .

Before proving the theorem, we show a general lemma.

Lemma 2.2. *Let (ω, ψ) be an $SU(3)$ -structure. Then, for every vector field X the following identity holds*

$$\iota_X \psi \wedge \psi = -2 * (\iota_X \omega),$$

where $*$ denotes the Hodge operator determined by the Riemannian metric g and the orientation $dV_g = \frac{1}{6}\omega^3$.

Proof. From the equation $\iota_X \Psi \wedge \Psi = 0$, which holds for every vector field X , we have

$$\iota_X \psi \wedge \psi = \iota_X \widehat{\psi} \wedge \widehat{\psi}, \quad \iota_X \psi \wedge \widehat{\psi} = -\iota_X \widehat{\psi} \wedge \psi.$$

Using the above identities and the relations $\iota_X \psi = \iota_{JX} \widehat{\psi}$, $\iota_{JX} \psi = -\iota_X \widehat{\psi}$, we get

$$\begin{aligned} \iota_X \psi \wedge \psi &= \iota_{JX} \widehat{\psi} \wedge \psi \\ &= \iota_{JX} (\widehat{\psi} \wedge \psi) + \widehat{\psi} \wedge \iota_{JX} \psi \\ &= \iota_{JX} (\widehat{\psi} \wedge \psi) - \widehat{\psi} \wedge \iota_X \widehat{\psi} \\ &= \iota_{JX} (\widehat{\psi} \wedge \psi) - \psi \wedge \iota_X \psi. \end{aligned}$$

Hence, $2\iota_X \psi \wedge \psi = \iota_{JX} (\widehat{\psi} \wedge \psi)$. Now, from condition (1.1) we know that $\widehat{\psi} \wedge \psi = -\frac{2}{3}\omega^3 = -4dV_g$. Thus,

$$\iota_X \psi \wedge \psi = -2\iota_{JX} dV_g = -2 * (JX)^\flat = -2 * (\iota_X \omega).$$

□

Proof of Theorem 2.1.

Let $X \in \mathfrak{g}$. Then, using the closedness of ω we have $0 = \mathcal{L}_X \omega = d(\iota_X \omega)$. Moreover, since $d\psi = 0$ and $\mathcal{L}_X \psi = 0$, then $d(\iota_X \psi \wedge \psi) = 0$ and Lemma 2.2 implies that $d * (\iota_X \omega) = 0$. Hence, the 1-form $\iota_X \omega$ is Δ_g -harmonic and \mathcal{F} coincides with the injective map $Z \mapsto \iota_Z \omega$ restricted to \mathfrak{g} . From this 1) follows.

In order to prove 2), we begin recalling that every Killing field on a compact manifold preserves every harmonic form. Consequently, for all $X, Y \in \mathfrak{g}$ we have

$$0 = \mathcal{L}_Y(\iota_X \omega) = \iota_{[Y, X]} \omega + \iota_X(\mathcal{L}_Y \omega) = \iota_{[Y, X]} \omega.$$

Since the map $Z \mapsto \iota_Z \omega$ is injective, we obtain that \mathfrak{g} is abelian. Now, G is compact abelian and it acts effectively on the compact manifold M . Therefore, the principal isotropy is trivial and $\dim(\mathfrak{g}) \leq 6$. When $\dim(\mathfrak{g}) = 6$, M can be identified with the 6-torus \mathbb{T}^6 endowed with a left-invariant metric, which is automatically flat. Hence, if (ω, ψ) is strict symplectic half-flat, then $\dim(\mathfrak{g}) \leq 5$.

As for 3), we fix a point p of M and we observe that the image of the isotropy representation $\rho : G_p \rightarrow O(6)$ is conjugate into $SU(3)$. Since $SU(3)$ has rank two and G_p is abelian, the dimension of \mathfrak{g}_p is at most two. If $\dim(\mathfrak{g}_p) = 2$, then the image of ρ is conjugate to a maximal torus of $SU(3)$ and its fixed point set in $T_p M$ is trivial. As $T_p(G \cdot p) \subseteq (T_p M)^{G_p}$, the orbit $G \cdot p$ is zero-dimensional, which implies that $\dim(\mathfrak{g}) = 2$.

Assertion 4) is equivalent to proving that G_p is trivial for every $p \in M$ whenever $\dim(\mathfrak{g}) \geq 4$. In this case, $\dim(\mathfrak{g}_p) \leq 1$ by 3), and therefore $\dim(G \cdot p) \geq 3$. If G_p contains a non-trivial element h , then $\rho(h)$ fixes every vector in $T_p(G \cdot p)$ and, consequently, its fixed point set in $T_p M$ must be non-trivial of dimension at least three. On the other hand, a non-trivial element of $SU(3)$ is easily seen to have a fixed point set of dimension at most two. This shows that $G_p = \{1_G\}$. The last assertion follows immediately from [14]. \square

Point 2) in the above theorem gives a direct proof of a result obtained in [16].

Corollary 2.3. *There are no compact homogeneous 6-manifolds endowed with an invariant strict symplectic half-flat structure.*

It is worth observing here that the non-compact case is less restrictive. For instance, it is possible to exhibit non-compact examples which are homogeneous under the action of a *semisimple* Lie group of automorphisms (see e.g. [16]). Moreover, in the next section we shall construct non-compact examples of cohomogeneity one with respect to a semisimple Lie group of automorphisms.

The next example was given in [8]. It shows that G can be non-trivial, that the upper bound on its dimension given in 2) can be attained, and that 4) is only a sufficient condition.

Example 2.4. On \mathbb{R}^6 with standard coordinates (x^1, \dots, x^6) consider three smooth functions $a(x^1)$, $b(x^2)$, $c(x^3)$ in such a way that

$$\lambda_1 := b(x^2) - c(x^3), \quad \lambda_2 := c(x^3) - a(x^1), \quad \lambda_3 := a(x^1) - b(x^2),$$

are \mathbb{Z}^6 -periodic. Then, the following pair of \mathbb{Z}^6 -invariant differential forms on \mathbb{R}^6 induces an $SU(3)$ -structure on $\mathbb{T}^6 = \mathbb{R}^6 / \mathbb{Z}^6$:

$$\omega = dx^{14} + dx^{25} + dx^{36}, \quad \psi = -e^{\lambda_3} dx^{126} + e^{\lambda_2} dx^{135} - e^{\lambda_1} dx^{234} + dx^{456},$$

where $dx^{ijk\dots}$ is a shorthand for the wedge product $dx^i \wedge dx^j \wedge dx^k \wedge \dots$. It is immediate to check that (ω, ψ) is strict symplectic half-flat whenever the functions λ_i are not all constant. The automorphism group of $(\mathbb{T}^6, \omega, \psi)$ is \mathbb{T}^3 when $a(x^1) b(x^2) c(x^3) \neq 0$, while it becomes \mathbb{T}^4 (\mathbb{T}^5) when one (two) of them vanishes identically.

Finally, we observe that there exist examples where the upper bound on the dimension of \mathfrak{g} given in 1) is more restrictive than the upper bound given in 2).

Example 2.5. In [5], the authors obtained the classification of six-dimensional nilpotent Lie algebras admitting symplectic half-flat structures. The only two non-abelian cases are described up to isomorphism by the following structure equations

$$(0, 0, 0, 0, e^{12}, e^{13}), \quad (0, 0, 0, e^{12}, e^{13}, e^{23}).$$

Denote by N the simply connected nilpotent Lie group corresponding to one of the above Lie algebras, and endow it with a left-invariant strict symplectic half-flat structure (ω, ψ) . By [13], there exists a co-compact discrete subgroup $\Gamma \subset N$ giving rise to a compact nilmanifold $\Gamma \backslash N$. Moreover, the left-invariant pair (ω, ψ) on N passes to the quotient defining an $SU(3)$ -structure of the same type on $\Gamma \backslash N$. By [15], we have that $b_1(\Gamma \backslash N)$ is either four or three.

3. NON-COMPACT COHOMOGENEITY ONE EXAMPLES

In this section, we construct complete examples of strict symplectic half-flat structures on a non-compact 6-manifold admitting a cohomogeneity one action of a semisimple Lie group of automorphisms. This points out the difference between the compact and the non-compact case, and together with the results in [16, §4.3] it suggests that the non-compact ambient provides a natural setting to obtain new examples.

From now on, we consider the natural cohomogeneity one action on $M = TS^3 \cong \mathbb{S}^3 \times \mathbb{R}^3$ induced by the transitive $SO(4)$ -action on \mathbb{S}^3 . Then, we have

$$TS^3 \cong SO(4) \times_{SO(3)} \mathbb{R}^3.$$

We refer the reader to [1, 14, 17, 18] for basic notions on cohomogeneity one isometric actions. Following the notation of [18], we consider the Lie algebra $\mathfrak{so}(4) \cong \mathfrak{su}(2) + \mathfrak{su}(2)$ and we fix the following basis of $\mathfrak{su}(2)$

$$H := \frac{1}{2} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad E := \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad V := \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

Let $\gamma : \mathbb{R} \rightarrow M$ be a normal geodesic such that $p := \gamma(0) \in \mathbb{S}^3$ and $\gamma_t := \gamma(t)$ is a regular point for all $t \neq 0$. The singular isotropy subalgebra is $\mathfrak{so}(4)_p = \mathfrak{su}(2)_{\text{diag}}$, while the principal isotropy subalgebra $\mathfrak{k} := \mathfrak{so}(4)_{\gamma_t}$, $t \neq 0$, is one-dimensional and spanned by (H, H) . We consider the following basis of $\mathfrak{so}(4) \cong \mathfrak{su}(2) + \mathfrak{su}(2)$

$$\begin{aligned} E_1 &:= (E, 0), & V_1 &:= (V, 0), & E_2 &:= (0, E), & V_2 &:= (0, V), \\ U &:= (H, H), & A &:= (H, -H). \end{aligned}$$

We let $\xi := \frac{\partial}{\partial t}$, and for any $Z \in \mathfrak{so}(4)$ we denote by \widehat{Z} the corresponding fundamental vector field on M . Then, a basis of $T_{\gamma_t}M$ for $t \neq 0$ is given by

$$(\xi, \widehat{A}, \widehat{E}_1, \widehat{V}_1, \widehat{E}_2, \widehat{V}_2)_{\gamma_t}.$$

We shall denote the dual coframe along γ_t by $(\xi^*, A^*, E_1^*, V_1^*, E_2^*, V_2^*)_{\gamma_t}$, where $\xi^* := dt$.

Let $K \subset \mathrm{SO}(4)$ be the principal isotropy subgroup corresponding to the Lie algebra \mathfrak{k} . The space of K -invariant 2-forms on $T_{\gamma_t}M$, $t \neq 0$, is spanned by

$$\begin{aligned} \omega_1 &:= \xi^* \wedge A^*, & \omega_2 &:= E_1^* \wedge V_1^*, & \omega_3 &:= E_2^* \wedge V_2^*, \\ \omega_4 &:= E_1^* \wedge E_2^* + V_1^* \wedge V_2^*, & \omega_5 &:= E_1^* \wedge V_2^* - V_1^* \wedge E_2^*. \end{aligned}$$

These forms extend as $\mathrm{SO}(4)$ -invariant 2-forms on the regular part $M_0 := \mathbb{S}^3 \times \mathbb{R}^+$. By [18], their differentials along γ_t are

$$(3.1) \quad \begin{aligned} d\omega_1|_{\gamma_t} &= \frac{1}{4} \xi^* \wedge (\omega_2 - \omega_3), & d\omega_2|_{\gamma_t} &= d\omega_3|_{\gamma_t} = 0, \\ d\omega_4|_{\gamma_t} &= -2 A^* \wedge \omega_5, & d\omega_5|_{\gamma_t} &= 2 A^* \wedge \omega_4. \end{aligned}$$

We now describe the general $\mathrm{SO}(4)$ -invariant symplectic 2-form ω on M . Along γ_t , $t \neq 0$, we have

$$\omega|_{\gamma_t} = \sum_{i=1}^5 f_i(t) \omega_i,$$

for suitable smooth functions $f_i \in \mathcal{C}^\infty(\mathbb{R}^+)$. By [18, Prop. 6.1], the $\mathrm{SO}(4)$ -invariant 2-form ω on M_0 corresponding to $\omega|_{\gamma_t}$ admits a smooth extension to the whole M if and only if the functions f_i extend smoothly on \mathbb{R} as follows:

- i) f_1 and f_4 are even and f_2, f_3, f_5 are odd;
- ii) $f_3'(0) = \frac{1}{2} f_1(0) + f_2'(0)$, $f_5'(0) = -\frac{1}{4} f_1(0) - f_2'(0)$, and $f_4(0) = 0$.

Moreover, $\omega|_p$ is non-degenerate if and only if $f_1(0) \neq 0$.

Using (3.1), we compute $d\omega$ and we see that ω is closed if and only if

$$f_4, f_5 \equiv 0, \quad \begin{cases} f_2' = -\frac{1}{4} f_1 \\ f_3' = \frac{1}{4} f_1 \end{cases}.$$

Combining this with the extendability conditions, we obtain that every $\mathrm{SO}(4)$ -invariant symplectic 2-form ω on M can be written as

$$(3.2) \quad \omega|_{\gamma_t} = f_1(t) \omega_1 + f_2(t) \omega_2 + f_3(t) \omega_3, \quad t \neq 0,$$

with $f_1 \in \mathcal{C}^\infty(\mathbb{R})$ even and nowhere vanishing, and

$$f_2(t) = -\frac{1}{4} \int_0^t f_1(s) ds = -f_3(t).$$

Notice that $\omega^3|_{\gamma_t} = -6f_1f_2^2\omega_1 \wedge \omega_2 \wedge \omega_3$ at every regular point of the geodesic γ_t . As f_1 is nowhere zero, we may assume that $f_1 < 0$, so that the volume form $\xi^* \wedge A^* \wedge E_1^* \wedge V_1^* \wedge E_2^* \wedge V_2^*$ defines the same orientation on $T_{\gamma_t}M$ as $\frac{1}{6} \omega^3|_{\gamma_t}$ for all $t \in \mathbb{R}^+$.

We now search for an $\text{SO}(4)$ -invariant closed 3-form $\psi \in \Omega^3(M)^{\text{SO}(4)}$ so that the pair (ω, ψ) defines an $\text{SO}(4)$ -invariant symplectic half-flat structure on M . For the sake of simplicity, we make the following Ansatz

$$\psi = du, \quad u \in \Omega^2(M)^{\text{SO}(4)}.$$

As before, along γ_t , $t \neq 0$, we can write

$$(3.3) \quad u|_{\gamma_t} = \sum_{i=1}^5 \phi_i(t) \omega_i,$$

for some smooth functions $\phi_i \in \mathcal{C}^\infty(\mathbb{R}^+)$ satisfying the same extendability conditions as the f_i 's. Then, omitting the dependence on t for brevity, we have

$$(3.4) \quad \psi|_{\gamma_t} = \psi_2 \xi^* \wedge \omega_2 + \psi_3 \xi^* \wedge \omega_3 + \phi'_4 \xi^* \wedge \omega_4 + \phi'_5 \xi^* \wedge \omega_5 + 2A^* \wedge (\phi_5 \omega_4 - \phi_4 \omega_5),$$

where $\psi_2 := \frac{1}{4} \phi_1 + \phi'_2$ and $\psi_3 := \phi'_3 - \frac{1}{4} \phi_1$.

By [11], the pair (ω, ψ) defines an $\text{SU}(3)$ -structure if and only if the following conditions hold:

- a) the compatibility condition $\omega \wedge \psi = 0$;
- b) the stability condition $P(\psi) < 0$, P being the characteristic quartic polynomial defined on 3-forms (see below for the definition);
- c) denoted by J the almost complex structure induced by (ω, ψ) , then the complex volume form $\Psi := \psi + i \widehat{\psi}$ with $\widehat{\psi} := J\psi$ fulfills the normalization condition (1.1);
- d) the symmetric bilinear form $g := \omega(\cdot, J\cdot)$ is positive definite.

The compatibility condition a) along γ_t reads $f_2 \psi_3 + f_3 \psi_2 = 0$. Since $f_2 = -f_3 \neq 0$, this implies

$$(3.5) \quad \psi_2 = \psi_3.$$

Recall that at each point $q \in M$ the 3-form ψ gives rise to an endomorphism $S \in \text{End}(T_q M)$ defined as follows for every $\theta \in T_q^* M$ and every $v \in T_q M$

$$\iota_v \psi \wedge \psi \wedge \theta = \theta(S(v)) \frac{\omega^3}{6}.$$

The endomorphism S satisfies $S^2 = P(\psi)\text{Id}$, and it gives rise to the almost complex structure $J := \frac{1}{\sqrt{|P(\psi)|}} S$ when $P(\psi) < 0$.

From the expressions

$$\begin{aligned} \iota_\xi \psi \wedge \psi|_{\gamma_t} &= 2(\psi_2^2 - (\phi'_4)^2 - (\phi'_5)^2) \xi^* \wedge \omega_2 \wedge \omega_3 - 4(\phi'_4 \phi_5 - \phi_4 \phi'_5) A^* \wedge \omega_2 \wedge \omega_3, \\ \iota_{\widehat{A}} \psi \wedge \psi|_{\gamma_t} &= 4(\phi_4 \phi'_5 - \phi'_4 \phi_5) \xi^* \wedge \omega_2 \wedge \omega_3 - 8(\phi_4^2 + \phi_5^2) A^* \wedge \omega_2 \wedge \omega_3, \end{aligned}$$

we see that the endomorphism $S \in \text{End}(T_{\gamma_t} M)$ maps the subspace of $T_{\gamma_t} M$ spanned by ξ and $\widehat{A}|_{\gamma_t}$ into itself with associated matrix given by

$$(3.6) \quad -\frac{1}{f_1 f_2^2} \begin{pmatrix} 4(\phi'_4 \phi_5 - \phi_4 \phi'_5) & 8(\phi_4^2 + \phi_5^2) \\ 2(\psi_2^2 - (\phi'_4)^2 - (\phi'_5)^2) & -4(\phi'_4 \phi_5 - \phi_4 \phi'_5) \end{pmatrix}.$$

Since the curve γ_t must be a normal geodesic for the metric g induced by (ω, ψ) , it follows that the tangent vector ξ is orthogonal to the orbit $\text{SO}(4) \cdot \gamma_t$ at every regular point of γ_t . In particular, we have

$$0 = g(\xi, \widehat{A}) = \omega(\xi, J(\widehat{A})) = \frac{1}{\sqrt{|P(\psi)|}} \omega(\xi, S(\widehat{A})) = \frac{4}{f_2^2 \sqrt{|P(\psi)|}} (\phi_4' \phi_5 - \phi_4 \phi_5'),$$

from which we get

$$(3.7) \quad \phi_4' \phi_5 = \phi_4 \phi_5'.$$

Using (3.5), (3.7) and the definition of $P(\psi)$, we obtain

$$(3.8) \quad P(\psi) = \frac{16}{f_1^2 f_2^4} (\phi_4^2 + \phi_5^2) (\psi_2^2 - (\phi_4')^2 - (\phi_5')^2).$$

Consequently, the stability condition b) reads

$$(3.9) \quad \psi_2^2 - (\phi_4')^2 - (\phi_5')^2 < 0, \quad \phi_4^2 + \phi_5^2 \neq 0,$$

for all $t \in \mathbb{R}^+$.

We now note that the vector field $J(\xi)$ is tangent to the $\text{SO}(4)$ -orbits and it belongs to the space of K -fixed vectors in $T_{\gamma_t}(\text{SO}(4) \cdot \gamma_t)^K$, which is spanned by $\widehat{A}|_{\gamma_t}$. Since the geodesic γ_t has unit speed, we see that

$$(3.10) \quad 1 = g(\xi, \xi) = \omega(\xi, J(\xi)) = -\frac{2}{f_2^2 \sqrt{|P(\psi)|}} (\psi_2^2 - (\phi_4')^2 - (\phi_5')^2).$$

Using (3.8), the relation (3.10) implies that

$$(3.11) \quad 4(\phi_4^2 + \phi_5^2) = f_1^2 ((\phi_4')^2 + (\phi_5')^2 - \psi_2^2).$$

Let us now focus on c). From (3.6) and (3.11), we obtain $J(\xi) = \frac{1}{f_1} \widehat{A}|_{\gamma_t}$. Using this and the identity $\widehat{\psi} = J\psi = -\psi(J \cdot, \cdot)$, we have

$$(3.12) \quad \widehat{\psi}|_{\gamma_t} = \xi^* \wedge \left(2 \frac{\phi_4}{f_1} \omega_5 - 2 \frac{\phi_5}{f_1} \omega_4 \right) + f_1 A^* \wedge (\psi_2 (\omega_2 + \omega_3) + \phi_4' \omega_4 + \phi_5' \omega_5).$$

Now, the normalization condition $\psi \wedge \widehat{\psi} = \frac{2}{3} \omega^3$ gives

$$4(\phi_4^2 + \phi_5^2) - f_1^2 (\psi_2^2 - (\phi_4')^2 - (\phi_5')^2) = 2 f_1^2 f_2^2.$$

Combining this with (3.11), we obtain

$$(3.13) \quad \phi_4^2 + \phi_5^2 = \frac{1}{4} (f_1 f_2)^2.$$

Note that (3.8), (3.11) and (3.13) imply

$$P(\psi) \equiv -4$$

along the geodesic γ_t . Thus, the stability of ψ holds also at $t = 0$.

Going back to (3.7), we see that either $\phi_4 = \lambda \phi_5$ or $\phi_5 = \lambda \phi_4$ for some $\lambda \in \mathbb{R} \setminus \{0\}$. Since ϕ_4 and ϕ_5 extend as an even and an odd function on \mathbb{R} , respectively, we see that either $\phi_4 \equiv 0$ or $\phi_5 \equiv 0$. As $f_1 f_2$ is an odd function on \mathbb{R} , (3.13) implies that

$$(3.14) \quad \phi_4 \equiv 0, \quad \phi_5 = \pm \frac{1}{2} f_1 f_2.$$

The matrix associated with the symmetric bilinear form $\omega(\cdot, J\cdot)$ along γ_t , $t \in \mathbb{R}^+$, is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & f_1^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 \frac{\phi_5' \phi_5}{f_1 f_2} & 0 & -2 \frac{\psi_2 \phi_5}{f_1 f_2} & 0 \\ 0 & 0 & 0 & -2 \frac{\phi_5' \phi_5}{f_1 f_2} & 0 & -2 \frac{\psi_2 \phi_5}{f_1 f_2} \\ 0 & 0 & -2 \frac{\psi_2 \phi_5}{f_1 f_2} & 0 & -2 \frac{\phi_5' \phi_5}{f_1 f_2} & 0 \\ 0 & 0 & 0 & -2 \frac{\psi_2 \phi_5}{f_1 f_2} & 0 & -2 \frac{\phi_5' \phi_5}{f_1 f_2} \end{pmatrix},$$

and condition d) can be written as

$$-2 \frac{\phi_5' \phi_5}{f_1 f_2} > 0, \quad \psi_2^2 < (\phi_5')^2.$$

The former condition is equivalent to $(f_2^2)'' > 0$, while the latter is satisfied whenever ψ is stable (cf. (3.9)).

Note that the metric g extends smoothly over the singular orbit \mathbb{S}^3 to a Hermitian symmetric bilinear form. The restriction of g on $T_p \mathbb{S}^3$ is positive definite as $g_p(\widehat{A}, \widehat{A}) = f_1^2(0) > 0$ and the orbit $\text{SO}(4) \cdot p$ is isotropy irreducible. Moreover, $T_p M = T_p \mathbb{S}^3 \oplus J(T_p \mathbb{S}^3)$, and from this we see that g_p is positive definite.

Summing up, the existence of a complete $\text{SO}(4)$ -invariant symplectic half-flat structure (ω, ψ) on M is equivalent to the existence of a smooth function $f_1 \in C^\infty(\mathbb{R})$ satisfying the following conditions:

- 1) f_1 is even and negative;
- 2) the function $f_2(t) := -\frac{1}{4} \int_0^t f_1(s) ds$ satisfies $(f_2^2)'' > 0$;
- 3) there exists an even smooth function $\psi_2 \in C^\infty(\mathbb{R})$ satisfying $\psi_2^2 = [(f_2^2)']^2 - f_2^2$.

Indeed, given f_1 we define the symplectic form ω on M as in (3.2), with $f_3 = -f_2$. As for ψ , we let $\psi_3 := \psi_2$, $\phi_4 := 0$, and $\phi_5 := \pm \frac{1}{2} f_1 f_2$ in (3.4). Then, (3.11) and (3.13) imply $\psi_2^2 = (\phi_5')^2 - f_2^2$, and we can choose the sign in the definition of ϕ_5 so that the extendability condition $\phi_5'(0) = -\psi_2(0)$ given in ii) is satisfied. It is also easy to see that we may choose ϕ_1, ϕ_2, ϕ_3 so that $\psi_2 = \frac{1}{4} \phi_1 + \phi_2'$ and $\psi_3 = \phi_3' - \frac{1}{4} \phi_1$, and the corresponding u as in (3.3) extends to a global 2-form on M . The resulting 3-form ψ is then stable by condition 3) and (3.8). The stability condition together with the inequality in 2) implies that the induced bilinear form g is everywhere positive definite. Hence, we have proved the following result.

Proposition 3.1. *The existence of a complete $\text{SO}(4)$ -invariant symplectic half-flat structure (ω, ψ) on $T\mathbb{S}^3 = \text{SO}(4) \times_{\text{SO}(3)} \mathbb{R}^3$ with $\psi \in d\Omega^2(M)$ is equivalent to the existence of a smooth function $f_1 \in C^\infty(\mathbb{R})$ satisfying conditions 1), 2), 3).*

Recall that the symplectic half-flat structure (ω, ψ) is strict if and only if the unique 2-form $\sigma \in [\Omega_0^{1,1}(M)]$ fulfilling $d\widehat{\psi} = \sigma \wedge \omega$ is not identically zero. Starting from (3.12),

using (3.1) and the identity $dA^*|_{\gamma_t} = \frac{1}{4}(\omega_3 - \omega_2)$ (cf. [18, (3.27)]), we obtain

$$d\widehat{\psi}|_{\gamma_t} = \left((f_1\phi'_5)' - 4\frac{\phi_5}{f_1} \right) \omega_1 \wedge \omega_5 + (f_1\psi_2)' \omega_1 \wedge (\omega_2 + \omega_3),$$

whence

$$\sigma|_{\gamma_t} = \frac{1}{f_1} (f_1\psi_2)' (\omega_2 + \omega_3) + \frac{1}{f_1} \left((f_1\phi'_5)' - 4\frac{\phi_5}{f_1} \right) \omega_5.$$

By [3], we know that the scalar curvature of the metric g induced by a symplectic half-flat structure is given by $\text{Scal}(g) = -\frac{1}{2}|\sigma|^2$. Hence, in our case we have

$$\text{Scal}(g)|_{\gamma_t} = -\frac{1}{f_1^2 f_2^2} \left[((f_1\psi_2)')^2 - \left((f_1\phi'_5)' - 4\frac{\phi_5}{f_1} \right)^2 \right] = -\left(\frac{(f_1\psi_2)'}{f_1\phi'_5} \right)^2,$$

where the second equality follows from the relations obtained so far.

We may construct plenty of complete $\text{SO}(4)$ -invariant strict symplectic half-flat structures on M by choosing a suitable f_1 as above. For instance, the function

$$f_1(t) := -\cosh(t), \quad t \in \mathbb{R},$$

fits in with conditions 1), 2), 3). With this choice, the scalar curvature is

$$\text{Scal}(g)|_{\gamma_t} = -\tanh^2(t) \frac{(6\cosh^2(t) - 5)^2}{4\cosh^4(t) - 8\cosh^2(t) + 5}.$$

This shows that the resulting symplectic half-flat structure is strict and non-homogeneous.

Note that the vanishing of σ is equivalent to the vanishing of $\text{Scal}(g)$. Hence, setting $(f_1\psi_2)' = 0$, the resulting $\text{SU}(3)$ -structure (ω, ψ) is Calabi-Yau and the associated metric is the well-known Stenzel's Ricci-flat metric on $T\mathbb{S}^3$ (cf. [19]).

Finally, we remark that the scalar curvature always vanishes at $t = 0$. Indeed, $(f_1\psi_2)'(0) = 0$, as $f_1\psi_2$ is even, while $f_1(0)\phi'_5(0) \neq 0$. This implies that an $\text{SO}(4)$ -invariant symplectic half-flat structure (ω, ψ) with ψ exact has constant scalar curvature if and only if it is Calabi-Yau.

REFERENCES

- [1] A. V. Alekseevsky, D. V. Alekseevsky. Riemannian G -manifolds with one dimensional orbit space. *Ann. Global Anal. Geom.*, **11**, 197–211, 1993.
- [2] D. Andriot. New supersymmetric flux vacua with intermediate $\text{SU}(2)$ structure. *J. High Energy Phys.*, **8**(096), 2008.
- [3] L. Bedulli and L. Vezzoni. The Ricci tensor of $\text{SU}(3)$ -manifolds. *J. Geom. Phys.*, **57**(4), 1125–1146, 2007.
- [4] S. Chiossi and S. Salamon. The intrinsic torsion of $\text{SU}(3)$ and G_2 structures. In *Differential geometry, Valencia, 2001*, 115–133. World Sci. Publ., River Edge, NJ, 2002.
- [5] D. Conti and A. Tomassini. Special symplectic six-manifolds. *Q. J. Math.*, **58**(3), 297–311, 2007.
- [6] P. de Bartolomeis. Geometric structures on moduli spaces of special Lagrangian submanifolds. *Ann. Mat. Pura Appl. (4)*, **179**, 361–382, 2001.
- [7] P. de Bartolomeis and A. Tomassini. On solvable generalized Calabi-Yau manifolds. *Ann. Inst. Fourier (Grenoble)*, **56**(5), 1281–1296, 2006.
- [8] P. de Bartolomeis and A. Tomassini. On the Maslov index of Lagrangian submanifolds of generalized Calabi-Yau manifolds. *Internat. J. Math.*, **17**(8), 921–947, 2006.

- [9] M. Fernández, V. Manero, A. Otal, and L. Ugarte. Symplectic half-flat solvmanifolds. *Ann. Global Anal. Geom.*, **43**(4), 367–383, 2013.
- [10] A. Fino and L. Ugarte. On the geometry underlying supersymmetric flux vacua with intermediate $SU(2)$ -structure. *Classical Quantum Gravity*, **28**(7), 075004, 21 pp., 2011.
- [11] N. Hitchin. Stable forms and special metrics. In *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, v. 288 of *Contemp. Math.*, pp. 70–89. Amer. Math. Soc., 2001.
- [12] S.-C. Lau, L.-S. Tseng, and S.-T. Yau. Non-Kähler SYZ mirror symmetry. *Comm. Math. Phys.*, **340**(1), 145–170, 2015.
- [13] A. I. Malčev. On a class of homogeneous spaces. *Amer. Math. Soc. Translation*, **1951**(39), 33 pp., 1951.
- [14] P. S. Mostert. On a compact Lie group action on a manifold. *Ann. of Math.*, **65**(2), 447–455, 1957; Errata, *ibid.* **66**(2), 589, 1957.
- [15] K. Nomizu. On the cohomology of compact homogeneous spaces of nilpotent Lie groups. *Ann. of Math. (2)*, **59**, 531–538, 1954.
- [16] F. Podestà, A. Raffero. Homogeneous symplectic half-flat 6-manifolds. *Ann. Global Anal. Geom.*, doi: 10.1007/s10455-018-9615-3.
- [17] F. Podestà and A. Spiro. Six-dimensional nearly Kähler manifolds of cohomogeneity one. *J. Geom. Phys.*, **60**(2), 156–164, 2010.
- [18] F. Podestà and A. Spiro. Six-dimensional nearly Kähler manifolds of cohomogeneity one (II). *Comm. Math. Phys.*, **312**(2), 477–500, 2012.
- [19] M. B. Stenzel. Ricci-flat metrics on the complexification of a compact rank one symmetric space. *Manuscripta Math.*, **80**(2), 151–163, 1993.
- [20] A. Tomassini and L. Vezzoni. On symplectic half-flat manifolds. *Manuscripta Math.*, **125**(4), 515–530, 2008.

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DEGLI STUDI DI FIRENZE,
VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY
E-mail address: `podesta@math.unifi.it`, `alberto.raffero@unifi.it`