

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Staphylococcus aureus undergoes major transcriptional reorganization during growth with Enterococcus faecalis in milk

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1684498> since 2018-12-13T09:07:09Z

Published version:

DOI:10.1016/j.fm.2018.01.007

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 ***Staphylococcus aureus* undergoes major transcriptional reorganization during growth**
2 **with *Enterococcus faecalis* in milk**

3

4 Gabriela Nogueira Viçosa^a, Cristian Botta^a, Ilario Ferrocino^a, Marta Bertolino^a, Marco
5 Ventura^b, Luís Augusto Nero^{c#}, Luca Cocolin^{a#}

6

7 ^aDepartment of Agricultural, Forest and Food Sciences, University of Turin, Grugliasco,
8 Italy; ^bLaboratory of Probiogenomics, Department of Life Sciences, University of Parma,
9 Italy; ^cDepartment of Veterinary, Federal University of Viçosa, Viçosa, Brazil.

10

11 #Address correspondence to: Luca Cocolin, lucasimone.cocolin@unito.it and Luís Augusto
12 Nero, nero@ufv.br

13

14

15

16

17

18

19

20

21

22

23

24

25 **Abstract**

26 Previous studies have demonstrated the antagonistic potential of lactic acid bacteria
27 (LAB) present in raw milk microbiota over *Staphylococcus aureus*, albeit the molecular
28 mechanisms underlying this inhibitory effect are not fully understood. In this study, we
29 compared the behavior of *S. aureus* ATCC 29213 alone and in the presence of a cheese-
30 isolated LAB strain, *Enterococcus faecalis* 41FL1 in skimmed milk at 30 °C for 24 h using
31 phenotypical and molecular approaches. Phenotypic analysis showed the absence of classical
32 staphylococcal enterotoxins in co-culture with a 1.2-log decrease in *S. aureus* final population
33 compared to single culture. Transcriptional activity of several exotoxins and global
34 regulators, including *agr*, was negatively impacted in co-culture, contrasting with the
35 accumulation of transcripts coding for surface proteins. After 24 h, the number of transcripts
36 coding for several metabolite responsive elements, as well as enzymes involved in glycolysis
37 and acetoin metabolism was increased in co-culture. The present study discusses the
38 complexity of the transcriptomic mechanisms possibly leading to *S. aureus* attenuated
39 virulence in the presence of *E. faecalis* and provides insights into this interspecies interaction
40 in a simulated food context.

41

42 **Keywords:** *Staphylococcus aureus*; *Enterococcus faecalis*; RNA sequencing; co-culture;
43 skimmed milk

44

45 1. Introduction

46 *Staphylococcus aureus* is a pathogen of major concern in foodstuffs due to the ability
47 of certain strains to produce thermoresistant staphylococcal enterotoxins (SE) that, once
48 ingested, may lead to gastrointestinal disorders symptoms, such as vomiting, nausea and
49 abdominal cramping (Argudín et al., 2010; Schelin et al., 2011). Despite remarkable
50 improvements in food safety procedures, SE still figure as a leading cause of foodborne
51 outbreaks in Europe, often related to the ingestion of contaminated milk and dairy products
52 (EFSA, 2016).

53 Lactic acid bacteria (LAB) are heavily present in raw milk microbiota and have been
54 extensively used in the food industry for the production of dairy and non-dairy fermented
55 products (Stiles and Holzapfel, 1997). In this context, LAB can exert a protective role against
56 the multiplication of spoilage and pathogenic microorganisms, mainly by modulating redox
57 potential, pH or through the production of inhibitory substances (Carr et al., 2002). Among
58 LAB, *Enterococcus faecalis* is one of the most frequent species recovered from milk and
59 cheese in the Mediterranean area, reaching levels of 10^5 to 10^7 CFU/g at the end of cheese
60 ripening and contributing to the development of organoleptic characteristics (Foulquié-
61 Moreno et al., 2006; Franciosi et al., 2009).

62 The conditions supporting *S. aureus* growth and enterotoxin production in foods are
63 extensive, since *S. aureus* is able to grow and survive in a wide range of temperatures, pH
64 and NaCl concentrations (Valero et al., 2009). The expression of several virulence
65 determinants in *S. aureus* is tightly coordinated to environmental conditions by a complex
66 gene regulatory network, which is mainly controlled by the accessory gene regulator (*agr*)
67 system (Novick et al., 1993). By responding to stress and external factors, *S. aureus* can
68 rapidly adapt its physiology and virulence in order to optimize growth and survival in
69 complex environments, such as foods (Novick et al., 1993).

70 Although LAB antagonistic potential over *S. aureus* in foods has been the subject of
71 research for more than fifty years, most of these early observations relied mainly on

72 phenotypical analysis to study this interaction (Daoud and Debevere, 1985; Haines and
73 Harmon, 1973; Hamama et al., 2002; Noleto et al., 1987; Perin et al., 2012). In recent years,
74 application of molecular techniques has proven to be a reliable strategy to explore the LAB-*S.*
75 *aureus* interaction in culture and food-mimicking media (Charlier et al., 2008; Cretenet et al.,
76 2011; Delpech et al., 2015; Even et al., 2009; Zdenkova et al., 2016). However, only a few
77 studies have been able to successfully determine the underlying mechanism leading to
78 attenuated *S. aureus* virulence in mixed bacterial cultures (Laughton et al., 2006; Li et al.,
79 2011).

80 In the present study, we monitored the behavior of *S. aureus* alone and in the presence
81 of *E. faecalis* during growth in skimmed milk. We used RNA sequencing analysis to explore
82 the adaptations in the *S. aureus* transcriptome that could, at least partly, support our
83 phenotypical findings, i.e., the reduction in *S. aureus* final population and the absence of SEs
84 in co-culture with *E. faecalis*. To the best of our knowledge, this study is the first
85 transcriptomic approach of the interaction between *S. aureus* and *E. faecalis* in skimmed milk
86 and provides interesting knowledge to for designing biocontrol-based strategies to improve
87 food safety.

88

89 **2. Material and methods**

90 *2.1. Bacterial strains and growth conditions*

91 *S. aureus* ATCC 29213 and *E. faecalis* 41FL1 (Dal Bello et al., 2010) were used in
92 this work. To evaluate bacterial interactions, *S. aureus* was inoculated alone and in
93 combination with *E. faecalis* in 100 mL of skimmed milk (Sigma-Aldrich, St. Louis, USA) in
94 250-mL Schott® flasks and statically incubated at 30° C for 24 h. Skimmed milk powder was
95 reconstituted in sterile distilled water (10% w/v) in aseptic conditions followed by treatment
96 at 121 °C for 5 min before bacterial inoculation. Both strains were subcultured at 30° C for
97 18 h and inoculated at a final concentration of 10³ CFU/mL for *S. aureus* and 10⁶ CFU/mL
98 for *E. faecalis*. Experiments were repeated three times to ensure reproducibility.

100 2.2. Microbiological analysis

101 Throughout the incubation period, bacterial growth was assessed at the following
102 intervals: 0 h (inoculation); 4 and 7 h (exponential phase); 12 and 24 h (post-exponential
103 phase). CFU of *S. aureus* population in single and co-cultures was determined on Baird-
104 Parker Rabbit Plasma Fibrinogen agar (bioMérieux, Marcy-l'Étoile, France) (De Buyser et
105 al., 2003), whereas *E. faecalis* population was enumerated on BHI agar (Sigma-Aldrich)
106 followed by a confirmatory step with the catalase test. All plates were incubated at 37° C for
107 48 h. Bacterial growth was determined in duplicate using the spread plate technique for each
108 sampling point.

109

110 2.3. Enterotoxin detection

111 At the same intervals set for *S. aureus* enumeration, samples for SE detection were
112 collected and immediately analyzed. SE production was qualitatively estimated in samples
113 using RIDASCREEN® SET TOTAL (R-Biopharm AG, Darmstadt, HE, GE) according to
114 manufacturer's instructions.

115

116 2.4. Determination of pH, sugars and organic acids content

117 Sugars and organic acids contents in milk were determined by high performance
118 liquid chromatography (HPLC) using the method described by Bertolino et al. (2011) with
119 minor modifications. Briefly, 5 mL of milk samples were added to 20 mL of 0.013 N H₂SO₄
120 (mobile phase) and mixed for 30 min with a horizontal shaker (Asal, Milan, Italy) at 100
121 oscillation/min. The samples were subsequently centrifuged for 5 min at 10000 X g and 10
122 °C, and the supernatant was filtered through a 0.2 µm polypropylene membrane filter (VWR,
123 Milan, Italy). The HPLC system (Thermo Finnigan Spectra System, San Jose, USA) was
124 equipped with an isocratic pump (P4000), a multiple autosampler (AS3000) fitted with a 20
125 µL loop, a UV detector (UV100) set at 210 and a refractive index detector RI-150. The

126 analysis were performed isocratically, at 0.8 ml min⁻¹ and 65 °C, with a 300 × 7.8 mm
127 i.d.cation exchange column (Aminex HPX-87H) equipped with a Cation H⁺ Microguard
128 cartridge (Bio-Rad Laboratories, Hercules, USA). Two replicates were analyzed for each
129 biological replicate. The data treatments were carried out using the Chrom QuestTM
130 chromatography data system (Thermo Finnigan Spectra System, San Jose, USA). Analytical
131 grade reagents were used as standards (Sigma-Aldrich). pH measurements were obtained
132 using a pH meter at the same intervals considered in the HPLC analysis (Crison, Modena,
133 Italy).

134

135 2.5. Whole genome sequencing

136 Genomic DNA (gDNA) was extracted from a *S. aureus* ATCC 29213 culture in BHI
137 broth grown to stationary phase using the Wizard[®] Genomic DNA Purification Kit
138 (Promega, Madison, WI, USA) according to manufacturer's instructions. Whole genome
139 sequencing (WGS) was performed at GenProbio srl (Parma, Italy) using the Illumina MiSeq
140 platform (Illumina, San Diego, CA, USA). From 2.5 µg of gDNA, a library for Illumina
141 paired-ended sequencing was prepared using the TruSeq DNA PCR-Free LT Kit (Illumina),
142 combined with fragmentation using a Bioruptor NGS ultrasonicator (Diagenode, LI,
143 Belgium) and size evaluation using Tape Station 2200 (Agilent Technologies, Palo Alto, CA,
144 USA). The library was sequenced (2 x 250 bp) using a Flow Cell V3 600 cycles (Illumina)
145 according to the manufacturer's instructions. *De novo* genome assembly was performed with
146 MIRA v.4.0.2 (Chevreux et al., 1999). Improvement quality of final contigs was performed
147 with Burrows-Wheeler Aligner (Li and Durbin, 2009), SAMtools suite (Li et al., 2009) and
148 GATK software package v.2.8-1 (McKenna et al., 2010). Reordering of the final contigs was
149 performed with Mauve v.2.3.1 (Darling et al., 2004) against *S. aureus* NCTC 8325 as
150 reference genome (NCBI Accession Number: NC_007795.1). Ribosomal RNA genes were
151 detected on the basis of RNAmmer v.1.2 (Lagesen et al., 2007) and transfer RNA genes were
152 identified using tRNAscan-SE v.1.21 (Lowe and Eddy, 1997). Protein-encoding open reading

153 frames (ORFs) were predicted using Prodigal v.2.6 (Hyatt et al., 2010). Automatic annotation
154 of the ORFs was performed with BLAST (Altschul et al., 1990) against NCBI database and
155 HMMER against the PFAM database (Sonnhammer et al., 1997). Manual corrections to
156 automated functional assignments were conducted to validate the presence or absence of
157 genes of interest. Based on the annotation, coding DNA sequences (CDS) were classified into
158 the Cluster of Orthologous Groups (COG) functional categories (Tatusov et al., 2000) using
159 EggNog v.4.0 database (Jensen et al., 2008).

160

161 *2.6. Transcriptomic analysis by RNA-Seq and gene expression analysis*

162 Bacterial cells from single and co-cultures in duplicates were harvested after 7 and 24
163 h of incubation at 30 °C, and total RNA was isolated using the RiboPure™ Bacteria kit
164 (Ambion, Life Technologies, Waltham MA). RNA was treated with DNase I (Turbo DNA
165 free; Ambion) to ensure complete removal of gDNA. RNA quantification and integrity were
166 determined by agarose gel electrophoresis and Agilent 2200 Tape Station Nucleic Acid
167 System (Agilent Technologies). Following quantification, rRNA was removed from 50 ng of
168 total RNA using Ribo-Zero rRNA removal kit for Gram-positive bacteria (Epicentre,
169 Madison, WI, USA) according to the supplier's instructions. The yield of rRNA depletion
170 was checked by Agilent 2200 Bioanalyzer (Agilent Technologies). rRNA-depleted samples
171 were then fragmented using RNaseIII (Life Technologies, CA, USA) followed by size
172 evaluation using Experion (Bio-Rad, CA, USA). RNA-Seq library was constructed with the
173 IonTorrent Total-RNAseq kit v2 (Life Technologies) according to the manufacturer's
174 protocol. Library templates were amplified on Ion Sphere Particles using Ion One Touch 200
175 Template Kit v2 (Life Technologies). Sequencing of libraries was loaded into IonTorrent 316
176 chips using an Ion PGM 200 sequencing kit (Life Technologies) at GenProbio srl (Parma,
177 Italy). Sequencing reads of each replicate were pooled and aligned to the *S. aureus* ATCC
178 29213 genome sequence through BWA39 with high stringency cut-offs (99% nucleotide
179 identity) to accurately assign co-culture reads to the correct genome. Alignment data were

180 analyzed using HTSeq (<http://www-huber.embl.de/users/anders/HTSeq/doc/overview.html>).
181 Raw mapped reads counts of each gene were normalized to the length of the gene itself and
182 to the number of reads mapped. The resulting data is presented as reads per kilobase of
183 transcript per million reads mapped (RPKM) and allows the comparison of genes in different
184 samples at different depths of sequencing. Based on RPKM values, we determined
185 transcriptional changes between *S. aureus* in single and co-culture by fold change analysis,
186 using the single culture as a reference. Genes were considered differentially expressed when
187 \log_2 fold change (LFC) was $\geq +1$ or ≤ -1 .

188

189 2.7. Real-time quantitative reverse transcription PCR (RT-qPCR)

190 Six *S. aureus* genes (*codY*, *alsD*, *agrC*, *spa*, *hld*, *sea*) up- or downregulated by more
191 than 2-fold at 7 h or 24 h in co-culture were selected to validate the data generated from
192 RNA-Seq experiments by RT-qPCR. *mgo2* was used as reference gene for RT-qPCR data
193 normalization since its expression profile remained invariant in all transcriptomes. The target
194 genes were selected based on their role as virulence factors and in central carbon metabolism.
195 The primer pairs used in RT-qPCR and reverse transcription reactions were designed using
196 the Primer-BLAST tool (Ye et al., 2012) based on the genome sequence of *S. aureus* ATCC
197 29213. Further confirmation of *in silico* specificity of selected primer sequences was
198 performed with UGENE software version 1.26.1 (Okonechnikov et al., 2012) and BLAST
199 (Altschul et al., 1990) against NCBI database. All primer sequences used in this study are
200 listed in Supplementary Table S5. Gene-specific reverse transcription was conducted as
201 follows: 600 ng of RNA were mixed with 1 μ L of reverse primer (100 μ M) and ultrapure
202 water in a reaction volume of 10 μ l. The mix was treated at 75 $^{\circ}$ C for 5 min for RNA
203 denaturation and immediately placed on ice for 10 min. Five microliters of M-MLV RT
204 Buffer (1 X), 5 μ l of dNTPs (10 μ M each), 1 μ l of M-MLV Reverse Transcriptase (8 U/ μ L)
205 and 0.6 μ L of RNasin ribonuclease inhibitor (20 U/ μ L) were added to the mix for a final
206 volume of 25 μ l by addition of ultrapure water. RT reaction was carried out at 42 $^{\circ}$ C for 1 h

207 in a Biorad DNA Engine thermal cycler (Bio-Rad) with subsequent storage of cDNA at -20
208 °C. RT-qPCR reactions were performed on corresponding cDNAs in a final volume of 20 µL
209 using SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad) as recommended by the
210 manufacturer in a MJ Research PTC-200 DNA Engine® Peltier Thermal Cycler (Bio-Rad)
211 with the following cycle conditions: initial denaturation at 98 °C for 30 sec, followed by 40
212 cycles of 95 °C for 15 sec and a 30-second step at 59.4 °C for *codY* and *alsD*, 54.8 °C for
213 *agrC* and *spa*, 51.4 °C for *hld* and *sea*. The specificity of primer sets used for qPCR
214 amplification was evaluated by melting curve analysis. All reactions were independently
215 conducted three times on two biological replicates. Gene expression data analysis using the 2⁻
216 $\Delta\Delta C_T$ method, correlation coefficients of oligonucleotides and efficiency of amplifications
217 were calculated as described previously (Livak and Schmittgen, 2001).

218

219 2.8. Sequencing data access

220 The annotated genome was submitted to the National Center for Biotechnology
221 Information (NCBI) database (Bioproject accession number: PRJNA344949). RNA-Seq data
222 generated in this experiment was deposited in the Sequence Read Archive (Leinonen et al.,
223 2011) of the NCBI and is publicly available through the accession number SRP092596.

224

225 2.9. Statistical analysis

226 Data from HPLC analysis were subjected to one-way analysis of variance (ANOVA),
227 while differences in mean counts of microbial populations in single and co-culture
228 experiments were assessed by a Student's t-test using Statistica software version 6 (Statsoft,
229 Tulsa, OK, USA). Statistical significance was set at $p < 0.05$.

230

231 3. Results

232 3.1. Growth kinetics, enterotoxin production and milk acidification

233 The mean counts of *S. aureus* ATCC 29213 and *E. faecalis* 41FL1 populations in
234 single and co-culture, as well as SE production, in sterile skimmed milk at 30 °C for 24 h are
235 reported in Table 1. pH measurements and water-soluble metabolites present in milk of single
236 and co-culture are reported in Table 2. *S. aureus* growth was hindered in co-culture with *E.*
237 *faecalis* compared to single culture, resulting in a 1.2-log inhibition after 24 h. The ability of
238 *S. aureus* to produce enterotoxins was verified only at 24 h in single culture, whereas no SE
239 was detected in co-culture throughout the incubation period (Table 1). Interestingly, *S. aureus*
240 ATCC 29213 showed virtually no impact on the growth dynamic of *E. faecalis* 41FL1, as the
241 latter largely followed the same growth pattern whether in single or co-culture, reaching a
242 final population density of approximately 1.0×10^9 CFU/mL after 24 h. Progressive
243 acidification of milk was observed in the presence of *E. faecalis* in single and co-culture, with
244 final pH values lower than 5 simultaneously with an increasing production of lactic acid and
245 consequently decrease in lactose content in milk (Table 2). Conversely, pH values, lactic
246 acid, and lactose contents remained virtually unaltered in *S. aureus* single culture.

247

248 3.2. *S. aureus* ATCC 29213 genome sequencing and transcriptomic response

249 The complete sequencing of *S. aureus* ATCC 29213 genome was obtained by
250 Illumina MiSeq Sequencing System technology for 250-bp paired-end sequencing. A total of
251 86 contigs were assembled using MIRA version 4.0.2 to provide a genome length of
252 2,847,591 bp with an average GC content of 32.81%. Sequencing of *S. aureus* ATCC 29213
253 genome revealed the presence of 2,676 CDS and 120 RNAs (19 rRNAs, 99 tRNAs and 2
254 pseudogenes). The number of sequences obtained in the RNA-Seq experiments ranged from
255 224,614 to 804,266 for each replicate, with a total of 4,017,675 read counts. For gene
256 expression studies, the transcriptome of *S. aureus* in single culture was used as a reference for
257 assessing changes in gene expression in co-culture by means of fold change analysis.
258 Therefore, we refer to upregulated and downregulated in comparing transcript levels between

259 the two conditions. Data was analyzed using an arbitrary cutoff value of LFC between -1 and
260 +1.

261 In Figure 1, the transcriptomic response of *S. aureus* ATCC 29213 in single and co-
262 culture was divided into 5 categories depicting the frequency of genes belonging to each
263 pattern over time. After 7 h of incubation, 303 genes were differentially expressed in co-
264 culture, of which 257 genes were upregulated and 46 genes were downregulated. After 24 h,
265 *S. aureus* transcriptome in co-culture showed an increased number of transcripts belonging to
266 597 genes and a reduced number of transcripts coding for 63 genes. In both time intervals, 9
267 genes were found to be transcribed in lower rates whereas transcripts of 59 genes were
268 significantly enriched in co-culture (Figure 1A and 1B; Tables S1 and S2). Considering
269 single and co-culture transcriptomes, transcripts coding for 748 genes were absent at 7 h,
270 whereas this number decreased to 197 genes at 24 h, with 146 shared genes (Figure 1C).
271 Moreover, transcripts coding for 1,375 and 1,512 genes was not detected in co-culture at 7
272 and 24 h, respectively, with 856 overlapping genes in this pattern (Figure 1D). Finally,
273 transcription of 35 and 39 genes occurred exclusively in co-culture at 7 and 24 h,
274 respectively, sharing no common representatives (Figure 1E; Table S3 e S4). Overall, these
275 observations indicate that the global transcriptomic response of *S. aureus* in co-culture with
276 *E. faecalis* diverges considerably from that observed in single culture.

277 To validate the RNA-Seq experiments, the expression of six *S. aureus* genes (*codY*,
278 *alsD*, *agrC*, *spa*, *hld*, *sea*) up- or downregulated in co-culture by more than twofold was
279 evaluated by RT-qPCR. Overall, trends in the expression of selected genes as determined by
280 RT-qPCR corroborated the RNA-Seq findings, with minor inconsistencies (Figure S1).
281 Notably, RT-qPCR showed an upregulation of *codY* by 6-fold in co-culture at 7 h, whereas
282 *alsD* showed a converse expression pattern from that obtained with RNA-Seq. The
283 expression of *agrC*, *spa*, *hld* and *sea* was found to be strongly decreased in co-culture at 7 h
284 and 24 h according to RT-qPCR analysis.

285 Gene expression analysis based on the relative abundance of transcripts assigned to
286 their respective COG categories revealed the occurrence of distinct transcriptional response
287 profiles in co-culture in each tested interval (Figure 2). To guide our analysis, we calculated
288 the relative percentage difference of total RPKM values obtained in each COG category at 7
289 and 24 h in co-culture versus single culture, and arbitrarily chosen to focus on COG
290 categories with changes of plus or minus 40%. After 7 h of incubation, a large amount of
291 transcripts in co-culture were assigned to categories ‘Transcription’ (+44.51%), ‘Replication,
292 recombination and repair’ (+182.94%) and ‘Posttranslational modification, protein turnover,
293 and chaperones’ (+72.81%), while the expression of genes related to the ‘Nucleotide
294 transport and metabolism’ (-45.93%) was negatively impacted, which strongly indicates the
295 triggering of stress response cascades in co-culture (Figure 2A). In contrast, after 24 hours of
296 incubation, the categories ‘Carbohydrate transport and metabolism’ (+108.78%), Coenzymes
297 transport and metabolism (+60.59%) and ‘Secondary metabolites biosynthesis, transport and
298 catabolism’ (+46.02%) were expressively more abundant in co-culture, whereas transcripts
299 encoding proteins assigned to the category ‘Translation, ribosomal structure, and biogenesis’
300 had considerably decreased expression compared to single culture (-45.23%), suggesting
301 major adaptations in *S aureus* central metabolism (Figure 2B).

302 Compared to single culture, the presence of transcripts encoding enzymes involved in
303 the glycolytic and tricarboxylic acid (TCA) cycle pathways was lower in co-culture at 7 h
304 (Figure 3). In particular, transcription of genes coding for six important glycolytic enzymes
305 (*fbaA*, *fda*, *tpiA*, *gapA1*, *pgm* and *pykA*) was 4.2, 2.0, 7.9, 2.7, 2.6 and 2.8-fold lower in co-
306 culture at 7 h compared to single culture, respectively. Transcripts encoding enzymes of the
307 pyruvate dehydrogenase complex (*pdhA*, *pdhB*, *pdhC*, *pdhD*) and citrate synthase (*citZ*) did
308 not accumulate in co-culture to the same extent as in single culture at 7 h. However, after 24
309 hours of incubation, we observed a noteworthy enrichment in the transcripts coding for
310 enzymes involved in glycolysis, acetoin and acetate metabolism (Figure 3). Transcripts of
311 genes coding for all glycolytic enzymes, except for *gapA1*, accumulated significantly in co-

312 culture compared to single culture at 24 h. Likewise, transcripts of acetolactate synthase
313 (*alsS*), acetolactate decarboxylase (*alsD*) and acetoin reductase (*butA*) involved in the 2,3-
314 butanediol pathway and acetoin production were present in significantly larger amounts in
315 co-culture at 24 h compared to single culture.

316 Considering the transcription of genes belonging to the *agr* locus (*agrBCDA*) in *S.*
317 *aureus* single and co-culture, we noted a transient upregulation of *agrC* by 10-fold in co-
318 culture at 7 h, but no transcripts of the *agrBCDA* locus were detected in co-culture at 24 h.
319 The RNAPIII transcript encoding the δ -hemolysin (Hld) was not detected in the co-culture
320 transcriptome at neither 7 h nor 24 h, while high amounts of this transcript were found in
321 single culture at 24 h accompanied by the presence of transcripts of *agrBCDA* (Table 3). In
322 addition, transcripts of other notable global transcription regulators of *S. aureus* virulence,
323 such as *mgrA* (Crosby et al., 2016), *saeS*, *saeP* (Giraud et al., 1997), and several *sarA*
324 homologs (Cheung et al., 2004) were absent in co-culture at 7 h or 24 h (Table 3).
325 Accordingly, transcripts coding for several *S. aureus* secreted toxins and exoenzymes were
326 absent or present in significantly reduced numbers in co-culture at 24 h, whereas transcripts
327 coding for several cell surface-associated proteins were upregulated in the same conditions
328 (Table 3).

329

330 **4. Discussion**

331 A deeper knowledge concerning bacterial interactions is needed to support new ways
332 of thinking the control of growth and virulence of pathogenic microorganisms in food
333 products. Although considerable research has been dedicated to investigating the negative
334 impact of certain LAB strains over the ability of *S. aureus* to grow and produce enterotoxins
335 in food-mimicking conditions (Alomar et al., 2008; Daoud and Devere, 1985; Kao and
336 Frazier, 1966; Noleto et al., 1987), rather less amount of knowledge has been pulled together
337 concerning the molecular mechanisms underlying this antagonistic phenomenon. In the
338 present study, we demonstrated that *S. aureus* ATCC 29213 growth and enterotoxigenic

339 ability were hindered in co-culture with *E. faecalis* 41FL1 in skimmed milk at 30 °C for 24 h
340 (Table 1). Our RNA-Seq data revealed considerable changes in the transcriptomic response
341 of *S. aureus* in co-culture compared to single culture in the tested conditions. To our
342 knowledge, this is the first study to present a comprehensive analysis of the alterations in the
343 *S. aureus* transcriptome during growth in the presence of *E. faecalis* in skimmed milk.

344 Considering *S. aureus* genes upregulated or whose transcripts were found exclusively
345 in co-culture at 7 h (Figure 2A; Table S1; Table S3), we identified an enrichment of
346 transcripts coding for repair and detoxifying enzymes, chaperones, and several stress-induced
347 transcriptional regulators. Of particular interest, we identified an upregulation of the
348 following *S. aureus* genes in co-culture: *arcA* (expressed exclusively in co-culture at 7 h and
349 upregulated by 2.63-fold at 24 h), *sigB* (7.99-fold), and *rex* (19.98-fold). The alternative
350 sigma factor σ^B is involved in the response to a variety of environmental stresses in *S. aureus*,
351 including pH-sensing transduction pathway (Gertz et al., 2000; Weinrick et al., 2004). The
352 arginine deiminase ArcA, as part of the arginine deiminase (ADI) operon (*arcABDC*),
353 catalyzes the hydrolysis of arginine into citrulline and ammonia, thus contributing to pH
354 homeostasis and survival under non-lethal acid stress (Makhlin et al., 2007). Remarkably,
355 *arcA* was shown to be upregulated in co-culture also at 24 h, as well as *arcD*, *arcB* and
356 *arcC1*, a homolog for carbamate kinase found in the *S. aureus* ATCC 29213 genome outside
357 of the ADI operon. Additionally, the redox sensing transcriptional factor Rex is known to be
358 involved in the regulation of metabolic pathways that mediate NAD^+ regeneration and ATP
359 synthesis in *S. aureus*, such as the ADI pathway, to assure survival and pathogenicity (Pagels
360 et al., 2010). It has been previously described the strong ability of *E. faecalis* to reduce the
361 redox potential and eliminate dissolved O_2 during growth in milk (Brasca et al., 2007;
362 Morandi et al., 2016), a characteristic likely to have contributed to the generation of an
363 oxygen-limiting environment in our co-culture conditions with profound implications over
364 *rex* expression in *S. aureus*. Collectively, the upregulation of such genes by *S. aureus* in co-
365 culture can be interpreted as part of a structured response to counteract the effects of pH- and

366 redox-induced stress caused by the presence of *E. faecalis*, also corroborated by the chemical
367 changes in milk samples where *E. faecalis* was inoculated (Table 2).

368 At 24 h, COG enrichment analysis revealed an accumulation of transcripts belonging
369 to the category of carbohydrate transport and metabolism in co-culture (Figure 2B). In
370 particular, transcripts encoding enzymes involved in glycolysis and acetoin production
371 (*alsSD* and *butA*) were significantly enriched in co-culture at 24 h (Figure 4), suggesting a
372 shift towards fermentative metabolism in *S. aureus* in co-culture with *E. faecalis*.
373 Interestingly, an upregulation of *alsSD* and *butA* in *S. aureus* grown in the presence of
374 another LAB species (*L. lactis*) in a cheese matrix has been reported (Cretenet et al., 2011).
375 In *S. aureus*, increased glycolytic activity and the activation of the acetoin biosynthetic
376 pathway in environments rich in organic acids were shown to be a strategy to enhance
377 survival by limiting metabolic-mediated cell death and lysis (Yang et al., 2006). In such
378 cases, the generation of acetoin by *S. aureus* could be useful for NAD⁺ regeneration, energy
379 conservation and maintenance of intracellular pH homeostasis (Thomas et al., 2014). In
380 support of these major alterations in the transcription of genes related to central carbon
381 metabolism in co-culture, we also identified that transcripts of metabolite-responsive
382 regulators known to be directly or indirectly involved in the regulation of the expression of
383 several virulence factors in *S. aureus*, namely *codY* (Pohl et al., 2009; Roux et al., 2014),
384 *rpiRA* and *rpiRB* (Zhu et al., 2011), accumulated in significantly larger amounts in co-culture
385 at 24 h, as well as *ccpA* and *ccpE* at both 7 h and 24 h (Sonenshein, 2007, 2005) (Table 3;
386 Table S1). In particular, CodY may act as a roadblock to transcription of *agr* P2 and P3
387 promoters (Majerczyk et al., 2010; Roux et al., 2014). Pathogenic bacteria use metabolite-
388 responsive regulators to link metabolic status, energy homeostasis and synthesis of virulence
389 determinants to the availability of biosynthetic intermediates derived from the glycolytic,
390 pentose phosphate and TCA pathways (Richardson et al., 2015). It has been shown that *S.*
391 *aureus* compromises its growth and pathogenic potential in favor of survival in conditions or
392 environments where energy generation is limited (Somerville et al., 2002; Zhu et al., 2009) or

393 in competition with other microorganisms (Filkins et al., 2015; Orazi and O'Toole, 2017;
394 Ramsey et al., 2016) mainly by shifting towards fermentative growth and shutting down the
395 activity of energy consuming processes, such as virulence and *agr*-based quorum-sensing.
396 This previous knowledge help us explain, at least partly, our findings related to *S. aureus*
397 decreased growth, failed *agr* activation and overall altered transcriptomic profile in co-culture
398 with *E. faecalis*.

399 The pathogenicity of *S. aureus* involves the production of a wide array of cell wall-
400 attached and secreted proteins, whose expression is regulated at multiple levels in response to
401 cell density, energy availability, environmental and intracellular signals (Cheung et al., 2004;
402 Novick and Geisinger, 2008). In this context, *trans*-acting factors, such as sigma factors, two-
403 component systems (TCS) and metabolite-responsive elements, act as a dynamic network that
404 allows *S. aureus* to fine-tune its metabolism and virulence (Ibarra et al., 2013). The most
405 well-described and widespread TCS in *S. aureus* is the *agr* locus, which comprises four genes
406 (*agrBDCA*) and also functions as a quorum-sensing system whose intracellular effector is the
407 bifunctional RNAIII, a small regulatory RNA transcribed from the P3 promoter of the *agr*
408 system at higher cell densities (Novick et al., 1993; Novick and Geisinger, 2008). As an
409 opposing effector of the *agr* locus in the double-selector switch that governs *S. aureus*
410 virulence is found another transcriptional regulator, i.e. the Repressor of toxins (Rot)
411 (Bronesky et al., 2016). Rot simultaneously activates the transcription of genes encoding
412 several surface proteins and immunomodulators while repressing the transcription of
413 exotoxins, pore-forming toxins and exoenzymes (Saïd-Salim et al., 2003). *rot* activity occurs
414 at low cell density and its translation is repressed by RNAIII (Geisinger et al., 2006; Hsieh et
415 al., 2008). Thus, Rot and RNAIII have opposing roles in enabling *S. aureus* to phenotypic
416 switch between defense (adhesion, immune evasion and biofilm formation) and offensive
417 modes (degradation of host tissues and subversion of host defenses) (Nitzan et al., 2015). In
418 our study, we observed that the classical post-exponential activation of *agr* and its regulon
419 did not occur in *S. aureus* during growth in co-culture with *E. faecalis*, an observation

420 seemingly not related to cell density, since *S. aureus* population in co-culture easily reached
421 the quorum sensing threshold ($\geq 10^6$ CFU/mL) (Table 1). Moreover, the transcription of *rot*
422 was found to be upregulated in co-culture at 7 h by 2.46-fold while no *rot* transcripts were
423 detected at 24 h (Table 3). Among Rot-regulated virulence genes (Saïd-Salim et al., 2003),
424 we found an upregulation of *coa* and *sdrC* as well as a lack of transcripts coding for the
425 serine protease operons (*splABCDEF* and *sspBC*) and urease complex (*ureABCDEF*) in co-
426 culture transcriptome at 7 h (Table 3), which is consistent with the finding of higher amounts
427 of *rot* transcripts under this condition. However, the number of transcripts coding for *hla* and
428 *geh*, genes whose transcription is supposedly negatively regulated by Rot (Saïd-Salim et al.,
429 2003), was found to be increased by 7.99-fold and 5.33-fold, respectively, in co-culture at 7
430 h, probably by *rot*- and/or *agr*-independent control mechanisms.

431 Enterotoxin production is one of the most important virulence-related traits of *S.*
432 *aureus* for food safety concerns, since SEs are often implicated in foodborne intoxication
433 outbreaks. We identified by means of whole genome sequencing the presence of genes
434 coding for staphylococcal enterotoxin A (*sea*), enterotoxin P (*selp*) and the enterotoxin gene
435 cluster (*seg*, *sen*, ψ *ent*₁, ψ *ent*₂, *sei*, *sem*, *seo*) in the genome of *S. aureus* ATCC 29213. Since
436 the assessment of SE production in our single- and co-culture experiments included only the
437 five classical enterotoxins, we thus focused on discussing the aspects concerning exclusively
438 *sea* regulation in the genetic background of *S. aureus* ATCC 29213. *sea* is located on the
439 genome of *Siphoviridae* temperate bacteriophages whose life cycle is characterized by two
440 phases: the lysogenic and the lytic phase (Deghorain and Van Melderen, 2012). The
441 expression of *sea* is not under the control of the *agr* system but related to the life cycle of the
442 prophage; transition to the phage's lytic phase potentially activates and/or even enhances *sea*
443 transcription (Sumby and Waldor, 2003; Tremaine et al., 1993). It has been shown that
444 prophage induction leads to an increase in the amount of phage replicative form, as well as
445 *sea* gene copies and transcripts, ultimately leading to an enhanced SEA production (Cao et
446 al., 2012; Zeaki et al., 2015). More recently, it has been demonstrated that the alternative

447 sigma factor σ^H (*sigH*) promotes and stabilizes the lysogenization of several prophages in *S.*
448 *aureus* by upregulating the mRNA levels of prophage integrases (Tao et al., 2010).
449 Paralleling these observations with our findings, it is worth citing that *sigH* transcripts
450 accumulated significantly (> 15-fold) in *S. aureus* co-culture at 7 h compared to single
451 culture (Table 3), which possibly contributed to the lack of SEA in co-culture throughout the
452 incubation period.

453

454 **5. Conclusions**

455 Many studies have attempted to explore the interaction between pathogenic bacteria
456 and intrinsic positive microbiota in foods. In the present work, we discussed the alterations in
457 the *S. aureus* transcriptome during growth with *E. faecalis* in skimmed milk and provided
458 novel contributions to the current understanding of the LAB-*S. aureus* interaction in food-
459 mimicking conditions. However, we failed to determine a single explanation for our
460 observations, which are likely to have arisen as a result of a combined effect acidic and redox
461 stress with utmost negative impact on the expression of *S. aureus* virulence. Further
462 assessment of our transcriptomic findings at a proteomic level would be expected to provide
463 additional insights into this bacterial interaction, since transcript levels do not necessarily
464 correlate with the levels of its respective protein. Additionally, it would be of great interest to
465 investigate whether *S. aureus* replicates the rearrangement of its gene expression in similar
466 ways as described herein in the presence of a more complex microbiota and/or different
467 surrounding matrix.

468

469 **Acknowledgements**

470 This research was funded by the “Conselho Nacional do Desenvolvimento Científico
471 e Tecnológico – CNPq”, Brazil [#485337/2012-6, 2012]. Gabriela Nogueira Viçosa is the
472 recipient of a Ph.D. fellowship from the “Coordenação de Aperfeiçoamento de Pessoal de
473 Nível Superior - CAPES” Foundation, Brazil [1529-13/7, 2013]. We also thank Francesca

474 Turrone and Gabriele Andrea Lugli from GenProbio, Parma, Italy for their support in the
475 analysis of RNA-Seq data.

476

477 **References**

478 Alomar, J., Loubiere, P., Delbes, C., Nouaille, S., Montel, M.C., 2008. Effect of
479 *Lactococcus garvieae*, *Lactococcus lactis* and *Enterococcus faecalis* on the behaviour of
480 *Staphylococcus aureus* in microfiltered milk. Food Microbiol. 25, 502–508.

481 Altschul, S., Gish, W., Miller, W., Myers, E., Lipman, D., 1990. Basic local
482 alignment search tool. J. Mol. Biol. 215, 403-410.

483 Argudín, M.Á., Mendoza, M.C., Rodicio, M.R., 2010. Food poisoning and
484 *Staphylococcus aureus* enterotoxins. Toxins (Basel). 2, 1751–73.

485 Brasca, M., Morandi, S., Lodi, R., Tamburini, A., 2007. Redox potential to
486 discriminate among species of lactic acid bacteria. J. Appl. Microbiol. 103, 1516–1524.

487 Bronesky, D., Wu, Z., Marzi, S., Walter, P., Geissmann, T., Moreau, K., Caldelari, I.,
488 Romby, P., 2016. *Staphylococcus aureus* RNAlII and its regulon link quorum sensing, stress
489 responses, metabolic adaptation, and regulation of virulence gene expression. Annu. Rev.
490 Microbiol. 70, 299–316.

491 Cao, R., Zeaki, N., Wallin-Carlquist, N., Skandamis, P.N., Schelin, J., Radstrom, P.,
492 2012. Elevated enterotoxin A expression and formation in *Staphylococcus aureus* and its
493 association with prophage induction. Appl. Environ. Microbiol. 78, 4942–4948.

494 Carr, F.J., Chill, D., Maida, N., 2002. The lactic acid bacteria: A literature survey.
495 Crit. Rev. Microbiol. 28, 281–370.

496 Charlier, C., Even, S., Gautier, M., Le Loir, Y., 2008. Acidification is not involved in
497 the early inhibition of *Staphylococcus aureus* growth by *Lactococcus lactis* in milk. Int.
498 Dairy J. 18, 197–203.

499 Cheung, A.L., Bayer, A.S., Zhang, G., Gresham, H., Xiong, Y.Q., 2004. Regulation
500 of virulence determinants in vitro and in vivo in *Staphylococcus aureus*. FEMS Immunol.

501 Med. Microbiol. 40, 1–9.

502 Chevreur, B., Wetter, T., Suhai, S., 1999. Genome Sequence Assembly Using Trace
503 Signals and Additional Sequence Information. Comput. Sci. Biol. Proc. Ger. Conf.
504 Bioinforma. '99, GCB, Hann. Ger. 45–56.

505 Cretenet, M., Nouaille, S., Thouin, J., Rault, L., Stenz, L., François, P., Hennekinne,
506 J.A., Piot, M., Maillard, M.B., Fauquant, J., Loubière, P., Loir, Y. Le, Even, S., 2011.
507 *Staphylococcus aureus* virulence and metabolism are dramatically affected by *Lactococcus*
508 *lactis* in cheese matrix. Environ. Microbiol. Rep. 3, 340–351.

509 Crosby, H.A., Schlievert, P.M., Merriman, J.A., King, J.M., Salgado-Pabón, W.,
510 Horswill, A.R., 2016. The *Staphylococcus aureus* global regulator MgrA modulates clumping
511 and virulence by controlling surface protein expression. PLOS Pathog. 12(5):e1005604.
512 doi:10.1371/journal.ppat.1005604

513 Dal Bello, B., Rantsiou, K., Bellio, A., Zeppa, G., Ambrosoli, R., Civera, T.,
514 Cocolin, L., 2010. Microbial ecology of artisanal products from North West of Italy and
515 antimicrobial activity of the autochthonous populations. LWT - Food Sci. Technol. 43, 1151–
516 1159.

517 Daoud, S.M., Debevere, J.M., 1985. The effect of *Bacillus subtilis* and *Streptococcus*
518 *faecalis* var. *liquefaciens* on staphylococcal A activity. Int. J. Food Microbiol. 2, 211–218.

519 Darling, A.C.E., Mau, B., Blattner, F.R., Perna, N.T., 2004. Mauve: Multiple
520 alignment of conserved genomic sequence with rearrangements. Genome Res. 14, 1394–
521 1403.

522 De Buyser, M.L., Lombard, B., Schulten, S.M., In't Veld, P.H., Scotter, S.L., Rollier,
523 P., Lahellec, C., Buyser, M.L. De, Lombard, B., Schulten, S.M., In, P.H., 2003. Validation of
524 EN ISO standard methods 6888 Part 1 and Part 2: 1999 - Enumeration of coagulase-positive
525 staphylococci in foods. Int. J. Food Microbiol. 83, 185–194.

526 Deghorain, M., Van Melderren, L., 2012. The staphylococci phages family: An
527 overview. Viruses 4, 3316–3335.

528 Delpech, P., Bornes, S., Alaterre, E., Bonnet, M., Gagne, G., Montel, M.-C., Delbès,
529 C., 2015. *Staphylococcus aureus* transcriptomic response to inhibition by H₂O₂-producing
530 *Lactococcus garvieae*. Food Microbiol. 51, 163–70.

531 European Food Safety Authority, 2016. The European Union summary report on
532 trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2015. EFSA J.
533 14(12):4634.

534 Even, S., Charlier, C., Nouaille, S., Ben Zakour, N.L., Cretenet, M., Cousin, F.J.,
535 Gautier, M., Cocaïgn-Bousquet, M., Loubière, P., Le Loir, Y., 2009. *Staphylococcus aureus*
536 virulence expression is impaired by *Lactococcus lactis* in mixed cultures. Appl. Environ.
537 Microbiol. 75, 4459–4472.

538 Filkins, L.M., Graber, J.A., Olson, D.G., Dolben, E.L., Lynd, L.R., Bhujju, S.,
539 O’Toole, G.A., 2015. Coculture of *Staphylococcus aureus* with *Pseudomonas aeruginosa*
540 drives *S. aureus* towards fermentative metabolism and reduced viability in a cystic fibrosis
541 model. J. Bacteriol. 197, 2252–2264.

542 Foulquié-Moreno, M.R., Sarantinopoulos, P., Tsakalidou, E., De Vuyst, L., 2006.
543 The role and application of enterococci in food and health. Int. J. Food Microbiol. 106, 1–24.

544 Franciosi, E., Settanni, L., Cavazza, A., Poznanski, E., 2009. Presence of enterococci
545 in raw cow’s milk and “Puzzone di Moena” cheese. J. Food Process. Preserv. 33, 204–217.

546 Geisinger, E., Adhikari, R.P., Jin, R., Ross, H.F., Novick, R.P., 2006. Inhibition of
547 *rot* translation by RNAIII, a key feature of *agr* function. Mol. Microbiol. 61, 1038–1048.

548 Gertz, S., Engelmann, S., Schmid, R., Ziebandt, A.-K., Tischer, K., Scharf, C.,
549 Hacker, J., Hecker, M., 2000. Characterization of the σ^B regulon in *Staphylococcus aureus*. J.
550 Bacteriol. 182, 6983–6991.

551 Giraud, A.T., Cheung, A.L., Nagel, R., 1997. The *sae* locus of *Staphylococcus*
552 *aureus* controls exoprotein synthesis at the transcriptional level. Arch. Microbiol. 168, 53–58.

553 Haines, W.C., Harmon, L.G., 1973. Effect of selected lactic acid bacteria on growth
554 of *Staphylococcus aureus* and production of enterotoxin. Appl. Microbiol. 25, 436–441.

555 Hamama, A., El Hankouri, N., El Ayadi, M., 2002. Fate of enterotoxigenic
556 *Staphylococcus aureus* in the presence of nisin-producing *Lactococcus lactis* strain during
557 manufacture of Jben, a Moroccan traditional fresh cheese. *Int. Dairy J.* 12, 933–938.

558 Hsieh, H.Y., Ching, W.T., Stewart, G.C., 2008. Regulation of *rot* expression in
559 *Staphylococcus aureus*. *J. Bacteriol.* 190, 546–554.

560 Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010.
561 Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC*
562 *Bioinformatics* 11(1):119. doi:10.1186/1471-2105-11-119

563 Ibarra, J.A., Perez-Rueda, E., Carroll, R.K., Shaw, L.N., 2013. Global analysis of
564 transcriptional regulators in *Staphylococcus aureus*. *BMC Genomics* Feb 26;14:126. doi:
565 10.1186/1471-2164-14-126

566 Jensen, L.J., Julien, P., Kuhn, M., von Mering, C., Muller, J., Doerks, T., Bork, P.,
567 2008. eggNOG: automated construction and annotation of orthologous groups of genes.
568 *Nucleic Acids Res.* 36, D250–D254.

569 Kao, C.T.C.T., Frazier, W.C.C., 1966. Effect of lactic acid bacteria on growth of
570 *Staphylococcus aureus*. *Appl. Microbiol.* 14, 251–255.

571 Lagesen, K., Hallin, P., Rødland, E.A., Stærfeldt, H.H., Rognes, T., Ussery, D.W.,
572 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids*
573 *Res.* 35, 3100–3108.

574 Laughton, J.M., Devillard, E., Heinrichs, D.E., Reid, G., McCormick, J.K., 2006.
575 Inhibition of expression of a staphylococcal superantigen-like protein by a soluble factor
576 from *Lactobacillus reuteri*. *Microbiology* 152, 1155–1167.

577 Leinonen, R., Sugawara, H., Shumway, M., 2011. The sequence read archive.
578 *Nucleic Acids Res.* 39, D19–D21.

579 Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-
580 Wheeler transform. *Bioinformatics* 25, 1754–1760.

581 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,

582 Abecasis, G., Durbin, R., 2009. The Sequence Alignment/Map format and SAMtools.
583 Bioinformatics 25, 2078–2079.

584 Li, J., Wang, W., Xu, S.X., Magarvey, N.A., McCormick, J.K., 2011. *Lactobacillus*
585 *reuteri*-produced cyclic dipeptides quench *agr*-mediated expression of toxic shock syndrome
586 toxin-1 in staphylococci. Proc. Natl. Acad. Sci. U S A 108(8):3360–3365.

587 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using
588 real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods 25, 402–408.

589 Lowe, T.M., Eddy, S.R., 1997. tRNAscan-SE: a program for improved detection of
590 transfer RNA genes in genomic sequence. Nucleic Acids Res. 25, 955–964.

591 Majerczyk, C.D., Dunman, P.M., Luong, T.T., Lee, C.Y., Sadykov, M.R.,
592 Somerville, G.A., Bodi, K., Sonenshein, A.L. 2010. Direct targets of CodY in *Staphylococcus*
593 *aureus*. J Bacteriol 192:2861–2877.

594 Makhlin, J., Kofman, T., Borovok, I., Kohler, C., Engelmann, S., Cohen, G.,
595 Aharonowitz, Y., 2007. *Staphylococcus aureus* ArcR controls expression of the arginine
596 deiminase operon. J. Bacteriol. 189, 5976–5986.

597 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytzky, A.,
598 Garimella, K., Altshuler, D., Gabriel, S., Daly, M., DePristo, M.A., 2010. The Genome
599 Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing
600 data. Genome Res. 20, 1297–1303.

601 Morandi, S., Silveti, T., Tamburini, A., Brasca, M., 2016. Changes in oxidation-
602 reduction potential during milk fermentation by wild lactic acid bacteria. J. Dairy Res. 83,
603 387–394.

604 Nitzan, M., Fechter, P., Peer, A., Altuvia, Y., Bronesky, D., Vandenesch, F., Romby,
605 P., Biham, O., Margalit, H., 2015. A defense-offense multi-layered regulatory switch in a
606 pathogenic bacterium. Nucleic Acids Res. 43, 1357–1369.

607 Noleto, A.L.S., Malburg, L.M., Bergdoll, M.S., 1987. Production of staphylococcal
608 enterotoxin in mixed cultures. Appl. Environ. Microbiol. 53, 2271–2274.

609 Novick, R.P., Geisinger, E., 2008. Quorum sensing in staphylococci.
610 Annu.Rev.Genet. 42, 541–564.

611 Novick, R.P., Ross, H.F., Projan, S.J., Kornblum, J., Kreiswirth, B., Moghazeh, S.,
612 1993. Synthesis of staphylococcal virulence factors is controlled by a regulatory RNA
613 molecule. EMBO J. 12, 3967–3975.

614 Okonechnikov, K., Golosova, O., Fursov, M., Varlamov, A., Vaskin, Y., Efremov, I.,
615 German Grehov, O.G., Kandrov, D., Rasputin, K., Syabro, M., Tleukenov, T., 2012. Unipro
616 UGENE: A unified bioinformatics toolkit. Bioinformatics 28, 1166–1167.

617 Orazi, G., O’Toole, G.A., 2017. *Pseudomonas aeruginosa* alters *Staphylococcus*
618 *aureus* sensitivity to vancomycin in a biofilm model of cystic fibrosis infection. MBio 8,
619 e00873-17. Pagels, M., Fuchs, S., Pané-Farré, J., Kohler, C., Menschner, L., Hecker, M.,
620 McNamarra, P.J., Bauer, M.C., Von Wachenfeldt, C., Liebeke, M., Lalk, M., Sander, G., Von
621 Eiff, C., Proctor, R.A., Engelmann, S., 2010. Redox sensing by a Rex-family repressor is
622 involved in the regulation of anaerobic gene expression in *Staphylococcus aureus*. Mol.
623 Microbiol. 76, 1142–1161.

624 Perin, L.M., Moraes, P.M., Viçosa, G.N., Silva Júnior, A., Nero, L.A., 2012.
625 Identification of bacteriocinogenic *Lactococcus* isolates from raw milk and cheese capable of
626 producing nisin A and nisin Z. Int. Dairy J. 25, 46–51.

627 Pohl, K., Francois, P., Stenz, L., Schlink, F., Geiger, T., Herbert, S., Goerke, C.,
628 Schrenzel, J., Wolz, C., 2009. CodY in *Staphylococcus aureus*: A regulatory link between
629 metabolism and virulence gene expression. J. Bacteriol. 191, 2953–2963.

630 Ramsey, M.M., Freire, M.O., Gabriliska, R.A., Rumbaugh, K.P., Lemon, K.P., 2016.
631 *Staphylococcus aureus* shifts toward commensalism in response to *Corynebacterium* species.
632 Front. Microbiol. 7, 1–15.

633 Richardson, A.R., Somerville, G.A., Sonenshein, A.L., 2015. Regulating the
634 intersection of metabolism and pathogenesis in Gram-positive bacteria. Microbiol. Spectr. 3,
635 1–27.

636 Roux, A., Todd, D.A., Velazquez, J. V., Cech, N.B., Sonenshein, A.L., 2014. CodY-
637 mediated regulation of the *Staphylococcus aureus* Agr system integrates nutritional and
638 population density signals. *J. Bacteriol.* 196, 1184–1196.

639 Saïd-Salim, B., Dunman, P.M., McAleese, F.M., Macapagal, D., Murphy, E.,
640 McNamara, P.J., Arvidson, S., Foster, T.J., Projan, S.J., Kreiswirth, B.N., 2003. Global
641 regulation of *Staphylococcus aureus* genes by Rot. *J. Bacteriol.* 185, 610–619.

642 Schelin, J., Wallin-Carlquist, N., Cohn, M.T., Lindqvist, R., Barker, G.C., Rådström,
643 P., 2011. The formation of *Staphylococcus aureus* enterotoxin in food environments and
644 advances in risk assessment. *Virulence* 2, 580–592.

645 Somerville, G.A., Chaussee, M.S., Morgan, C.I., Fitzgerald, J.R., Dorward, D.W.,
646 Reitzer, L.J., Musser, J.M., 2002. *Staphylococcus aureus* aconitase inactivation unexpectedly
647 inhibits post-exponential-phase growth and enhances stationary-phase survival. *Infect.*
648 *Immun.* 70, 6373–6382.

649 Sonenshein, A.L., 2007. Control of key metabolic intersections in *Bacillus subtilis*.
650 *Nat. Rev. Microbiol.* 5, 917–927.

651 Sonenshein, A.L., 2005. CodY, a global regulator of stationary phase and virulence
652 in Gram-positive bacteria. *Curr. Opin. Microbiol.* 8, 203–207.

653 Sonnhammer, E.L.L., Eddy, S.R., Durbin, R., 1997. Pfam: A comprehensive
654 database of protein domain families based on seed alignments. *Proteins Struct. Funct. Genet.*
655 28, 405–420.

656 Stiles, M.E., Holzapel, W.H., 1997. Lactic acid bacteria of foods and their current
657 taxonomy. *Int. J. Food Microbiol.* 36, 1–29.

658 Sumbly, P., Waldor, M.K., 2003. Transcription of the toxin genes present within the
659 *Staphylococcal* phage $\phi Sa3ms$ is intimately linked with the phage's life cycle. *J. Bacteriol.*
660 185, 6841–6851.

661 Tao, L., Wu, X., Sun, B., 2010. Alternative sigma factor σ^H modulates prophage
662 integration and excision in *Staphylococcus aureus*. *PLoS Pathog.* 6(5): e1000888.

663 Tatusov, R.L., Galperin, M.Y., Natale, D.A., Koonin, E. V, 2000. The COG
664 database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids*
665 *Res.* 28, 33–36.

666 Thomas, V.C., Sadykov, M.R., Chaudhari, S.S., Jones, J., Endres, J.L., Widhelm,
667 T.J., Ahn, J.S., Jawa, R.S., Zimmerman, M.C., Bayles, K.W., 2014. A central role for carbon-
668 overflow pathways in the modulation of bacterial cell death. *PLoS Pathog.* 10(6):e1004205.
669 doi:10.1371/journal.ppat.1004205

670 Tremaine, M.T., Brockman, D.K., Betley, M.J., 1993. Staphylococcal enterotoxin A
671 gene (*sea*) expression is not affected by the accessory gene regulator (*agr*). *Infect. Immun.*
672 61, 356–359.

673 Valero, A., Pérez-Rodríguez, F., Carrasco, E., Fuentes-Alventosa, J.M., García-
674 gimeno, R.M., Zurera, G., 2009. Modelling the growth boundaries of *Staphylococcus aureus*:
675 Effect of temperature, pH and water activity. *Int. J. Food Microbiol.* 133, 186–194.

676 Weinrick, B., Dunman, P.M., McAleese, F., Projan, S.J., Fang, Y., Novick, R.P.,
677 Murphy, E., Projan, S.J., Fang, Y., Novick, R.P., 2004. Effect of mild acid on gene
678 expression in *Staphylococcus aureus*. *J. Bacteriol.* 186, 8407–8423.

679 Yang, S.J., Dunman, P.M., Projan, S.J., Bayles, K.W., 2006. Characterization of the
680 *Staphylococcus aureus* CidR regulon: Elucidation of a novel role for acetoin metabolism in
681 cell death and lysis. *Mol. Microbiol.* 60, 458–468.

682 Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., Madden, T.L., 2012.
683 Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC*
684 *Bioinformatics* 13:134. doi:10.1186/1471-2105-13-134

685 Zdenkova, K., Alibayov, B., Karamonova, L., Purkrtova, S., Karpiskova, R.,
686 Demnerova, K., 2016. Transcriptomic and metabolic responses of *Staphylococcus aureus* in
687 mixed culture with *Lactobacillus plantarum*, *Streptococcus thermophilus* and *Enterococcus*
688 *durans* in milk. *J. Ind. Microbiol. Biotechnol.* 43, 1237–1247.

689 Zeaki, N., Susilo, Y.B., Pregiel, A., Rådström, P., Schelin, J., 2015. Prophage-

690 encoded staphylococcal enterotoxin A: Regulation of production in *Staphylococcus aureus*
691 strains representing different *sea* regions. *Toxins (Basel)*. 7, 5359–5376.

692 Zhu, Y., Nandakumar, R., Sadykov, M.R., Madayiputhiya, N., Luong, T.T., Gaupp,
693 R., Lee, C.Y., Somerville, G.A., 2011. RpiR homologues may link *Staphylococcus aureus*
694 RNAIII synthesis and pentose phosphate pathway regulation. *J. Bacteriol.* 193, 6187–6196.

695 Zhu, Y., Xiong, Y.Q., Sadykov, M.R., Fey, P.D., Lei, M.G., Lee, C.Y., Bayer, A.S.,
696 Somerville, G.A., 2009. Tricarboxylic acid cycle-dependent attenuation of *Staphylococcus*
697 *aureus in vivo* virulence by selective inhibition of amino acid transport. *Infect. Immun.* 77,
698 4256–4264.

699

700 Table 1. Mean counts and standard deviation (MC \pm SD) of *S. aureus* ATCC 29213 and *E.*
 701 *faecalis* 41FL1 populations (log₁₀ CFU/mL) and staphylococcal enterotoxin production in
 702 single and co-culture experiments in sterile skimmed milk at 30 °C.

Incubation time (hours)	<i>S. aureus</i>				<i>E. faecalis</i>	
	Single culture		Co-culture		Single culture	Co-culture
	MC \pm SD	SE production	MC \pm SD	SE production		
0	3.33 \pm 0.17	nd	3.10 \pm 0.55	nd	6.26 \pm 0.12	6.30 \pm 0.06
4	3.91 \pm 0.62	nd	4.65 \pm 0.30	nd	7.29 \pm 0.24	7.29 \pm 0.16
7	7.17 \pm 1.28	nd	6.01 \pm 0.28	nd	8.12 \pm 0.11	8.09 \pm 0.18
12	8.28 \pm 0.53 ^a	nd	7.15 \pm 0.13 ^b	nd	8.82 \pm 0.15	9.02 \pm 0.23
24	8.66 \pm 0.32 ^a	detected*	7.49 \pm 0.16 ^b	nd	9.17 \pm 0.11	9.22 \pm 0.18


703 nd: not detected (below detection limit); *combined detection of all five classical
 704 staphylococcal enterotoxins (SEA, SEB, SEC, SED, SEE); ^{a,b}Different letters in each line
 705 indicate difference at 95% level of significance.

16 Table 2. pH, organic acids and sugar contents in skimmed milk during growth of *S. aureus* ATCC 29213 and *E. faecalis* 41FL1 alone or in combination
 17 at 30 °C for 24 h.

Chemical parameter	Treatment	Hours				
		0	4	7	12	24
pH	<i>E. faecalis</i>	6.53 ± 0.00 ^a	6.36 ± 0.02 ^a	6.19 ± 0.01 ^a	5.77 ± 0.02 ^a	4.94 ± 0.02 ^a
	Co-culture	6.54 ± 0.00 ^a	6.37 ± 0.00 ^a	6.21 ± 0.09 ^a	5.78 ± 0.02 ^a	4.93 ± 0.04 ^a
	<i>S. aureus</i>	6.57 ± 0.05 ^b	6.79 ± 0.08 ^b	6.72 ± 0.00 ^b	6.73 ± 0.04 ^b	6.81 ± 0.02 ^b
	Significance	*	*	*	*	*
Lactose	<i>E. faecalis</i>	42.16 ± 0.19	42.01 ± 0.22	41.18 ± 0.38 ^a	40.46 ± 0.38 ^a	35.16 ± 0.20 ^a
	Co-culture	41.77 ± 0.59	41.60 ± 0.50	40.69 ± 0.64 ^a	39.64 ± 0.65 ^a	35.13 ± 0.50 ^a
	<i>S. aureus</i>	42.05 ± 0.21	41.91 ± 0.06	41.53 ± 0.17 ^b	41.12 ± 0.09 ^b	41.01 ± 0.37 ^b
	Significance	ns	ns	*	*	*
Citric acid	<i>E. faecalis</i>	0.37 ± 0.01	0.36 ± 0.01	0.30 ± 0.01 ^b	0.18 ± 0.01 ^b	0.11 ± 0.06 ^b
	Co-culture	0.37 ± 0.01	0.38 ± 0.01	0.26 ± 0.01 ^a	0.14 ± 0.01 ^a	0.09 ± 0.01 ^a
	<i>S. aureus</i>	0.38 ± 0.01	0.37 ± 0.01	0.35 ± 0.01 ^c	0.34 ± 0.01 ^c	0.33 ± 0.01 ^c
	Significance	ns	ns	*	*	*


Chemical parameter	Treatment	Hours				
		0	4	7	12	24
Pyruvic acid	<i>E. faecalis</i>	nd	nd	0.01 ± 0.00	0.02 ± 0.00 ^b	0.03 ± 0.00 ^b
	Co-culture	nd	nd	0.01 ± 0.00	0.02 ± 0.00 ^b	0.03 ± 0.00 ^b
	<i>S. aureus</i>	nd	nd	0.01 ± 0.00	0.01 ± 0.00 ^a	0.01 ± 0.00 ^a
	Significance	-	-	ns	*	*
Lactic acid	<i>E. faecalis</i>	0.04 ± 0.02 ^b	0.14 ± 0.01 ^b	0.33 ± 0.03 ^b	1.73 ± 0.09 ^b	5.45 ± 0.05 ^b
	Co-culture	0.04 ± 0.01 ^b	0.15 ± 0.01 ^b	0.39 ± 0.01 ^c	1.88 ± 0.02 ^c	5.52 ± 0.02 ^b
	<i>S. aureus</i>	nd ^a	nd ^a	nd ^a	nd ^a	0.08 ± 0.01 ^a
	Significance	*	*	*	*	*
Acetic acid	<i>E. faecalis</i>	nd	0.02 ± 0.01 ^b	0.13 ± 0.01 ^b	0.28 ± 0.01 ^b	0.40 ± 0.01 ^b
	Co-culture	nd	0.05 ± 0.01 ^c	0.13 ± 0.01 ^b	0.30 ± 0.01 ^b	0.40 ± 0.01 ^b
	<i>S. aureus</i>	nd	nd ^a	nd ^a	0.02 ± 0.01 ^a	0.03 ± 0.01 ^a
	Significance	ns	*	*	*	*

^{a,b,c} Different letters in the same column indicate difference at 95% level of significance; ns: not significant; nd: not detected (below detection limit).



710
 711 Figure 1. Venn diagrams depicting the number of genes and their respective tr
 712 response at 7 and 24 h of incubation at 30 °C in skimmed milk. A) Downregul
 713 co-culture ($LFC \leq -1.0$); B) Upregulated genes in co-culture ($LFC \geq +1.0$); C)
 714 transcripts not detected in single and co-culture (below detection limit); D)
 715 transcripts absent only in co-culture (below detection limit); E) Genes wit
 716 detected only in co-culture. LFC: \log_2 fold change.


717



718

719 Figure 2. Overview of total abundance of *S. aureus* transcripts assigned to COG functional
 720 categories in single (orange) and co-culture (yellow) in skimmed milk at 7h and 24h of
 721 incubation. RPKM: reads per kilobase of transcript per million reads mapped. J: translation,
 722 ribosomal structure and biogenesis; K: transcription; L: replication, recombination and repair;
 723 D: cell cycle control, cell division, chromosome partitioning; V: defense mechanisms; T:
 724 signal transduction mechanisms; M: cell wall, membrane, envelope biogenesis; U:
 725 intracellular trafficking, secretion and vesicular transport; O: posttranslational modification,
 726 protein turnover, chaperones; C: energy production and conversion; G: carbohydrate
 727 transport and metabolism; E: amino acid transport and metabolism; F: nucleotide transport

728 and metabolism; H: coenzyme transport and metabolism; I: lipid transport and metabolism; P:
729 inorganic ion transport and metabolism; Q: secondary metabolites biosynthesis, transport and
730 catabolism; R: general function prediction only; S: function unknown; NOF: no orthologous
731 found.



732 Figure 3. Transcriptional response of *S. aureus* genes related to central metabolic pathways
 733 during growth with *E. faecalis* at 7 and 24 h of incubation. Genes involved in glucose and
 734 lactose transport, glycolysis, gluconeogenesis, TCA cycle, pentose phosphate pathway,
 735 acetate, lactate, ethanol and acetoin metabolism.

16 Table 3. Transcriptional response profile of virulence genes and transcriptional regulators of major relevance in *S. aureus* during growth in co-culture
 17 with *E. faecalis* at 7 and 24 h of incubation.

Function	ORF denomination	Gene product	Gene	RPKM values at			RPKM values at		
				7 h		LFC	24 h		LFC
				SC	CC		SC	CC	
Adherence and attachment to host tissues	ATCC29213_13_64	clumping factor A	<i>clfA</i>	1215.8	1733.0	0.5	1720.3	2469.0	0.5
	ATCC29213_46_136	clumping factor B	<i>clfB</i>	483.6	281.1	-0.8	238.3	934.4	2.0
	ATCC29213_46_5	fibronectin-binding protein FnbB	<i>fnbB</i>	31.9	254.8	3.0	406.6	529.4	0.4
	ATCC29213_46_6	fibronectin-binding protein FnbA	<i>fnbA</i>	88.5	0.0	NDCC	305.9	0.0	NDCC
	ATCC29213_21_1	extracellular matrix-binding protein Ebh	<i>ebh</i>	17.3	0.0	NDCC	58.3	134.2	1.2
	ATCC29213_21_45	elastin-binding protein EbpS	<i>ebpS</i>	503.9	1006.9	1.0	502.2	2092.0	2.1
	ATCC29213_14_90	extracellular adherence protein Eap	<i>eap</i>	105.9	0.0	NDCC	450.5	1407.5	1.6
	ATCC29213_10_37	serine-aspartate repeat-containing	<i>sdrC</i>	115.7	264.1	1.2	131.7	219.5	0.7

		protein C								
		serine-aspartate	rich							
	ATCC29213_10_38	fibrinogen/bone	sialoprotein-	<i>sdrD</i>	141.2	451.4	1.7	168.8	375.1	1.2
		binding protein D								
	ATCC29213_47_1	serine-aspartate	repeat-containing	<i>sdrE</i>	15.3	0.0	NDCC	28.5	305.8	3.4
		protein E								
	ATCC29213_11_1	serine-aspartate	repeat-containing	<i>sdrE</i>	31.3	0.0	NDCC	66.6	416.1	2.6
		protein E								
	ATCC29213_46_152	putative cell-wall-anchored protein		<i>sasF</i>	506.4	385.4	-0.4	243.5	961.0	2.0
		SasF (LPXAG motif)								
	ATCC29213_45_10	surface protein G		<i>sasG</i>	0.0	0.0	NDBC	165.1	589.6	1.8
	ATCC29213_46_1	surface protein G		<i>sasG</i>	24.5	0.0	NDCC	195.6	489.0	1.3
	ATCC29213_51_1	surface protein G		<i>sasG</i>	0.0	0.0	NDBC	124.4	194.4	0.6
	ATCC29213_63_1	surface protein G		<i>sasG</i>	51.0	407.9	3.0	142.4	508.5	1.8
Immune evasion and	ATCC29213_17_21	alpha-hemolysin		<i>hla</i>	95.9	766.4	3.0	2242.5	0.0	NDCC
host damaging	ATCC29213_44_61	phospholipase C beta-hemolysin		<i>hlb</i>	0.0	0.0	NDBC	0.0	0.0	NDBC

enzymes	ATCC29213_42_85	gamma-hemolysin component A	<i>hlgA</i>	0.0	0.0	NDBC	164.6	0.0	NDCC
	ATCC29213_42_86	gamma-hemolysin component C	<i>hlgC</i>	0.0	0.0	NDBC	283.9	0.0	NDCC
	ATCC29213_42_87	gamma-hemolysin component B	<i>hlgB</i>	94.1	0.0	NDCC	125.1	0.0	NDCC
	ATCC29213_44_72	RNAIII (delta-hemolysin)	<i>hld</i>	0.0	0.0	NDBC	3738.7	0.0	NDCC
	ATCC29213_4_76	staphylocoagulase	<i>coa</i>	23.3	372.0	4.0	142.2	0.0	NDCC
	ATCC29213_13_68	thermonuclease	<i>nuc</i>	0.0	0.0	NDBC	124.7	0.0	NDCC
	ATCC29213_39_45	secretory antigen SsaA	<i>ssaA</i>	183.9	1470.0	3.0	464.3	0.0	NDCC
	ATCC29213_46_71	staphylococcal secretory antigen SsaA	<i>ssaA</i>	179.9	0.0	NDCC	15.9	2389.3	7.2
	ATCC29213_2_3	immunoglobulin G binding protein A precursor	<i>spa</i>	5373.7	1631.0	-1.7	153.6	1807.3	3.6
	ATCC29213_42_84	immunoglobulin G binding protein Sbi precursor	<i>sbi</i>	35.1	0.0	NDCC	1361.8	466.3	-1.55
	ATCC29213_31_2	serine protease SplF	<i>splF</i>	32.0	0.0	NDCC	68.0	0.0	NDCC
	ATCC29213_31_3	serine protease SplC	<i>splC</i>	0.0	0.0	NDBC	0.0	0.0	NDBC
	ATCC29213_31_4	serine protease SplB	<i>splB</i>	0.0	0.0	NDBC	84.6	0.0	NDCC

		ATCC29213_31_5	serine protease SplA	<i>splA</i>	65.0	0.0	NDCC	51.8	0.0	NDCC
		ATCC29213_70_1	serine protease SplD	<i>splD</i>	0.0	0.0	NDCC	102.0	0.0	NDCC
		ATCC29213_70_2	serine protease SplF	<i>splF</i>	32.5	0.0	NDCC	51.8	0.0	NDCC
		ATCC29213_14_153	staphostatin B	<i>sspC</i>	0.0	0.0	NDBC	111.4	0.0	NDCC
		ATCC29213_14_154	staphopain B	<i>sspB</i>	0.0	0.0	NDBC	41.4	517.3	3.64
		ATCC29213_33_49	staphopain A	<i>sspP</i>	0.0	0.0	NDBC	21.0	523.9	4.64
		ATCC29213_33_50	staphostatin A	<i>sspA_2</i>	0.0	0.0	NDBC	0.0	0.0	NDBC
		ATCC29213_39_30	urease subunit gamma	<i>ureA</i>	0.0	0.0	NDBC	80.9	0.0	NDCC
		ATCC29213_39_31	urease subunit beta	<i>ureB</i>	0.0	0.0	NDBC	89.4	0.0	NDCC
		ATCC29213_39_32	urease subunit alpha	<i>ureC</i>	53.6	0.0	NDCC	106.9	356.2	1.74
		ATCC29213_39_33	urease accessory protein UreE	<i>ureE</i>	0.0	0.0	NDBC	0.0	0.0	NDBC
		ATCC29213_39_34	urease accessory protein UreF	<i>ureF</i>	33.4	0.0	NDCC	35.5	443.3	3.64
		ATCC29213_39_35	urease accessory protein UreG	<i>ureG</i>	112.3	1197.1	3.41	179.1	0.0	NDCC
		ATCC29213_39_36	urease accessory protein UreD	<i>ureD</i>	55.0	0.0	NDCC	43.8	0.0	NDCC
		ATCC29213_4_165	glycerol ester hydrolase	<i>geh</i>	66.5	354.2	2.41	512.3	0.0	NDCC
ESAT-6	secretion	ATCC29213_4_127	virulence factor EsxA	<i>esxA</i>	34761.	15051.7	-1.2	11009.7	2084.9	-2.40

system				4					
	ATCC29213_4_128	type VII secretion protein EsaA	<i>esaA</i>	75.9	0.0	NDCC	84.7	201.7	1.25
	ATCC29213_4_129	secretion protein EssA	<i>essA</i>	100.4	0.0	NDCC	266.8	0.0	NDCC
	ATCC29213_4_130	type VII secretion protein EsaB	<i>esaB</i>	0.0	0.0	NDBC	0.0	0.0	NDBC
	ATCC29213_4_131	type VII secretion protein EssB	<i>essB</i>	86.2	0.0	NDCC	82.4	0.0	NDCC
	ATCC29213_4_132	type VII secretion protein EssC	<i>essC</i>	207.2	165.6	-0.3	96.3	0.0	NDCC
	ATCC29213_4_133	protein EsaC	<i>esaC</i>	117.3	0.0	NDCC	280.5	0.0	NDCC
	ATCC29213_4_134	virulence factor EsxB	<i>esxB</i>	146.4	0.0	NDCC	77.8	0.0	NDCC
Superantigens	ATCC29213_17_24	staphylococcal exotoxin 1	<i>set1</i>	0.0	0.0	NDBC	0.0	0.0	NDBC
	ATCC29213_17_25	staphylococcal exotoxin 4	<i>set4</i>	31.7	0.0	NDCC	33.7	0.0	NDCC
	ATCC29213_17_26	staphylococcal exotoxin 3	<i>set3</i>	95.1	0.0	NDCC	33.7	0.0	NDCC
	ATCC29213_5_5	staphylococcal exotoxin 6	<i>set6</i>	0.0	0.0	NDBC	35.9	0.0	NDCC
	ATCC29213_5_6	staphylococcal exotoxin 7	<i>set7</i>	66.2	1057.6	4.0	70.3	0.0	NDCC
	ATCC29213_5_7	staphylococcal exotoxin 8	<i>set8</i>	0.0	0.0	NDBC	137.0	0.0	NDCC
	ATCC29213_6_2	staphylococcal exotoxin 9	<i>set9</i>	0.0	0.0	NDBC	139.2	347.9	1.3
	ATCC29213_6_3	staphylococcal exotoxin 10	<i>set10</i>	0.0	0.0	NDBC	86.8	0.0	NDCC

ATCC29213_6_4	staphylococcal exotoxin 11	<i>set11</i>	0.0	0.0	NDBC	140.7	0.0	NDCC
ATCC29213_6_5	staphylococcal exotoxin 12	<i>set12</i>	0.0	0.0	NDBC	35.0	0.0	NDCC
ATCC29213_6_6	staphylococcal exotoxin 13	<i>set13</i>	0.0	0.0	NDBC	35.0	0.0	NDCC
ATCC29213_6_7	staphylococcal exotoxin 14	<i>set14</i>	0.0	0.0	NDBC	107.3	0.0	NDCC
ATCC29213_7_3	staphylococcal exotoxin 15	<i>set15</i>	0.0	0.0	NDBC	107.3	0.0	NDCC
ATCC29213_31_17	enterotoxin type G	<i>seg</i>	177.8	0.0	NDCC	78.7	0.0	NDCC
ATCC29213_31_18	enterotoxin type N	<i>sem</i>	0.0	0.0	NDBC	16.2	0.0	NDCC
ATCC29213_31_19	pseudoenterotoxin 2, ent2	<i>ψ ent 2</i>	116.4	0.0	NDCC	30.9	0.0	NDCC
ATCC29213_31_20	pseudoenterotoxin 1, ent1	<i>ψ ent 1</i>	229.3	0.0	NDCC	0.0	0.0	NDBC
ATCC29213_31_21	enterotoxin type I	<i>sei</i>	126.3	0.0	NDCC	50.4	839.1	4.1
ATCC29213_31_22	enterotoxin type M	<i>sem</i>	64.0	0.0	NDCC	34.0	0.0	NDCC
ATCC29213_31_23	enterotoxin type O	<i>seo</i>	0.0	0.0	NDBC	32.0	0.0	NDCC
ATCC29213_34_23	enterotoxin type A	<i>sea</i>	416.4	0.0	NDCC	1027.5	790.3	-0.4

Global regulators

ATCC29213_44_73	accessory gene regulator protein AgrB	<i>agrB</i>	186.3	0.0	NDCC	435.7	0.0	NDCC
ATCC29213_44_74	accessory gene regulator	<i>agrD</i>	321.6	0.0	NDCC	1025.4	0.0	NDCC

	autoinducing peptide AgrD								
ATCC29213_44_75	accessory gene regulator sensor histidine kinase AgrC	<i>agrC</i>	53.8	572.9	3.4	514.3	0.0	NDCC	
ATCC29213_44_76	accessory gene regulator DNA- binding response regulator AgrA	<i>agrA</i>	321.1	0.0	NDCC	955.7	0.0	NDCC	
ATCC29213_12_25	histidine protein kinase SaeS	<i>saeS</i>	109.0	0.0	NDCC	2212.2	0.0	NDCC	
ATCC29213_12_26	two-component response regulator SaeR	<i>saeR</i>	335.1	0.0	NDCC	2796.5	890.5	-1.7	
ATCC29213_11_54	HTH-type transcriptional regulator SarA	<i>sarA</i>	2274.5	3930.6	0.8	4018.6	3266.7	-0.3	
ATCC29213_39_37	HTH-type transcriptional regulator SarR	<i>sarR</i>	861.3	4236.4	2.3	2394.5	0.0	NDCC	
ATCC29213_2_4	HTH-type transcriptional regulator SarS	<i>sarS</i>	305.7	0.0	NDCC	130.0	0.0	NDCC	
ATCC29213_42_51	HTH-type transcriptional regulator SarZ	<i>sarZ</i>	876.3	0.0	NDCC	1150.7	0.0	NDCC	

	ATCC29213_46_2	HTH-type transcriptional regulator SarT	<i>sarT</i>	0.0	0.0	NDCC	68.6	0.0	NDCC
	ATCC29213_46_3	HTH-type transcriptional regulator SarU	<i>sarU</i>	0.0	0.0	NDCC	65.8	0.0	NDCC
	ATCC29213_39_10	HTH-type transcriptional regulator SarV	<i>sarV</i>	394.1	0.0	NDCC	768.1	0.0	NDCC
	ATCC29213_39_39	HTH-type transcriptional regulator SarY	<i>sarY</i>	61.9	0.0	NDCC	164.5	0.0	NDCC
	ATCC29213_11_103	HTH-type transcriptional regulator SarX	<i>sarX</i>	0.0	0.0	NDCC	0.0	0.0	NDBC
	ATCC29213_12_6	HTH-type transcriptional regulator MgrA	<i>mgrA</i>	1608.9	0.0	NDCC	1737.7	0.0	NDCC
	ATCC29213_28_117	HTH-type transcriptional regulator repressor of toxin Rot	<i>rot</i>	1490.7	3666.0	1.30	1645.5	0.0	NDCC
Metabolite- responsive regulators	ATCC29213_18_43	GTP-sensing transcriptional pleiotropic repressor CodY	<i>codY</i>	237.9	0.0	NDCC	284.5	790.3	1.47

	ATCC29213_28_91	catabolite control protein A	<i>ccpA</i>	162.7	743.2	2.1913	667.2	1853.0	1.4737
	ATCC29213_11_107	carbon catabolite responsive regulator CcpE	<i>ccpE</i>	159.3	848.8	2.4137	296.3	705.4	1.2513
	ATCC29213_4_162	RpiR family transcriptional regulator	<i>rpiRA</i>	57.5	0.0	NDCC	30.5	381.8	3.6
	ATCC29213_4_38	RpiR family transcriptional regulator	<i>rpiRB</i>	26.2	0.0	NDCC	153.1	695.8	2.2
	ATCC29213_40_7	RpiR family transcriptional regulator	<i>rpiRC</i>	210.9	0.0	NDCC	686.6	0.0	NDCC
Sigma factors	ATCC29213_24_10	RNA polymerase sigma factor A	<i>sigA</i>	124.7	996.9	2.9986	430.9	1933.1	2.1656
	ATCC29213_36_4	RNA polymerase sigma factor B	<i>sigB</i>	238.9	1909.2	2.9986	587.2	793.3	0.4342
	ATCC29213_36_5	serine-protein kinase RsbW	<i>rsbW</i>	96.0	1534.5	3.9986	153.1	0.0	NDCC
	ATCC29213_36_6	Anti-sigma factor B antagonist RsbV	<i>rsbV</i>	70.5	0.0	NDCC	562.2	0.0	NDCC
	ATCC29213_36_7	Sigma factor B regulation protein RsbU	<i>rsbU</i>	91.9	0.0	NDCC	134.3	0.0	NDCC

ATCC29213_10_10	RNA polymerase sigma factor H	<i>sigH</i>	80.8	1291.8	15.984	193.3	0.0	NDCC
ATCC29213_28_129	RNA polymerase sigma factor S	<i>rpoS</i>	48.9	0.0	NDCC	0.0	1299.7	NDPC

18 **RPKM:** reads per kilobase of transcript per million reads mapped

19 **LFC:** log₂ fold change

20 **SC:** single culture

21 **CC:** co-culture

22 **NDBC:** not detected in both conditions (below detection limit)

23 **NDCC:** not detected in co-culture (below detection limit)

24 **NDPC:** not detected in pure culture (below detection limit)