

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Emerging roles of Fgf14 in behavioral control

### **This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1686134> since 2019-01-08T15:28:25Z

*Published version:*

DOI:10.1016/j.bbr.2018.08.034

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

<b>Sexual behavior</b>	<b>Wild type</b>	<b><i>Fgf14</i> -/-</b>	<b><i>P</i> value</b>
<b>Latency to mount (sec)</b>	393.5±73.36	854.3±45.71	<b>***0.0001</b>
<b>Time mounting (sec)</b>	6.36±2.34	0.45±0.45	<b>*0.020</b>
<b>Number of mounts (sec)</b>	8.14±2.44	0.21±0.21	<b>**0.003</b>
<b>Duration of each episode (sec)</b>	0.51±0.096	0.079±0.079	<b>**0.002</b>
<b>Social approach (sec)</b>	447.7±26.84	406.1±34.42	0.35
<b>Avoidance (sec)</b>	439.8±26.48	417.5±31.63	0.59
<b>Passive behavior (sec)</b>	4.61±1.72	75.26±14.64	<b>***0.0001</b>

**Table II.** Analysis of 15 minutes of sexual behavior (n = 14 for both genotypes). \* p < 0.05, \*\* p < 0.01, \*\*\* p < 0.001.