

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effectiveness of artificial floods for benthic community recovery after sediment flushing from a dam

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1687326> since 2019-01-30T14:35:29Z

Published version:

DOI:10.1007/s10661-019-7232-7

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Effectiveness of artificial floods for benthic community recovery after sediment flushing from**
2 **a dam**

3
4 Alberto Doretto^{1*}, Tiziano Bo², Francesca Bona¹, Mattia Apostolo¹, Davide Bonetto³, Stefano
5 Fenoglio⁴
6

7 ¹DBIOS, University of Torino, Via Accademia Albertina 13, I-10123, Torino, Italy

8 ²Naturastaff, Via Lunga 11, I-14040, Mongardino (AT), Italy

9 ³Settore Presidio del Territorio – Ufficio polizia locale faunistico ambientale, Provincia di Cuneo,
10 Corso Nizza 21, I-12100, Italy

11 ⁴DISIT, University of Piemonte Orientale, Viale Teresa Michel 25, I-15121, Alessandria, Italy
12
13

14 *Corresponding author. E-mail address: alberto.doretto@unito.it

15 **Abstract**

16

17 The number of dams is predicted to increase worldwide under the current global change scenario. A
18 major environmental problem associated with dams is the release of large quantities of fine sediment
19 downstream. Therefore, future studies in river conservation will largely be focused on the
20 management of sediments trapped by reservoirs. The aim of this study was to investigate the
21 downstream ecological impacts of sediment flushing from a dam and the effectiveness of artificial
22 flash floods as a recovery strategy. Artificial flash floods have often been employed to remove large
23 amounts of sediment from riverbeds, but their importance in improving the biological quality of lotic
24 environments is almost unknown. We carried out a series of quantitative macroinvertebrate samplings
25 over a 2-year period that started before sediment release and included the artificial flushing events.
26 We characterized the macroinvertebrate community in its structural and functional aspects, and tested
27 the performance of two biomonitoring indexes, comparing their diagnostic ability. Our results
28 demonstrated that sediment flushing significantly altered the structure and composition of benthic
29 communities for more than 1 year. Flash floods exacerbated the overall biological quality, but we
30 believe that this treatment was useful because, by removing large amounts of sediment, the biological
31 recovery process was accelerated. Finally, regarding the water quality assessment, we found that the
32 biomonitoring index for siltation, composed of a selection of taxonomical and functional metrics, was
33 more reliable than the generic one.

34

35

36 **Keywords:** macroinvertebrates, siltation, dam, multi-metric index, Alpine stream, restoration.

37

38 **Introduction**

39

40 Dam and reservoir management is likely to represent one of the biggest challenges in river
41 conservation over the next years. Dams are increasing both in number and in geographical extent
42 (Fox et al. 2016), but numerous aspects of their correct management remain unresolved. In some
43 countries there is a growing tendency to remove large dams (Gangloff 2013; Grant and Lewis 2015),
44 while the current and anticipated changes in hydrological regimes with climate change are causing
45 an increase in dam construction worldwide (O'Connor et al. 2015; Zarfl et al. 2015).

46 Dams can have a significant ecological impact on lotic ecosystems, disrupting natural flow regimes
47 and altering naturally evolved processes (Van Cappellen and Maavara 2016; Arheimer et al. 2017).
48 For example, dams inhibit natural dispersal pathways and movements of aquatic organisms, causing
49 population isolation and extirpation of migratory fauna (Olden 2016). Dams transform upstream lotic
50 habitats into lentic environments, with a strong impact on structural and functional composition of
51 aquatic coenoses (Gray and Ward 1982; Guareschi et al. 2014; Rothenberger et al. 2017), and
52 interrupt the natural downstream transport of sediments. Moreover, the quality of water released from
53 reservoirs is often significantly altered during reservoir residence (WCD 2000). Dams generally trap
54 all the bedload, i.e. gravel and coarse sand particles, and much of the suspended load, which has
55 detrimental effects on many natural processes, both in downstream and impounded reaches (Brandt
56 2000). Downstream sections are usually sediment starved causing substrate armoring and increased
57 bank erosion and riverbed incision (Kondolf 1997; Graf 2006), with many detrimental ecological
58 effects. By contrast, in the upstream impounded reach, sediment trapping in reservoirs dramatically
59 alters the morphological and ecological features of the systems, decreasing substrate particle size and
60 transforming the river into a lentic body (Tang et al. 2016; Rothenberger et al. 2017). Moreover,
61 sediment accumulation within impoundments reduces storage capacity of reservoirs, interfering with
62 the functional purpose of the dam and reducing the efficiency of associated hydroelectric power plants
63 (Annandale 2013).

64 Unfortunately, loss of storage due to sedimentation was rarely considered in the design of many dams,
65 resulting in a widespread problem of reservoir sediment management (Morris and Fan 1998; Owens
66 et al. 2015), with an estimated 28% of the global sediment flux being potentially trapped in reservoirs
67 (Vorosmarty et al. 2003). There are three main strategies for managing sedimentation in reservoirs.
68 The first - sediment yield reduction - aims to reduce sediment inflow to reservoirs by controlling
69 erosion in the upstream river network (Quiñonero-Rubio et al. 2016). The second - sediment routing
70 - a modern approach whereby inflow sediments are allowed to pass through or around the reservoir

71 (Sumi and Kantoush 2010; Auel and Boes 2011, Martín et al. 2017). Unfortunately, these approaches
72 are not suitable for many existing dams. The last, most common strategy is sediment removal by
73 eliminating accumulated sediment through excavation, dredging, or drawdown flushing (Espa et al.
74 2013; Kondolf et al. 2014).

75 Sediment flushing can restore or preserve reservoir storage but often has dramatic negative impacts
76 on freshwater ecosystems. In some cases, sediment flushing causes the release of an excessive
77 quantity of fine material, forming thick layers of sediments, which can settle in the river for years
78 before being transported downstream by the flow, affecting the diversity and abundance of the river
79 biota. The adverse effects associated with the unnatural deposition of fine sediment in the river bed
80 have been documented for different aquatic groups, including macroinvertebrates (Wood and
81 Armitage 1997; Mebane 2001; Bo et al. 2007; Bilotta and Brazier 2008; Jones et al. 2012), fish (Kemp
82 et al. 2011; Sear et al. 2016) and plants (Izagirre et al. 2009; Jones et al. 2014). To accelerate removal
83 of these deposits, dam managers can implement artificial flash floods, i.e. sudden releases of water
84 from the dam for washing away and moving the settled sediments downstream. Artificial floods have
85 been applied for achieving restoration goals, for example, to reproduce the natural flow regime in
86 heavily regulated streams affected by multiple dams in their basins (Robinson et al. 2003, 2004,
87 2012).

88 Given the complexity and unpredictability of the flushing operations, data on their impact and
89 effectiveness are scarce. Several operational strategies have been proposed and applied in the last
90 years for successfully desilting reservoirs; for example, it is recommended to perform flushing
91 operations during the period of maximum seasonal run-off to maintain the natural pattern of sediment
92 transport (White 2001; Kondolf et al. 2014). Moreover, flushing-related sedimentation can be
93 mitigated by providing additional water from nearby reservoirs, and stopping sediment release at
94 night to limit the chronic stress associated with these activities (Espa et al. 2013; Quadroni et al.
95 2016). However, the effects of sediment flushing on aquatic communities are often underestimated
96 due to the lack of pre-impact data and specific biomonitoring tools. We therefore aimed to analyze
97 the impact of sediment flushing from an Alpine river dam on downstream benthic macroinvertebrate
98 communities in terms of richness, density and structure; and to monitor the temporal recovery of these
99 coenoses after two artificial flash floods. Finally, we compared the diagnostic power of two
100 biomonitoring indexes based on macroinvertebrates, one generic and the other designed for detecting
101 siltation effects.

102

103 **Materials and methods**

104

105 *Study Area*

106 The study was carried out in the Stura di Demonte river (hereinafter referred to as Stura), which is
107 considered one of the least impacted rivers in the entire Italian Alpine area. In fact, the Stura hosts
108 populations of endangered fish species, such as marble trout (*Salmo marmoratus*), grayling
109 (*Thymallus thymallus*) and European bullhead (*Cottus gobio*). For this reason, the SCI (Sites of
110 Community Importance) Stura di Demonte (Code IT1160036) was established according to the EU
111 43/1992 Directive (“Habitat Directive”) to protect the most conserved river stretch, which includes
112 four municipalities between the villages of Demonte and Roccasparvera. The study area was located
113 downstream of the Roccasparvera Dam (44.340256°, 7.440539°; Fig. 1), which was built in 1957 and
114 subsequently acquired by the Italian power company, Enel Green Power, in March 1963.
115 Roccasparvera is a 13 m concrete overflow dam, with a slightly arched planimetric course and a total
116 capacity of $0.580 \times 10^6 \text{ m}^3$. The land use of the watershed is dominated by forests (48%), followed
117 by pastures (39%) and rocky areas (10.8%), while the urbanized and agricultural areas account for
118 only for 1.4% and 0.8%, respectively.

119

120 *Sedimentation and dam operations*

121 Due to a safety-related emergency, in January 2016, a sudden and rapid depletion in the volume of
122 water in the reservoir was necessary. On 7th January at 10:15 AM, the sluice gates on the dam were
123 opened, allowing the water to flush at $5.4 \text{ m}^3 \text{ s}^{-1}$; the discharge was progressively increased to 13.7
124 $\text{m}^3 \text{ s}^{-1}$ (at 02:45 PM), and was successively reduced until the reservoir was empty at the end of the
125 flushing event on 8th January 2016 at 02:00 AM ($4.5 \text{ m}^3 \text{ s}^{-1}$). During flushing, water turbidity,
126 suspended solids concentration (SSC), water temperature, dissolved oxygen (DO), pH, and electric
127 conductivity were monitored continually by Enel Green Power at the station ROC1 (44.338820°,
128 7.448782°), located 0.7 km downstream. This event produced a significant environmental impact,
129 with a massive layer of sediments being accumulated for almost 8 km downstream of the dam
130 (personal observations). These sediments filled the river bed of the Stura and numerous irrigation
131 canals, completely transforming their morphology. Enel Green Power and the local environmental
132 authority (Province of Cuneo) agreed on a recovery strategy aimed at accelerating the removal of
133 sediment from the impacted reach. Since such a large amount of sediment would have taken many
134 years to disperse with normal flows, sudden water releases (artificial flash floods) were implemented
135 on two occasions (18th and 25th May 2016; Table 1), which increased erosion and transport in the

136 downstream section. In both circumstances, they started at 10:30 AM and consisted of a progressive
137 increase of released water every 5 minutes: from $10 \text{ m}^3 \text{ s}^{-1}$ to $41 \text{ m}^3 \text{ s}^{-1}$. This latter condition was kept
138 constant until 01:00 PM and then the volume of sluiced water was progressively reduced.

139

140 *Sampling design*

141 Benthic macroinvertebrate samples were collected in ten surveys from December 2015 (i.e., pre-
142 impact) to December 2017 (Table 1). Ten quantitative samples were collected randomly in each
143 survey, using a Surber sampler (250 μm mesh size; 0.062 m^2 area). Organisms were counted in the
144 laboratory and identified to the family level using taxonomic keys (Campaioli et al. 1994; 1999).

145

146 *Statistical analysis*

147 Statistical analysis was performed using R (R Development Core Team 2017). Changes in the
148 composition and structure of benthic invertebrate communities throughout the study were visually
149 evaluated using Principal Coordinate Analysis (PCoA). A Bray-Curtis similarity matrix was used for
150 multivariate ordination of the samples, while a Permutational Analysis of Variance (PERMANOVA)
151 was applied to test significant effects of the “sampling session” on the composition and structure of
152 benthic invertebrate communities. These analyses were carried out with the functions “vegdist”,
153 “betadisper” and “adonis” of the R package *vegan* (Oksanen et al. 2015). Univariate analysis was
154 then performed to detect the specific impacts of the sediment release and the temporal recovery of
155 macroinvertebrate communities. Four different community metrics were considered, namely total
156 taxa richness, total density of macroinvertebrates (Ind m^{-2}), EPT (Ephemeroptera, Plecoptera and
157 Trichoptera) richness and EPT density. Each Surber sample was considered as a replicate, and prior
158 to performing regression models outliers were removed using the method of Zuur et al. (2010) for
159 data exploration. Generalized Additive Models (GAMs) were employed to test the non-linear
160 response of the selected community compared to time in days from the sediment release; for this
161 operation, negative values were assigned to the pre-impact data (9th December 2015). All the GAMs
162 were carried out using the “gam” function of the *mgcv* R package (Wood and Wood 2015) and
163 applying a Poisson distribution. A negative binomial distribution was then applied in case of
164 overdispersion.

165 Finally, our data were analyzed in the context of river quality assessment, comparing the performance
166 of two selected indexes, namely the STAR_ICMi and the multi-metric index proposed by Doretto et
167 al. (2018). The former was originally developed as a calibration tool in the context of the Water

168 Framework Directive (EU 60/2000 Directive), and is currently the official biomonitoring index
169 provided by Italian regulations to classify the ecological status of running waters. STAR_ICMi is a
170 multi-metric index composed of six community metrics, belonging to three categories (i.e. diversity,
171 abundance, and sensitivity/tolerance). The weighted values of these metrics are aggregated in the
172 STAR_ICMi index, and the final score is obtained by the ratio between the observed and the reference
173 value (for details see Bo et al. 2017). The latter instead represents a recently proposed multi-metric
174 index using benthic invertebrates designed to detect the impacts of river siltation. This index includes
175 two diversity metrics (i.e., total taxa and EPT richness) and a functional metric (i.e., the abundance
176 of taxa preferring shallow, fast-flowing water with coarse substrates: Ecological Group A *sensu*
177 Usseglio-Polatera et al. 2000). Compared to the STAR_ICMi, which is generic, the multi-metric
178 index is a stressor-specific tool designed to assess the effects of siltation (for further details see
179 Doretto et al. 2018). These indexes were calculated for each sampling occasion, pooling together the
180 ten macroinvertebrate samples collected on that date. The temporal trends of the selected indexes
181 were thereby compared to evaluate which method provided the best performance in relation to the
182 sediment release.

183

184 **Results**

185

186 *Sediment and water chemistry parameters*

187 The flushing operations dramatically altered the physical and chemical parameters of the water, with
188 the greatest changes occurring during the night between 7th and 8th January 2016 (Fig. 2). An
189 exponential increase in SSC was documented; the SSC varied from 0.00 before the flushing to 10 mg
190 L⁻¹ at the end of the operation (mean = 12.07 mg L⁻¹), with intermediate peaks. The highest level of
191 SSC occurred on 8th January from 00:00 AM to 10:00 AM; during this period the SSC was
192 consistently higher than 20 mg L⁻¹, reaching a peak of 31.05 mg L⁻¹ (Fig. 2a). A very similar trend
193 was also observed for the turbidity, as this parameter was highly correlated with the SSC. From the
194 onset to the end of the flushing operations, the water turbidity increased from 0.80 to 2160 NTU,
195 reaching a maximum value of 6209 NTU (mean = 2413 NTU). During the most intense phase of
196 flushing, the turbidity was higher than 4300 NTU (Fig. 2b). Water temperature varied from 3.80 to
197 5.28 °C (mean = 4.26°C) but a sharp decline was recorded between 7:15 PM and 09:15 PM, when
198 the water temperature dropped to 0.26 °C, but then abruptly increased again (Fig. 2c) possibly due to
199 a thermal stratification of the impounded reach. The DO concentration markedly reduced during the

200 monitored period from 12.50 mg L⁻¹ at the beginning of the operations to 5.01 mg L⁻¹ at 02:00 AM
201 (Fig. 2d). DO then slowly increased again, reaching a value of 7.09 mg L⁻¹ at the end of the operations
202 (mean = 8.90 mg L⁻¹). A similar temporal trend was also observed for the pH (Fig. 2e), which varied
203 from 8.20 to 7.14 (mean = 7.70). Around 01:00 AM, the pH dropped to 7.17 and then remained
204 relatively constant until the end of the flushing activity. Finally, marked oscillations were observed
205 in relation to the electric conductivity over the monitored period (Fig. 2f). This parameter varied from
206 447 μS cm⁻¹ at the beginning of the operations to 483 μS cm⁻¹ at the end (mean = 421 μS cm⁻¹).
207 However, the lowest and highest values were 225 and 502 μS cm⁻¹, respectively.

208

209 *Benthic invertebrates*

210 We collected 7,860 organisms belonging to 38 families from nine different systematic groups. Diptera
211 and Trichoptera were the orders with the highest number of families, nine and eight, respectively,
212 whereas Gastropoda, Crustacea and Triclada were each represented by only one family. The number
213 of taxa per sample ranged from 1 to 21 (mean = 9), while the total density of benthic invertebrates
214 ranged from 40 to 8440 Ind m⁻² (mean = 1638 Ind m⁻²). Mayflies - Heptageniidae and Baetidae - as
215 well as dipteran larvae of the families Chironomidae and Simuliidae were the most abundant taxa.

216 PCoA ordination of the samples depicted a clear temporal shift in the composition and structure of
217 benthic invertebrate communities (Fig. 3) and PERMANOVA results showed a significant effect of
218 the sampling occasion ($F_{9,86} = 9.9; P < 0.001$). Compared to the pre-impact situation (i.e. 2015.12.09),
219 the communities just after the sediment release were significantly different in terms of composition
220 and structure, and oriented at the right side of the plot. The macroinvertebrate communities slowly
221 approached the pre-impact community levels only after both of the flash floods (from 2016.07.22
222 onward), reaching full overlap with the pre-flushing community on the last sampling occasion (i.e.
223 2017.12.12, nearly 2 years later).

224 Results of the multivariate analysis were confirmed and further strengthened by those of the GAMs
225 (Table 2). The sediment release and consequent flash floods significantly reduced the total taxa
226 richness, which showed a unimodal pattern with a negative peak around 150 days after the sediment
227 release (Fig. 4a). Taxa richness then progressively recovered over time, reaching values comparable
228 to pre-impact levels after 600 days. The combined effects of sedimentation and flash floods also
229 resulted in a similar temporal pattern for the EPT richness (Fig. 4b). Prior to impact, the average
230 number of EPT families was ten, which subsequently dropped to three after the sedimentation event
231 and to one after the flash floods. Again, the results of the model showed a progressive increase in the

232 EPT richness over time, with an almost complete recovery after 600 days. Conversely, GAMs
233 illustrated very fluctuating patterns in the density of benthic invertebrates. Both the total (Fig. 4c) and
234 the EPT density (Fig. 4d) showed a dramatic depletion after the sediment release as well as the flash
235 floods. However, compared to the taxa and EPT richness, both these two events were followed by
236 peaks in density. A less pronounced decrease was also observed after 400 days, and then both the
237 total and EPT density increased again to nearly pre-impact values. It is interesting to note that these
238 marked temporal fluctuations were mainly driven by the extremely high abundance of Baetidae (Fig.
239 5a), Chironomidae (Fig. 5b) and Simuliidae (Fig. 5c) specimens during the first post-release sampling
240 occasions (Table 2). Thus, the specific response of these few taxa accounted for the observed results
241 for the total and EPT density.

242

243 *Biomonitoring indexes*

244 The two biomonitoring indexes for assessing the impact of the sediment release from the
245 Roccapervera dam gave different responses (Fig. 6). The STAR_ICMi (Fig. 6a) on the pre-impact
246 sampling occasion was 1.1 and then decreased as a consequence of the sedimentation event and flash
247 floods. However, these changes only corresponded to a slight deterioration in the water quality
248 classification, which shifted from High to Good and only dropped to Scarce on the sampling occasion
249 2016.06.06 (i.e., after the second flash flood, STAR_ICMi = 0.25). After the artificial floods, the
250 index increased, ranging from 0.80 to 1.15, and it reached the High quality class on the last two
251 sampling occasions (i.e. after 475 days). Overall, the STAR_ICMi detected a change in the water
252 quality class, but this only resulted in a shift from High to Good for most of the recovery period, with
253 the exception of the sampling occasion 2016.06.06. Conversely, the temporal variation of the multi-
254 metric index for siltation proposed by Doretto et al. (2018) depicted a different temporal pattern (Fig.
255 6b). From the pre-impact value of 0.83, this index dropped markedly after the sediment release (0.33)
256 and the flash floods, achieving a minimum value of 0 on the sampling occasion 2016.06.06. The index
257 then increased over time, reaching full recovery on the last sample date. Compared to the
258 STAR_ICMi, this stressor-specific index showed a more pronounced variation between the pre-
259 impact value and that recorded just after the sediment release. Moreover, the temporal pattern of this
260 index was more similar to the community metrics, especially taxa richness.

261

262 **Discussion**

263 Although several studies have documented the effects of sediment releases from dams, long-term
264 series encompassing pre-impact data and multi-year sampling campaigns, like this study, are rare.
265 Usually, flushing operations are programmed over long periods of time, and carried out by alternating
266 sediment and water pulses between the day-time and night-time, or by providing volumes of diluting
267 water from adjacent reservoirs (Crosa et al. 2010; Espa et al. 2013; 2015). These practices allow
268 minimization of the ecological impacts of excessive releases of sediments and facilitate the recovery
269 of aquatic communities.

270 Unfortunately, adopting these approaches is not always possible according to the specific features of
271 the dam and the magnitude of the related impacts. The sediment flushing from the Roccasparvera
272 dam, documented in this study, was concentrated in a short period (from 7th January at 10:15 AM to
273 8th January at 02:00 AM). In just a few hours, drastic alterations of all the main physical and chemical
274 attributes of water were detected; for example, the SSC and turbidity were in the range of three orders
275 of magnitude greater than before the sediment release. At the same time, a dramatic reduction in the
276 concentration of DO and pH was observed. Similar changes have been documented by other authors
277 in relation to flushing events (Gray and Ward 1982; Peter et al. 2014; Khakzad and Elfimov 2015,
278 Espa et al. 2016) and have led to a strong impact on benthic invertebrate assemblages. We found a
279 temporal shift in the composition of communities; pre-impact macroinvertebrate communities were
280 mainly composed of EPT taxa, while after the sediment release and flash floods, the number of taxa
281 belonging to these orders was significantly reduced. Our results corroborate other studies carried out
282 in the Alpine area (Espa et al. 2013; Peter et al. 2014; Martín et al. 2017). More profound effects were
283 evident when macroinvertebrate density was considered, with a collapse of the total and EPT density
284 due to the sediment release, which was exacerbated with the flash floods. Similar results were also
285 observed by other authors (Doeg and Khoen 1994; Quadroni et al. 2016); for example, Crosa et al.
286 (2010) studied the ecological consequences of a reservoir flushing in an Italian alpine stream. They
287 found the density of invertebrates dropped, on average, from 2000 to 100 Ind m⁻². In our study, 2
288 years were necessary to achieve full recovery of the benthic communities to their pre-impact taxa
289 richness and density of macroinvertebrates. However, taxa richness recovered slowly and
290 progressively, while there was a fluctuating response of density. The extremely high abundance and
291 dominance of few generalist taxa, such as Baetidae, Chiromomidae and Simuliidae, may account for
292 the temporal pattern observed for density. Several authors have demonstrated that these families are
293 early colonists and tend to be dominant during the first phases of post-disturbance recolonization,
294 including experimental floods from reservoirs (Otermin et al. 2002; Robinson et al. 2004). Compared
295 to our results, a faster post-flushing recovery of macroinvertebrate communities has been documented
296 in other studies (Crosa et al. 2010; Espa et al. 2013; 2016). This delayed recovery pattern could be

297 explained by the fact that the flash floods were performed 5 months after the sedimentation event.
298 However, we postulate that, in the long-term, these artificial treatments contribute to restoring the
299 natural habitat for benthic invertebrates, facilitating the recovery of communities.

300 Another possible factor that could account for the full community recovery is the overall pristine
301 conditions of the investigated stream, which was reflected by the excellent pre-impact ecological
302 status of the sampled reach. As expected, the sedimentation process produced a deterioration of the
303 water quality, but this mainly consisted in a shift from High to Good, even when the river bottom was
304 buried by a large amount of fine sediment (personal observation) and the density and richness of
305 benthic invertebrates were highly reduced. The water quality further dropped to Scarce as an
306 immediate consequence of the flash floods. Although these operations were successful in removing
307 the fine sediment and promoting long-term recovery, they initially induced stressful conditions for
308 benthic invertebrates, dislodging them from the substrate. This may account for the low water quality
309 on that occasion, after which the water quality quickly re-established to Good and then High. In
310 particular, despite the temporary worsening of the biological quality after the flash floods, we believe
311 that this treatment had a considerable importance overall, because large amounts of sediment were
312 removed, which would have taken much longer to disperse naturally, thus the biological recovery
313 process was accelerated. The limited performance of STAR_ICMi was not completely unexpected
314 since it is a generic multi-metric index, as confirmed by other authors who obtained similar results
315 when assessing the impacts of sedimentation from dams (Espa et al. 2015; 2016). On the contrary,
316 the performance of the index for siltation recently proposed by Doretto et al. (2018) offered
317 interesting results, although no ecological classification has been provided yet. Nevertheless, when
318 looking at the temporal trend of the index, it was more reliable than the STAR_ICMi in detecting the
319 effects of disturbance in relation to both the sedimentation release and the flash floods. Quadroni et
320 al. (2016) evaluated the consequences of sediment flushing in an Alpine stream and stated that the
321 STAR_ICMi can underestimate impacts, resulting in an imprecise water quality assessment. This is
322 the first time that our stressor-specific index has been tested using an independent dataset. Overall,
323 its performance offered promising applications and confirmed the useful role of macroinvertebrates
324 as bioindicators for running waters (Rosenberg and Resh 1993; Buss et al. 2015). Moreover, the
325 combination of taxonomical and functional metrics considered here appears very successful for
326 detecting the specific effects of siltation, and this supports the adoption of functional groups for river
327 biomonitoring (Merritt et al. 2015).

328 While several solutions have been proposed in relation to the timing of the sluiced water and the
329 amount of flushed fine sediment, such as thresholds in the SSC and discharge, the development of
330 stressor-specific methods based on the biota response still remains unexplored (Crosa et al. 2010;

331 Quadroni et al. 2016). Our report therefore offers the application of a diagnostic biomonitoring tool
332 designed to evaluate the specific impacts associated with siltation. Since the number of reservoirs and
333 dams is predicted to increase in the near future (Fox et al. 2016; Van Cappellen and Maavara 2016),
334 developing science-based biomonitoring and management tools is an important challenge for river
335 ecologists and managers.

336

337 **Acknowledgements**

338 The authors wish to thank AECOM and Enel Green Power for their collaboration and sharing of data
339 about the sedimentation event. We are very grateful to Elena Piano and Radhika Srinivasan for their
340 assistance during data analysis and linguistic revision respectively. We would also like to
341 acknowledge the Province of Cuneo (Settore Presidio del Territorio – Ufficio polizia locale faunistico
342 ambientale) for their useful assistance. This work was supported by “Italian Mountain Lab”.

343

344 **References**

345 Annandale, G. (2013). Quenching the thirst, sustainable water supply and climate change. North
346 Charleston, SC: CreateSpace Independent Publishing Platform.

347 Arheimer, B., Donnelly, C., & Lindström, G. (2017). Regulation of snow-fed rivers affects flow
348 regimes more than climate change. *Nature communications*, 8(1), 62.

349 Auel, C., & Boes, R. M. (2011). Sediment bypass tunnel design–review and outlook. In A. J.

350 Bilotta, G. S., & Brazier, R. E. (2008). Understanding the influence of suspended solids on water
351 quality and aquatic biota. *Water Research* 42(12), 2849–2861.

352 Bo, T., Fenoglio, S., Malacarne, G., Pessino, M., & Sgariboldi, F. (2007). Effects of clogging on
353 stream macroinvertebrates: an experimental approach. *Limnologica* 37(2), 186–192.

354 Bo, T., Doretto, A., Laini, A., Bona, F., & Fenoglio, S. (2017). Biomonitoring with macroinvertebrate
355 communities in Italy: what happened to our past and what's the future? *Journal of Limnology*, 76(s1),
356 21–28.

357 Brandt, S. A. (2000). Classification of geomorphological effects downstream of dams. *Catena*, 40(4),
358 375–401.

359 Buss, D. F., Carlisle, D. M., Chon, T. S., Culp, J., Harding, J. S., Keizer-Vlek, H. E., Robinson, W.
360 A., Strachan, S., Thirion, C., & Hughes, R. M. (2015). Stream biomonitoring using
361 macroinvertebrates around the globe: a comparison of large-scale programs. *Environmental*
362 *Monitoring and Assessment* 187(1), 4132.

363 Campaioli, S., Ghetti, P. F., Minelli, A., & Ruffo, S. (1994). Manuale per il riconoscimento dei
364 macroinvertebrati delle acque dolci italiane (Vol. I). Trento, IT: Provincia Autonoma di Trento.

365 Campaioli, S., Ghetti, P. F., Minelli, A., & Ruffo, S. (1999). Manuale per il riconoscimento dei
366 macroinvertebrati delle acque dolci italiane (Vol. II). Trento, IT: Provincia Autonoma di Trento.

367 Crosa, G., Castelli, E., Gentili, G., & Espa, P. (2010). Effects of suspended sediments from reservoir
368 flushing on fish and macroinvertebrates in an alpine stream. *Aquatic Sciences*, 72(1), 85–95.

369 Doretto, A., Piano, E., Bona, F., & Fenoglio, S. (2018). How to assess the impact of fine sediments
370 on the macroinvertebrate communities of alpine streams? A selection of the best metrics. *Ecological*
371 *Indicators*, 84, 60–69.

372 Espa, P., Castelli, E., Crosa, G., & Gentili, G. (2013). Environmental effects of storage preservation
373 practices: controlled flushing of fine sediment from a small hydropower reservoir. *Environmental*
374 *management*, 52(1), 261–276.

375 Espa, P., Crosa, G., Gentili, G., Quadroni, S., & Petts, G. (2015). Downstream ecological impacts of
376 controlled sediment flushing in an Alpine valley river: a case study. *River Research and Applications*,
377 31(8), 931–942.

378 Espa, P., Brignoli, M. L., Crosa, G., Gentili, G., & Quadroni, S. (2016). Controlled sediment flushing
379 at the Cancano Reservoir (Italian Alps): management of the operation and downstream environmental
380 impact. *Journal of Environmental Management*, 182, 1–12.

381 Fox, G. A., Sheshukov, A., Cruse, R., Kolar, R. L., Guertault, L., Gesch, K. R., & Dutnell, R. C.
382 (2016). Reservoir sedimentation and upstream sediment sources: Perspectives and future research
383 needs on streambank and gully erosion. *Environmental Management*, 57(5), 945-955.

384 Gangloff, M. M. (2013). Taxonomic and ecological tradeoffs associated with small dam removals.
385 *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23(4), 475–480.

386 Graf, W. L. (2006). Downstream hydrologic and geomorphic effects of large dams on American
387 rivers. *Geomorphology*, 79(3), 336–360.

388 Grant, G. E., & Lewis, S. L. (2015). The remains of the dam: What have we learned from 15 years of
389 US dam removals? In Lollino G., Marconi A., Locat J., Huang Y., Canals Artigas M. (Ed.)
390 Engineering Geology for Society and Territory-Volume 3 (pp. 31–35). Cham: Springer.

391 Gray, L. J., & Ward, J. V. (1982). Effects of sediment releases from a reservoir on stream
392 macroinvertebrates. *Hydrobiologia*, 96(2), 177-184.

393 Guareschi S., Laini A., Racchetti E., Bo T., Fenoglio S., & Bartoli M. (2014). How
394 hydromorphological constraints and regulated flows govern macroinvertebrate communities along an
395 entire lowland river. *Ecohydrology*, 7, 366–377.

396 Izagirre, O., Serra, A., Guasch, H., & Elosegi, A. (2009). Effects of sediment deposition on periphytic
397 biomass, photosynthetic activity and algal community structure. *Science of the Total Environment*,
398 407(21), 5694–5700.

399 Jones, J. I., Murphy, J. F., Collins, A. L., Sear, D. A., Naden, P. S., & Armitage, P. D. (2012). The
400 impact of fine sediment on macroinvertebrates. *River Research and Applications*, 28(8), 1055– 1071.

401 Jones, J. I., Duerdoth, C. P., Collins, A. L., Naden, P. S., & Sear, D. A. (2014). Interactions between
402 diatoms and fine sediment. *Hydrological Processes*, 28(3), 1226–1237.

403 Kemp, P., Sear, D., Collins, A., Naden, P., & Jones, I. (2011). The impacts of fine sediment on
404 riverine fish. *Hydrological Processes*, 25(11), 1800–1821.

405 Khakzad, H., & Elfimov, V. I. (2015). A Review of Environmental Characteristics and Effects of Dez
406 Dam Flushing Operation on Downstream. *Environmental Practice*, 17(3), 211–232.

407 Kondolf, G. M. (1997). Hungry water: effects of dams and gravel mining on river channels.
408 *Environmental Management*, 21(4), 533–551.

409 Kondolf, G. M., Gao, Y., Annandale, G. W., Morris, G. L., Jiang, E., Zhang, J., Carling, P., Fu., K.,
410 Guo, Q., Hotchkiss, R., Peteuil, C., Sumi, T., Wang, H. W., Wang, Z., Wei, Z., Wu, B., & Yang, C.
411 T. (2014). Sustainable sediment management in reservoirs and regulated rivers: Experiences from
412 five continents. *Earth's Future*, 2(5), 256–280.

413 Martín, E. J., Doering, M., & Robinson, C. T. (2017). Ecological Assessment of a Sediment By-pass
414 Tunnel on a Receiving Stream in Switzerland. *River Research and Applications*, 33(6), 925– 936.

415 Mebane, C. A., 2001. Testing bioassessment metrics: macroinvertebrate, sculpin, and salmonid
416 responses to stream habitat, sediment, and metals. *Environmental Monitoring and Assessment*, 67(3),
417 293–322.

418 Merritt, R. W., Fenoglio, S., & Cummins, K. W. (2016). Promoting a functional macroinvertebrate
419 approach in the biomonitoring of Italian lotic systems. *Journal of Limnology*, 76(s1), 5–8.

420 Morris, G. L., & Fan, J. (1998). Reservoir sedimentation handbook: design and management of dams,
421 reservoirs, and watersheds for sustainable use. New York: McGraw Hill Professional.

422 Olden, J. D. (2016). Challenges and opportunities for fish conservation in dam-impacted waters.
423 Cambridge: Cambridge University Press.

424 O'Connor, J. E., Duda, J. J., & Grant, G. E. (2015). 1000 dams down and counting. *Science*,
425 348(6234), 496–497.

426 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R.,
427 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, & E., Wagner, H. (2015).
428 Vegan: Community Ecology Package. *R Package Version 2.2-1*.

429 Otermin, A., Basaguren, A., & Pozo, J. (2002). Re-colonization by the macroinvertebrate community
430 after a drought period in a first-order stream (Aguera Basin, Northern Spain). *Limnetica*, 21, 117–
431 128.

432 Owens, P. N., Batalla, R. J., Collins, A. J., Gomez, B., Hicks, D. M., Horowitz, A. J., Kondolf, G.
433 M., Marden, M., Page, M. J., Peacock, D. H., Peticrew, E. L., Salomons, W., & Trustrum, N. A.
434 (2005). Fine-grained sediment in river systems: environmental significance and management issues.
435 *River Research and Applications*, 21(7), 693–717.

436 Peter, D. H., Castella, E., & Slaveykova, V. I. (2014). Effects of a reservoir flushing on trace metal
437 partitioning, speciation and benthic invertebrates in the floodplain. *Environmental Science: Processes
438 & Impacts*, 16(12), 2692–2702.

439 Quadroni, S., Brignoli, M. L., Crosa, G., Gentili, G., Salmaso, F., Zaccara, S., & Espa, P. (2016).
440 Effects of sediment flushing from a small Alpine reservoir on downstream aquatic fauna.
441 *Ecohydrology*, 9(7), 1276–1288.

442 Quiñonero_Rubio, J. M., Nadeu, E., Boix_Fayos, C., & Vente, J. (2016). Evaluation of the
443 Effectiveness of Forest Restoration and Check_Dams to Reduce Catchment Sediment Yield. *Land
444 Degradation & Development*, 27(4), 1018–1031.

445 R Development Core Team, 2017. R: a language and environment for statistical computing. R
446 Foundation for Statistical Computing. Vienna, Austria.

447 Robinson, C. T., Uehlinger, U., & Monaghan, M. T. (2003). Effects of a multi-year experimental
448 flood regime on macroinvertebrates downstream of a reservoir. *Aquatic Sciences*, 65(3), 210–222.

449 Robinson, C. T., Uehlinger, U. R. S., & Monaghan, M. T. (2004). Stream ecosystem response to
450 multiple experimental floods from a reservoir. *River Research and Applications*, 20(4), 359–377.

451 Robinson, C. T. (2012). Long-term changes in community assembly, resistance, and resilience
452 following experimental floods. *Ecological Applications*, 22(7), 1949–1961.

453 Rosenberg, D. M., & Resh, V. H. (1993). *Freshwater Biomonitoring and Benthic Macroinvertebrates*.
454 Boston: Kluwer Academic Publishers.

455 Rothenberger, M. B., Hoyt, V., Germanoski, D., Conlon, M., Wilson, J., & Hitchings, J. (2017). A
456 risk assessment study of water quality, biota, and legacy sediment prior to small dam removal in a
457 tributary to the Delaware River. *Environmental Monitoring and Assessment*, 189(7), 344.

458 Sear, D. A., Jones, J. I., Collins, A. L., Hulin, A., Burke, N., Bateman, S., Pattison, I., & Naden, P. S.
459 (2016). Does fine sediment source as well as quantity affect salmonid embryo mortality and
460 development? *Science of the Total Environment*, 541, 957–968.

461 Sumi, T., & Kantoush, S. A. (2010). Integrated management of reservoir sediment routing by
462 flushing, replenishing, and bypassing sediments in Japanese river basins. *Proceedings of the 8th*
463 *International Symposium on Ecohydraulics, Seoul, Korea*, 831–838.

464 Tang, Q., Bao, Y., He, X., Fu, B., Collins, A. L., & Zhang, X. (2016). Flow regulation manipulates
465 contemporary seasonal sedimentary dynamics in the reservoir fluctuation zone of the Three Gorges
466 Reservoir, China. *Science of the Total Environment*, 548, 410–420.

467 Usseglio-Polatera, P., Bournaud, M., Richoux, P., & Tachet, H. (2000). Biological and ecological
468 traits of benthic freshwater macroinvertebrates: relationships and definition of groups with similar
469 traits. *Freshwater Biology*, 43(2), 175–205.

470 Van Cappellen, P., & Maavara, T. (2016). Rivers in the Anthropocene: global scale modifications of
471 riverine nutrient fluxes by damming. *Ecohydrology & Hydrobiology*, 16(2), 106–111.

472 Vorosmarty, C. J., Meybeck, B., Fekete, K., Sharma, P., Green, P., & Syvitski, J. P. M. (2003),
473 Anthropogenic sediment retention: major global impact from registered river impoundments, *Global*
474 *and Planetary Change*, 39, 169–190.

475 WCD (2000). *World Commission on Dams - Dams and Development: A New Framework for*
476 *Decision-making: the Report of the World Commission on Dams*. London: Earthscan.

- 477 White, R. (2001). Evacuation of sediments from reservoirs. London: Thomas Telford Ltd.
- 478 Wood, P. J., & Armitage, P. D. (1997). Biological effects of fine sediment in the lotic environment.
479 *Environmental Management*, 21(2), 203–217.
- 480 Wood, S., & Wood, M. S. (2015). Package ‘mgcv’. *R package version*, 1–7.
- 481 Zarfl, C., Lumsdon, A. E., Berlekamp, J., Tydecks, L., & Tockner, K. (2015). A global boom in
482 hydropower dam construction. *Aquatic Sciences*, 77(1), 161–170.
- 483 Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common
484 statistical problems. *Methods in Ecology and Evolution*, 1(1), 3–14.
- 485

486 **Tables**

487

488 **Table 1** Sampling scheme: Date = date, Label = labels used in the statistical analysis
489 (year.month.day), Activity = data and information associated to each date

Date	Label	Activity
9 th December 2015	2015.12.09	Pre-impact data
7 th - 8 th January 2016	-	Sediment flushing
23 rd March 2016	2016.03.23	Benthic invertebrates sampling
17 th May 2016	2017.05.17	Benthic invertebrates sampling
18 th May 2016	-	First flash flood event
24 th May 2016	2016.05.24	Benthic invertebrates sampling
25 th May 2016	-	Second flash flood event
06 th June 2016	2016.06.06	Benthic invertebrates sampling
22 nd July 2016	2016.07.22	Benthic invertebrates sampling
09 th December 2016	2016.12.09	Benthic invertebrates sampling
15 th March 2017	2017.03.15	Benthic invertebrates sampling
27 th April 2017	2017.04.27	Benthic invertebrates sampling
12 th December 2017	2017.12.12	Benthic invertebrates sampling

490

491

492 **Table 2** Statistics of the Generalized Additive Models: Est = estimate, SE = standard error, z = z-
 493 value, P = p-value, X^2 = Chi-square

Metric	Intercept				Smooth	
	Est	SE	z	P	X^2	P
Taxa richness	2.124	0.037	57.080	< 0.001	146.900	< 0.001
Total density	6.721	0.047	143.900	< 0.001	698.100	< 0.001
EPT richness	1.426	0.054	26.530	< 0.001	122.900	< 0.001
EPT density	5.781	0.047	121.900	< 0.001	893.600	< 0.001
Baetidae density	4.118	0.054	76.180	< 0.001	821	< 0.001
Chironomidae density	4.980	0.048	102.900	< 0.001	915	< 0.001
Simuliidae density	3.667	0.053	68.790	< 0.001	1173	< 0.001

494

495

496 **Figure captions**

497

498 **Fig. 1** Area of study: the Stura di Demonte watershed and the investigated stream reach

499

500 **Fig. 2** Values of: **a** concentration of suspended solids (SSC), **b** turbidity, **c** water temperature, **d**
501 concentration of dissolved oxygen (DO), **e** pH and **f** conductivity measured at the ROC1 station
502 during the sedimentation release from the Roccasparvera dam

503

504 **Fig. 3** Principal Coordinate Analysis (PCoA) ordination plots based on the dissimilarity matrices
505 (Bray-Curtis). Symbols represent the benthic community samples collected in each sampling
506 occasion (year.month.day). The grey lines link each sample with its corresponding centroid (black
507 dots)

508

509 **Fig. 4** Representation of the GAM smoother applied to time interacting with the community metrics:
510 **a** total taxa richness, **b** EPT richness, **c** total density and **d** EPT density. The black line represents the
511 smoothed function (i.e. $s(\text{Time}, \text{degrees of freedom})$), while the grey area indicates the 95%
512 confidence interval. Black vertical line indicates the sediment flushing, while the dotted vertical lines
513 represent the two artificial floods. Grey ticks on the x-axis represent the sampling occasions

514

515 **Fig. 5** Representation of the GAM smoother applied to time interacting with the density of: **a**
516 Baetidae, **b** Chironomidae and **c** Simuliidae. The black line represents the smoothed function (i.e.
517 $s(\text{Time}, \text{degrees of freedom})$), while the grey area indicates the 95% confidence interval. Black
518 vertical line indicates the sediment flushing, while the dotted vertical lines represent the two artificial
519 floods. Grey ticks on the x-axis represent the sampling occasions

520

521 **Fig. 6** Bars indicate the values of **a** the STAR_ICM index and **b** the sedimentation index (Doretto et
522 al. 2018) calculated for each sampling occasion (year.month.day)

523