

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Emerging roles of Fgf14 in behavioral control

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1686134> since 2019-01-08T15:28:25Z

Published version:

DOI:10.1016/j.bbr.2018.08.034

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Resident intruder parameters	Wild type	<i>Fgf14</i> ^{-/-}	<i>P</i> value
Time threatening (sec)	84.93±21.4	4.08±2.64	***0.0004
Number of threat episodes	17.93±4.39	1.0±0.65	***0.0003
Mean time of each threat episode (sec)	4.40±0.68	1.45±1.02	*0.026
Latency to first attack (sec)	630.1±82.4	769.3±71.4	0.21
Time attacking (sec)	19.05±8.8	2.51±1.98	*0.045
Number of attack episodes	7.0±3.03	1.8±1.29	0.066
Mean time of each attack episode (sec)	1.32±0.41	0.20±0.12	*0.0095
Social behavior (sec)	78.25±5.05	445.2±38.81	***<0.0001

Table III. Analysis of 15 minutes of resident intruder test (n = 14 for wild type and n = 16 for *Fgf14*^{-/-} mice.* p < 0.05, ** p < 0.01, *** p < 0.001.