



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Emerging roles of Fgf14 in behavioral control

inis is the author's manuscript	
Original Citation:	
Availability:	
This version is available http://hdl.handle.net/2318/1686134	since 2019-01-08T15:28:25Z
Published version:	
DOI:10.1016/j.bbr.2018.08.034	
Terms of use:	
Open Access	
Anyone can freely access the full text of works made available as under a Creative Commons license can be used according to the tof all other works requires consent of the right holder (author or p protection by the applicable law.	terms and conditions of said license. Use

(Article begins on next page)

Resident intruder parameters	Wild type	Fgf14 -/-	P value
Time threatening (sec)	84.93±21.4	4.08±2.64	***0.0004
Number of threat episodes	17.93±4.39	1.0 ± 0.65	***0.0003
Mean time of each threat episode (sec)	4.40±0.68	1.45±1.02	*0.026
Latency to first attack (sec)	630.1±82.4	769.3±71.4	0.21
Time attacking (sec)	19.05±8.8	2.51±1.98	*0.045
Number of attack episodes	7.0 ± 3.03	1.8±1.29	0.066
Mean time of each attack episode (sec)	1.32±0.41	0.20±0.12	*0.0095
Social behavior (sec)	78.25±5.05	445.2±38.81	***<0.0001

Table III. Analysis of 15 minutes of resident intruder test (n = 14 for wild type and n = 16 for $Fgf14^{-/-}$ mice.* p < 0.05, ** p < 0.01, *** p < 0.001.