

Fatty acid profile, meat quality and flavour acceptability of beef from double-muscling Piemontese young bulls fed ground flaxseed

Manuela Renna, Alberto Brugiapaglia, Emanuela Zanardi, Gianluigi Destefanis, Aldo Prandini, Maurizio Moschini, Samantha Sigolo & Carola Lussiana

To cite this article: Manuela Renna, Alberto Brugiapaglia, Emanuela Zanardi, Gianluigi Destefanis, Aldo Prandini, Maurizio Moschini, Samantha Sigolo & Carola Lussiana (2019): Fatty acid profile, meat quality and flavour acceptability of beef from double-muscling Piemontese young bulls fed ground flaxseed, Italian Journal of Animal Science, DOI: [10.1080/1828051X.2018.1530958](https://doi.org/10.1080/1828051X.2018.1530958)

To link to this article: <https://doi.org/10.1080/1828051X.2018.1530958>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 01 Feb 2019.



Submit your article to this journal [↗](#)



Article views: 16



View Crossmark data [↗](#)

Fatty acid profile, meat quality and flavour acceptability of beef from double-muscled Piemontese young bulls fed ground flaxseed

Manuela Renna^a , Alberto Brugiapaglia^a , Emanuela Zanardi^b , Gianluigi Destefanis^{a*}, Aldo Prandini^c , Maurizio Moschini^c , Samantha Sigolo^c  and Carola Lussiana^a 

^aDipartimento di Scienze Agrarie, Forestali e Alimentari, University of Torino, Grugliasco, TO, Italy; ^bDipartimento di Scienze degli Alimenti e del Farmaco, University of Parma, Parma, Italy; ^cDipartimento di Scienze Animali, della Nutrizione e degli Alimenti, Catholic University of the Sacred Heart, Piacenza, Italy

ABSTRACT

This study was designed to assess the effects of dietary flaxseed on the proximate composition, fatty acid profile, lipid oxidation, colour and flavour of the *longissimus thoracis* muscle of double-muscled Piemontese young bulls. Compared to a control diet, the flaxseed diet had no significant effect on the moisture, protein or ether extract content. Flaxseed modified the fatty acid composition of beef. A more than doubled n-3 polyunsaturated fatty acids (PUFA) content (from 21.1 to 46.7 g/kg of total fatty acids – TFA – in control and flaxseed beef, respectively; $p < .001$) and a lowered n-6/n-3 PUFA ratio (13.13 versus 5.60; $p < .001$) were observed. The total *trans*-octadecadienoic acids were also increased by flaxseed (7.8 vs 12.1 g/kg TFA; $p = .001$). The total saturated fatty acids (SFA), monounsaturated fatty acids and PUFA contents, as well as the PUFA/SFA ratio, remained unaffected by the treatment. Lipid oxidation stability tended to be negatively affected by flaxseed ($p = .096$). Such a negative effect was more pronounced as the ageing period advanced. The malondialdehyde content was ≤ 0.72 mg MDA/kg meat, and was therefore below the threshold values for rancidity. Meat colour was not influenced by dietary treatment. The inclusion of flaxseed in the diet increased beef flavour intensity perceived by consumers and did not negatively affect flavour acceptability. Properly combining the choice of animal breed and diet allows a slight improvement of the nutritional value of meat for human consumption.

HIGHLIGHTS

- Proximate composition, colour and flavour of Piemontese beef are not affected by dietary flaxseed
- Dietary flaxseed increases ALA and long-chain n-3 PUFA and lowers the n-6/n-3 PUFA ratio of Piemontese beef
- Flaxseed unprotected from ruminal biohydrogenation does not allow labelling Piemontese beef as source of n-3 FA in the European Union

ARTICLE HISTORY

Received 15 February 2018
Revised 19 June 2018
Accepted 1 September 2018

KEYWORDS

Linum usitatissimum L;
mh/mh genotype;
meat quality; lipids; TBARS

Introduction

The food scandals involving beef, the new eating habits of younger generations, the environmental and ethical issues related to animal production, the high prices of beef and the contemporaneous economic recession, are the main reasons why consumers have turned away from beef (Henchion et al. 2014). However, health concerns, mainly those related to the total fat content and to the supposed negative effects due to saturated fatty acids (SFA) and *trans*

fatty acids, have the strongest incidence on reducing beef consumption (Mapiye et al. 2015; Ruiz-Núñez et al. 2016).

In late-maturing beef cattle breeds, such as Continental European breeds, the amount of intramuscular fat in young bulls is about 2.5% and is even lower (about 1%) in double-muscled animals (*mh/mh* genotype). Owing to the low fat content, the beef from double-muscled animals has a lower SFA content and a more favourable polyunsaturated/saturated fatty

CONTACT Prof. Alberto Brugiapaglia  alberto.brugiapaglia@unito.it  Dipartimento di Scienze Agrarie, Forestali e Alimentari, University of Torino, Largo Paolo Braccini 2, Grugliasco 10095, Torino, Italy

*This paper is dedicated to the memory of Professor Gianluigi Destefanis who passed away on April 8, 2018.

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

acids (PUFA/SFA) ratio than meat from other genotypes (Fiems 2012).

The double-musclé Piedmontese breed is the most numerous Italian beef breed. Almost 270,000 cattle (of which about 128,000 are cows) are registered in the Herd Book, the latter being the largest one among the European meat breeds (ANABORAPI 2016). Piedmontese meat is greatly appreciated by Italian consumers, especially because of its very low intramuscular fat content.

The results of a study conducted by Brugiapaglia et al. (2014) confirmed the positive characteristics of Piedmontese meat (low ether extract content, low concentration of SFA and high PUFA/SFA ratio). However, the traditional feeding system of the Piedmontese breed, which is based on high concentrate diets (e.g. a large use of corn and corn by-products), has led to an unfavourable increase in the n-6/n-3 PUFA ratio in the meat.

An excess of n-6 PUFA can interfere with the metabolism of n-3 PUFA, reducing their incorporation into tissue lipids (Daley et al. 2010), thus representing a risk factor for a number of widespread human pathologies (Simopoulos 2011). In recent years, several trials on beef cattle have highlighted that different feeding strategies can be adopted to favour the deposition of n-3 PUFA in muscle tissues in order to obtain a healthier meat (Baba et al. 2016). Flaxseed, which provides large amounts of α -linolenic acid (C18:3 n-3, ALA), has already been successfully used to improve the nutritional value of the PUFA fraction of beef (Scollan et al. 2014).

However, an enrichment in PUFA could induce a greater susceptibility to lipid oxidation (Guyon et al. 2016), a process that could reduce the organoleptic quality of meat, with particular reference to colour and flavour.

To the best of our knowledge, no studies have been carried out to determine the effects of flaxseed on the meat characteristics of Piedmontese young bulls. Therefore, the aim of this trial was to evaluate the effect of dietary inclusion of flaxseed on the proximate composition, fatty acid (FA) profile, lipid oxidation, colour and flavour of meat from Piedmontese young bulls.

Materials and methods

Animals and dietary treatments

Animal care, handling and experimental procedures were in compliance with the EU Directive 2010/63/EU on the protection of animals used for scientific purposes.

The trial was carried out under practical conditions on a private farm specialised for beef production (Fontanellato, Parma, Italy; latitude: 44°51' 08"N; longitude: 10°07'34"E; altitude: 65 m a.s.l.). Eighteen male Piedmontese calves (4.5 ± 0.59 months) were purchased and transported to the farm, divided into two groups ($n = 9$) and housed in adjacent pens indoors. The two groups were not balanced on a live weight basis, considering the high risks associated with a dominance hierarchy at the feeding rack.

In the pre-experimental period (172 days), all the animals were given the same mixed diet, consisting of a commercial concentrate for fattening cattle (concentrate A: corn, wheat middlings, sunflower meal, roasted dehulled soybean meal, wheat bran, roasted soybean meal, calcium carbonate, cane molasses, canola meal, dicalcium phosphate and sodium chloride), ryegrass hay, corn meal, distillers dried grains with solubles, dried beet pulp and soybean meal (Table 1).

The animals were weighed individually at the beginning of the experimental period. The experimental period lasted 135 days and was divided into two fattening periods (FP1, 66 days; FP2, 69 days) to better fulfil the nutritional requirements of the growing animals. The groups were randomly assigned to two isoenenergetic and isonitrogenous diets (Table 1): a control diet (C; initial live weight: 381 ± 18.5 kg) or an experimental diet containing ground flaxseed (*Linum usitatissimum* L.) (FS; initial live weight: 423 ± 23.5 kg) unprotected from ruminal biohydrogenation. The amount of ground flaxseed (dry matter – DM: 917 g/kg; ether extract – EE: 360 g/kg DM; α -linolenic acid: 200 g/kg DM) was set at 10.0% of the diet DM so as not to exceed the maximum content of cyanogenic substances tolerated by ruminants (Petit 2010). The daily feed intake was monitored by group.

Feed analyses

The AOAC International (2000) procedures were used to determine the dry matter (DM: method no. 930.15), ash (method no. 942.05), crude protein (CP: method no. 984.13) and acid detergent fibre (ADF: method no. 973.18). The ether extract (EE) was determined according to method no. 920.39 of AOAC International (2003). The neutral detergent fibre (NDF) was analysed according to Van Soest et al. (1991); α -amylase (Sigma, Sigma-Aldrich Milano, Milano, Italy) was added, but no sodium sulphite, and the results were corrected for the residual ash content. The chemical composition of the feed (g/kg DM) and the FA proportion of the feed

Table 1. Ingredients, proximate composition and fatty acid profile of the experimental diets.

	FP1		FP2	
	C	FS	C	FS
<i>Ingredients (g/kg DM)</i>				
Concentrate A ^a	353	271	313	244
Corn meal	298	316	209	272
Dried sugar beet pulp	181	134	139	70
Ryegrass hay	125	159	116	143
Barley meal	0	0	155	121
Ground flaxseed	0	99	0	100
Soybean meal	44	22	25	5
Concentrate B ^b	0	0	43	46
<i>Proximate composition (g/kg DM)</i>				
DM (g/kg)	861	865	864	868
CP	174	170	172	169
EE	40	76	36	74
Ash	68	62	64	57
NDF	357	356	355	340
ADF	192	192	179	172
NEF (MJ/kg DM)	7.94	8.19	7.89	8.18
<i>Fatty acid composition (g/kg of TFA)</i>				
C14:0	1.60	0.90	0.19	0.10
C16:0	168.50	108.70	17.78	10.96
C18:0	23.10	25.00	2.21	2.48
C18:1 c9	268.40	242.30	25.29	23.96
C18:2 n-6	490.00	318.60	49.50	31.97
C18:3 n-3	29.60	292.00	3.27	29.36
Others	18.80	12.40	1.75	1.17

FP: fattening period; C: control diet; FS: flaxseed diet; DM: dry matter; CP: crude protein; EE: ether extract; NDF: neutral detergent fibre; ADF: acid detergent fibre; NEF: net energy for fattening; TFA: total fatty acids; c: *cis*; Others: C12:0 + C16:1 *t3* + C16:1 *c9* + C18:1 *c11* + C20:0 + C18:3 n-6.

^aContaining: corn 26%, wheat bran shorts middlings 20%, sunflower meal 15%, rice bran 12%, roasted dehulled soybean meal 9%, wheat bran 7%, roasted soybean meal 4.2%, wheat flour middlings 2%, calcium carbonate 1.5%, cane molasses 1%, canola meal 1%, dicalcium phosphate 0.5%, sodium chloride 0.5%, vitamins and minerals 0.3%. Chemical composition (as fed basis): 18% CP; 5.5% CF (crude fibre); 8.0% cellulose; 7.1% ash; 0.2% Na. Vitamin-mineral composition (given values supplied per kg of concentrate): 88,938 U of vitamin A; 1,338 U of vitamin D₃; 68 mg of vitamin E; 2.5 mg of vitamin B₁; 1.3 mg of vitamin B₂; 1.3 mg of vitamin B₆; 0.011 mg of vitamin B₁₂; 22 mg of vitamin PP; 0.011 mg of biotin; 2.5 mg of pantothenic acid; 750 mg of choline chloride; 54 mg of Mn; 57 mg of Fe; 194 mg of Zn; 12.6 mg of Cu; 0.54 mg of Co; 2.2 mg of I; 0.18 mg of Se.

^bContaining: corn germ meal 45%, wheat middlings 25%, wheat bran 15%, corn 14.7%, vitamins and minerals 0.3%. Chemical composition (as fed basis): 17.5% CP; 3.4% CF (crude fibre); 7.8% cellulose; 5.6% ash; 0.1% Na. Vitamin-mineral composition (given values supplied per kg of concentrate): 4,000 mg of vitamin E; 5.0 mg of Se.

(g/kg of the total detected FA, analysed as detailed in Renna et al. 2014), are reported in Table 1.

Slaughtering and meat sampling

At an age of 15 months, the animals were transported to the slaughterhouse, individually weighed and then slaughtered, in accordance with the European guidelines (EC No. 1099/2009). The carcasses were split into two sides and chilled at 4 °C. Twenty-four hours *post mortem*, the portion between the 9th and 13th thoracic vertebra of *longissimus thoracis* (LT) muscle was taken from the right side. Each sample was divided into three subsamples of equal length.

Two subsamples were randomly assigned an ageing period of 7 or 10 days. The subsamples were placed in a plastic tray, over-wrapped with a polyvinyl chloride oxygen permeable film and stored in the dark at 4 °C for the designated ageing period.

The remaining subsample was cut into three 2-cm thick steaks which were randomly assigned ageing

periods of 2, 7 or 10 days. The steaks were placed in a polystyrene tray, over-wrapped as described before and stored in the dark at 4 °C until lipid oxidation analysis was carried out. The presence of the myostatin gene mutation that is responsible for the double-muscling phenotype (*mh/mh* genotype) in the Piemontese breed was detected in the meat samples using the method described by Di Stasio and Rolando (2005).

Meat analyses

Proximate composition and fatty acid profile

A 2.5 cm thick steak was cut from each 7-day aged subsample, homogenised and divided into two parts. A portion was used to determine the moisture content, according to AOAC method 950.46 (AOAC International 2000). The remaining part was freeze-dried (Edwards MF 1000, Milano, Italy) and then analysed to establish the protein and ether extract contents, and the fatty acid composition.

The nitrogen content was determined by means of the Kjeldahl method (method 928.08; AOAC International 2000) using a Büchi Distillation Unit K-355 (Flawil, Switzerland), and a nitrogen-to-protein conversion factor of 6.25 was used to calculate the protein content.

The ether extract content was determined by means of Soxhlet extraction, without prior acid hydrolysis, using a Büchi Extraction System B-811 (Flawil, Switzerland) with petroleum ether, according to AOAC method 991.36 (AOAC International 2000). The proximate composition results were expressed as g/kg.

Meat fatty acid composition was assessed as reported by Schmid et al. (2009). The fatty acid methyl esters (FAME) were separated and quantified by a high resolution gas chromatograph (Shimadzu GC 2010 Plus, Shimadzu Corporation Analytical Instruments Division, Kyoto, Japan) equipped with a flame-ionisation detector and a CP-Sil 88 capillary column (100 m × 0.25 mm ID, 0.20 µm film thickness; Varian Inc., Lake Forest, CA, USA). The column temperature was held at 45 °C for 5 min, then raised 20 °C min⁻¹ up to 195 °C and maintained for 65 min. The temperature of the injector and the flame-ionization detector was maintained at 250 and 280 °C, respectively; the injection volume was 0.1 µl; nitrogen constant linear flow rate was set at 40 ml min⁻¹. Peaks were identified by comparing the retention times with pure FAME standards (Fluka, Sigma and Supelco, Sigma-Aldrich Milano, Milano, Italy; Matreya Inc., Pleasant Gap, PA, USA), as detailed in Renna et al. (2012), and through comparisons with published chromatograms (Juárez et al. 2011; Alves and Bessa 2014; Vahmani et al. 2016). Quantification was assessed using tridecanoic acid (C13:0; Fluka, Sigma-Aldrich Milano, Milano, Italy) as internal standard. The fatty acid content was determined as mg/100 g of LT muscle, and was then expressed as g/100 g of total detected fatty acids.

Lipid oxidation

After each ageing period (2, 7 or 10 days), a sample of 10 g was taken from each steak. Lipid oxidation was assessed by means of measurement of the thiobarbituric acid-reactive substances (TBARS) using the method of Tarladgis et al. (1960), modified by Novelli et al. (1998). The TBARS values were expressed as mg malondialdehyde/kg of muscle tissue (mg MDA/kg).

Colour

Meat colour was measured using a Minolta CR-331C colourimeter with a 25 mm diameter measured area, D65 illuminant and 2° standard observer in the CIELAB space (CIE 1976). The colour measurements were obtained on a freshly cut surface of subsamples aged 7 or 10 days, after 1 h of blooming at 4 °C. CIELAB coordinates, Lightness (L*), redness (a*) and yellowness (b*) were recorded and chroma (C*) and hue angle (H*) were calculated as $C^* = (a^{*2} + b^{*2})^{0.5}$ and $H^* = \tan^{-1} (b^*/a^*)$.

Consumer sensory evaluation

Two 2.0-cm-thick steaks, cut from each subsample aged 10 days, were vacuum-packaged and stored at -25 °C. Twenty-four hours prior to sensory analysis, the steaks were thawed at 4 °C.

Consumer tests were carried out in individual booths at the Sensory Analysis Laboratory of the Department of Agricultural, Forest and Food Sciences, University of Torino, according to AMSA procedures (AMSA 2015). A total of 63 untrained and regular beef consumers, 39 males and 24 females, ranging in age from 21 to 60 years, were recruited. Informed consents were obtained from all the consumers.

The steaks were cooked on a preheated double plate grill to an internal temperature of 70 °C. After cooking, each steak was cut into 1.3 × 1.3 × 1.9 cm samples which were placed in a foil pouch labelled with a three-digit random number.

The consumers were requested to smell and taste the meat and, then, to rate the perceived flavour intensity (aroma and taste) of each sample using a 9-point, end anchored, intensity scale, where 1 = 'none or extremely bland' and 9 = 'extremely intense' (Miller 1998). The midpoint of the scale (5) was considered a neutral response (i.e. 'neither bland nor intense').

In addition, consumers were asked to rate flavour of each sample as either acceptable or unacceptable. Each consumer evaluated 4 samples (two samples per dietary treatment).

Statistical analyses

The statistical data analyses were carried out using IBM SPSS (IBM Corp. Released 2011; IBM SPSS Statistics for Windows, Version 20.0, Armonk, NY, USA).

The animal was considered as the experimental unit. It was assumed that a general pen effect was negligible due to (i) identical structural characteristics of the pens, (ii) exposition of the animals within the

Table 2. Proximate composition (g/kg meat) of *longissimus thoracis* muscle of Piemontese bulls fed the control (C) and flaxseed (FS) diets ($n = 9$).

	Dietary treatment		SEM	<i>p</i> -value
	C	FS		
Moisture	752.9	754.7	1.43	.422
Protein	229.5	231.5	1.35	.343
Ether Extract	5.2	4.6	0.58	.532

SEM: standard error of the mean; *p*: probability.

two pens to identical management practices, (iii) high uniformity of animals, in terms of genotype, physiology and health status, between the pens, (iv) no changes in animal numbers during the trial and no moving of the animals in and out the pens (St-Pierre 2007). We also assumed that, for the considered dependent variables, the individual responses of bulls within the same pen were not correlated as: (i) the same rearing density was used in both pens, no differential feed access due to social behaviour and no competition for space among the animals within the pens occurred; (ii) the correlation of individual responses of animals within pens has been demonstrated to be of low entity for dependent variables measuring meat quality (Jauhiainen et al. 2008).

The proximate and FA compositions were analysed using the GLM procedure, considering the dietary treatment as fixed effect and the cold carcass weight as covariate. The TBARS values were analysed using the GLM procedure for repeated measures over period, with the dietary treatment as fixed factor, the ageing period (days) as repeated factor, and the dietary treatment \times ageing period interaction. Colour data were analysed by a split-plot design, considering the dietary treatment effect in the main plot and the day effect in the sub-plot, as well as their interaction. Flavour intensity was analysed using the GLM procedure considering the dietary treatment as fixed effect and the consumer as random effect. Acceptability from flavour data was analysed by chi-square test.

The results were reported as least square means and standard error of the mean. Significance was declared at $p < .05$.

Results and discussion

Proximate composition

The final weight and the feed conversion ratio of the bulls averaged 549 kg and 7.0, respectively. The dietary treatment had no significant effect on the LT proximate composition (Table 2). The very low ether extract values found for Piemontese meat in this trial are due to the lack of acid hydrolysis preceding Soxhlet

extraction, and therefore to a lack of phospholipids extraction. The ether extract content of beef from both C-and FS-fed groups was in agreement with previous results of our research group using identical analytical methods (Brugiapaglia and Destefanis 2012; Brugiapaglia et al. 2014) and comparable with the values found on LT muscle of double-muscling Belgian Blue bulls (De Smet et al. 2000).

Some authors have found that the ether extract (Corazzin et al. 2012) and protein (Corazzin et al. 2012; Barahona et al. 2016) increase with the inclusion of flaxseed in the diet, even for the inclusion levels considered in the current trial. However, most of the published literature has shown, in agreement with our findings, that flaxseed does not exert any significant effect on the meat proximate constituents (Juárez et al. 2012; Mapiye, Aalhus et al. 2013; Albertí et al. 2014).

Fatty acid profile

The dietary inclusion of flaxseed significantly modified the FA profile of meat (Table 3), without altering the total intramuscular FA proportion, in agreement with other studies including ALA-rich oilseeds in rations for cattle (Juárez et al. 2011; Mapiye, Aalhus et al. 2013; Mapiye, Turner et al. 2013).

No significant changes were observed for the total SFA, MUFA (monounsaturated fatty acids) and PUFA contents, as previously observed in young bulls of other double-muscling breeds when fed flaxseed (Raes, De Smet et al. 2004; Albertí et al. 2014). The PUFA/SFA ratio was not affected by diet and, for both groups, fell within the recommended values for human nutrition (0.4–1.0) (Jiménez-Colmenero et al. 2012), as generally observed for double-muscling breeds (Scollan et al. 2014; Sevane et al. 2014).

The percentage of PUFA was about double in comparison with that observed in flaxseed-fed young bulls belonging to early-maturing breeds (Mach et al. 2006), but similar to the values reported for Belgian Blue young bulls fed diets rich in n-3 PUFA (Raes et al. 2003). Such a result has to be ascribed to muscle hyperplasia, which determines an increase in the number of cells and, consequently, a high proportion of phospholipids over total lipids (Fiems 2012).

The flaxseed-fed Piemontese bulls showed a higher ALA percentage than Pirenaica bulls (Albertí et al. 2014), a result that is in agreement with the typical high n-3 PUFA proportion over total FA in the meat of this breed, and of leaner animals in general (Aldai et al. 2006; Sevane et al. 2014). Among n-3 PUFA, the proportions of C18:3 n-3 ($p < .001$), C20:5 n-3 ($p < .01$) and C22:6 n-3

Table 3. Fatty acid profile of total lipids in *longissimus thoracis* muscle of Piemontese bulls fed the control (C) and flaxseed (FS) diets ($n = 9$).

	Dietary treatment		SEM	<i>p</i> -value
	C	FS		
<i>Individual SFA (g/100 g of TFA)</i>				
C10:0	0.06	0.06	0.009	.855
C12:0	0.05	0.05	0.005	.766
C14:0	1.53	1.45	0.086	.569
C15:0	2.36	2.19	0.298	.701
C15 anteiso	0.10	0.11	0.009	.414
C16:0	20.45	18.92	0.531	.078
C17:0	1.23	1.09	0.104	.382
C17 iso	0.35	0.35	0.020	.925
C18:0	13.23	15.31	0.560	.028
C18 iso	0.02	0.04	0.005	.072
C20:0	0.07	0.10	0.007	.014
C22:0	0.10	0.11	0.012	.635
<i>Individual MUFA (g/100 g of TFA)</i>				
C14:1 c9	0.56	0.55	0.050	.818
C16:1 t9	0.11	0.11	0.018	.758
C16:1 c7	0.45	0.37	0.026	.072
C16:1 c9	1.30	0.99	0.080	.124
C17:1 c9	0.34	0.38	0.049	.687
C18:1 t6-9	0.25	0.29	0.021	.245
C18:1 t10-11	2.74	2.47	0.283	.544
C18:1 t12	0.57	1.21	0.057	.000
C18:1 c9+t13-14	22.48	19.21	1.622	.206
C18:1 c11+t15	1.78	1.59	0.077	.130
C18:1 c12	0.36	0.50	0.032	.011
C18:1 c15+t16	0.36	0.40	0.042	.511
C20:1 c11	0.11	0.09	0.006	.024
<i>Individual PUFA (g/100 g of TFA)</i>				
C18:2 c9,t13+c9,t14	0.13	0.24	0.028	.016
C18:2 c9,t12+t8,c13	0.15	0.23	0.016	.004
C18:2 t9,c12	0.12	0.20	0.009	.000
C18:2 t10,c15+t11,c15	0.24	0.40	0.032	.005
C18:2 n-6 (LA)	21.52	21.86	1.431	.877
C18:2 c9,c15	0.11	0.09	0.012	.293
C18:2 c9,t11+t7,c9+t8,c10 (CLA)	0.15	0.14	0.008	.514
C18:3 n-6 (GLA)	0.08	0.09	0.008	.822
C18:3 n-3 (ALA)	0.88	2.98	0.140	.000
C20:2 n-6	0.12	0.08	0.008	.005
C20:3 n-9	0.04	0.06	0.004	.003
C20:3 n-6	0.65	0.62	0.051	.729
C20:3 n-3	0.02	0.03	0.006	.362
C20:4 n-6	3.60	3.62	0.255	.974
C20:5 n-3 (EPA)	0.22	0.41	0.039	.007
C22:4 n-6	0.39	0.28	0.028	.020
C22:5 n-3 (DPA)	0.62	0.72	0.086	.424
C22:6 n-3 (DHA)	0.02	0.04	0.005	.010
<i>Groups of FA (g/100 g of TFA)</i>				
Σ SFA	39.56	39.78	0.698	.839
Σ MUFA	31.40	28.14	1.664	.218
Σ C18:1	28.52	25.67	1.647	.274
Σ C18:1 t	3.55	3.97	0.310	.379
Σ PUFA	29.04	32.08	1.910	.313
Σ C18:2	22.40	23.16	1.426	.732
Σ C18:2 t	0.78	1.21	0.065	.001
Σ n-6 PUFA	26.62	26.97	1.710	.895
Σ n-3 PUFA	2.11	4.67	0.237	.000
<i>Selected FA (mg/100 g meat)</i>				
C16:0	120.75	91.27	10.389	.083
C18:0	76.94	73.87	6.739	.770
C18:1 c9+t13-14	136.47	94.28	17.186	.128
C18:2 n-6 (LA)	122.53	102.16	5.370	.025
C18:3 n-3 (ALA)	4.90	14.24	0.657	.000
Σ SFA	231.26	191.26	17.535	.156
Σ MUFA	189.00	137.36	21.200	.131
Σ C18:1	172.14	125.38	19.880	.144
Σ C18:1 t	20.78	19.11	2.494	.667

(continued)

Table 3. Continued.

	Dietary treatment		SEM	<i>p</i> -value
	C	FS		
Σ PUFA	165.30	150.36	7.031	.184
Σ C18:2	127.78	108.39	5.673	.041
Σ C18:2 t	4.61	5.81	0.507	.142
Σ n-6 PUFA	151.62	126.13	6.277	.018
Σ n-3 PUFA	11.85	22.13	0.890	.000
<i>FA ratios and DI</i>				
PUFA/SFA	0.74	0.81	0.057	.408
n-6/n-3 PUFA	13.13	5.60	0.511	.000
DI ₁₄ *	0.38	0.38	0.047	.920
DI ₁₆ **	0.06	0.05	0.006	.193
DI ₁₈ ***	1.72	1.28	0.134	.048

TFA: total fatty acids; SEM: standard error of the mean; *p*: probability; SFA: saturated fatty acids; MUFA: monounsaturated fatty acids; *c*: *cis*; *t*: *trans*; PUFA: polyunsaturated fatty acids; LA: linoleic acid; CLA: conjugated linoleic acid; GLA: γ -linolenic acid; ALA: α -linolenic acid; EPA: eicosapentaenoic acid; DPA: docosapentaenoic acid; DHA: docosahexaenoic acid; FA: fatty acids; DI: desaturase index.

*DI₁₄ = C14:1 c9/C14:0.

**DI₁₆ = C16:1 c9/C16:0.

***DI₁₈ = C18:1 c9/C18:0.

($p = .01$) were higher in the FS group than in the C group. The overall increase in the concentrations of ALA and its long-chain metabolites was statistically significant, but quantitatively limited, as a result of the typical well-known high biohydrogenation rate of dietary ALA, which lowers the duodenal availability of n-3 PUFA for incorporation in the tissues (Doreau and Ferlay 1994). The docosahexaenoic acid (DHA) content was enhanced by the dietary flaxseed, but its level remained very low, thus confirming that consistent enhancements of DHA are almost exclusively achieved when this fatty acid is supplied directly with the diet. The concentrations of ALA (4.90 and 14.24 mg/100 g of meat in C- and FS-fed bulls, respectively) and EPA + DHA (1.35 and 2.14 mg/100 g of meat in C- and FS-fed bulls, respectively) in Piemontese meat were significantly higher in the FS group than in the C group, but remained in both cases very low. Also in case of FS-fed bulls, such values represented only <1% of the intake recommended by EFSA (2009; 2012) for adult people.

The n-6/n-3 PUFA ratio in the FS group was considerably lower than that of the C group ($p < .001$), and was closer to the recommended value (≤ 4) for human nutrition (Simopoulos 2011). As previously observed in flaxseed-supplemented Belgian Blue cattle (Raes, Haak et al. 2004), the n-6/n-3 PUFA ratio was found to be above 5. Lower values can be achieved in cattle fed flaxseed (Herdmann et al. 2010), but are difficult to obtain in lean animals, due to the large amount of LA generally found in the phospholipid fraction of their muscles (Raes, Haak et al. 2004).

The sum of conjugated linoleic acid (CLA) isomers C18:2 c9,t11+t7,c9+t8,c10 was unaffected by the

treatment. Its concentration was low, most probably as a consequence of both dietary and genetic reasons. The type of basal diet used, rich in concentrates as commonly occurs in the fattening phase of ruminants, probably led to a ι 10-shifted rumen biohydrogenation pathway, that is usually associated to low rumenic acid (C18:2 c_9,ι 11) levels (Bessa et al. 2015). Low rumenic acid contents are also usually observed in Piemontese meat (Brugiapaglia et al. 2014) due both to a higher incorporation of CLA in the neutral lipids than in the phospholipids and to a low Stearoyl-CoA desaturase (SCD) activity, which is typical of leaner animals (Sevane et al. 2014).

Many of the detected non-conjugated octadecadienoic isomers are known or thought to be intermediate products of the ruminal biohydrogenation of dietary ALA, which explains their significant increase in the meat of flaxseed-fed cattle in the current and other trials (Juárez et al. 2011; Mapiye, Turner et al. 2013). The sum of *trans* octadecadienoic isomers was higher in the FS meat than in the C meat ($p = .001$), while no differences in the sum content of *trans*-octadecenoic acids were found between groups. It is well-known that *trans* fatty acids increase the risk of cardiovascular disease, but the effect of animal *trans* fatty acids (1–8% of total fats) is still unclear. However, their potential impact on human health is weaker than that of industrial *trans* fats, due to a lower *trans* fatty acids intake from the diet (Brouwer et al. 2013). During GC analysis, the major C18:1 *trans* peak contained a co-elution of C18:1 ι 10 and C18:1 ι 11. As C18:1 ι 10-11 are typically major products of PUFA biohydrogenation, it is apparent that ALA in ground flaxseed was extensively biohydrogenated, which is consistent with the increased C18:0 content when feeding flaxseed. It has been found that C18:1 ι 13–14 can be the major C18:1 *trans* isomer in beef when feeding flaxseed in a barley grain based diet (Juarez et al. 2011). Unfortunately during GC analysis for the current study, C18:1 ι 13–14 co-eluted with C18:1 c_9 , and any changes could not be detected. Interestingly, however, there was an increase in C18:1 ι 12 when feeding flaxseed, and this is consistent with Juarez et al. (2011) where feeding flaxseed in a barley grain based diet increased the content of C18:1 *trans* isomers with double bonds from carbons 12 to 16. Considering the findings of Vahmani et al. (2016) and Juárez et al. (2011), the increased contents of C18:2 c_9,ι 12+ ι 8, c 13 and C18:2 c_9,ι 13+ c_9,ι 14 in the meat from FS-fed bulls can be also respectively imputed to the higher contents of C18:1 ι 12 and C18:1 ι 13/ ι 14 as precursors for Δ 9-desaturase activity. The obtained results further confirm

feeding ALA enriched oilseeds in concentrate based diets may shift biohydrogenation away from pathways with intermediates containing ι 10 and ι 11 double bonds.

Among the SFA, only C18:0 and C20:0 were affected by the treatment ($p < .05$), and they were found to be higher in the FS group than in the C one. The contents of *de novo* synthesised SFA (C10:0, C12:0, C14:0 and C16:0) did not differ significantly between the groups in our study. Some authors have reported a significant reduction in SFA (Raes et al. 2003; Mach et al. 2006; Corazzin et al. 2012), which has been ascribed to the inhibitory effects of the lipid supplementation on *de novo* FA synthesis, while others have found no influence on *de novo* synthesised SFA (Raes, De Smet et al. 2004; Albertí et al. 2014). The dietary treatment did not influence the amount of odd- or branched-chain fatty acids to any great extent, in accordance with the general low inhibitory effect of ALA on rumen microbial synthesis (Mapiye, Aalhus et al. 2013).

Lipid oxidation

The average MDA content tended (0.28 vs 0.42 mg MDA/kg; $p = .096$) to be lower in the meat of the C group than that of the FS group (Table 4). As expected, the MDA levels increased as the ageing period increased ($p < .001$). A significant interaction between dietary treatment and ageing period was also observed ($p < .05$). In fact, the TBARS values in the C group only increased significantly from d2 to d7, while they increased over the whole ageing period in the FS group.

Some authors have reported that the susceptibility of meat to oxidation mainly depends on the n-3 PUFA content, and in particular on ALA, in the cell membranes (Realini et al. 2004; Daly et al. 2007; Kouba & Mourot 2011). In the current trial, the higher n-3 PUFA content in the meat from the FS-fed animals

Table 4. Thiobarbituric acid reactive substances (TBARS; mg MDA/kg of muscle tissue) values of *longissimus thoracis* muscle of Piemontese bulls fed the control (C) and flaxseed (FS) diets ($n = 9$), during ageing (2, 7 and 10 days at 4 ± 1 °C).

Ageing	Dietary treatment		SEM	p-value		
	C	FS		D	AG	D × AG
d2	0.11 ^a	0.15 ^a	0.008			
d7	0.33 ^b	0.39 ^b	0.096	.096	.000	.038
d10	0.41 ^b	0.72 ^c	0.101			

SEM: standard error of the mean; p: probability; D: dietary treatment; AG: ageing period; D × AG: interaction between dietary treatment and ageing period. Means with different superscripts (a–c) within each column indicate significant differences among days.

contributed to a greater susceptibility of the muscle to lipid oxidation.

The toxicity of MDA in human nutrition is well known (Watts 1962); however, its level was below the threshold value for rancidity (1–2 mg MDA/kg) in the meat of both groups.

Colour

After 7 and 10 days of ageing, when meat is generally sold at retail, the dietary treatment had no significant effect on CIELAB colour parameters (Table 5).

This result is in agreement with that of the study of Mach et al. (2006), where different lipid sources, including whole flaxseed, at different concentration in the diet, did not influence L*, a* and b* values in meat from Holstein bulls. Similarly, Alberti et al. (2014) reported that adding whole linseed to a concentrate

Table 5. Colour parameters of *longissimus thoracis* muscle of Piemontese bulls fed the control (C) and flaxseed (FS) diets ($n = 9$) during ageing (7 and 10 days at $4 \pm 1^\circ\text{C}$).

	Dietary treatment				SEM	<i>p</i> -value		
	C		FS			D	AG	D × AG
	d7	d10	d7	d10				
L*	41.24	41.69	41.63	41.55	0.195	.924	.346	.195
a*	27.65	27.64	27.24	27.41	0.377	.633	.841	.817
b*	9.64	9.77	9.29	9.41	0.263	.471	.628	.990
C*	29.29	29.32	28.80	29.00	0.433	.597	.798	.851
H*	19.20	19.48	18.70	18.84	0.304	.375	.499	.815

SEM: standard error of the mean; *p*: probability; D: dietary treatment; AG: ageing period; D × AG: interaction between dietary treatment and ageing period; L*: lightness; a*: redness; b*: yellowness; C*: chroma; H*: hue angle.

fed to young bulls did not significantly modify meat colour.

Consumer sensory evaluation

The frequency distribution of 252 ratings for the two dietary treatments is reported in Figure 1. Both mode and median values of flavour intensity of FS group were 5, which corresponds to 'neither bland nor intense' on the 9-point intensity scale. For C group the mode and the median values were 3 and 4 respectively, and no ratings were found in the 9th category.

The perceived flavour intensity of FS group was significantly higher than that of C group (Table 6). On the contrary, flavour acceptability was not affected by treatment (Table 7).

Table 6. Flavour intensity of *longissimus thoracis* muscle of Piemontese bulls fed the control (C) and flaxseed (FS) diets ($n = 9$).

	C	FS	SEM	<i>p</i> -value
Flavour intensity	4.09	4.69	0.098	.039

SEM: standard error of the mean; *p*: probability. Flavour intensity was evaluated on a 9-point, end-anchored, intensity scale, where 1='none or extremely bland'; 9='extremely intense'.

Table 7. Flavour acceptability of *longissimus thoracis* muscle of Piemontese bulls fed the control (C) and flaxseed (FS) diets.

	Number of ratings (%)		χ^2	<i>p</i> -value
	C	FS		
Acceptable	104 (83)	96 (76)	1.19	.275
Unacceptable	22 (17)	30 (24)		
Total	126 (100)	126 (100)		

χ^2 : Chi square; *p*: probability.




Figure 1. Sensory ratings distribution of flavour intensity. Flavour was evaluated on a 9-point, end-anchored, intensity scale, where 1 = 'none or extremely bland'; 9 = 'extremely intense'.

Beef was designated as acceptable if two-thirds or more of consumers sampling each steak indicated that they were satisfied (Platter et al. 2003). In our study both C group (83%) and FS group (76%) beef resulted in an acceptable rating.

Flavour is a complex attribute of meat palatability and in part depends on meat fat content and composition (Calkins & Hodgen 2007). According to Van Ba et al. (2012), meat with high PUFA content produces unpleasant odours and flavours associated with oxidation.

The higher flavour intensity of FS group may be linked to the differences between treatments in the n-3 PUFA content and most probably to the higher amounts of lower molecular weight unsaturated aldehydes derived from fat oxidation, especially from long chain PUFA (Van Ba et al. 2012).

Literature on how feeding flax to cattle affects palatability is conflicting. Barahona et al. (2016) found that a linseed enriched diet produced meat that had a greater intensity of beef flavour. In contrast, Drouillard et al. (2004) reported no differences in sensory panel observations for juiciness, tenderness or flavour. According to Scollan and Huws (2005) feeding flaxseed, which resulted in meat ALA content of 1.2% of total lipids, had no negative effect on meat organoleptic characteristics. On the contrary, with an increase of the ALA content to 2.8% of the total lipids, an 'abnormal' and 'rancid' beef was recorded (Scollan et al. 2004). These results support the conclusion of Wood et al. (2004) that only when the concentrations of ALA approach 3% of total lipids there are adverse effects on lipid stability, colour or flavour.

After 10 days of ageing, meat of FS group underwent low oxidative deterioration (0.72 mg MDA/kg). Greene and Cumuze (1982) found that detection of MDA by untrained panellists varied from 0.2 to 2 mg MDA/kg and did not necessarily relate to flavour acceptability.

Examining eating quality in 73 beef loins produced using different feeding treatments and containing different concentrations of n-6 and n-3 PUFA, Campo et al. (2006) found that when TBARS values increased, the scores for beef flavour intensity and abnormal flavour increased. This study identifies a TBARS value of 2.3 mg MDA/kg as the point where rancid and other abnormal flavours overpower beef flavour to produce an unacceptable flavour profile in beef.

Conclusions

Our results demonstrate that the inclusion of 10% DM of ground flaxseed can be used in the diets of

Piemontese young bulls, without any modifications of the beef proximate composition, colour and flavour acceptability. The dietary flaxseed significantly modified the FA profile of the meat, increasing the *trans*-octadecadienoic acids and the n-3 PUFA contents, and lowering the n-6/n-3 PUFA ratio. The supplies of ALA and of EPA + DHA on a per-serving basis remained very low (i.e. <1% of the recommended intake for adult humans) even in the meat obtained from the FS-fed bulls. Increasing the n-3 PUFA content in the diet of Piemontese young bulls could be a strategic way of enhancing the nutritional value of the derived beef, without forcing consumers to change their eating habits. Alpha-linolenic acid from flaxseed is extensively biohydrogenated; finding ways to increase its ruminal bypass would be a priority when trying to increase the n-3 PUFA content of Piemontese beef, especially if such a strategy would like be adopted by the cattle industry as a way of promoting healthy Piemontese beef.

Acknowledgments

The authors would like to thank Dr. Claudio Pini for having formulated the diets and the AN.FO.RA. farm (Fontanellato, PR) for its technical assistance and care of the animals. We also acknowledge the staff of UNIPEG slaughterhouse in Pegognaga (MN) for having organised the samples collection.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

ORCID

Manuela Renna  <http://orcid.org/0000-0003-4296-7589>

Alberto Brugiapaglia  <http://orcid.org/0000-0002-0619-7137>

Emanuela Zanardi  <http://orcid.org/0000-0002-2253-3647>

Aldo Prandini  <http://orcid.org/0000-0002-8650-8766>

Maurizio Moschini  <http://orcid.org/0000-0002-7167-709X>

Samantha Sigolo  <http://orcid.org/0000-0001-9081-0565>

Carola Lussiana  <http://orcid.org/0000-0002-0641-5966>

References

- Albertí P, Beriain MJ, Ripoll G, Sarriés V, Panea B, Mendizabal JA, Purroy A, Olleta JL, Sañudo C. 2014. Effect of including linseed in a concentrate fed to young bulls on intramuscular fatty acids and beef color. *Meat Sci.* 96:1258–1265.
- Aldai N, Murray BE, Oliván M, Martínez A, Troy DJ, Osoro K, Nájera AI. 2006. The influence of breed and *mh*-genotype on carcass conformation, meat physico-chemical

- characteristics, and the fatty acid profile of muscle from yearling bulls. *Meat Sci.* 72:486–495.
- Alves SP, Bessa RJB. 2014. The trans-10,cis-15 18:2: a missing intermediate of *trans*-10 shifted rumen biohydrogenation pathway? *Lipids.* 49:527–541.
- AMSA. 2015. Research guidelines for cookery, sensory evaluation and instrumental tenderness measurements of meat. Chicago (IL): American Meat Science Association.
- ANABORAPI. 2016. Relazione Tecnica e Statistiche [Technical Report and Statistics]. [accessed 2017 Dec 11]. http://www.anaborapi.it/index.php?option=com_content&view=category&layout=blog&id=26&Itemid=28. Italian.
- AOAC International. 2000. Official methods of analysis. 17th ed. Gaithersburg (MD): Association of Official Analytical Chemists.
- AOAC International. 2003. Official methods of analysis. 17th ed., 2nd revision. Gaithersburg (MD): Association of Official Analytical Chemists.
- Baba Y, Kallas Z, Costa-Font M, Gil JM, Realini CE. 2016. Impact of hedonic evaluation on consumers' preferences for beef attributes including its enrichment with n-3 and CLA fatty acids. *Meat Sci.* 111:9–17.
- Barahona M, Olleta JL, Sañudo C, Albertí P, Panea B, Pérez-Juan M, Realini CE, Campo MM. 2016. Effects of whole linseed and rumen-protected conjugated linoleic acid enriched diets on beef quality. *Animal.* 10:709–717.
- Bessa RJB, Alves SP, Santos-Silva J. 2015. Constraints and potentials for the nutritional modulation of the fatty acid composition of ruminant meat. *Eur J Lipid Sci Technol.* 117:1325–1344.
- Brouwer I, Wanders A, Katan M. 2013. Trans fatty acids and cardiovascular health: research completed? *Eur J Clin Nutr.* 67:541–547.
- Brugiapaglia A, Destefanis G. 2012. Influence of the housing system on meat quality of double muscled Piemontese young bulls. *Livest Sci.* 145:73–78.
- Brugiapaglia A, Lussiana C, Destefanis G. 2014. Fatty acid profile and cholesterol content of beef at retail of Piemontese, Limousin and Friesian breeds. *Meat Sci.* 96: 568–573.
- Calkins CR, Hodgen JM. 2007. A fresh look at meat flavor. *Meat Sci.* 77:63–80.
- Campo MM, Nute GR, Hughes SI, Enser M, Wood JD, Richardson RI. 2006. Flavour perception of oxidation in beef. *Meat Sci.* 72:303–311.
- CIE. 1976. Recommendations on uniform colour spaces-colour difference equations, psychometric colour terms (Supplement no. 2 to CIE publication No. 15). Paris, France: Commission Internationale de l'Éclairage.
- Corazzin M, Bovolenta S, Sepulcri A, Piasentier E. 2012. Effect of whole linseed addition on meat production and quality of Italian Simmental and Holstein young bulls. *Meat Sci.* 90:99–105.
- Daley CA, Abbott A, Doyle PS, Nader GA, Larson S. 2010. A review of fatty acid profiles and antioxidant content in grass-fed and grain-fed beef. *Nutr J.* 9:10.
- Daly C, Moloney A, Monahan F. 2007. Lipid and colour stability of beef from grazing heifers supplemented with sunflower oil alone or with fish oil. *Meat Sci.* 77:634–642.
- De Smet S, Webb E, Claeys E, Uytterhaegen L, Demeyer D. 2000. Effect of dietary energy and protein levels on fatty acid composition of intramuscular fat in double-muscled Belgian Blue bulls. *Meat Sci.* 56:73–79.
- Di Stasio L, Rolando A. 2005. A PCR-RFLP method for genotyping the myostatin locus in Piemontese cattle. *Anim Genet.* 36:521.
- Doreau M, Ferlay A. 1994. Digestion and utilisation of fatty acids by ruminants. *Anim Feed Sci Technol.* 45:379–396.
- Drouillard J, Seyfert M, Good E, Loe E, Depenbusch B, Daubert R. 2004. Flaxseed for finishing beef cattle: effects on animal performance, carcass quality, and meat composition. Proceedings 60th Flax Institute; Mar 17–19; Fargo, Fargo, ND: ND Flax Institute, Department of Plant Sciences; p. 108–117.
- European Food Safety Authority (EFSA) 2009. Opinion on the substantiation of health claims related to alpha linolenic acid and maintenance of normal blood cholesterol concentrations (ID 493) and maintenance of normal blood pressure (ID 625) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA Journal.* 7:1252. p. 1–17.
- European Food Safety Authority (EFSA) 2012. Scientific opinion on the tolerable upper intake level of eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and docosapentaenoic acid (DPA). EFSA panel on dietetic products, nutrition and allergies (NDA). *EFSA Journal.* 10: 2815. p. 1–48.
- European Union. 2009. Council Regulation (EC) No 1099/2009 on the protection of animals at the time of killing. *Official Journal of the European Union, L.* 303:1–30.
- European Union. 2010. Directive 2010/63/EU on the protection of animals used for scientific purposes. *Official Journal of the European Union, L.* 279:33–79.
- Fiems LO. 2012. Double muscling in cattle: genes, husbandry, carcasses and meat. *Animals (Basel).* 2:472–506.
- Greene B, Cumuze T. 1982. Relationship between TBA numbers and inexperienced panelists' assessments of oxidized flavor in cooked beef. *J Food Sci.* 47:52–54.
- Guyon C, Meynier A, De Lamballerie M. 2016. Protein and lipid oxidation in meat: a review with emphasis on high-pressure treatments. *Trends Food Sci Technol.* 50:131–143.
- Henchion M, McCarthy M, Resconi VC, Troy D. 2014. Meat consumption: trends and quality matters. *Meat Sci.* 98: 561–568.
- Herdmann A, Martin JR, Nuernberg G, Dannenberger D, Nuernberg K. 2010. Effect of dietary n-3 and n-6 PUFA on lipid composition of different tissues of German Holstein bulls and the fate of bioactive fatty acids during processing. *J Agric Food Chem.* 58:8314–8321.
- Jauhiainen L, Manninen M, Öfversten J. 2008. Effect of a pen in group-feeding trials: modelling data from suckler cows. *J Agric Sci.* 146:381–387.
- Jiménez-Colmenero F, Herrero A, Cofrades S, Ruiz-Capillas C. 2012. Meat and functional foods. In: Hui YH, editor. *Handbook of meat and meat processing.* Boca Raton (FL): CRC Press; p. 225–248.
- Juárez M, Dugan ME, Aalhus JL, Aldai N, Basarab JA, Baron VS, McAllister TA. 2011. Effects of vitamin E and flaxseed on rumen-derived fatty acid intermediates in beef intramuscular fat. *Meat Sci.* 88:434–440.
- Juárez M, Dugan MER, Aldai N, Basarab JA, Baron VS, McAllister TA, Aalhus JL. 2012. Beef quality attributes as affected by increasing the intramuscular levels of vitamin E and omega-3 fatty acids. *Meat Sci.* 90:764–769.

- Kouba M, Mourot J. 2011. A review of nutritional effects on fat composition of animal products with special emphasis on n-3 polyunsaturated fatty acids. *Biochimie*. 93:13–17.
- Mach N, Devant M, Díaz I, Font-Furnols M, Oliver M, García J, Bach A. 2006. Increasing the amount of n-3 fatty acid in meat from young Holstein bulls through nutrition. *J Anim Sci*. 84:3039–3048.
- Mapiye C, Aalhus JL, Turner TD, Rolland DC, Basarab JA, Baron VS, McAllister TA, Block HC, Uttaro B, Lopez-Campos O, et al. 2013. Effects of feeding flaxseed or sunflower-seed in high-forage diets on beef production, quality and fatty acid composition. *Meat Sci*. 95:98–109.
- Mapiye C, Turner T, Rolland D, Basarab J, Baron V, McAllister T, Block H, Uttaro B, Aalhus J, Dugan M. 2013b. Adipose tissue and muscle fatty acid profiles of steers fed red clover silage with and without flaxseed. *Livest Sci*. 151:11–20.
- Mapiye C, Vahmani P, Mlambo V, Muchenje V, Dzama K, Hoffman L, Dugan M. 2015. The *trans*-octadecenoic fatty acid profile of beef: implications for global food and nutrition security. *Food Res Int*. 76:992–1000.
- Miller R. 1998. Sensory evaluation of pork. Fact Sheet. Des Moines (IA): National Pork Board/American Meat Science Association. [accessed 2017 Dec 11]. <http://www.extension.org/pages/27315/sensory-evaluation-of-pork>.
- Novelli E, Zanardi E, Ghiretti G, Campanini G, Dazzi G, Madarena G, Chizzolini R. 1998. Lipid and cholesterol oxidation in frozen stored pork, salame Milano and mortadella. *Meat Sci*. 48:29–40.
- Petit H. 2010. Feed intake, milk production and milk composition of dairy cows fed flaxseed. *Can J Anim Sci*. 90: 115–127.
- Platter W, Tatum J, Belk K, Chapman P, Scanga J, Smith G. 2003. Relationships of consumer sensory ratings, marbling score, and shear force value to consumer acceptance of beef strip loin steaks. *J Anim Sci*. 81:2741–2750.
- Raes K, De Smet S, Balcaen A, Claeys E, Demeyer D. 2003. Effect of diets rich in N-3 polyunsaturated fatty acids on muscle lipids and fatty acids in Belgian Blue double-muscléd young bulls. *Reprod Nutr Dev*. 43:331–345.
- Raes K, De Smet S, Demeyer D. 2004. Effect of dietary fatty acids on incorporation of long chain polyunsaturated fatty acids and conjugated linoleic acid in lamb, beef and pork meat: a review. *Anim Feed Sci Technol*. 113:199–221.
- Raes K, Haak L, Balcaen A, Claeys E, Demeyer D, De Smet S. 2004. Effect of linseed feeding at similar linoleic acid levels on the fatty acid composition of double-muscléd Belgian Blue young bulls. *Meat Sci*. 66:307–315.
- Realini C, Duckett S, Brito G, Dalla Rizza M, De Mattos D. 2004. Effect of pasture vs. concentrate feeding with or without antioxidants on carcass characteristics, fatty acid composition, and quality of Uruguayan beef. *Meat Sci*. 66: 567–577.
- Renna M, Cornale P, Lussiana C, Malfatto V, Fortina R, Mimosi A, Battaglini LM. 2012. Use of *Pisum sativum* (L.) as alternative protein resource in diets for dairy sheep: effects on milk yield, gross composition and fatty acid profile. *Small Rumin Res*. 102:142–150.
- Renna M, Gasmi-Boubaker A, Lussiana C, Battaglini LM, Belfayez K, Fortina R. 2014. Fatty acid composition of the seed oils of selected *Vicia* L. taxa from Tunisia. *Ital J Anim Sci*. 13:308–316.
- Ruiz-Núñez B, Dijck-Brouwer DJ, Muskiet FA. 2016. The relation of saturated fatty acids with low-grade inflammation and cardiovascular disease. *J Nutr Biochem*. 36:1–20.
- Schmid A, Collomb M, Hadorn R. 2009. Fatty acid composition of Swiss cooked sausages. *Fleischwirtschaft*. 89: 101–104.
- Scollan ND, Dannenberger D, Nuernberg K, Richardson I, MacKintosh S, Hocquette J-F, Moloney AP. 2014. Enhancing the nutritional and health value of beef lipids and their relationship with meat quality. *Meat Sci*. 97: 384–394.
- Scollan ND, Enser M, Richardson RI, Gulati S, Hallett KG, Nute GR, Wood JD. 2004. The effects of ruminally protected dietary lipid on the fatty acid composition and quality of beef muscle. In: Puolanne E, Ruusunen M, editors. *Proceedings 50th International Congress of Meat Science and Technology*; Aug 8–13; Helsinki, Finland; p. 116.
- Scollan ND, Huws SA. 2005. Enrichment of ruminant products with nutritionally beneficial fatty acids. *Recent Adv Animal Nutr Aust*. 15:21–31.
- Sevane N, Nute G, Sañudo C, Cortes O, Cañon J, Williams J, Dunner S the GemQual Consortium. 2014. Muscle lipid composition in bulls from 15 European breeds. *Livest Sci*. 160:1–11.
- Simopoulos AP. 2011. Importance of the omega-6/omega-3 balance in health and disease: evolutionary aspects of diet. In: Simopoulos AP, editor. *Healthy agriculture, healthy nutrition, healthy people*. Basel, Switzerland: Karger Medical and Scientific Publishers; p. 10–21.
- St-Pierre NR. 2007. Design and analysis of pen studies in the animal sciences. *J Dairy Sci*. 90:E87–E99.
- Tarladgis BG, Watts BM, Younathan MT, Dugan L. 1960. A distillation method for the quantitative determination of malonaldehyde in rancid foods. *J Am Oil Chem Soc*. 37: 44–48.
- Vahmani P, Rolland DC, Gzyl KE, Dugan MER. 2016. Non-conjugated *cis/trans* 18:2 in beef fat are mainly Δ -9 desaturation products of *trans*-18:1 isomers. *Lipids*. 51:1427–1433.
- Van Ba H, Hwang I, Jeong D, Touseef A. 2012. Principle of meat aroma flavors and future prospect. In: Akyar I, editor. *Latest research into quality control*. London (UK): InTechOpen; p. 145–176.
- Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci*. 74: 3583–3597.
- Watts BM. 1962. Meat products. In: Day A and Simhulber RPR, editors. *Symposium on food: lipids and their oxidation*. Westport (CT): AVI Publ. Co.; p. 202–219.
- Wood JD, Richardson RI, Nute GR, Fisher AV, Campo MM, Kasapidou E, Sheard PR, Enser M. 2004. Effects of fatty acids on meat quality: a review. *Meat Sci*. 66:21–32.